Psychedelic treatments; questions from clinical research and implementation

Robert Schoevers Dept. of Psychiatry University Medical Center Groningen

EMA Multi Stakeholder Workshop











Conflict of interest

Research funding	Netherlands Organisation Health Research & Development, EU Horizon 2020,
Speakers fee	NIMH (USA), EU Horizon 2023, UMCG, Stichting tot Steun, Nationaal programma
Scientific advisory	Groningen, Healthcare insurers innovation funds, Janssen Pharmaceuticals,
Clinical trials	Novartis, Compass Pathways, Clexio Biosciences, GH research
Shareholder	None
Other	None

How to maintain initial positive effects?

IMPORTANCE OF ONGOING SUPPORT



THE LANCET Psychiatry

Maintenance ketamine treatment for depression: a systematic review of efficacy, safety, and tolerability

Sanne Y Smith-Apeldoorn, Jolien KE Veraart, Jan Spijker, Jeanine Kamphuis, Robert A Schoevers

Can we combine with other medications?

Pharmacodynamic Interactions Between Ketamine and Psychiatric Medications Used in the Treatment of Depression: A Systematic Review

Jolien K. E. Veraart°, Sanne Y. Smith-Apeldoorn, Iris M. Bakker, Berber A. E. Visser, Jeanine Kamphuis, Robert A. Schoevers, Daan J. Touw

Safety of Ketamine Augmentation to Monoamine Oxidase Inhibitors in Treatment-Resistant Depression:

A Systematic Literature Review and Case Series

Jolien K. E. Veraart, MD^{a,b,+}; Sanne Y. Smith-Apeldoorn, MD^a; Mats Kutscher, MSc^a; Maurice Vischjager, MSc^a; Annemarie van der Meij, MD^c; Jeanine Kamphuis, MD, PhD^a; and Robert A. Schoevers, MD, PhD^{ad}



Can we prescribe in patients with comorbidities?

Ketamine Treatment for Depression in Patients With a History of Psychosis or Current Psychotic Symptoms: A Systematic Review

Jolien K. E. Veraart, MD^{a,b,+}; Sanne Y. Smith-Apeldoorn, MD^a; Jan Spijker, MD, PhD^{c,d}; Jeanine Kamphuis, MD, PhD^a, and Robert A. Schoevers, MD, PhD^{a,e}



Research suggests that mental illnesses lie along a spectrum – but the field's latest diagnostic manual still splits them apart. What is optimal support?

Patient perspectives and experiences with psilocybin treatment for treatment-resistant depression: a qualitative study

Joost J. Breeksema^{1,2×}, Alistair Niemeijer³, Erwin Krediet⁴, Tilman Karsten⁵, Jeanine Kamphuis¹, Eric Vermetten⁶⁴, Wim van den Brink^{6,7} & Robert Schoevers¹

scientific reports



What are adverse effects of classic psychedelics and how often do they occur?



Adverse events in clinical treatments with serotonergic psychedelics and MDMA: A mixed-methods systematic review

Joost J Breeksema^{1,2,3}, Bouwe W Kuin¹, Jeanine Kamphuis¹, Wim van den Brink⁴, Eric Vermetten² and Robert A Schoevers¹



Journal of Psychopharmacology 1-18 © The Author(s) 2022 © • • • •

Article reuse guidelines: sagepub.com/journals-permission DOI: 10.1177/0269881122111694 journals.sagepub.com/home/jop

AE's are poorly defined and remain underreported;

 lack of systematic assessment of AEs
 sample selection

 Acute challenging experiences may be therapeutically

Acute challenging experiences may be therapeutically meaningful
Systematic and detailed assessment & reporting required in all future studies

Who participated in published studies?

Similar effects across psychedelics & diagnoses?

Psychedelic Treatments for Psychiatric Disorders: A Systematic Review and Thematic Synthesis of Patient Experiences in Qualitative Studies

Joost J. Breeksema^{1,2} · Alistair R. Niemeijer³ · Erwin Krediet^{2,5} · Eric Vermetten^{2,4,5} · Robert A. Schoevers¹

Therapeutic processes

1.Insight (e.g. illness, causes, own role)

2. Changing perspective of self

3. Connectedness self/others

4. Transcendent experiences

5. More access to emotions

Patient outcomes

1.Symptom decrease

2.Self image change

3.Connected

4. Larger emotional repertoire

5. Quality of life

CNS Drugs https://doi.org/10.1007/s40263-020-00748-y


SYSTEMATIC REVIEW

What form of psychotherapy works best?

Psychedelic assisted & ketamine assisted psychotherapies; two reviews in progress;

- Wide range of psychotherapeutic approaches
- Form of psychotherapy represents prevailing interests of each era
- Studies mostly descriptive, low methodological quality
- Massive heterogeneity in:
 - Drug formulation and dosing regimens
 - Patient populations
 - Description of specific psychotherapeutic interventions
 - Proposed mechanisms: treatment adherence, cognitive flexibility, reduced defensiveness, access to traumatic memories, transformational experiences
 - Psychotherapy provided during or after administration
 - Psycholytic, psychedelic?

How many sessions are needed?





Questions from research & implementation

- Knowledge gaps
- Upcoming projects
- Next steps



PRESS RELEASE



European Union funds groundbreaking research into psychedelic therapy



Funded by the European Union

A European consortium of 19 partners has been awarded over €6.5M by the European Union to study psilocybin to treat psychological distress in people with progressive incurable illnesses requiring palliative care. This is the first time the EU has funded a multi-site clinical study into psychedelic-assisted therapy.

Psilocybin therapy for psychological distress in palliative care patients

Being confronted with a terminal illness

Psychological and existential distress

- Profound changes in physical, emotional, social and spiritual well-being
- Depression and anxiety (up to 80%)
- Demoralization
- Decreases in quality of life

Poor treatment outcome

• Evidence for psychotherapy and antidepressants remains inconclusive





Multi-site double-blind placebo-controlled clinical trial

• Four medical conditions

- 1. Chronic obstructive pulmonary disorder (COPD) in Groningen
- 2. Amyotrophic lateral sclerosis (ALS) in Copenhagen
- 3. Multiple sclerosis (MS) in Prague
- 4. Atypical Parkinson disorder (APD) in Lisbon


• Inclusion screening

- 100+ patients
- Age > 18
- Current depressive episode (ICD-10)
- Life expectancy of at least 6 months
- Randomisation and blinding (2:1 active vs placebo)





Psilocybin Therapy





Classified as public by the European Medicines Agency

PsyPal key elements

All participants receive psilocybin Ascending dose (2x) Therapy manual and training standardized Mixed methods - including spiritual, existential Explore phenomena eg 'End of life Distress' Different chronic deteriorating medical conditions Combinations with other medications Continued Care and Public and Patient Involvement

- 1) Peer support networks
- 2) Digital Care platform









Questions from research & implementation

- Knowledge gaps
- Upcoming projects
- Next steps



Signalement

Therapeutische toepassingen van psychedelica

Kansen, voor beł -psychiat





RIE

MEEST GELEZEN ARTIKELEN

Sekssomnia tijdens behandeling met ee...

a worldwide point of reference for research, innovation and implementation of novel treatments in mental health care



Improve the health and well-being of patients suffering from mental disorders by establishing a country-wide research and implementation infrastructure

25% of the general population faces a mental health problem

€ 22 billion economic costs per year in the Netherlands alone

highest population disease burden and cause the greatest loss of productivity

of patients develop a serious 'treatment-resistant' mental 30% disorder

existential distress

OCD addiction

> eating disorders chronic pain

post-traumatic stress disorder

depression



Nationwide coordination and collaboration of R&D&I efforts.

Follow-up care to monitor

Large-scale research answering the gaps

Translation to clinical practice

High-quality training and certification for health care professionals Patient-centred approach

Controlled. sustainable, ethical. safe OVERVIEW

long-term effects

Development of specific therapeutic skills

> Creating clear guidelines

Thank you for your attention!

Research group: <u>Psychedelic Treatment and Mechanisms (umcgresearch.org)</u> Contact: <u>r.a.schoevers@umcg.nl</u>

NIVERSITAIR MEDISCH CENTRUM BRONNEN