

SUMMARY OF THE RISK MANAGEMENT PLAN

Table of Contents

I.	THE MEDICINE AND WHAT IT IS USED FOR	2
II.	RISKS ASSOCIATED WITH THE MEDICINE AND ACTIVITIES TO MINIMIZE OR FURTHER CHARACTERIZE THE RISKS	2
II.A	List of important risks and missing information	3
II.B	Summary of important risks	3
II.C	Postauthorization development plan	8
II.C.1	Studies which are conditions of the marketing authorization	8
II.C.2	Other studies in postauthorization development plan	8
References	9

Summary of risk management plan for XOSPATA (Gilteritinib)

This is a summary of the risk management plan (RMP) for XOSPATA. The RMP details important risks of XOSPATA, how these risks can be minimized, and how more information will be obtained about XOSPATA's risks.

XOSPATA's summary of product characteristics (SmPC) and its package leaflet give essential information to healthcare professionals and patients on how XOSPATA should be used.

This summary of the RMP for XOSPATA should be read in the context of all this information including the assessment report of the evaluation and its plain-language summary, all which is part of the European Public Assessment Report (EPAR).

Important new concerns or changes to the current ones will be included in updates of XOSPATA's RMP.

I. THE MEDICINE AND WHAT IT IS USED FOR

XOSPATA is authorized for treatment of adult patients who have relapsed or refractory acute myeloid leukemia (AML) with an FMS-like tyrosine kinase (FLT3) mutation (see the SmPC for the full indication). It contains gilteritinib as the active substance and it is available as a film-coated 40 mg tablet.

Further information about the evaluation of XOSPATA's benefits can be found in XOSPATA's EPAR, including in its plain-language summary, available on the European Medicines Agency (EMA) website, under the medicine's webpage:

<https://www.ema.europa.eu/en/medicines/human/EPAR/xospata>

II. RISKS ASSOCIATED WITH THE MEDICINE AND ACTIVITIES TO MINIMIZE OR FURTHER CHARACTERIZE THE RISKS

Important risks of XOSPATA, together with measures to minimize such risks and the proposed studies for learning more about XOSPATA's risks, are outlined below.

Measures to minimize the risks identified for medicinal products can be:

- Specific information, such as warnings, precautions, and advice on correct use, in the package leaflet and SmPC addressed to patients and healthcare professionals;
- Important advice on the medicine's packaging;
- The authorized pack size — the amount of medicine in a pack is chosen so to ensure that the medicine is used correctly;
- The medicine's legal status — the way a medicine is supplied to the patient (e.g., with or without prescription) can help to minimize its risks.

Together, these measures constitute *routine risk minimization* measures.

In the case of XOSPATA, these measures are supplemented with additional risk minimization measures mentioned under relevant important risks below.

In addition to these measures, information about adverse reactions is collected continuously and regularly analyzed, including PSUR assessment, so that immediate action can be taken as necessary. These measures constitute routine pharmacovigilance activities.

II.A List of important risks and missing information

Important risks of XOSPATA are risks that need special risk management activities to further investigate or minimize the risk, so that the medicinal product can be safely taken. Important risks can be regarded as identified or potential. Identified risks are concerns for which there is sufficient proof of a link with the use of XOSPATA. Potential risks are concerns for which an association with the use of this medicine is possible based on available data, but this association has not been established yet and needs further evaluation. Missing information refers to information on the safety of the medicinal product that is currently missing and needs to be collected (e.g., on the long-term use of the medicine).

List of Important Risks	
Important identified risks	<ul style="list-style-type: none"> • Posterior reversible encephalopathy syndrome • Differentiation syndrome
Important potential risks	<ul style="list-style-type: none"> • Torsades de Pointes • Serious Gastrointestinal disorders • Eye disorders • Pulmonary adverse events • Pancreatitis • Embryo-fetal lethality, suppressed fetal growth, and teratogenicity

II.B Summary of important risks

Important identified risk: Posterior reversible encephalopathy syndrome (PRES)	
Evidence for linking the risk to the medicine	<p>PRES is a rare neurological disorder that can present with rapidly evolving symptoms including seizure, headache, confusion, blindness, and other visual and neurological disturbances, with or without associated hypertension. A diagnosis of PRES requires confirmation by brain imaging, preferably magnetic resonance imaging (MRI). In clinical trials, there have been uncommon reports of PRES with gilteritinib with symptoms including seizure and altered mental status. Symptoms have resolved after discontinuation of gilteritinib. Early recognition of PRES may mitigate secondary complications that can lead to morbidity/mortality. Discontinuation of gilteritinib in patients who develop PRES is recommended.</p>
Risk factors and risk groups	<p>In general, there appears to be a predominance of female PRES cases to male cases; the syndrome occurs across many age groups (pediatric, adult, and elderly) [Fugate et al, 2010; Li et al, 2012; Li et al, 2013].</p> <p>PRES has been linked to a number of medical conditions including hypertensive encephalopathy, renal disease, systemic inflammatory syndrome, autoimmune disease, and vasculitis [Le & Loghin, 2014]. PRES has also been linked to medications including chemotherapies (e.g., anthracyclines and alkylating agents), monoclonal antibodies, and immunosuppressant drugs [Le & Loghin, 2014]. PRES has also been associated with use of several multikinase inhibitors (e.g., sunitinib, sorafenib) [Costa et al, 2014; Foerster et al, 2013].</p>

Important identified risk: Posterior reversible encephalopathy syndrome (PRES)

Risk minimization measures	<p>Routine risk communication:</p> <ul style="list-style-type: none">• SmPC sections 4.2, 4.4 and 4.8;• PL sections 2 and 4. <p>Routine risk minimization activities recommending specific clinical measures to address the risk:</p> <ul style="list-style-type: none">• Recommendation to discontinue gilteritinib in patients who develop PRES is provided in SmPC sections 4.2 and 4.4.
----------------------------	--

MRI: Magnetic Resonance Imaging; PL: Package Leaflet; PRES: Posterior Reversible Encephalopathy Syndrome; SmPC: Summary of Product Characteristics

Important identified risk: Differentiation Syndrome

Evidence for linking the risk to the medicine	<p>Differentiation syndrome is associated with rapid proliferation and differentiation of myeloid cells and may be life-threatening or fatal if not treated. Of the 319 patients treated with gilteritinib in the clinical trials, 3% experienced differentiation syndrome. Symptoms of differentiation syndrome in patients treated with gilteritinib included fever, dyspnea, pleural effusion, pericardial effusion, pulmonary edema, hypotension, rapid weight gain, peripheral edema, rash, and renal dysfunction. Some cases had concomitant acute febrile neutrophilic dermatosis. Differentiation syndrome occurred as early as 1 day and up to 82 days after gilteritinib initiation and has been observed with or without concomitant leukocytosis. Of the 11 patients who experienced differentiation syndrome, 9 (82%) recovered after treatment or after dose interruption of gilteritinib.</p>
Risk factors and risk groups	<p>Patients treated with chemotherapeutic agents and targeted therapies that induce myeloid differentiation are at risk for developing differentiation syndrome as a rare treatment associated complication.</p>
Risk minimization measures	<p>Routine risk communication:</p> <ul style="list-style-type: none">• SmPC sections 4.2; 4.4; 4.8;• PL sections 2 and 4. <p>Routine risk minimization activities recommending specific clinical measures to address the risk:</p> <ul style="list-style-type: none">• Recommendation for monitoring is provided in SmPC section 4.4;• Recommendation for treatment interruption of gilteritinib is provided in SmPC section 4.2 and 4.4. <p>Additional risk minimization measures:</p> <ul style="list-style-type: none">• Healthcare Professional Information• Patient alert card

PL: Package Leaflet; SmPC: Summary of Product Characteristics

Important potential risk: Torsades de Pointes

Evidence for linking the risk to the medicine

Gilteritinib has been associated with prolonged cardiac ventricular repolarization (QT interval). A concentration-related increase in change from baseline of QTcF was observed across gilteritinib doses ranging from 20 to 450 mg. The effect of gilteritinib 120 mg once a day on QTc has been evaluated in patients, which showed an absence of large mean increases (i.e., 20 msec) in the QTc interval.

Of 319 patients treated with gilteritinib at 120 mg tested in clinical trials, 4 patients (1.3%) experienced a QTcF > 500 msec. Additionally, across all doses, 2.3% of relapse/refractory subjects had a maximum post-baseline QTcF interval > 500 msec.

In clinical trials, there have been reports of QT prolongation with gilteritinib, occurring in approximately 7% of patients treated with gilteritinib (36/496 patients). Another approximately 7% (36/496 patients) experienced related events such as syncope, cardiac arrest, ventricular tachycardia, ventricular arrhythmia, loss of consciousness, ventricular fibrillation, and sudden death. There were no patients that discontinued gilteritinib due to QT prolongation.

The clinical trials excluded patients with baseline QTcF \geq 450 msec, long QT syndrome, hypokalemia or hypomagnesemia, or significant cardiovascular disease.

Monitoring for arrhythmias and early recognition of QT prolongation may mitigate the development of serious outcomes. Interruption and/or dose reduction of gilteritinib in patients who develop QT prolongation (QTcF > 500 msec) is recommended. Consideration for gilteritinib dose reduction in patients with QTc > 30 msec change from baseline on Day 8, if confirmed.

Risk factors and risk groups

Patients with pre-existing QTcF > 450 msec, long QT syndrome, electrolyte abnormalities (hypokalemia or hypomagnesemia), or those who are taking medications known to prolong the QT interval (e.g., anti-arrhythmic medicines, fluoroquinolones, triazole antifungals, and 5-HT₃ [serotonin type 3] receptor antagonists) may be at a higher risk for QT prolongation.

A concentration-related increase in changes from baseline for QTcF was observed across gilteritinib doses ranging from 20 to 450 mg. Therefore, strong inhibitors of CYP3A or P-gp, such as, antifungals (e.g., voriconazole, posaconazole), antibiotics (e.g., clarithromycin, azithromycin), angiotensin-converting enzyme inhibitors (e.g., captopril), antivirals (e.g., ritonavir), and beta blockers (e.g., carvedilol) may increase the risk of QTc interval prolongation because they can increase gilteritinib plasma concentrations.

Risk minimization measures

Routine risk communication:

- Description of QT prolongation is included in:
 - SmPC sections 4.2, 4.4 and 4.8;
 - PL sections 2 and 4.

Routine risk minimization activities recommending specific clinical measures to address the risk:

Important potential risk: Torsades de Pointes

	<ul style="list-style-type: none">• Recommendation for ECG monitoring is provided in SmPC sections 4.2 and 4.4;• Recommendation for treatment interruption of gilteritinib in patients with QTc > 500 msec with resumption at a reduced dose when QTc interval returns to baseline or ≤480 msec is provided in SmPC section 4.2 and 4.4;• Consideration for gilteritinib dose reduction in patients with QTc > 30 msec change from baseline on Day 8, if confirmed is provided in SmPC section 4.2 and 4.4.• Recommendation to correct hypokalemia or hypomagnesemia prior to and during gilteritinib administration is provided in SmPC section 4.4., as hypokalemia or hypomagnesemia may increase QT prolongation.• Caution in patients with relevant cardiac history is noted in SmPC section 4.4.
--	--

CYP: Cytochrome P450; ECG: Electrocardiogram; P-gp: P-glycoprotein; PL: Package Leaflet; QTc: QT Interval Corrected for Heart Rate; QTcF: QT Interval Corrected for Heart Rate Using Fridericia's Formula; SmPC: Summary of Product Characteristics.

Important potential risk: Serious GI disorders

Evidence for linking the risk to the medicine	Serious GI disorders have been observed in gilteritinib clinical trials; however, an association with gilteritinib has not been confirmed.
Risk factors and risk groups	Risk factors for serious GI disorders in the general population include hyperlipidemia, pre-existing conditions such as diverticulitis, alcohol use, obesity, and cigarette smoking. In AML patients, serious GI disorders can occur secondary to sepsis and infection.
Risk minimization measures	None

AML: Acute Myeloid Leukemia; GI: Gastrointestinal.

Important potential risk: Eye disorders

Evidence for linking the risk to the medicine	Eye disorders have been observed in gilteritinib clinical trials; however, an association with gilteritinib has not been confirmed.
Risk factors and risk groups	Risk factors for eye disorders in the general population include age, hypertension, diabetes, pre-existing retinal disease, bleeding disorders, ultraviolet exposure and cigarette smoking.
Risk minimization measures	None

Important potential risk: Pulmonary adverse events	
Evidence for linking the risk to the medicine	Pulmonary events have been observed in gilteritinib clinical trials; however, an association with gilteritinib has not been confirmed.
Risk factors and risk groups	Risk factors for pulmonary toxicity in the general population include diabetes, alcohol use, environmental pollution, genetic/autoimmune diseases such as asthma and cystic fibrosis, and cigarette smoking. In AML patients, pulmonary events can occur secondary to sepsis and infection, thrombocytopenia with associated bleeding complications and also in association with differentiation syndrome.
Risk minimization measures	None

AML: Acute Myeloid Leukemia

Important potential risk: Pancreatitis	
Evidence for linking the risk to the medicine	Acute pancreatitis and pancreatitis have been observed in gilteritinib clinical trials; however, an association with gilteritinib has not been confirmed.
Risk factors and risk groups	Risk factors for pancreatitis in the general population include hyperlipidemia, diabetes, alcohol use, and cigarette smoking. In AML patients, pancreatitis can occur secondary to sepsis and infection.
Risk minimization measures	<p>Routine risk communication:</p> <ul style="list-style-type: none"> • SmPC section 4.2 and 4.4; • PL section 2. <p>Routine risk minimization activities recommending specific clinical measures to address the risk:</p> <ul style="list-style-type: none"> • Recommendation to evaluate and monitor patients who develop signs and symptoms suggestive of pancreatitis is provided in SmPC section 4.4. • Recommendation for treatment interruption of gilteritinib is provided in SmPC section 4.2 and 4.4. Gilteritinib can be resumed at a reduced dose when the signs and symptoms of pancreatitis have resolved.

AML: Acute Myeloid Leukemia; PL: Package Leaflet; SmPC: Summary of Product Characteristics.

Important potential risk: Embryo-fetal lethality, suppressed fetal growth, and teratogenicity	
Evidence for linking the risk to the medicine	Gilteritinib can cause fetal harm when administered to pregnant women. There are no data from the use of gilteritinib in pregnant women. Embryo-fetal development studies in rats with gilteritinib have shown suppressed fetal growth, embryo-fetal deaths, and teratogenicity.
Risk factors and risk groups	Persons at risk include female patients of child-bearing potential, as well as male patients with female partners who are of child-bearing potential.
Risk minimization measures	<p>Routine risk communication:</p> <ul style="list-style-type: none"> • SmPC sections 4.4 and 4.6; • PL section 2. <p>Routine risk minimization activities recommending specific clinical measures to address the risk:</p> <ul style="list-style-type: none"> • Recommendation for females of reproductive potential to use contraception during treatment and for at least 6 months after the last dose of gilteritinib is provided in SmPC sections 4.4 and 4.6; • Recommendation for women using hormonal contraceptives to add a barrier method of contraception is provided in SmPC sections 4.4 and 4.6; • Recommendation for males with female partners of reproductive potential to use effective contraception during treatment and for at least 4 months after the last dose is provided in SmPC sections 4.4 and 4.6.

PL: Package Leaflet; SmPC: Summary of Product Characteristics.

II.C Postauthorization development plan

II.C.1 Studies which are conditions of the marketing authorization

There are no studies that are conditions of the marketing authorization or specific obligation of XOSPATA.

II.C.2 Other studies in postauthorization development plan

None

References

- Costa R, Costa R, Costa R, Junior GM, Cartaxo HQ, de Barros AC. Reversible posterior encephalopathy syndrome secondary to sunitinib. *Case Rep Oncol Med*. 2014;2014:952624. Epub 13-May-2014.
- Foerster R, Welzel T, Debus J, Gruellich C, Jaeger D, Potthoff K. Posterior reversible leukoencephalopathy syndrome associated with pazopanib. *Case Rep Oncol*. 2013;6(1):204-8.
- Fugate JE, Claassen DO, Cloft HJ, Kallmes DF, Kozak OS, Rabinstein AA. Posterior reversible encephalopathy syndrome: associated clinical and radiologic findings. *Mayo Clin Proc*. 2010;85(5):427-32.
- Le EM, Loghin ME. Posterior reversible encephalopathy syndrome: a neurologic phenomenon in cancer patients. *Curr Oncol Rep*. 2014;16(5):383.
- Li R, Mitchell P, Dowling R, Yan B. Is hypertension predictive of clinical recurrence in posterior reversible encephalopathy syndrome? *J Clin Neurosci*. 2013;20(2):248-52.
- Li Y, Gor D, Walicki D, Jenny D, Jones D, Barbour P, et al. Spectrum and potential pathogenesis of reversible posterior leukoencephalopathy syndrome. *J Stroke Cerebrovasc Dis*. 2012;21(8):873-82.