

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Tracleer 62.5 mg film-coated tablets

Tracleer 125 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Tracleer 62.5 mg film-coated tablets

Each film-coated tablet contains 62.5 mg bosentan (as monohydrate).

Tracleer 125 mg film-coated tablets

Each film-coated tablet contains 125 mg bosentan (as monohydrate).

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet (tablets):

Tracleer 62.5 mg film-coated tablets

Orange-white, round, biconvex, film-coated tablets, embossed with “62,5” on one side.

Tracleer 125 mg film-coated tablets

Orange-white, oval, biconvex, film-coated tablets, embossed with “125” on one side.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Treatment of pulmonary arterial hypertension (PAH) to improve exercise capacity and symptoms in patients with WHO functional class III. Efficacy has been shown in:

- Primary (idiopathic and heritable) pulmonary arterial hypertension
- Pulmonary arterial hypertension secondary to scleroderma without significant interstitial pulmonary disease
- Pulmonary arterial hypertension associated with congenital systemic-to-pulmonary shunts and Eisenmenger’s physiology

Some improvements have also been shown in patients with pulmonary arterial hypertension WHO functional class II (see section 5.1).

Tracleer is also indicated to reduce the number of new digital ulcers in patients with systemic sclerosis and ongoing digital ulcer disease (see section 5.1).

### 4.2 Posology and method of administration

Method of administration

Tablets are to be taken orally morning and evening, with or without food. The film-coated tablets are to be swallowed with water.

Patients should be advised not to swallow the desiccant found in the white high-density polyethylene bottles.

## Posology

### Pulmonary arterial hypertension

Treatment should only be initiated and monitored by a physician experienced in the treatment of PAH.

#### Adults

In adult patients, Tracleer treatment should be initiated at a dose of 62.5 mg twice daily for 4 weeks and then increased to the maintenance dose of 125 mg twice daily. The same recommendations apply to re-introduction of Tracleer after treatment interruption (see section 4.4).

#### Paediatric population

Paediatric pharmacokinetic data have shown that bosentan plasma concentrations in children with PAH aged from 1 year to 15 years were on average lower than in adult patients and were not increased by increasing the dose of Tracleer above 2 mg/kg body weight or by increasing the dosing frequency from twice daily to three times daily (see section 5.2). Increasing the dose or the dosing frequency will likely not result in additional clinical benefit.

Based on these pharmacokinetic results, when used in children with PAH aged 1 year and older, the recommended starting and maintenance dose is 2 mg/kg morning and evening.

In neonates with persistent pulmonary hypertension of the newborn (PPHN), the benefit of bosentan has not been shown in the standard-of-care treatment. No recommendation on a posology can be made (see sections 5.1 and 5.2).

#### *Management in the event of clinical deterioration of PAH*

In the event of clinical deterioration (e.g., decrease in 6-minute walk test distance by at least 10% compared with pre-treatment measurement) despite Tracleer treatment for at least 8 weeks (target dose for at least 4 weeks), alternative therapies should be considered. However, some patients who show no response after 8 weeks of treatment with Tracleer may respond favourably after an additional 4 to 8 weeks of treatment.

In the event of late clinical deterioration despite treatment with Tracleer (i.e., after several months of treatment), the treatment should be re-assessed. Some patients not responding well to 125 mg twice daily of Tracleer may slightly improve their exercise capacity when the dose is increased to 250 mg twice daily. A careful benefit/risk assessment should be made, taking into consideration that the liver toxicity is dose dependent (see sections 4.4 and 5.1).

#### *Discontinuation of treatment*

There is limited experience with abrupt discontinuation of Tracleer in patients with PAH. No evidence for acute rebound has been observed. However, to avoid the possible occurrence of harmful clinical deterioration due to potential rebound effect, gradual dose reduction (halving the dose for 3 to 7 days) should be considered. Intensified monitoring is recommended during the discontinuation period. If the decision to withdraw Tracleer is taken, it should be done gradually while an alternative therapy is introduced.

### Systemic sclerosis with ongoing digital ulcer disease

Treatment should only be initiated and monitored by a physician experienced in the treatment of systemic sclerosis.

#### Adults

Tracleer treatment should be initiated at a dose of 62.5 mg twice daily for 4 weeks and then increased to the maintenance dose of 125 mg twice daily. The same recommendations apply to re-introduction of Tracleer after treatment interruption (see section 4.4).

Controlled clinical study experience in this indication is limited to 6 months (see section 5.1).

The patient's response to treatment and need for continued therapy should be re-evaluated on a regular basis. A careful benefit/risk assessment should be made, taking into consideration the liver toxicity of bosentan (see sections 4.4 and 4.8).

#### Paediatric population

There are no data on the safety and efficacy in patients under the age of 18 years. Pharmacokinetic data are not available for Tracleer in young children with this disease.

### Special populations

#### *Hepatic impairment*

Tracleer is contraindicated in patients with moderate to severe liver dysfunction (see sections 4.3, 4.4 and 5.2). No dose adjustment is needed in patients with mild hepatic impairment (i.e., Child-Pugh class A) (see section 5.2).

#### *Renal impairment*

No dose adjustment is required in patients with renal impairment. No dose adjustment is required in patients undergoing dialysis (see section 5.2).

#### *Elderly*

No dose adjustment is required in patients over the age of 65 years.

### **4.3 Contraindications**

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1
- Moderate to severe hepatic impairment, i.e., Child-Pugh class B or C (see section 5.2)
- Baseline values of liver aminotransferases, i.e., aspartate aminotransferase (AST) and/or alanine aminotransferase (ALT), greater than 3 × the upper limit of normal (ULN; see section 4.4)
- Concomitant use of cyclosporine A (see section 4.5)
- Pregnancy (see sections 4.4 and 4.6)
- Women of childbearing potential who are not using reliable methods of contraception (see sections 4.4, 4.5 and 4.6)

### **4.4 Special warnings and precautions for use**

The efficacy of Tracleer has not been established in patients with severe PAH. Transfer to a therapy that is recommended at the severe stage of the disease (e.g., epoprostenol) should be considered if the clinical condition deteriorates (see section 4.2).

The benefit/risk balance of bosentan has not been established in patients with WHO class I functional status of PAH.

Tracleer should only be initiated if the systemic systolic blood pressure is higher than 85 mmHg.

Tracleer has not been shown to have a beneficial effect on the healing of existing digital ulcers.

### Liver function

Elevations in liver aminotransferases, i.e., aspartate and alanine aminotransferases (AST and/or ALT), associated with bosentan are dose dependent. Liver enzyme changes typically occur within the first 26 weeks of treatment but may also occur late in treatment (see section 4.8). These increases may be partly due to competitive inhibition of the elimination of bile salts from hepatocytes but other mechanisms, which have not been clearly established, are probably also involved in the occurrence of liver dysfunction. The accumulation of bosentan in hepatocytes leading to cytolysis with potentially severe damage of the liver, or an immunological mechanism, are not excluded. Liver dysfunction risk may also be increased when medicinal products that are inhibitors of the bile salt export pump, e.g., rifampicin, glibenclamide and cyclosporine A (see sections 4.3 and 4.5), are co-administered with bosentan, but limited data are available.

**Liver aminotransferase levels must be measured prior to initiation of treatment and subsequently at monthly intervals for the duration of treatment with Tracleer. In addition, liver aminotransferase levels must be measured 2 weeks after any dose increase.**

### Recommendations in the event of ALT/AST elevations

#### **ALT/AST levels**

#### **Treatment and monitoring recommendations**

> 3 and  $\leq 5 \times$  ULN

The result should be confirmed by a second liver test; if confirmed, a decision should be made on an individual basis to continue Tracleer, possibly at a reduced dose, or to stop Tracleer administration (see section 4.2). Monitoring of aminotransferase levels should be continued at least every 2 weeks. If the aminotransferase levels return to pre-treatment values continuing or re-introducing Tracleer according to the conditions described below should be considered.

> 5 and  $\leq 8 \times$  ULN

The result should be confirmed by a second liver test; if confirmed, treatment should be stopped and aminotransferase levels monitored at least every 2 weeks. If the aminotransferase levels return to pre-treatment values re-introducing Tracleer according to the conditions described below should be considered.

>  $8 \times$  ULN

Treatment must be stopped and re-introduction of Tracleer is not to be considered.

**In the event of associated clinical symptoms of liver injury, i.e., nausea, vomiting, fever, abdominal pain, jaundice, unusual lethargy or fatigue, flu-like syndrome (arthralgia, myalgia, fever), treatment must be stopped and re-introduction of Tracleer is not to be considered.**

### Re-introduction of treatment

Re-introduction of treatment with Tracleer should only be considered if the potential benefits of treatment with Tracleer outweigh the potential risks and when liver aminotransferase levels are within pre-treatment values. The advice of a hepatologist is recommended. Re-introduction must follow the guidelines detailed in section 4.2. **Aminotransferase levels must then be checked within 3 days**

**after re-introduction, then again after a further 2 weeks, and thereafter according to the recommendations above.**

ULN = upper limit of normal

### Haemoglobin concentration

Treatment with bosentan has been associated with dose-related decreases in haemoglobin concentration (see section 4.8). In placebo-controlled studies, bosentan-related decreases in haemoglobin concentration were not progressive, and stabilised after the first 4–12 weeks of treatment. It is recommended that haemoglobin concentrations be checked prior to initiation of treatment, every month during the first 4 months, and quarterly thereafter. If a clinically relevant decrease in haemoglobin concentration occurs, further evaluation and investigation should be undertaken to determine the cause and need for specific treatment. In the post-marketing period, cases of anaemia requiring red blood cell transfusion have been reported (see section 4.8).

### Women of childbearing potential

As Tracleer may render hormonal contraceptives ineffective, and taking into account the risk that pulmonary hypertension deteriorates with pregnancy as well as the teratogenic effects observed in animals:

- Tracleer treatment must not be initiated in women of childbearing potential unless they practise reliable contraception and the result of the pre-treatment pregnancy test is negative
- Hormonal contraceptives cannot be the sole method of contraception during treatment with Tracleer
- Monthly pregnancy tests are recommended during treatment to allow early detection of pregnancy

For further information see sections 4.5 and 4.6.

### Pulmonary veno-occlusive disease

Cases of pulmonary oedema have been reported with vasodilators (mainly prostacyclins) when used in patients with pulmonary veno-occlusive disease. Consequently, should signs of pulmonary oedema occur when Tracleer is administered in patients with PAH, the possibility of associated veno-occlusive disease should be considered. In the post-marketing period there have been rare reports of pulmonary oedema in patients treated with Tracleer who had a suspected diagnosis of pulmonary veno-occlusive disease.

### Pulmonary arterial hypertension patients with concomitant left ventricular failure

No specific study has been performed in patients with pulmonary hypertension and concomitant left ventricular dysfunction. However, 1,611 patients (804 bosentan- and 807 placebo-treated patients) with severe chronic heart failure (CHF) were treated for a mean duration of 1.5 years in a placebo-controlled study (study AC-052-301/302 [ENABLE 1 & 2]). In this study there was an increased incidence of hospitalisation due to CHF during the first 4–8 weeks of treatment with bosentan, which could have been the result of fluid retention. In this study, fluid retention was manifested by early weight gain, decreased haemoglobin concentration and increased incidence of leg oedema. At the end of this study, there was no difference in overall hospitalisations for heart failure nor in mortality between bosentan- and placebo-treated patients. Consequently, it is recommended that patients be monitored for signs of fluid retention (e.g., weight gain), especially if they concomitantly suffer from severe systolic dysfunction. Should this occur, starting treatment with diuretics is recommended, or the

dose of existing diuretics should be increased. Treatment with diuretics should be considered in patients with evidence of fluid retention before the start of treatment with Tracleer.

#### Pulmonary arterial hypertension associated with HIV infection

There is limited clinical study experience with the use of Tracleer in patients with PAH associated with HIV infection, treated with antiretroviral medicinal products (see section 5.1). An interaction study between bosentan and lopinavir+ritonavir in healthy subjects showed increased plasma concentrations of bosentan, with the maximum level during the first 4 days of treatment (see section 4.5). When treatment with Tracleer is initiated in patients who require ritonavir-boosted protease inhibitors, the patient's tolerability of Tracleer should be closely monitored with special attention, at the beginning of the initiation phase, to the risk of hypotension and to liver function tests. An increased long-term risk of hepatic toxicity and haematological adverse events cannot be excluded when bosentan is used in combination with antiretroviral medicinal products. Due to the potential for interactions related to the inducing effect of bosentan on CYP450 (see section 4.5), which could affect the efficacy of antiretroviral therapy, these patients should also be monitored carefully regarding their HIV infection.

#### Pulmonary hypertension secondary to chronic obstructive pulmonary disease (COPD)

Safety and tolerability of bosentan was investigated in an exploratory, uncontrolled 12-week study in 11 patients with pulmonary hypertension secondary to severe COPD (stage III of GOLD classification). An increase in minute ventilation and a decrease in oxygen saturation were observed, and the most frequent adverse event was dyspnoea, which resolved with discontinuation of bosentan.

#### Concomitant use with other medicinal products

Concomitant use of Tracleer and cyclosporine A is contraindicated (see sections 4.3 and 4.5).

Concomitant use of Tracleer with glibenclamide, fluconazole and rifampicin is not recommended. For further details please refer to section 4.5.

Concomitant administration of both a CYP3A4 inhibitor and a CYP2C9 inhibitor with Tracleer should be avoided (see section 4.5).

### **4.5 Interaction with other medicinal products and other forms of interaction**

Bosentan is an inducer of the cytochrome P450 (CYP) isoenzymes CYP2C9 and CYP3A4. *In vitro* data also suggest an induction of CYP2C19. Consequently, plasma concentrations of substances metabolised by these isoenzymes will be decreased when Tracleer is co-administered. The possibility of altered efficacy of medicinal products metabolised by these isoenzymes should be considered. The dosage of these products may need to be adjusted after initiation, dose change or discontinuation of concomitant Tracleer treatment.

Bosentan is metabolised by CYP2C9 and CYP3A4. Inhibition of these isoenzymes may increase the plasma concentration of bosentan (see ketoconazole). The influence of CYP2C9 inhibitors on bosentan concentration has not been studied. The combination should be used with caution.

*Fluconazole and other inhibitors of both CYP2C9 and CYP3A4:* Co-administration with fluconazole, which inhibits mainly CYP2C9, but to some extent also CYP3A4, could lead to large increases in plasma concentrations of bosentan. The combination is not recommended. For the same reason, concomitant administration of both a potent CYP3A4 inhibitor (such as ketoconazole, itraconazole or ritonavir) and a CYP2C9 inhibitor (such as voriconazole) with Tracleer is not recommended.

*Cyclosporine A:* Co-administration of Tracleer and cyclosporine A (a calcineurin inhibitor) is contraindicated (see section 4.3). When co-administered, initial trough concentrations of bosentan were approximately 30-fold higher than those measured after bosentan alone. At steady state, bosentan plasma concentrations were 3- to 4-fold higher than with bosentan alone. The mechanism of this interaction is most likely inhibition of transport protein-mediated uptake of bosentan into hepatocytes by cyclosporine. The blood concentrations of cyclosporine A (a CYP3A4 substrate) decreased by approximately 50%. This is most likely due to induction of CYP3A4 by bosentan.

*Tacrolimus, sirolimus:* Co-administration of tacrolimus or sirolimus and Tracleer has not been studied in man but co-administration of tacrolimus or sirolimus and Tracleer may result in increased plasma concentrations of bosentan in analogy to co-administration with cyclosporine A. Concomitant Tracleer may reduce the plasma concentrations of tacrolimus and sirolimus. Therefore, concomitant use of Tracleer and tacrolimus or sirolimus is not advisable. Patients in need of the combination should be closely monitored for adverse events related to Tracleer and for tacrolimus and sirolimus blood concentrations.

*Glibenclamide:* Co-administration of bosentan 125 mg twice daily for 5 days decreased the plasma concentrations of glibenclamide (a CYP3A4 substrate) by 40%, with potential significant decrease of the hypoglycaemic effect. The plasma concentrations of bosentan were also decreased by 29%. In addition, an increased incidence of elevated aminotransferases was observed in patients receiving concomitant therapy. Both glibenclamide and bosentan inhibit the bile salt export pump, which could explain the elevated aminotransferases. This combination should not be used. No drug-drug interaction data are available with the other sulfonylureas.

*Rifampicin:* Co-administration in 9 healthy subjects for 7 days of bosentan 125 mg twice daily with rifampicin, a potent inducer of CYP2C9 and CYP3A4, decreased the plasma concentrations of bosentan by 58%, and this decrease could achieve almost 90% in an individual case. As a result, a significantly reduced effect of bosentan is expected when it is co-administered with rifampicin. Concomitant use of rifampicin and Tracleer is not recommended. Data on other CYP3A4 inducers, e.g., carbamazepine, phenobarbital, phenytoin and St. John's wort are lacking, but their concomitant administration is expected to lead to reduced systemic exposure to bosentan. A clinically significant reduction of efficacy cannot be excluded.

*Lopinavir+ritonavir (and other ritonavir-boosted protease inhibitors):* Co-administration of bosentan 125 mg twice daily and lopinavir+ritonavir 400+100 mg twice daily for 9.5 days in healthy volunteers resulted in initial trough plasma concentrations of bosentan that were approximately 48-fold higher than those measured after bosentan administered alone. On day 9, plasma concentrations of bosentan were approximately 5-fold higher than with bosentan administered alone. Inhibition by ritonavir of transport protein-mediated uptake into hepatocytes and of CYP3A4, thereby reducing the clearance of bosentan, most likely causes this interaction. When administered concomitantly with lopinavir+ritonavir, or other ritonavir-boosted protease inhibitors, the patient's tolerability of Tracleer should be monitored.

After co-administration of bosentan for 9.5 days, the plasma exposures to lopinavir and ritonavir decreased to a clinically non-significant extent (by approximately 14% and 17%, respectively). However, full induction by bosentan might not have been reached and a further decrease of protease inhibitors cannot be excluded. Appropriate monitoring of the HIV therapy is recommended. Similar effects would be expected with other ritonavir-boosted protease inhibitors (see section 4.4).

*Other antiretroviral agents:* No specific recommendation can be made with regard to other available antiretroviral agents due to the lack of data. Due to the marked hepatotoxicity of nevirapine, which could add to bosentan liver toxicity, this combination is not recommended.



*Hormonal contraceptives:* Co-administration of bosentan 125 mg twice daily for 7 days with a single dose of oral contraceptive containing norethisterone 1 mg + ethinyl estradiol 35 mcg decreased the AUC of norethisterone and ethinyl estradiol by 14% and 31%, respectively. However, decreases in exposure were as much as 56% and 66%, respectively, in individual subjects. Therefore, hormone-based contraceptives alone, regardless of the route of administration (i.e., oral, injectable, transdermal or implantable forms), are not considered as reliable methods of contraception (see sections 4.4 and 4.6).

*Warfarin:* Co-administration of bosentan 500 mg twice daily for 6 days decreased the plasma concentrations of both S-warfarin (a CYP2C9 substrate) and R-warfarin (a CYP3A4 substrate) by 29% and 38%, respectively. Clinical experience with concomitant administration of bosentan with warfarin in patients with PAH did not result in clinically relevant changes in International Normalized Ratio (INR) or warfarin dose (baseline versus end of the clinical studies). In addition, the frequency of changes in warfarin dose during the studies due to changes in INR or due to adverse events was similar among bosentan- and placebo-treated patients. No dose adjustment is needed for warfarin and similar oral anticoagulant agents when bosentan is initiated, but intensified monitoring of INR is recommended, especially during bosentan initiation and the up-titration period.

*Simvastatin:* Co-administration of bosentan 125 mg twice daily for 5 days decreased the plasma concentrations of simvastatin (a CYP3A4 substrate) and its active  $\beta$ -hydroxy acid metabolite by 34% and 46%, respectively. The plasma concentrations of bosentan were not affected by concomitant simvastatin. Monitoring of cholesterol levels and subsequent dosage adjustment should be considered.

*Ketoconazole:* Co-administration for 6 days of bosentan 62.5 mg twice daily with ketoconazole, a potent CYP3A4 inhibitor, increased the plasma concentrations of bosentan approximately 2-fold. No dose adjustment of Tracleer is considered necessary. Although not demonstrated through *in vivo* studies, similar increases in bosentan plasma concentrations are expected with the other potent CYP3A4 inhibitors (such as itraconazole or ritonavir). However, when combined with a CYP3A4 inhibitor, patients who are poor metabolisers of CYP2C9 are at risk of increases in bosentan plasma concentrations that may be of higher magnitude, thus leading to potential harmful adverse events.

*Epoprostenol:* Limited data obtained from a study (AC-052-356 [BREATHE-3]) in which 10 paediatric patients received the combination of bosentan and epoprostenol indicate that after both single- and multiple-dose administration, the  $C_{max}$  and AUC values of bosentan were similar in patients with or without continuous infusion of epoprostenol (see section 5.1).

*Sildenafil:* Co-administration of bosentan 125 mg twice daily (steady state) with sildenafil 80 mg three times a day (at steady state) concomitantly administered during 6 days in healthy volunteers resulted in a 63% decrease in the sildenafil AUC and a 50% increase in the bosentan AUC. Caution is recommended in the case of co-administration.

*Digoxin:* Co-administration for 7 days of bosentan 500 mg twice daily with digoxin decreased the AUC,  $C_{max}$  and  $C_{min}$  of digoxin by 12%, 9% and 23%, respectively. The mechanism for this interaction may be induction of P-glycoprotein. This interaction is unlikely to be of clinical relevance.

### Paediatric population

Interaction studies have only been performed in adults.

## **4.6 Fertility, pregnancy and lactation**

### Pregnancy

Studies in animals have shown reproductive toxicity (teratogenicity, embryotoxicity; see section 5.3). There are no reliable data on the use of Tracleer in pregnant women. The potential risk for humans is still unknown. Tracleer is contraindicated in pregnancy (see section 4.3).

### Women of childbearing potential

Before the initiation of Tracleer treatment in women of childbearing potential, the absence of pregnancy should be checked, appropriate advice on reliable methods of contraception provided, and reliable contraception initiated. Patients and prescribers must be aware that due to potential pharmacokinetic interactions, Tracleer may render hormonal contraceptives ineffective (see section 4.5). Therefore, women of childbearing potential must not use hormonal contraceptives (including oral, injectable, transdermal or implantable forms) as the sole method of contraception but must use an additional or an alternative reliable method of contraception. If there is any doubt about what contraceptive advice should be given to the individual patient, consultation with a gynaecologist is recommended. Because of possible hormonal contraception failure during Tracleer treatment, and also bearing in mind the risk that pulmonary hypertension severely deteriorates with pregnancy, monthly pregnancy tests during treatment with Tracleer are recommended to allow early detection of pregnancy.

### Breast-feeding

It is not known whether bosentan is excreted into human breast milk. Breast-feeding is not recommended during treatment with Tracleer.

### Fertility

Animal studies showed testicular effects (see section 5.3). In a study investigating the effects of bosentan on testicular function in male PAH patients, 8 out of 24 patients showed a decreased sperm concentration from baseline of at least 42% after 3 or 6 months of treatment with bosentan. Based on these findings and preclinical data, it cannot be excluded that bosentan may have a detrimental effect on spermatogenesis in men. In male children, a long-term impact on fertility after treatment with bosentan cannot be excluded.

## **4.7 Effects on ability to drive and use machines**

No specific studies have been conducted to assess the direct effect of Tracleer on the ability to drive and use machines. However, Tracleer may induce hypotension, with symptoms of dizziness, blurred vision or syncope that could affect the ability to drive or use machines.

## **4.8 Undesirable effects**

In 20 placebo-controlled studies, conducted in a variety of therapeutic indications, a total of 2,486 patients were treated with bosentan at daily doses ranging from 100 mg to 2000 mg and 1,838 patients were treated with placebo. The mean treatment duration was 45 weeks. Adverse reactions were defined as events occurring in at least 1% of patients on bosentan and at a frequency at least 0.5% more than on placebo. The most frequent adverse reactions are headache (11.5%), oedema / fluid retention (13.2%), abnormal liver function test (10.9%) and anaemia / haemoglobin decrease (9.9%).

Treatment with bosentan has been associated with dose-dependent elevations in liver aminotransferases and decreases in haemoglobin concentration (see section 4.4).

Adverse reactions observed in 20 placebo-controlled studies and post-marketing experience with bosentan are ranked according to frequency using the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data).

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness. No clinically relevant differences in adverse reactions were observed between the overall dataset and the approved indications.

<b>System organ class</b>	<b>Frequency</b>	<b>Adverse reaction</b>
Blood and lymphatic system disorders	Common	Anaemia, haemoglobin decrease, (see section 4.4)
	Not known	Anaemia or haemoglobin decreases requiring red blood cell transfusion <sup>1</sup>
	Uncommon	Thrombocytopenia <sup>1</sup>
	Uncommon	Neutropenia, leukopenia <sup>1</sup>
Immune system disorders	Common	Hypersensitivity reactions (including dermatitis, pruritus and rash) <sup>2</sup>
	Rare	Anaphylaxis and/or angioedema <sup>1</sup>
Nervous system disorders	Very common	Headache <sup>3</sup>
	Common	Syncope <sup>1,4</sup>
Eye disorders	Not known	Blurred vision <sup>1</sup>
Cardiac disorders	Common	Palpitations <sup>1,4</sup>
Vascular disorders	Common	Flushing
	Common	Hypotension <sup>1,4</sup>
Respiratory, thoracic and mediastinal disorders	Common	Nasal congestion <sup>1</sup>
Gastrointestinal disorders	Common	Gastrooesophageal reflux disease Diarrhoea
Hepatobiliary disorders	Very common	Abnormal liver function test (see section 4.4)
	Uncommon	Aminotransferase elevations associated with hepatitis (including possible exacerbation of underlying hepatitis) and/or jaundice <sup>1</sup> (see section 4.4)
	Rare	Liver cirrhosis, liver failure <sup>1</sup>
Skin and subcutaneous disorders	Common	Erythema
General disorders and administration site conditions	Very common	Oedema, fluid retention <sup>5</sup>

<sup>1</sup> Data derived from post-marketing experience, frequencies based on statistical modelling of placebo-controlled clinical trial data.

<sup>2</sup> Hypersensitivity reactions were reported in 9.9% of patients on bosentan and 9.1% of patients on placebo.

<sup>3</sup> Headache was reported in 11.5% of patients on bosentan and 9.8% of patients on placebo.

<sup>4</sup> These types of reactions can also be related to the underlying disease.

<sup>5</sup> Oedema or fluid retention was reported in 13.2% of patients on bosentan and 10.9% of patients on placebo.

In the post-marketing period rare cases of unexplained hepatic cirrhosis were reported after prolonged therapy with Tracleer in patients with multiple co-morbidities and therapies with medicinal products. There have also been rare reports of liver failure. These cases reinforce the importance of strict adherence to the monthly schedule for monitoring of liver function for the duration of treatment with Tracleer (see section 4.4).

### Paediatric population

#### *Uncontrolled clinical studies in paediatric patients*

The safety profile in the first paediatric uncontrolled study performed with the film-coated tablet (BREATHE-3: n = 19, median age 10 years [range 3–15 years], open-label bosentan 2 mg/kg twice daily; treatment duration 12 weeks) was similar to that observed in the pivotal trials in adult patients with PAH. In BREATHE-3, the most frequent adverse reactions were flushing (21%), headache, and abnormal liver function test (each 16%).

A pooled analysis of uncontrolled paediatric studies conducted in PAH with the bosentan 32 mg dispersible tablet formulation (FUTURE 1/2, FUTURE 3/Extension) included a total of 100 children treated with bosentan 2 mg/kg twice daily (n = 33), 2 mg/kg three times daily (n = 31), or 4 mg/kg twice daily (n = 36). At enrolment, six patients were between 3 months and 1 year old, 15 children were between 1 and less than 2 years old, and 79 were between 2 and 12 years old. The median treatment duration was 71.8 weeks (range 0.4–258 weeks).

The safety profile in this pooled analysis of uncontrolled paediatric studies was similar to that observed in the pivotal trials in adult patients with PAH except for infections, which were more frequently reported than in adults (69.0% vs 41.3%). This difference in infection frequency may in part be due to the longer median treatment exposure in the paediatric set (median 71.8 weeks) compared with the adult set (median 17.4 weeks). The most frequent adverse events were upper respiratory tract infections (25%), pulmonary (arterial) hypertension (20%), nasopharyngitis (17%), pyrexia (15%), vomiting (13%), bronchitis (10%), abdominal pain (10%), and diarrhoea (10%). There was no relevant difference in adverse event frequencies between patients above and below the age of 2 years; however, this is based on only 21 children less than 2 years, including 6 patients between 3 months to 1 year of age. Adverse events of liver abnormalities and anaemia/haemoglobin decrease occurred in 9% and 5% of patients, respectively.

In a randomised placebo-controlled study, conducted in PPHN patients (FUTURE-4), a total of 13 neonates were treated with the bosentan dispersible tablet formulation at a dose of 2 mg/kg twice daily (8 patients were on placebo). The median bosentan and placebo treatment duration was, respectively, 4.5 days (range 0.5–10.0 days) and 4.0 days (range 2.5–6.5 days). The most frequent adverse events in the bosentan- and placebo-treated patients were, respectively, anaemia or haemoglobin decrease (7 and 2 patients), generalised oedema (3 and 0 patients), and vomiting (2 and 0 patients).

### Laboratory abnormalities

#### *Liver test abnormalities*

In the clinical programme, dose-dependent elevations in liver aminotransferases generally occurred within the first 26 weeks of treatment, usually developed gradually, and were mainly asymptomatic. In the post-marketing period rare cases of liver cirrhosis and liver failure have been reported.

The mechanism of this adverse effect is unclear. These elevations in aminotransferases may reverse spontaneously while continuing treatment with the maintenance dose of Tracleer or after dose reduction, but interruption or cessation may be necessary (see section 4.4).

In the 20 integrated placebo-controlled studies, elevations in liver aminotransferases  $\geq 3 \times \text{ULN}$  were observed in 11.2% of the bosentan-treated patients as compared to 2.4% of the placebo-treated patients. Elevations to  $\geq 8 \times \text{ULN}$  were seen in 3.6% of the bosentan-treated patients and 0.4% of the placebo-treated patients. Elevations in aminotransferases were associated with elevated bilirubin ( $\geq 2 \times \text{ULN}$ ) without evidence of biliary obstruction in 0.2% (5 patients) on bosentan and 0.3% (6 patients) on placebo.

In the pooled analysis of 100 PAH children from uncontrolled paediatric studies FUTURE 1/2 and FUTURE 3/Extension, elevations in liver aminotransferases  $\geq 3 \times \text{ULN}$  were observed in 2% of patients.

In the FUTURE-4 study including 13 neonates with PPHN treated with bosentan 2 mg/kg twice daily for less than 10 days (range 0.5–10.0 days), there were no cases of liver aminotransferases  $\geq 3 \times \text{ULN}$  during treatment, but one case of hepatitis occurred 3 days after the end of bosentan treatment.

### *Haemoglobin*

In the adult placebo-controlled studies, a decrease in haemoglobin concentration to below 10 g/dL from baseline was reported in 8.0% of bosentan-treated patients and 3.9% of placebo-treated patients (see section 4.4).

In the pooled analysis of 100 PAH children from uncontrolled paediatric studies FUTURE 1/2 and FUTURE 3/Extension, a decrease in haemoglobin concentration from baseline to below 10 g/dL was reported in 10.0% of patients. There was no decrease to below 8 g/dL.

In the FUTURE-4 study, 6 out of 13 bosentan-treated neonates with PPHN experienced a decrease in haemoglobin from within the reference range at baseline to below the lower limit of normal during the treatment.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via **the national reporting system** listed in [Appendix V](#).

## **4.9 Overdose**

Bosentan has been administered as a single dose of up to 2,400 mg to healthy subjects and up to 2,000 mg/day for 2 months in patients with a disease other than pulmonary hypertension. The most common adverse reaction was headache of mild to moderate intensity.

Massive overdose may result in pronounced hypotension requiring active cardiovascular support. In the post-marketing period there was one reported overdose of 10,000 mg of Tracleer taken by an adolescent male patient. He had symptoms of nausea, vomiting, hypotension, dizziness, sweating and blurred vision. He recovered completely within 24 hours with blood pressure support. Note: bosentan is not removed through dialysis.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: other antihypertensives, ATC code: C02KX01

## Mechanism of action

Bosentan is a dual endothelin receptor antagonist (ERA) with affinity for both endothelin A and B (ET<sub>A</sub> and ET<sub>B</sub>) receptors. Bosentan decreases both pulmonary and systemic vascular resistance resulting in increased cardiac output without increasing heart rate.

The neurohormone endothelin-1 (ET-1) is one of the most potent vasoconstrictors known and can also promote fibrosis, cell proliferation, cardiac hypertrophy and remodelling, and is pro-inflammatory. These effects are mediated by endothelin binding to ET<sub>A</sub> and ET<sub>B</sub> receptors located in the endothelium and vascular smooth muscle cells. ET-1 concentrations in tissues and plasma are increased in several cardiovascular disorders and connective tissue diseases, including PAH, scleroderma, acute and chronic heart failure, myocardial ischaemia, systemic hypertension and atherosclerosis, suggesting a pathogenic role of ET-1 in these diseases. In PAH and heart failure, in the absence of endothelin receptor antagonism, elevated ET-1 concentrations are strongly correlated with the severity and prognosis of these diseases.

Bosentan competes with the binding of ET-1 and other ET peptides to both ET<sub>A</sub> and ET<sub>B</sub> receptors, with a slightly higher affinity for ET<sub>A</sub> receptors ( $K_i = 4.1\text{--}43$  nanomolar) than for ET<sub>B</sub> receptors ( $K_i = 38\text{--}730$  nanomolar). Bosentan specifically antagonises ET receptors and does not bind to other receptors.

## Efficacy

### *Animal models*

In animal models of pulmonary hypertension, chronic oral administration of bosentan reduced pulmonary vascular resistance and reversed pulmonary vascular and right ventricular hypertrophy. In an animal model of pulmonary fibrosis, bosentan reduced collagen deposition in the lungs.

### *Efficacy in adult patients with pulmonary arterial hypertension*

Two randomised, double-blind, multi-centre, placebo-controlled studies have been conducted in 32 (study AC-052-351) and 213 (study AC-052-352 [BREATHE-1]) adult patients with WHO functional class III–IV PAH (primary pulmonary hypertension or pulmonary hypertension secondary mainly to scleroderma). After 4 weeks of bosentan 62.5 mg twice daily, the maintenance doses studied in these studies were 125 mg twice daily in AC-052-351, and 125 mg twice daily and 250 mg twice daily in AC-052-352.

Bosentan was added to patients' current therapy, which could include a combination of anticoagulants, vasodilators (e.g., calcium channel blockers), diuretics, oxygen and digoxin, but not epoprostenol. Control was placebo plus current therapy.

The primary endpoint for each study was change in 6-minute walk distance at 12 weeks for the first study and 16 weeks for the second study. In both studies, treatment with bosentan resulted in significant increases in exercise capacity. The placebo-corrected increases in walk distance compared with baseline were 76 metres ( $p = 0.02$ ; t-test) and 44 metres ( $p = 0.0002$ ; Mann-Whitney U test) at the primary endpoint of each study, respectively. The differences between the two groups, 125 mg twice daily and 250 mg twice daily, were not statistically significant but there was a trend towards improved exercise capacity in the group treated with 250 mg twice daily.

The improvement in walk distance was apparent after 4 weeks of treatment, was clearly evident after 8 weeks of treatment and was maintained for up to 28 weeks of double-blind treatment in a subset of the patient population.

In a retrospective responder analysis based on change in walking distance, WHO functional class and dyspnoea of the 95 patients randomised to bosentan 125 mg twice daily in the placebo-controlled studies, it was found that at week 8, 66 patients had improved, 22 were stable and 7 had deteriorated. Of the 22 patients stable at week 8, 6 improved at week 12/16 and 4 deteriorated compared with baseline. Of the 7 patients who deteriorated at week 8, 3 improved at week 12/16 and 4 deteriorated compared with baseline.

Invasive haemodynamic parameters were assessed in the first study only. Treatment with bosentan led to a significant increase in cardiac index associated with a significant reduction in pulmonary artery pressure, pulmonary vascular resistance and mean right atrial pressure.

A reduction in symptoms of PAH was observed with bosentan treatment. Dyspnoea measurement during walk tests showed an improvement in bosentan-treated patients. In the AC-052-352 study, 92% of the 213 patients were classified at baseline as WHO functional class III and 8% as class IV. Treatment with bosentan led to a WHO functional class improvement in 42.4% of patients (placebo 30.4%). The overall change in WHO functional class during both studies was significantly better among bosentan-treated patients as compared with placebo-treated patients. Treatment with bosentan was associated with a significant reduction in the rate of clinical worsening compared with placebo at 28 weeks (10.7% vs 37.1%, respectively;  $p = 0.0015$ ).

In a randomised, double-blind, multi-centre, placebo-controlled study (AC-052-364 [EARLY]), 185 PAH patients in WHO functional class II (mean baseline 6-minute walk distance of 435 metres) received bosentan 62.5 mg twice daily for 4 weeks followed by 125 mg twice daily ( $n = 93$ ), or placebo ( $n = 92$ ) for 6 months. Enrolled patients were PAH-treatment-naïve ( $n = 156$ ) or on a stable dose of sildenafil ( $n = 29$ ). The co-primary endpoints were percentage change from baseline in pulmonary vascular resistance (PVR) and change from baseline in 6-minute walk distance to Month 6 versus placebo. The table below illustrates the pre-specified protocol analyses.

	PVR (dyn.sec/cm <sup>5</sup> )		6-Minute Walk Distance (m)	
	Placebo (n=88)	Bosentan (n=80)	Placebo (n=91)	Bosentan (n=86)
Baseline (BL); mean (SD)	802 (365)	851 (535)	431 (92)	443 (83)
Change from BL; mean (SD)	128 (465)	-69 (475)	-8 (79)	11 (74)
Treatment effects	-22.6%		19	
95% CL	-34, -10		-4, 42	
P-value	< 0.0001		0.0758	

CL = confidence limit; PVR = pulmonary vascular resistance; SD = standard deviation.

Treatment with bosentan was associated with a reduction in the rate of clinical worsening, defined as a composite of symptomatic progression, hospitalisation for PAH and death, compared with placebo (proportional risk reduction 77%, 95% confidence interval [CI] 20–94%,  $p = 0.0114$ ). The treatment effect was driven by improvement in the component symptomatic progression. There was one hospitalisation related to PAH worsening in the bosentan group and three hospitalisations in the placebo group. Only one death occurred in each treatment group during the 6-month double-blind study period, therefore no conclusion can be drawn on survival.

Long-term data were generated from all 173 patients who were treated with bosentan in the controlled phase and/or were switched from placebo to bosentan in the open-label extension phase of the EARLY study. The mean duration of exposure to bosentan treatment was  $3.6 \pm 1.8$  years (up to 6.1 years), with 73% of patients treated for at least 3 years and 62% for at least 4 years. Patients could receive additional PAH treatment as required in the open-label extension. The majority of patients were diagnosed with idiopathic or heritable PAH (61%). Overall, 78% of patients remained in WHO functional class II. Kaplan-Meier estimates of survival were 90% and 85% at 3 and 4 years after the start of treatment, respectively. At the same timepoints, 88% and 79% of patients remained free from

PAH worsening (defined as all-cause death, lung transplantation, atrial septostomy or start of intravenous or subcutaneous prostanoid treatment). The relative contributions of previous placebo treatment in the double-blind phase and of other medications started during the open-label extension period are unknown.

In a prospective, multi-centre, randomised, double-blind, placebo-controlled study (AC-052-405 [BREATHE-5]), patients with PAH WHO functional class III and Eisenmenger physiology associated with congenital heart disease received bosentan 62.5 mg twice daily for 4 weeks, then 125 mg twice daily for a further 12 weeks (n = 37, of whom 31 had a predominantly right to left, bidirectional shunt). The primary objective was to show that bosentan did not worsen hypoxaemia. After 16 weeks, the mean oxygen saturation was increased in the bosentan group by 1.0% (95% CI -0.7%–2.8%) as compared to the placebo group (n = 17), showing that bosentan did not worsen hypoxaemia. The mean pulmonary vascular resistance was significantly reduced in the bosentan group (with a predominant effect observed in the subgroup of patients with bidirectional intracardiac shunt). After 16 weeks, the mean placebo-corrected increase in 6-minute walk distance was 53 metres (p = 0.0079), reflecting improvement in exercise capacity. Twenty-six patients continued to receive bosentan in the 24-week open-label extension phase (AC-052-409) of the BREATHE-5 study (mean duration of treatment = 24.4 ± 2.0 weeks) and, in general, efficacy was maintained.

An open-label, non-comparative study (AC-052-362 [BREATHE-4]) was performed in 16 patients with WHO functional class III PAH associated with HIV infection. Patients were treated with bosentan 62.5 mg twice daily for 4 weeks followed by 125 mg twice daily for a further 12 weeks. After 16 weeks' treatment, there were significant improvements from baseline in exercise capacity: the mean increase in 6-minute walk distance was 91.4 metres from 332.6 metres on average at baseline (p < 0.001). No formal conclusion can be drawn regarding the effects of bosentan on antiretroviral drug efficacy (see also section 4.4).

There are no studies to demonstrate beneficial effects of Tracleer treatment on survival. However, long-term vital status was recorded for all 235 patients who were treated with bosentan in the two pivotal placebo-controlled studies (AC-052-351 and AC-052-352) and/or their two uncontrolled, open-label extensions. The mean duration of exposure to bosentan was 1.9 years ± 0.7 years (min: 0.1 years; max: 3.3 years) and patients were observed for a mean of 2.0 ± 0.6 years. The majority of patients were diagnosed as primary pulmonary hypertension (72%) and were in WHO functional class III (84%). In this total population, Kaplan-Meier estimates of survival were 93% and 84% 1 and 2 years after the start of treatment with bosentan, respectively. Survival estimates were lower in the subgroup of patients with PAH secondary to systemic sclerosis. The estimates may have been influenced by the initiation of epoprostenol treatment in 43/235 patients.

#### *Studies performed in children with pulmonary arterial hypertension*

##### **BREATHE-3 (AC-052-356)**

Bosentan film-coated tablets were evaluated in an open-label uncontrolled study in 19 paediatric patients with PAH aged 3 to 15 years. This study was primarily designed as a pharmacokinetic study (see section 5.2). Patients had primary pulmonary hypertension (10 patients) or PAH related to congenital heart diseases (9 patients) and were in WHO functional class II (n = 15, 79%) or class III (n = 4, 21%) at baseline. Patients were divided into three body-weight groups and dosed with bosentan at approximately 2 mg/kg twice daily for 12 weeks. Half of the patients in each group were already being treated with intravenous epoprostenol and the dose of epoprostenol remained constant for the duration of the study.

Haemodynamics were measured in 17 patients. The mean increase from baseline in cardiac index was 0.5 L/min/m<sup>2</sup>, the mean decrease in mean pulmonary arterial pressure was 8 mmHg, and the mean decrease in PVR was 389 dyn·sec·cm<sup>-5</sup>. These haemodynamic improvements from baseline were



similar with or without co-administration of epoprostenol. Changes in exercise test parameters at week 12 from baseline were highly variable and none were significant.

#### FUTURE 1/2 (AC-052-365/AC-052-367)

FUTURE 1 was an open-label, uncontrolled study that was conducted with the dispersible tablet formulation of bosentan administered at a maintenance dose of 4 mg/kg twice daily to 36 patients from 2 to 11 years of age. It was primarily designed as a pharmacokinetic study (see section 5.2). At baseline, patients had idiopathic (31 patients [86%]) or familial (5 patients [14%]) PAH, and were in WHO functional class II (n = 23, 64%) or class III (n = 13, 36%). In the FUTURE 1 study, the median exposure to study treatment was 13.1 weeks (range: 8.4 to 21.1). 33 of these patients were provided with continued treatment with bosentan dispersible tablets at a dose of 4 mg/kg twice daily in the FUTURE 2 uncontrolled extension phase for a median overall treatment duration of 2.3 years (range: 0.2 to 5.0 years). At baseline in FUTURE 1, 9 patients were taking epoprostenol. 9 patients were newly initiated on PAH-specific medication during the study. The Kaplan-Meier event-free estimate for worsening of PAH (death, lung transplantation, or hospitalisation for PAH worsening) at 2 years was 78.9%. The Kaplan-Meier estimate of overall survival at 2 years was 91.2%.

#### FUTURE 3 (AC-052-373)

In this open-label randomised study with the bosentan 32 mg dispersible tablet formulation, 64 children with stable PAH from 3 months to 11 years of age were randomised to 24 weeks' bosentan treatment 2 mg/kg twice daily (n = 33) or 2 mg/kg three times daily (n = 31). 43 (67.2%) were ≥ 2 years to 11 years old, 15 (23.4%) were between 1 and 2 years old, and 6 (9.4%) were between 3 months and 1 year old. The study was primarily designed as a pharmacokinetic study (see section 5.2), and efficacy endpoints were only exploratory. The aetiology of PAH, according to Dana Point classification, included idiopathic PAH (46%), heritable PAH (3%), associated PAH after corrective cardiac surgery (38%), and PAH related to congenital heart disease associated with systemic-to-pulmonary shunts, including Eisenmenger syndrome (13%). Patients were in WHO functional class I (n = 19, 29%), class II (n = 27, 42%) or class III (n = 18, 28%) at start of study treatment. At study entry, patients were treated with PAH medications (most frequently phosphodiesterase type-5 inhibitor [sildenafil] alone [35.9%], bosentan alone [10.9%], and a combination of bosentan, iloprost, and sildenafil [10.9%]) and continued their PAH treatment during the study.

At study start, less than half of the patients included (45.3% [29/64]) had bosentan treatment alone not combined with other PAH medication. 40.6% (26/64) remained on bosentan monotherapy during the 24 weeks of study treatment without experiencing PAH worsening. The analysis on the global population included (64 patients) showed that the majority had remained at least stable (i.e., without deterioration) based on non-paediatric-specific WHO functional class assessment (97% twice daily, 100% three times daily) and physician's global clinical impression (94% twice daily, 93% three times daily) during the treatment period. The Kaplan-Meier event-free estimate for worsening of PAH (death, lung transplantation, or hospitalisation for PAH worsening) at 24 weeks was 96.9% and 96.7% in the twice daily and three times daily groups, respectively.

There was no evidence of any clinical benefit with 2 mg/kg three times daily as compared to 2 mg/kg twice daily dosing.

#### *Study performed in neonates with persistent pulmonary hypertension of the newborn (PPHN):*

#### FUTURE 4 (AC-052-391)

This was a double-blind, placebo-controlled, randomised study in pre-term or term neonates (gestational age 36–42 weeks) with PPHN. Patients with suboptimal response to inhaled nitric oxide (iNO) despite at least 4 hours of continuous treatment were treated with bosentan dispersible tablets at 2 mg/kg twice daily (N = 13) or placebo (N = 8) via nasogastric tube as add-on therapy on top of iNO

until complete weaning of iNO or until treatment failure (defined as need for extra-corporeal membrane oxygenation [ECMO] or initiation of alternative pulmonary vasodilator), and for a maximum of 14 days.

The median exposure to study treatment was 4.5 (range: 0.5–10.0) days in the bosentan group and 4.0 (range: 2.5–6.5) days in the placebo group.

The results did not indicate an additional benefit of bosentan in this population:

- The median time to complete weaning from iNO was 3.7 days (95% confidence limits [CLs] 1.17, 6.95) on bosentan and 2.9 days (95% CLs 1.26, 4.23) on placebo ( $p = 0.34$ ).
- The median time to complete weaning from mechanical ventilation was 10.8 days (95% CLs 3.21, 12.21 days) on bosentan and 8.6 days (95% CLs 3.71, 9.66 days) on placebo ( $p = 0.24$ ).
- One patient in the bosentan group had treatment failure (need for ECMO as per protocol definition), which was declared based on increasing Oxygenation Index values within 8 h after the first study drug dose. This patient recovered within the 60-day follow-up period.

#### *Combination with epoprostenol*

The combination of bosentan and epoprostenol has been investigated in two studies: AC-052-355 (BREATHE-2) and AC-052-356 (BREATHE-3). AC-052-355 was a multi-centre, randomised, double-blind, parallel-group study of bosentan versus placebo in 33 patients with severe PAH who were receiving concomitant epoprostenol therapy. AC-052-356 was an open-label, uncontrolled study; 10 of the 19 paediatric patients were on concomitant bosentan and epoprostenol therapy during the 12-week study. The safety profile of the combination was not different from the one expected with each component and the combination therapy was well tolerated in children and adults. The clinical benefit of the combination has not been demonstrated.

#### *Systemic sclerosis with digital ulcer disease*

Two randomised, double-blind, multi-centre, placebo-controlled studies have been conducted in 122 (study AC-052-401 [RAPIDS-1]) and 190 (study AC-052-331 [RAPIDS-2]) adult patients with systemic sclerosis and digital ulcer disease (either ongoing digital ulcers or a history of digital ulcers within the previous year). In study AC-052-331, patients had to have at least one digital ulcer of recent onset, and across the two studies 85% of patients had ongoing digital ulcer disease at baseline. After 4 weeks of bosentan 62.5 mg twice daily, the maintenance dose studied in both these studies was 125 mg twice daily. The duration of double-blind therapy was 16 weeks in study AC-052-401, and 24 weeks in study AC-052-331.

Background treatments for systemic sclerosis and digital ulcers were permitted if they remained constant for at least 1 month prior to the start of treatment and during the double-blind study period.

The number of new digital ulcers from baseline to study endpoint was a primary endpoint in both studies. Treatment with bosentan resulted in fewer new digital ulcers for the duration of therapy, compared with placebo. In study AC-052-401, during 16 weeks of double-blind therapy, patients in the bosentan group developed a mean of 1.4 new digital ulcers vs 2.7 new digital ulcers in the placebo group ( $p = 0.0042$ ). In study AC-052-331, during 24 weeks of double-blind therapy, the corresponding figures were 1.9 vs 2.7 new digital ulcers, respectively ( $p = 0.0351$ ). In both studies, patients on bosentan were less likely to develop multiple new digital ulcers during the study and took longer to develop each successive new digital ulcer than did those on placebo. The effect of bosentan on reduction of the number of new digital ulcers was more pronounced in patients with multiple digital ulcers.

No effect of bosentan on time to healing of digital ulcers was observed in either study.

## 5.2 Pharmacokinetic properties

The pharmacokinetics of bosentan have mainly been documented in healthy subjects. Limited data in patients show that the exposure to bosentan in adult PAH patients is approximately 2-fold greater than in healthy adult subjects.

In healthy subjects, bosentan displays dose- and time-dependent pharmacokinetics. Clearance and volume of distribution decrease with increased intravenous doses and increase with time. After oral administration, the systemic exposure is proportional to dose up to 500 mg. At higher oral doses,  $C_{\max}$  and AUC increase less than proportionally to the dose.

### Absorption

In healthy subjects, the absolute bioavailability of bosentan is approximately 50% and is not affected by food. The maximum plasma concentrations are attained within 3–5 hours.

### Distribution

Bosentan is highly bound (> 98%) to plasma proteins, mainly albumin. Bosentan does not penetrate into erythrocytes.

A volume of distribution ( $V_{ss}$ ) of about 18 litres was determined after an intravenous dose of 250 mg.

### Biotransformation and elimination

After a single intravenous dose of 250 mg, the clearance was 8.2 L/h. The terminal elimination half-life ( $t_{1/2}$ ) is 5.4 hours.

Upon multiple dosing, plasma concentrations of bosentan decrease gradually to 50–65% of those seen after single dose administration. This decrease is probably due to auto-induction of metabolising liver enzymes. Steady-state conditions are reached within 3–5 days.

Bosentan is eliminated by biliary excretion following metabolism in the liver by the cytochrome P450 isoenzymes, CYP2C9 and CYP3A4. Less than 3% of an administered oral dose is recovered in urine.

Bosentan forms three metabolites and only one of these is pharmacologically active. This metabolite is mainly excreted unchanged via the bile. In adult patients, the exposure to the active metabolite is greater than in healthy subjects. In patients with evidence of the presence of cholestasis, the exposure to the active metabolite may be increased.

Bosentan is an inducer of CYP2C9 and CYP3A4 and possibly also of CYP2C19 and the P-glycoprotein. *In vitro*, bosentan inhibits the bile salt export pump in hepatocyte cultures.

*In vitro* data demonstrated that bosentan had no relevant inhibitory effect on the CYP isoenzymes tested (CYP1A2, 2A6, 2B6, 2C8, 2C9, 2D6, 2E1, 3A4). Consequently, bosentan is not expected to increase the plasma concentrations of medicinal products metabolised by these isoenzymes.

## Pharmacokinetics in special populations

Based on the investigated range of each variable, it is not expected that the pharmacokinetics of bosentan will be influenced by gender, body weight, race, or age in the adult population to any relevant extent.

### *Children*

Pharmacokinetics were studied in paediatric patients in 4 clinical studies (BREATHE-3, FUTURE 1, FUTURE-3 and FUTURE-4; see section 5.1). Due to limited data in children below 2 years of age, pharmacokinetics remain not well characterised in this age category.

Study AC-052-356 (BREATHE-3) evaluated the pharmacokinetics of single and multiple oral doses of the film-coated tablet formulation of bosentan in 19 children aged from 3 to 15 years with PAH who were dosed on the basis of body weight with 2 mg/kg twice daily. In this study, the exposure to bosentan decreased with time in a manner consistent with the known auto-induction properties of bosentan. The mean AUC (CV%) values of bosentan in paediatric patients treated with 31.25, 62.5 or 125 mg twice daily were 3,496 (49), 5,428 (79), and 6,124 (27) ng·h/mL, respectively, and were lower than the value of 8,149 (47) ng·h/mL observed in adult patients with PAH receiving 125 mg twice daily. At steady state, the systemic exposures in paediatric patients weighing 10–20 kg, 20–40 kg and > 40 kg were 43%, 67% and 75%, respectively, of the adult systemic exposure.

In study AC-052-365 (FUTURE 1), dispersible tablets were administered in 36 PAH children aged from 2 to 11 years. No dose proportionality was observed, as steady-state bosentan plasma concentrations and AUCs were similar at oral doses of 2 and 4 mg/kg (AUC<sub>τ</sub>: 3,577 ng·h/mL and 3,371 ng·h/mL for 2 mg/kg twice daily and 4 mg/kg twice daily, respectively). The average exposure to bosentan in these paediatric patients was about half the exposure in adult patients at the 125 mg twice daily maintenance dose but showed a large overlap with the exposures in adults.

In study AC-052-373 (FUTURE 3), using dispersible tablets, the exposure to bosentan in the patients treated with 2 mg/kg twice daily was comparable to that in the FUTURE 1 study. In the overall population (n = 31), 2 mg/kg twice daily resulted in a daily exposure of 8,535 ng·h/mL; AUC<sub>τ</sub> was 4,268 ng·h/mL (CV: 61%). In patients between 3 months and 2 years the daily exposure was 7,879 ng·h/mL; AUC<sub>τ</sub> was 3,939 ng·h/mL (CV: 72%). In patients between 3 months and 1 year (n = 2) AUC<sub>τ</sub> was 5,914 ng·h/mL (CV: 85%), and in patients between 1 and 2 years (n = 7) AUC<sub>τ</sub> was 3,507 ng·h/mL (CV: 70%). In the patients above 2 years (n = 22) the daily exposure was 8,820 ng·h/mL; AUC<sub>τ</sub> was 4,410 ng·h/mL (CV: 58%). Dosing bosentan 2 mg/kg three times daily did not increase exposure; daily exposure was 7,275 ng·h/mL (CV: 83%, n = 27).

Based on the findings in studies BREATHE-3, FUTURE 1, and FUTURE-3, it appears that the exposure to bosentan reaches a plateau at lower doses in paediatric patients than in adults, and that doses higher than 2 mg/kg twice daily (4 mg/kg twice daily or 2 mg/kg three times daily) will not result in greater exposure to bosentan in paediatric patients.

In study AC-052-391 (FUTURE 4) conducted in neonates, bosentan concentrations increased slowly and continuously over the first dosing interval, resulting in low exposure (AUC<sub>0-12</sub> in whole blood: 164 ng·h/mL, n = 11). At steady state, AUC<sub>τ</sub> was 6,165 ng·h/mL (CV: 133%, n = 7), which is similar to the exposure observed in adult PAH patients receiving 125 mg twice daily and taking into account a blood/plasma distribution ratio of 0.6.

The consequences of these findings regarding hepatotoxicity are unknown. Gender and concomitant use of intravenous epoprostenol had no significant effect on the pharmacokinetics of bosentan.

### *Hepatic impairment*

In patients with mildly impaired liver function (Child-Pugh class A) no relevant changes in the pharmacokinetics have been observed. The steady-state AUC of bosentan was 9% higher and the AUC of the active metabolite, Ro 48-5033, was 33% higher in patients with mild hepatic impairment than in healthy volunteers.

The impact of moderately impaired liver function (Child-Pugh class B) on the pharmacokinetics of bosentan and its primary metabolite Ro 48-5033 was investigated in a study including 5 patients with pulmonary hypertension associated with portal hypertension and Child-Pugh class B hepatic impairment, and 3 patients with PAH from other causes and normal liver function. In the patients with Child-Pugh class B liver impairment, the mean (95% CI) steady-state AUC of bosentan was 360 (212–613) ng·h/mL, i.e., 4.7 times higher, and the mean (95% CI) AUC of the active metabolite Ro 48-5033 was 106 (58.4–192) ng·h/mL, i.e., 12.4 times higher than in the patients with normal liver function (bosentan: mean [95% CI] AUC: 76.1 [9.07–638] ng·h/mL; Ro 48-5033: mean [95% CI] AUC 8.57 [1.28–57.2] ng·h/ml). Though the number of patients included was limited and with high variability, these data indicate a marked increase in the exposure to bosentan and its primary metabolite Ro 48-5033 in patients with moderate liver function impairment (Child-Pugh class B).

The pharmacokinetics of bosentan have not been studied in patients with Child-Pugh class C hepatic impairment. Tracleer is contraindicated in patients with moderate to severe hepatic impairment, i.e., Child-Pugh class B or C (see section 4.3).

### *Renal impairment*

In patients with severe renal impairment (creatinine clearance 15–30 mL/min), plasma concentrations of bosentan decreased by approximately 10%. Plasma concentrations of bosentan metabolites increased about 2-fold in these patients as compared with subjects with normal renal function. No dose adjustment is required in patients with renal impairment. There is no specific clinical experience in patients undergoing dialysis. Based on physicochemical properties and the high degree of protein binding, bosentan is not expected to be removed from the circulation by dialysis to any significant extent (see section 4.2).

## **5.3 Preclinical safety data**

A 2-year carcinogenicity study in mice showed an increased combined incidence of hepatocellular adenomas and carcinomas in males, but not in females, at plasma concentrations about 2 to 4 times the plasma concentrations achieved at the therapeutic dose in humans. In rats, oral administration of bosentan for 2 years produced a small, significant increase in the combined incidence of thyroid follicular cell adenomas and carcinomas in males, but not in females, at plasma concentrations about 9 to 14 times the plasma concentrations achieved at the therapeutic dose in humans. Bosentan was negative in tests for genotoxicity. There was evidence of a mild thyroid hormonal imbalance induced by bosentan in rats. However, there was no evidence of bosentan affecting thyroid function (thyroxine, TSH) in humans.

The effect of bosentan on mitochondrial function is unknown.

Bosentan has been shown to be teratogenic in rats at plasma levels higher than 1.5 times the plasma concentrations achieved at the therapeutic dose in humans. Teratogenic effects, including malformations of the head and face and of the major vessels, were dose dependent. The similarities of the pattern of malformations observed with other ET receptor antagonists and in ET knock-out mice indicate a class effect. Appropriate precautions must be taken for women of childbearing potential (see sections 4.3, 4.4 and 4.6).

Development of testicular tubular atrophy and impaired fertility has been linked with chronic administration of endothelin receptor antagonists in rodents.

In fertility studies in male and female rats, no effects on sperm count, motility and viability, or on mating performance or fertility were observed at exposures that were 21 and 43 times the expected therapeutic level in humans, respectively; nor was there any adverse effect on the development of the pre-implantation embryo or on implantation.

Slightly increased incidence of testicular tubular atrophy was observed in rats given bosentan orally at doses as low as 125 mg/kg/day (about 4 times the maximum recommended human dose [MRHD] and the lowest doses tested) for two years but not at doses as high as 1,500 mg/kg/day (about 50 times the MRHD) for 6 months. In a juvenile rat toxicity study, where rats were treated from Day 4 *post partum* up to adulthood, decreased absolute weights of testes and epididymides, and reduced number of sperm in epididymides were observed after weaning. The NOAEL was 21 times (at Day 21 *post partum*) and 2.3 times (Day 69 *post partum*) the human therapeutic exposure, respectively.

However, no effects on general development, growth, sensory, cognitive function and reproductive performance were detected at 7 (males) and 19 (females) times the human therapeutic exposure at Day 21 *post partum*. At adult age (Day 69 *post partum*), no effects of bosentan were detected at 1.3 (males) and 2.6 (females) times the therapeutic exposure in children with PAH.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core:

Maize starch  
Pregelatinised starch  
Sodium starch glycollate  
Povidone  
Glycerol dibehenate  
Magnesium stearate

#### Film coat:

Hypromellose  
Glycerol triacetate  
Talc  
Titanium dioxide (E171)  
Iron oxide yellow (E172)  
Iron oxide red (E172)  
Ethylcellulose

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

4 years

For white high-density polyethylene bottles, use within 30 days after the first opening.

#### **6.4 Special precautions for storage**

For PVC/PE/PVDC/aluminium-blisters:  
Do not store above 30 °C.

For white high-density polyethylene bottles:  
This medicinal product does not require any special storage conditions.  
For storage conditions after first opening of the medicinal product, see section 6.3.

#### **6.5 Nature and contents of container**

##### Tracleer 62.5 mg film-coated tablets

PVC/PE/PVDC/aluminium-blisters containing 14 film-coated tablets.  
Cartons contain 14, 56 or 112 film-coated tablets.

White high-density polyethylene bottles with a silica gel desiccant containing 56 film-coated tablets.  
Cartons contain 56 film-coated tablets.

##### Tracleer 125 mg film-coated tablets

PVC/PE/PVDC/aluminium-blisters containing 14 film-coated tablets.  
Cartons contain 56 or 112 film-coated tablets.

White high-density polyethylene bottles with a silica gel desiccant containing 56 film-coated tablets.  
Cartons contain 56 film-coated tablets.

Not all pack sizes may be marketed.

#### **6.6 Special precautions for disposal and other handling**

No special requirements for disposal.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

### **7. MARKETING AUTHORISATION HOLDER**

Actelion Registration Ltd  
Chiswick Tower, 13<sup>th</sup> Floor  
389 Chiswick High Road  
London W4 4AL  
United Kingdom

### **8. MARKETING AUTHORISATION NUMBERS**

##### Tracleer 62.5 mg film-coated tablets

EU/1/02/220/001  
EU/1/02/220/002  
EU/1/02/220/003  
EU/1/02/220/007

Tracleer 125 mg film-coated tablets

EU/1/02/220/004

EU/1/02/220/005

EU/1/02/220/008

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 15 May 2002

Date of renewal: 20 April 2012

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>.



## **1. NAME OF THE MEDICINAL PRODUCT**

Tracleer 32 mg dispersible tablets

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each dispersible tablet contains 32 mg bosentan (as monohydrate).  
Excipient: 3.7 mg of Aspartame (E951) are present in each dispersible tablet.

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Dispersible tablet:

Pale yellow to off-white, clover-shape tablets, quadrisected on one side and debossed with “32” on the other side. The dispersible tablet can be divided into four equal parts.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Treatment of pulmonary arterial hypertension (PAH) to improve exercise capacity and symptoms in patients with WHO functional class III. Efficacy has been shown in:

- Primary (idiopathic and heritable) pulmonary arterial hypertension
- Pulmonary arterial hypertension secondary to scleroderma without significant interstitial pulmonary disease
- Pulmonary arterial hypertension associated with congenital systemic-to-pulmonary shunts and Eisenmenger’s physiology

Some improvements have also been shown in patients with pulmonary arterial hypertension WHO functional class II (see section 5.1).

Tracleer is also indicated to reduce the number of new digital ulcers in patients with systemic sclerosis and ongoing digital ulcer disease (see section 5.1).

### **4.2 Posology and method of administration**

#### Method of administration

Tablets are to be taken orally morning and evening, with or without food.

The dispersible tablets should be added to a little water on a spoon, and the liquid stirred to aid dissolution, before swallowing. A little more water should be added to the spoon and swallowed by the patient, to make sure all of the medicine has been administered. If possible, a glass of water should be taken to ensure that all the medicine has been ingested. If necessary the dispersible tablet can be divided by breaking it along the lines cut into the surface (see section 6.6).

The dispersible tablet has been studied only in paediatric patients. A bioavailability comparison between dispersible tablets and film-coated tablets performed in adult subjects indicated lower exposure to bosentan with the dispersible tablet (see section 5.2). Thus its use in adults should be reserved for patients who cannot take the film-coated tablet.

## Posology

### Pulmonary arterial hypertension

Treatment should only be initiated and monitored by a physician experienced in the treatment of PAH.

#### Adults

In adult patients, Tracleer treatment should be initiated at a dose of 62.5 mg twice daily for 4 weeks and then increased to the maintenance dose of 125 mg twice daily. The same recommendations apply to re-introduction of Tracleer after treatment interruption (see section 4.4).

#### Paediatric population

Paediatric pharmacokinetic data have shown that bosentan plasma concentrations in children with PAH aged from 1 year to 15 years were on average lower than in adult patients and were not increased by increasing the dose of Tracleer above 2 mg/kg body weight or by increasing the dosing frequency from twice daily to three times daily (see section 5.2). Increasing the dose or the dosing frequency will likely not result in additional clinical benefit.

Based on these pharmacokinetic results, when used in children with PAH aged 1 year and older, the recommended starting and maintenance dose is 2 mg/kg morning and evening.

In neonates with persistent pulmonary hypertension of the newborn (PPHN), the benefit of bosentan has not been shown in the standard-of-care treatment. No recommendation on a posology can be made (see sections 5.1 and 5.2).

#### *Management in the event of clinical deterioration of PAH*

In the event of clinical deterioration (e.g., decrease in 6-minute walk test distance by at least 10% compared with pre-treatment measurement) despite Tracleer treatment for at least 8 weeks (target dose for at least 4 weeks), alternative therapies should be considered. However, some patients who show no response after 8 weeks of treatment with Tracleer may respond favourably after an additional 4 to 8 weeks of treatment.

In the event of late clinical deterioration despite treatment with Tracleer (i.e., after several months of treatment), the treatment should be re-assessed. Some patients not responding well to 125 mg twice daily of Tracleer may slightly improve their exercise capacity when the dose is increased to 250 mg twice daily. A careful benefit/risk assessment should be made, taking into consideration that the liver toxicity is dose dependent (see sections 4.4 and 5.1).

#### *Discontinuation of treatment*

There is limited experience with abrupt discontinuation of Tracleer in patients with PAH. No evidence for acute rebound has been observed. However, to avoid the possible occurrence of harmful clinical deterioration due to potential rebound effect, gradual dose reduction (halving the dose for 3 to 7 days) should be considered. Intensified monitoring is recommended during the discontinuation period. If the decision to withdraw Tracleer is taken, it should be done gradually while an alternative therapy is introduced.

### Systemic sclerosis with ongoing digital ulcer disease

Treatment should only be initiated and monitored by a physician experienced in the treatment of systemic sclerosis.

#### Adults

Tracleer treatment should be initiated at a dose of 62.5 mg twice daily for 4 weeks and then increased to the maintenance dose of 125 mg twice daily. The same recommendations apply to re-introduction of Tracleer after treatment interruption (see section 4.4).

Controlled clinical study experience in this indication is limited to 6 months (see section 5.1).

The patient's response to treatment and need for continued therapy should be re-evaluated on a regular basis. A careful benefit/risk assessment should be made, taking into consideration the liver toxicity of bosentan (see sections 4.4 and 4.8).

#### Paediatric population

There are no data on the safety and efficacy in patients under the age of 18 years. Pharmacokinetic data are not available for Tracleer in young children with this disease.

### Special populations

#### *Hepatic impairment*

Tracleer is contraindicated in patients with moderate to severe liver dysfunction (see sections 4.3, 4.4 and 5.2). No dose adjustment is needed in patients with mild hepatic impairment (i.e., Child-Pugh class A) (see section 5.2).

#### *Renal impairment*

No dose adjustment is required in patients with renal impairment. No dose adjustment is required in patients undergoing dialysis (see section 5.2).

#### *Elderly*

No dose adjustment is required in patients over the age of 65 years.

### **4.3 Contraindications**

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1
- Moderate to severe hepatic impairment, i.e., Child-Pugh class B or C (see section 5.2)
- Baseline values of liver aminotransferases, i.e., aspartate aminotransferase (AST) and/or alanine aminotransferase (ALT), greater than 3 × the upper limit of normal (ULN; see section 4.4)
- Concomitant use of cyclosporine A (see section 4.5)
- Pregnancy (see sections 4.4 and 4.6)
- Women of childbearing potential who are not using reliable methods of contraception (see sections 4.4, 4.5 and 4.6)

### **4.4 Special warnings and precautions for use**

The efficacy of Tracleer has not been established in patients with severe PAH. Transfer to a therapy that is recommended at the severe stage of the disease (e.g., epoprostenol) should be considered if the clinical condition deteriorates (see section 4.2).

The benefit/risk balance of bosentan has not been established in patients with WHO class I functional status of PAH.

Tracleer should only be initiated if the systemic systolic blood pressure is higher than 85 mmHg. Tracleer has not been shown to have a beneficial effect on the healing of existing digital ulcers.

#### Liver function

Elevations in liver aminotransferases, i.e., aspartate and alanine aminotransferases (AST and/or ALT), associated with bosentan are dose dependent. Liver enzyme changes typically occur within the first 26 weeks of treatment but may also occur late in treatment (see section 4.8). These increases may be partly due to competitive inhibition of the elimination of bile salts from hepatocytes but other mechanisms, which have not been clearly established, are probably also involved in the occurrence of liver dysfunction. The accumulation of bosentan in hepatocytes leading to cytolysis with potentially severe damage of the liver, or an immunological mechanism, are not excluded. Liver dysfunction risk may also be increased when medicinal products that are inhibitors of the bile salt export pump, e.g., rifampicin, glibenclamide and cyclosporine A (see sections 4.3 and 4.5), are co-administered with bosentan, but limited data are available.

**Liver aminotransferase levels must be measured prior to initiation of treatment and subsequently at monthly intervals for the duration of treatment with Tracleer. In addition, liver aminotransferase levels must be measured 2 weeks after any dose increase.**

#### Recommendations in the event of ALT/AST elevations

##### **ALT/AST levels**

##### **Treatment and monitoring recommendations**

> 3 and  $\leq 5 \times$  ULN

The result should be confirmed by a second liver test; if confirmed, a decision should be made on an individual basis to continue Tracleer, possibly at a reduced dose, or to stop Tracleer administration (see section 4.2). Monitoring of aminotransferase levels should be continued at least every 2 weeks. If the aminotransferase levels return to pre-treatment values continuing or re-introducing Tracleer according to the conditions described below should be considered.

> 5 and  $\leq 8 \times$  ULN

The result should be confirmed by a second liver test; if confirmed, treatment should be stopped and aminotransferase levels monitored at least every 2 weeks. If the aminotransferase levels return to pre-treatment values re-introducing Tracleer according to the conditions described below should be considered.

> 8  $\times$  ULN

Treatment must be stopped and re-introduction of Tracleer is not to be considered.

**In the event of associated clinical symptoms of liver injury, i.e., nausea, vomiting, fever, abdominal pain, jaundice, unusual lethargy or fatigue, flu-like syndrome (arthralgia, myalgia, fever), treatment must be stopped and re-introduction of Tracleer is not to be considered.**

#### Re-introduction of treatment

Re-introduction of treatment with Tracleer should only be considered if the potential benefits of treatment with Tracleer outweigh the potential risks and when liver aminotransferase levels are within pre-treatment values. The advice of a hepatologist is recommended. Re-introduction must follow the guidelines detailed in section 4.2. **Aminotransferase levels must then be checked within 3 days after re-introduction, then again after a further 2 weeks, and thereafter according to the recommendations above.**

ULN = upper limit of normal

### Haemoglobin concentration

Treatment with bosentan has been associated with dose-related decreases in haemoglobin concentration (see section 4.8). In placebo-controlled studies, bosentan-related decreases in haemoglobin concentration were not progressive, and stabilised after the first 4–12 weeks of treatment. It is recommended that haemoglobin concentrations be checked prior to initiation of treatment, every month during the first 4 months, and quarterly thereafter. If a clinically relevant decrease in haemoglobin concentration occurs, further evaluation and investigation should be undertaken to determine the cause and need for specific treatment. In the post-marketing period, cases of anaemia requiring red blood cell transfusion have been reported (see section 4.8).

### Women of childbearing potential

As Tracleer may render hormonal contraceptives ineffective, and taking into account the risk that pulmonary hypertension deteriorates with pregnancy as well as the teratogenic effects observed in animals:

- Tracleer treatment must not be initiated in women of childbearing potential unless they practise reliable contraception and the result of the pre-treatment pregnancy test is negative
- Hormonal contraceptives cannot be the sole method of contraception during treatment with Tracleer
- Monthly pregnancy tests are recommended during treatment to allow early detection of pregnancy

For further information see sections 4.5 and 4.6.

### Pulmonary veno-occlusive disease

Cases of pulmonary oedema have been reported with vasodilators (mainly prostacyclins) when used in patients with pulmonary veno-occlusive disease. Consequently, should signs of pulmonary oedema occur when Tracleer is administered in patients with PAH, the possibility of associated veno-occlusive disease should be considered. In the post-marketing period there have been rare reports of pulmonary oedema in patients treated with Tracleer who had a suspected diagnosis of pulmonary veno-occlusive disease.

### Pulmonary arterial hypertension patients with concomitant left ventricular failure

No specific study has been performed in patients with pulmonary hypertension and concomitant left ventricular dysfunction. However, 1,611 patients (804 bosentan- and 807 placebo-treated patients) with severe chronic heart failure (CHF) were treated for a mean duration of 1.5 years in a placebo-controlled study (study AC-052-301/302 [ENABLE 1 & 2]). In this study there was an increased incidence of hospitalisation due to CHF during the first 4–8 weeks of treatment with bosentan, which could have been the result of fluid retention. In this study, fluid retention was manifested by early weight gain, decreased haemoglobin concentration and increased incidence of leg oedema. At the end of this study, there was no difference in overall hospitalisations for heart failure nor in mortality between bosentan- and placebo-treated patients. Consequently, it is recommended that patients be monitored for signs of fluid retention (e.g., weight gain), especially if they concomitantly suffer from severe systolic dysfunction. Should this occur, starting treatment with diuretics is recommended, or the dose of existing diuretics should be increased. Treatment with diuretics should be considered in patients with evidence of fluid retention before the start of treatment with Tracleer.

### Pulmonary arterial hypertension associated with HIV infection

There is limited clinical study experience with the use of Tracleer in patients with PAH associated with HIV infection, treated with antiretroviral medicinal products (see section 5.1). An interaction study between bosentan and lopinavir+ritonavir in healthy subjects showed increased plasma concentrations of bosentan, with the maximum level during the first 4 days of treatment (see section 4.5). When treatment with Tracleer is initiated in patients who require ritonavir-boosted protease inhibitors, the patient's tolerability of Tracleer should be closely monitored with special attention, at the beginning of the initiation phase, to the risk of hypotension and to liver function tests. An increased long-term risk of hepatic toxicity and haematological adverse events cannot be excluded when bosentan is used in combination with antiretroviral medicinal products. Due to the potential for interactions related to the inducing effect of bosentan on CYP450 (see section 4.5), which could affect the efficacy of antiretroviral therapy, these patients should also be monitored carefully regarding their HIV infection.

### Pulmonary hypertension secondary to chronic obstructive pulmonary disease (COPD)

Safety and tolerability of bosentan was investigated in an exploratory, uncontrolled 12-week study in 11 patients with pulmonary hypertension secondary to severe COPD (stage III of GOLD classification). An increase in minute ventilation and a decrease in oxygen saturation were observed, and the most frequent adverse event was dyspnoea, which resolved with discontinuation of bosentan.

### Concomitant use with other medicinal products

Concomitant use of Tracleer and cyclosporine A is contraindicated (see sections 4.3 and 4.5).

Concomitant use of Tracleer with glibenclamide, fluconazole and rifampicin is not recommended. For further details please refer to section 4.5.

Concomitant administration of both a CYP3A4 inhibitor and a CYP2C9 inhibitor with Tracleer should be avoided (see section 4.5).

Tracleer 32 mg dispersible tablets contain a source of phenylalanine (Aspartame – E951). This may be harmful for people with phenylketonuria.

## **4.5 Interaction with other medicinal products and other forms of interaction**

Bosentan is an inducer of the cytochrome P450 (CYP) isoenzymes CYP2C9 and CYP3A4. *In vitro* data also suggest an induction of CYP2C19. Consequently, plasma concentrations of substances metabolised by these isoenzymes will be decreased when Tracleer is co-administered. The possibility of altered efficacy of medicinal products metabolised by these isoenzymes should be considered. The dosage of these products may need to be adjusted after initiation, dose change or discontinuation of concomitant Tracleer treatment.

Bosentan is metabolised by CYP2C9 and CYP3A4. Inhibition of these isoenzymes may increase the plasma concentration of bosentan (see ketoconazole). The influence of CYP2C9 inhibitors on bosentan concentration has not been studied. The combination should be used with caution.

*Fluconazole and other inhibitors of both CYP2C9 and CYP3A4:* Co-administration with fluconazole, which inhibits mainly CYP2C9, but to some extent also CYP3A4, could lead to large increases in plasma concentrations of bosentan. The combination is not recommended. For the same reason, concomitant administration of both a potent CYP3A4 inhibitor (such as ketoconazole, itraconazole or ritonavir) and a CYP2C9 inhibitor (such as voriconazole) with Tracleer is not recommended.

*Cyclosporine A:* Co-administration of Tracleer and cyclosporine A (a calcineurin inhibitor) is contraindicated (see section 4.3). When co-administered, initial trough concentrations of bosentan were approximately 30-fold higher than those measured after bosentan alone. At steady state, bosentan plasma concentrations were 3- to 4-fold higher than with bosentan alone. The mechanism of this interaction is most likely inhibition of transport protein-mediated uptake of bosentan into hepatocytes by cyclosporine. The blood concentrations of cyclosporine A (a CYP3A4 substrate) decreased by approximately 50%. This is most likely due to induction of CYP3A4 by bosentan.

*Tacrolimus, sirolimus:* Co-administration of tacrolimus or sirolimus and Tracleer has not been studied in man but co-administration of tacrolimus or sirolimus and Tracleer may result in increased plasma concentrations of bosentan in analogy to co-administration with cyclosporine A. Concomitant Tracleer may reduce the plasma concentrations of tacrolimus and sirolimus. Therefore, concomitant use of Tracleer and tacrolimus or sirolimus is not advisable. Patients in need of the combination should be closely monitored for adverse events related to Tracleer and for tacrolimus and sirolimus blood concentrations.

*Glibenclamide:* Co-administration of bosentan 125 mg twice daily for 5 days decreased the plasma concentrations of glibenclamide (a CYP3A4 substrate) by 40%, with potential significant decrease of the hypoglycaemic effect. The plasma concentrations of bosentan were also decreased by 29%. In addition, an increased incidence of elevated aminotransferases was observed in patients receiving concomitant therapy. Both glibenclamide and bosentan inhibit the bile salt export pump, which could explain the elevated aminotransferases. This combination should not be used. No drug-drug interaction data are available with the other sulfonylureas.

*Rifampicin:* Co-administration in 9 healthy subjects for 7 days of bosentan 125 mg twice daily with rifampicin, a potent inducer of CYP2C9 and CYP3A4, decreased the plasma concentrations of bosentan by 58%, and this decrease could achieve almost 90% in an individual case. As a result, a significantly reduced effect of bosentan is expected when it is co-administered with rifampicin. Concomitant use of rifampicin and Tracleer is not recommended. Data on other CYP3A4 inducers, e.g., carbamazepine, phenobarbital, phenytoin and St. John's wort are lacking, but their concomitant administration is expected to lead to reduced systemic exposure to bosentan. A clinically significant reduction of efficacy cannot be excluded.

*Lopinavir+ritonavir (and other ritonavir-boosted protease inhibitors):* Co-administration of bosentan 125 mg twice daily and lopinavir+ritonavir 400+100 mg twice daily for 9.5 days in healthy volunteers resulted in initial trough plasma concentrations of bosentan that were approximately 48-fold higher than those measured after bosentan administered alone. On day 9, plasma concentrations of bosentan were approximately 5-fold higher than with bosentan administered alone. Inhibition by ritonavir of transport protein-mediated uptake into hepatocytes and of CYP3A4, thereby reducing the clearance of bosentan, most likely causes this interaction. When administered concomitantly with lopinavir+ritonavir, or other ritonavir-boosted protease inhibitors, the patient's tolerability of Tracleer should be monitored.

After co-administration of bosentan for 9.5 days, the plasma exposures to lopinavir and ritonavir decreased to a clinically non-significant extent (by approximately 14% and 17%, respectively). However, full induction by bosentan might not have been reached and a further decrease of protease inhibitors cannot be excluded. Appropriate monitoring of the HIV therapy is recommended. Similar effects would be expected with other ritonavir-boosted protease inhibitors (see section 4.4).

*Other antiretroviral agents:* No specific recommendation can be made with regard to other available antiretroviral agents due to the lack of data. Due to the marked hepatotoxicity of nevirapine, which could add to bosentan liver toxicity, this combination is not recommended.

*Hormonal contraceptives:* Co-administration of bosentan 125 mg twice daily for 7 days with a single dose of oral contraceptive containing norethisterone 1 mg + ethinyl estradiol 35 mcg decreased the AUC of norethisterone and ethinyl estradiol by 14% and 31%, respectively. However, decreases in exposure were as much as 56% and 66%, respectively, in individual subjects. Therefore, hormone-based contraceptives alone, regardless of the route of administration (i.e., oral, injectable, transdermal or implantable forms), are not considered as reliable methods of contraception (see sections 4.4 and 4.6).

*Warfarin:* Co-administration of bosentan 500 mg twice daily for 6 days decreased the plasma concentrations of both S-warfarin (a CYP2C9 substrate) and R-warfarin (a CYP3A4 substrate) by 29% and 38%, respectively. Clinical experience with concomitant administration of bosentan with warfarin in patients with PAH did not result in clinically relevant changes in International Normalized Ratio (INR) or warfarin dose (baseline versus end of the clinical studies). In addition, the frequency of changes in warfarin dose during the studies due to changes in INR or due to adverse events was similar among bosentan- and placebo-treated patients. No dose adjustment is needed for warfarin and similar oral anticoagulant agents when bosentan is initiated, but intensified monitoring of INR is recommended, especially during bosentan initiation and the up-titration period.

*Simvastatin:* Co-administration of bosentan 125 mg twice daily for 5 days decreased the plasma concentrations of simvastatin (a CYP3A4 substrate) and its active  $\beta$ -hydroxy acid metabolite by 34% and 46%, respectively. The plasma concentrations of bosentan were not affected by concomitant simvastatin. Monitoring of cholesterol levels and subsequent dosage adjustment should be considered.

*Ketoconazole:* Co-administration for 6 days of bosentan 62.5 mg twice daily with ketoconazole, a potent CYP3A4 inhibitor, increased the plasma concentrations of bosentan approximately 2-fold. No dose adjustment of Tracleer is considered necessary. Although not demonstrated through *in vivo* studies, similar increases in bosentan plasma concentrations are expected with the other potent CYP3A4 inhibitors (such as itraconazole or ritonavir). However, when combined with a CYP3A4 inhibitor, patients who are poor metabolisers of CYP2C9 are at risk of increases in bosentan plasma concentrations that may be of higher magnitude, thus leading to potential harmful adverse events.

*Epoprostenol:* Limited data obtained from a study (AC-052-356 [BREATHE-3]) in which 10 paediatric patients received the combination of bosentan and epoprostenol indicate that after both single- and multiple-dose administration, the  $C_{max}$  and AUC values of bosentan were similar in patients with or without continuous infusion of epoprostenol (see section 5.1).

*Sildenafil:* Co-administration of bosentan 125 mg twice daily (steady state) with sildenafil 80 mg three times a day (at steady state) concomitantly administered during 6 days in healthy volunteers resulted in a 63% decrease in the sildenafil AUC and a 50% increase in the bosentan AUC. Caution is recommended in the case of co-administration.

*Digoxin:* Co-administration for 7 days of bosentan 500 mg twice daily with digoxin decreased the AUC,  $C_{max}$  and  $C_{min}$  of digoxin by 12%, 9% and 23%, respectively. The mechanism for this interaction may be induction of P-glycoprotein. This interaction is unlikely to be of clinical relevance.

### Paediatric population

Interaction studies have only been performed in adults.



## **4.6 Fertility, pregnancy and lactation**

### Pregnancy

Studies in animals have shown reproductive toxicity (teratogenicity, embryotoxicity; see section 5.3). There are no reliable data on the use of Tracleer in pregnant women. The potential risk for humans is still unknown. Tracleer is contraindicated in pregnancy (see section 4.3).

### Women of childbearing potential

Before the initiation of Tracleer treatment in women of childbearing potential, the absence of pregnancy should be checked, appropriate advice on reliable methods of contraception provided, and reliable contraception initiated. Patients and prescribers must be aware that due to potential pharmacokinetic interactions, Tracleer may render hormonal contraceptives ineffective (see section 4.5). Therefore, women of childbearing potential must not use hormonal contraceptives (including oral, injectable, transdermal or implantable forms) as the sole method of contraception but must use an additional or an alternative reliable method of contraception. If there is any doubt about what contraceptive advice should be given to the individual patient, consultation with a gynaecologist is recommended. Because of possible hormonal contraception failure during Tracleer treatment, and also bearing in mind the risk that pulmonary hypertension severely deteriorates with pregnancy, monthly pregnancy tests during treatment with Tracleer are recommended to allow early detection of pregnancy.

### Breast-feeding

It is not known whether bosentan is excreted into human breast milk. Breast-feeding is not recommended during treatment with Tracleer.

### Fertility

Animal studies showed testicular effects (see section 5.3). In a study investigating the effects of bosentan on testicular function in male PAH patients, 8 out of 24 patients showed a decreased sperm concentration from baseline of at least 42% after 3 or 6 months of treatment with bosentan. Based on these findings and preclinical data, it cannot be excluded that bosentan may have a detrimental effect on spermatogenesis in men. In male children, a long-term impact on fertility after treatment with bosentan cannot be excluded.

## **4.7 Effects on ability to drive and use machines**

No specific studies have been conducted to assess the direct effect of Tracleer on the ability to drive and use machines. However, Tracleer may induce hypotension, with symptoms of dizziness, blurred vision or syncope that could affect the ability to drive or use machines.

## **4.8 Undesirable effects**

In 20 placebo-controlled studies, conducted in a variety of therapeutic indications, a total of 2,486 patients were treated with bosentan at daily doses ranging from 100 mg to 2000 mg and 1,838 patients were treated with placebo. The mean treatment duration was 45 weeks. Adverse reactions were defined as events occurring in at least 1% of patients on bosentan and at a frequency at least 0.5% more than on placebo. The most frequent adverse reactions are headache (11.5%), oedema / fluid retention (13.2%), abnormal liver function test (10.9%) and anaemia / haemoglobin decrease (9.9%).

Treatment with bosentan has been associated with dose-dependent elevations in liver aminotransferases and decreases in haemoglobin concentration (see section 4.4).

Adverse reactions observed in 20 placebo-controlled studies and post-marketing experience with bosentan are ranked according to frequency using the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data).

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness. No clinically relevant differences in adverse reactions were observed between the overall dataset and the approved indications.

<b>System organ class</b>	<b>Frequency</b>	<b>Adverse reaction</b>
Blood and lymphatic system disorders	Common	Anaemia, haemoglobin decrease, (see section 4.4)
	Not known	Anaemia or haemoglobin decreases requiring red blood cell transfusion <sup>1</sup>
	Uncommon	Thrombocytopenia <sup>1</sup>
	Uncommon	Neutropenia, leukopenia <sup>1</sup>
Immune system disorders	Common	Hypersensitivity reactions (including dermatitis, pruritus and rash) <sup>2</sup>
	Rare	Anaphylaxis and/or angioedema <sup>1</sup>
Nervous system disorders	Very common	Headache <sup>3</sup>
	Common	Syncope <sup>1,4</sup>
Eye disorders	Not known	Blurred vision <sup>1</sup>
Cardiac disorders	Common	Palpitations <sup>1,4</sup>
Vascular disorders	Common	Flushing
	Common	Hypotension <sup>1,4</sup>
Respiratory, thoracic and mediastinal disorders	Common	Nasal congestion <sup>1</sup>
Gastrointestinal disorders	Common	Gastrooesophageal reflux disease Diarrhoea
Hepatobiliary disorders	Very common	Abnormal liver function test (see section 4.4)
	Uncommon	Aminotransferase elevations associated with hepatitis (including possible exacerbation of underlying hepatitis) and/or jaundice <sup>1</sup> (see section 4.4)
	Rare	Liver cirrhosis, liver failure <sup>1</sup>
Skin and subcutaneous disorders	Common	Erythema
General disorders and administration site conditions	Very common	Oedema, fluid retention <sup>5</sup>

<sup>1</sup> Data derived from post-marketing experience, frequencies based on statistical modelling of placebo-controlled clinical trial data.

<sup>2</sup> Hypersensitivity reactions were reported in 9.9% of patients on bosentan and 9.1% of patients on placebo.

<sup>3</sup> Headache was reported in 11.5% of patients on bosentan and 9.8% of patients on placebo.

<sup>4</sup> These types of reactions can also be related to the underlying disease.

<sup>5</sup> Oedema or fluid retention was reported in 13.2% of patients on bosentan and 10.9% of patients on placebo.

In the post-marketing period rare cases of unexplained hepatic cirrhosis were reported after prolonged therapy with Tracleer in patients with multiple co-morbidities and therapies with medicinal products. There have also been rare reports of liver failure. These cases reinforce the importance of strict adherence to the monthly schedule for monitoring of liver function for the duration of treatment with Tracleer (see section 4.4).

### Paediatric population

#### *Uncontrolled clinical studies in paediatric patients*

The safety profile in the first paediatric uncontrolled study performed with the film-coated tablet (BREATHE-3: n = 19, median age 10 years [range 3–15 years], open-label bosentan 2 mg/kg twice daily; treatment duration 12 weeks) was similar to that observed in the pivotal trials in adult patients with PAH. In BREATHE-3, the most frequent adverse reactions were flushing (21%), headache, and abnormal liver function test (each 16%).

A pooled analysis of uncontrolled paediatric studies conducted in PAH with the bosentan 32 mg dispersible tablet formulation (FUTURE 1/2, FUTURE 3/Extension) included a total of 100 children treated with bosentan 2 mg/kg twice daily (n = 33), 2 mg/kg three times daily (n = 31), or 4 mg/kg twice daily (n = 36). At enrolment, six patients were between 3 months and 1 year old, 15 children were between 1 and less than 2 years old, and 79 were between 2 and 12 years old. The median treatment duration was 71.8 weeks (range 0.4–258 weeks).

The safety profile in this pooled analysis of uncontrolled paediatric studies was similar to that observed in the pivotal trials in adult patients with PAH except for infections, which were more frequently reported than in adults (69.0% vs 41.3%). This difference in infection frequency may in part be due to the longer median treatment exposure in the paediatric set (median 71.8 weeks) compared with the adult set (median 17.4 weeks). The most frequent adverse events were upper respiratory tract infections (25%), pulmonary (arterial) hypertension (20%), nasopharyngitis (17%), pyrexia (15%), vomiting (13%), bronchitis (10%), abdominal pain (10%), and diarrhoea (10%). There was no relevant difference in adverse event frequencies between patients above and below the age of 2 years; however, this is based on only 21 children less than 2 years including 6 patients between 3 months to 1 year of age. Adverse events of liver abnormalities and anaemia/haemoglobin decrease occurred in 9% and 5% of patients, respectively.

In a randomised placebo-controlled study, conducted in PPHN patients (FUTURE-4), a total of 13 neonates were treated with the bosentan dispersible tablet formulation at a dose of 2 mg/kg twice daily (8 patients were on placebo). The median bosentan and placebo treatment duration was, respectively, 4.5 days (range 0.5–10.0 days) and 4.0 days (range 2.5–6.5 days). The most frequent adverse events in the bosentan- and placebo-treated patients were, respectively, anaemia or haemoglobin decrease (7 and 2 patients), generalised oedema (3 and 0 patients), and vomiting (2 and 0 patients).

### Laboratory abnormalities

#### *Liver test abnormalities*

In the clinical programme, dose-dependent elevations in liver aminotransferases generally occurred within the first 26 weeks of treatment, usually developed gradually, and were mainly asymptomatic. In the post-marketing period rare cases of liver cirrhosis and liver failure have been reported.

The mechanism of this adverse effect is unclear. These elevations in aminotransferases may reverse spontaneously while continuing treatment with the maintenance dose of Tracleer or after dose reduction, but interruption or cessation may be necessary (see section 4.4).

In the 20 integrated placebo-controlled studies, elevations in liver aminotransferases  $\geq 3 \times \text{ULN}$  were observed in 11.2% of the bosentan-treated patients as compared to 2.4% of the placebo-treated patients. Elevations to  $\geq 8 \times \text{ULN}$  were seen in 3.6% of the bosentan-treated patients and 0.4% of the placebo-treated patients. Elevations in aminotransferases were associated with elevated bilirubin ( $\geq 2 \times \text{ULN}$ ) without evidence of biliary obstruction in 0.2% (5 patients) on bosentan and 0.3% (6 patients) on placebo.

In the pooled analysis of 100 PAH children from uncontrolled paediatric studies FUTURE 1/2 and FUTURE 3/Extension, elevations in liver aminotransferases  $\geq 3 \times \text{ULN}$  were observed in 2% of patients.

In the FUTURE-4 study including 13 neonates with PPHN treated with bosentan 2 mg/kg twice daily for less than 10 days (range 0.5–10.0 days), there were no cases of liver aminotransferases  $\geq 3 \times \text{ULN}$  during treatment, but one case of hepatitis occurred 3 days after the end of bosentan treatment.

### *Haemoglobin*

In the adult placebo-controlled studies, a decrease in haemoglobin concentration to below 10 g/dL from baseline was reported in 8.0% of bosentan-treated patients and 3.9% of placebo-treated patients (see section 4.4).

In the pooled analysis of 100 PAH children from uncontrolled paediatric studies FUTURE 1/2 and FUTURE 3/Extension, a decrease in haemoglobin concentration from baseline to below 10 g/dL was reported in 10.0% of patients. There was no decrease to below 8 g/dL.

In the FUTURE-4 study, 6 out of 13 bosentan-treated neonates with PPHN experienced a decrease in haemoglobin from within the reference range at baseline to below the lower limit of normal during the treatment.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via **the national reporting system** listed in [Appendix V](#).

## **4.9 Overdose**

Bosentan has been administered as a single dose of up to 2,400 mg to healthy subjects and up to 2,000 mg/day for 2 months in patients with a disease other than pulmonary hypertension. The most common adverse reaction was headache of mild to moderate intensity.

Massive overdose may result in pronounced hypotension requiring active cardiovascular support. In the post-marketing period there was one reported overdose of 10,000 mg of Tracleer taken by an adolescent male patient. He had symptoms of nausea, vomiting, hypotension, dizziness, sweating and blurred vision. He recovered completely within 24 hours with blood pressure support. Note: bosentan is not removed through dialysis.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: other antihypertensives, ATC code: C02KX01

#### Mechanism of action

Bosentan is a dual endothelin receptor antagonist (ERA) with affinity for both endothelin A and B (ET<sub>A</sub> and ET<sub>B</sub>) receptors. Bosentan decreases both pulmonary and systemic vascular resistance resulting in increased cardiac output without increasing heart rate.

The neurohormone endothelin-1 (ET-1) is one of the most potent vasoconstrictors known and can also promote fibrosis, cell proliferation, cardiac hypertrophy and remodelling, and is pro-inflammatory. These effects are mediated by endothelin binding to ET<sub>A</sub> and ET<sub>B</sub> receptors located in the endothelium and vascular smooth muscle cells. ET-1 concentrations in tissues and plasma are increased in several cardiovascular disorders and connective tissue diseases, including PAH, scleroderma, acute and chronic heart failure, myocardial ischaemia, systemic hypertension and atherosclerosis, suggesting a pathogenic role of ET-1 in these diseases. In PAH and heart failure, in the absence of endothelin receptor antagonism, elevated ET-1 concentrations are strongly correlated with the severity and prognosis of these diseases.

Bosentan competes with the binding of ET-1 and other ET peptides to both ET<sub>A</sub> and ET<sub>B</sub> receptors, with a slightly higher affinity for ET<sub>A</sub> receptors ( $K_i = 4.1\text{--}43$  nanomolar) than for ET<sub>B</sub> receptors ( $K_i = 38\text{--}730$  nanomolar). Bosentan specifically antagonises ET receptors and does not bind to other receptors.

#### Efficacy

##### *Animal models*

In animal models of pulmonary hypertension, chronic oral administration of bosentan reduced pulmonary vascular resistance and reversed pulmonary vascular and right ventricular hypertrophy. In an animal model of pulmonary fibrosis, bosentan reduced collagen deposition in the lungs.

##### *Efficacy in adult patients with pulmonary arterial hypertension*

Two randomised, double-blind, multi-centre, placebo-controlled studies have been conducted in 32 (study AC-052-351) and 213 (study AC-052-352 [BREATHE-1]) adult patients with WHO functional class III–IV PAH (primary pulmonary hypertension or pulmonary hypertension secondary mainly to scleroderma). After 4 weeks of bosentan 62.5 mg twice daily, the maintenance doses studied in these studies were 125 mg twice daily in AC-052-351, and 125 mg twice daily and 250 mg twice daily in AC-052-352.

Bosentan was added to patients' current therapy, which could include a combination of anticoagulants, vasodilators (e.g., calcium channel blockers), diuretics, oxygen and digoxin, but not epoprostenol. Control was placebo plus current therapy.

The primary endpoint for each study was change in 6-minute walk distance at 12 weeks for the first study and 16 weeks for the second study. In both studies, treatment with bosentan resulted in significant increases in exercise capacity. The placebo-corrected increases in walk distance compared with baseline were 76 metres ( $p = 0.02$ ; t-test) and 44 metres ( $p = 0.0002$ ; Mann-Whitney U test) at the primary endpoint of each study, respectively. The differences between the two groups, 125 mg twice

daily and 250 mg twice daily, were not statistically significant but there was a trend towards improved exercise capacity in the group treated with 250 mg twice daily.

The improvement in walk distance was apparent after 4 weeks of treatment, was clearly evident after 8 weeks of treatment and was maintained for up to 28 weeks of double-blind treatment in a subset of the patient population.

In a retrospective responder analysis based on change in walking distance, WHO functional class and dyspnoea of the 95 patients randomised to bosentan 125 mg twice daily in the placebo-controlled studies, it was found that at week 8, 66 patients had improved, 22 were stable and 7 had deteriorated. Of the 22 patients stable at week 8, 6 improved at week 12/16 and 4 deteriorated compared with baseline. Of the 7 patients who deteriorated at week 8, 3 improved at week 12/16 and 4 deteriorated compared with baseline.

Invasive haemodynamic parameters were assessed in the first study only. Treatment with bosentan led to a significant increase in cardiac index associated with a significant reduction in pulmonary artery pressure, pulmonary vascular resistance and mean right atrial pressure.

A reduction in symptoms of PAH was observed with bosentan treatment. Dyspnoea measurement during walk tests showed an improvement in bosentan-treated patients. In the AC-052-352 study, 92% of the 213 patients were classified at baseline as WHO functional class III and 8% as class IV. Treatment with bosentan led to a WHO functional class improvement in 42.4% of patients (placebo 30.4%). The overall change in WHO functional class during both studies was significantly better among bosentan-treated patients as compared with placebo-treated patients. Treatment with bosentan was associated with a significant reduction in the rate of clinical worsening compared with placebo at 28 weeks (10.7% vs 37.1%, respectively;  $p = 0.0015$ ).

In a randomised, double-blind, multi-centre, placebo-controlled study (AC-052-364 [EARLY]), 185 PAH patients in WHO functional class II (mean baseline 6-minute walk distance of 435 metres) received bosentan 62.5 mg twice daily for 4 weeks followed by 125 mg twice daily ( $n = 93$ ), or placebo ( $n = 92$ ) for 6 months. Enrolled patients were PAH-treatment-naïve ( $n = 156$ ) or on a stable dose of sildenafil ( $n = 29$ ). The co-primary endpoints were percentage change from baseline in pulmonary vascular resistance (PVR) and change from baseline in 6-minute walk distance to Month 6 versus placebo. The table below illustrates the pre-specified protocol analyses.

	PVR (dyn.sec/cm <sup>5</sup> )		6-Minute Walk Distance (m)	
	Placebo (n=88)	Bosentan (n=80)	Placebo (n=91)	Bosentan (n=86)
Baseline (BL); mean (SD)	802 (365)	851 (535)	431 (92)	443 (83)
Change from BL; mean (SD)	128 (465)	-69 (475)	-8 (79)	11 (74)
Treatment effects	-22.6%		19	
95% CL	-34, -10		-4, 42	
P-value	< 0.0001		0.0758	

CL = confidence limit; PVR = pulmonary vascular resistance; SD = standard deviation.

Treatment with bosentan was associated with a reduction in the rate of clinical worsening, defined as a composite of symptomatic progression, hospitalisation for PAH and death, compared with placebo (proportional risk reduction 77%, 95% confidence interval [CI] 20–94%,  $p = 0.0114$ ). The treatment effect was driven by improvement in the component symptomatic progression. There was one hospitalisation related to PAH worsening in the bosentan group and three hospitalisations in the placebo group. Only one death occurred in each treatment group during the 6-month double-blind study period, therefore no conclusion can be drawn on survival.

Long-term data were generated from all 173 patients who were treated with bosentan in the controlled phase and/or were switched from placebo to bosentan in the open-label extension phase of the EARLY study. The mean duration of exposure to bosentan treatment was  $3.6 \pm 1.8$  years (up to 6.1 years), with 73% of patients treated for at least 3 years and 62% for at least 4 years. Patients could receive additional PAH treatment as required in the open-label extension. The majority of patients were diagnosed with idiopathic or heritable PAH (61%). Overall, 78% of patients remained in WHO functional class II. Kaplan-Meier estimates of survival were 90% and 85% at 3 and 4 years after the start of treatment, respectively. At the same timepoints, 88% and 79% of patients remained free from PAH worsening (defined as all-cause death, lung transplantation, atrial septostomy or start of intravenous or subcutaneous prostanoid treatment). The relative contributions of previous placebo treatment in the double-blind phase and of other medications started during the open-label extension period are unknown.

In a prospective, multi-centre, randomised, double-blind, placebo-controlled study (AC-052-405 [BREATHE-5]), patients with PAH WHO functional class III and Eisenmenger physiology associated with congenital heart disease received bosentan 62.5 mg twice daily for 4 weeks, then 125 mg twice daily for a further 12 weeks ( $n = 37$ , of whom 31 had a predominantly right to left, bidirectional shunt). The primary objective was to show that bosentan did not worsen hypoxaemia. After 16 weeks, the mean oxygen saturation was increased in the bosentan group by 1.0% (95% CI  $-0.7\%$ – $2.8\%$ ) as compared to the placebo group ( $n = 17$ ), showing that bosentan did not worsen hypoxaemia. The mean pulmonary vascular resistance was significantly reduced in the bosentan group (with a predominant effect observed in the subgroup of patients with bidirectional intracardiac shunt). After 16 weeks, the mean placebo-corrected increase in 6-minute walk distance was 53 metres ( $p = 0.0079$ ), reflecting improvement in exercise capacity. Twenty-six patients continued to receive bosentan in the 24-week open-label extension phase (AC-052-409) of the BREATHE-5 study (mean duration of treatment =  $24.4 \pm 2.0$  weeks) and, in general, efficacy was maintained.

An open-label, non-comparative study (AC-052-362 [BREATHE-4]) was performed in 16 patients with WHO functional class III PAH associated with HIV infection. Patients were treated with bosentan 62.5 mg twice daily for 4 weeks followed by 125 mg twice daily for a further 12 weeks. After 16 weeks' treatment, there were significant improvements from baseline in exercise capacity: the mean increase in 6-minute walk distance was 91.4 metres from 332.6 metres on average at baseline ( $p < 0.001$ ). No formal conclusion can be drawn regarding the effects of bosentan on antiretroviral drug efficacy (see also section 4.4).

There are no studies to demonstrate beneficial effects of Tracleer treatment on survival. However, long-term vital status was recorded for all 235 patients who were treated with bosentan in the two pivotal placebo-controlled studies (AC-052-351 and AC-052-352) and/or their two uncontrolled, open-label extensions. The mean duration of exposure to bosentan was  $1.9 \text{ years} \pm 0.7 \text{ years}$  (min: 0.1 years; max: 3.3 years) and patients were observed for a mean of  $2.0 \pm 0.6$  years. The majority of patients were diagnosed as primary pulmonary hypertension (72%) and were in WHO functional class III (84%). In this total population, Kaplan-Meier estimates of survival were 93% and 84% 1 and 2 years after the start of treatment with bosentan, respectively. Survival estimates were lower in the subgroup of patients with PAH secondary to systemic sclerosis. The estimates may have been influenced by the initiation of epoprostenol treatment in 43/235 patients.

#### *Studies performed in children with pulmonary arterial hypertension*

##### **BREATHE-3 (AC-052-356)**

Bosentan film-coated tablets were evaluated in an open-label uncontrolled study in 19 paediatric patients with PAH aged 3 to 15 years. This study was primarily designed as a pharmacokinetic study (see section 5.2). Patients had primary pulmonary hypertension (10 patients) or PAH related to congenital heart diseases (9 patients) and were in WHO functional class II ( $n = 15$ , 79%) or class III ( $n$

= 4, 21%) at baseline. Patients were divided into three body-weight groups and dosed with bosentan at approximately 2 mg/kg twice daily for 12 weeks. Half of the patients in each group were already being treated with intravenous epoprostenol and the dose of epoprostenol remained constant for the duration of the study.

Haemodynamics were measured in 17 patients. The mean increase from baseline in cardiac index was 0.5 L/min/m<sup>2</sup>, the mean decrease in mean pulmonary arterial pressure was 8 mmHg, and the mean decrease in PVR was 389 dyn·sec·cm<sup>-5</sup>. These haemodynamic improvements from baseline were similar with or without co-administration of epoprostenol. Changes in exercise test parameters at week 12 from baseline were highly variable and none were significant.

#### FUTURE 1/2 (AC-052-365/AC-052-367)

FUTURE 1 was an open-label, uncontrolled study that was conducted with the dispersible tablet formulation of bosentan administered at a maintenance dose of 4 mg/kg twice daily to 36 patients from 2 to 11 years of age. It was primarily designed as a pharmacokinetic study (see section 5.2). At baseline, patients had idiopathic (31 patients [86%]) or familial (5 patients [14%]) PAH, and were in WHO functional class II (n = 23, 64%) or class III (n = 13, 36%). In the FUTURE 1 study, the median exposure to study treatment was 13.1 weeks (range: 8.4 to 21.1). 33 of these patients were provided with continued treatment with bosentan dispersible tablets at a dose of 4 mg/kg twice daily in the FUTURE 2 uncontrolled extension phase for a median overall treatment duration of 2.3 years (range: 0.2 to 5.0 years). At baseline in FUTURE 1, 9 patients were taking epoprostenol. 9 patients were newly initiated on PAH-specific medication during the study. The Kaplan-Meier event-free estimate for worsening of PAH (death, lung transplantation, or hospitalisation for PAH worsening) at 2 years was 78.9%. The Kaplan-Meier estimate of overall survival at 2 years was 91.2%.

#### FUTURE 3 (AC-052-373)

In this open-label randomised study with the bosentan 32 mg dispersible tablet formulation, 64 children with stable PAH from 3 months to 11 years of age were randomised to 24 weeks' bosentan treatment 2 mg/kg twice daily (n = 33) or 2 mg/kg three times daily (n = 31). 43 (67.2%) were ≥ 2 years to 11 years old, 15 (23.4%) were between 1 and 2 years old, and 6 (9.4%) were between 3 months and 1 year old. The study was primarily designed as a pharmacokinetic study (see section 5.2), and efficacy endpoints were only exploratory. The aetiology of PAH, according to Dana Point classification, included idiopathic PAH (46%), heritable PAH (3%), associated PAH after corrective cardiac surgery (38%), and PAH related to congenital heart disease associated with systemic-to-pulmonary shunts, including Eisenmenger syndrome (13%). Patients were in WHO functional class I (n = 19, 29%), class II (n = 27, 42%) or class III (n = 18, 28%) at start of study treatment. At study entry, patients were treated with PAH medications (most frequently phosphodiesterase type-5 inhibitor [sildenafil] alone [35.9%], bosentan alone [10.9%], and a combination of bosentan, iloprost, and sildenafil [10.9%]) and continued their PAH treatment during the study.

At study start, less than half of the patients included (45.3% [29/64]) had bosentan treatment alone not combined with other PAH medication. 40.6% (26/64) remained on bosentan monotherapy during the 24 weeks of study treatment without experiencing PAH worsening. The analysis on the global population included (64 patients) showed that the majority had remained at least stable (i.e., without deterioration) based on non-paediatric-specific WHO functional class assessment (97% twice daily, 100% three times daily) and physician's global clinical impression (94% twice daily, 93% three times daily) during the treatment period. The Kaplan-Meier event-free estimate for worsening of PAH (death, lung transplantation, or hospitalisation for PAH worsening) at 24 weeks was 96.9% and 96.7% in the twice daily and three times daily groups, respectively.

There was no evidence of any clinical benefit with 2 mg/kg three times daily as compared to 2 mg/kg twice daily dosing.



*Study performed in neonates with persistent pulmonary hypertension of the newborn (PPHN):*

#### FUTURE 4 (AC-052-391)

This was a double-blind, placebo-controlled, randomised study in pre-term or term neonates (gestational age 36–42 weeks) with PPHN. Patients with suboptimal response to inhaled nitric oxide (iNO) despite at least 4 hours of continuous treatment were treated with bosentan dispersible tablets at 2 mg/kg twice daily (N = 13) or placebo (N = 8) via nasogastric tube as add-on therapy on top of iNO until complete weaning of iNO or until treatment failure (defined as need for extra-corporeal membrane oxygenation [ECMO] or initiation of alternative pulmonary vasodilator), and for a maximum of 14 days.

The median exposure to study treatment was 4.5 (range: 0.5–10.0) days in the bosentan group and 4.0 (range: 2.5–6.5) days in the placebo group.

The results did not indicate an additional benefit of bosentan in this population:

- The median time to complete weaning from iNO was 3.7 days (95% confidence limits [CLs] 1.17, 6.95) on bosentan and 2.9 days (95% CLs 1.26, 4.23) on placebo (p = 0.34).
- The median time to complete weaning from mechanical ventilation was 10.8 days (95% CLs 3.21, 12.21 days) on bosentan and 8.6 days (95% CLs 3.71, 9.66 days) on placebo (p = 0.24).
- One patient in the bosentan group had treatment failure (need for ECMO as per protocol definition), which was declared based on increasing Oxygenation Index values within 8 h after the first study drug dose. This patient recovered within the 60-day follow-up period.

#### *Combination with epoprostenol*

The combination of bosentan and epoprostenol has been investigated in two studies: AC-052-355 (BREATHE-2) and AC-052-356 (BREATHE-3). AC-052-355 was a multi-centre, randomised, double-blind, parallel-group study of bosentan versus placebo in 33 patients with severe PAH who were receiving concomitant epoprostenol therapy. AC-052-356 was an open-label, uncontrolled study; 10 of the 19 paediatric patients were on concomitant bosentan and epoprostenol therapy during the 12-week study. The safety profile of the combination was not different from the one expected with each component and the combination therapy was well tolerated in children and adults. The clinical benefit of the combination has not been demonstrated.

#### *Systemic sclerosis with digital ulcer disease*

Two randomised, double-blind, multi-centre, placebo-controlled studies have been conducted in 122 (study AC-052-401 [RAPIDS-1]) and 190 (study AC-052-331 [RAPIDS-2]) adult patients with systemic sclerosis and digital ulcer disease (either ongoing digital ulcers or a history of digital ulcers within the previous year). In study AC-052-331, patients had to have at least one digital ulcer of recent onset, and across the two studies 85% of patients had ongoing digital ulcer disease at baseline. After 4 weeks of bosentan 62.5 mg twice daily, the maintenance dose studied in both these studies was 125 mg twice daily. The duration of double-blind therapy was 16 weeks in study AC-052-401, and 24 weeks in study AC-052-331.

Background treatments for systemic sclerosis and digital ulcers were permitted if they remained constant for at least 1 month prior to the start of treatment and during the double-blind study period.

The number of new digital ulcers from baseline to study endpoint was a primary endpoint in both studies. Treatment with bosentan resulted in fewer new digital ulcers for the duration of therapy, compared with placebo. In study AC-052-401, during 16 weeks of double-blind therapy, patients in the bosentan group developed a mean of 1.4 new digital ulcers vs 2.7 new digital ulcers in the placebo group (p = 0.0042). In study AC-052-331, during 24 weeks of double-blind therapy, the corresponding figures were 1.9 vs 2.7 new digital ulcers, respectively (p = 0.0351). In both studies, patients on

bosentan were less likely to develop multiple new digital ulcers during the study and took longer to develop each successive new digital ulcer than did those on placebo. The effect of bosentan on reduction of the number of new digital ulcers was more pronounced in patients with multiple digital ulcers.

No effect of bosentan on time to healing of digital ulcers was observed in either study.

## 5.2 Pharmacokinetic properties

The pharmacokinetics of bosentan have mainly been documented in healthy subjects. Limited data in patients show that the exposure to bosentan in adult PAH patients is approximately 2-fold greater than in healthy adult subjects.

In healthy subjects, bosentan displays dose- and time-dependent pharmacokinetics. Clearance and volume of distribution decrease with increased intravenous doses and increase with time. After oral administration, the systemic exposure is proportional to dose up to 500 mg. At higher oral doses,  $C_{max}$  and AUC increase less than proportionally to the dose.

### Absorption

In healthy subjects, the absolute bioavailability of bosentan is approximately 50% and is not affected by food. The maximum plasma concentrations are attained within 3–5 hours.

### Distribution

Bosentan is highly bound (> 98%) to plasma proteins, mainly albumin. Bosentan does not penetrate into erythrocytes.

A volume of distribution ( $V_{ss}$ ) of about 18 litres was determined after an intravenous dose of 250 mg.

### Biotransformation and elimination

After a single intravenous dose of 250 mg, the clearance was 8.2 L/h. The terminal elimination half-life ( $t_{1/2}$ ) is 5.4 hours.

Upon multiple dosing, plasma concentrations of bosentan decrease gradually to 50–65% of those seen after single dose administration. This decrease is probably due to auto-induction of metabolising liver enzymes. Steady-state conditions are reached within 3–5 days.

Bosentan is eliminated by biliary excretion following metabolism in the liver by the cytochrome P450 isoenzymes, CYP2C9 and CYP3A4. Less than 3% of an administered oral dose is recovered in urine.

Bosentan forms three metabolites and only one of these is pharmacologically active. This metabolite is mainly excreted unchanged via the bile. In adult patients, the exposure to the active metabolite is greater than in healthy subjects. In patients with evidence of the presence of cholestasis, the exposure to the active metabolite may be increased.

Bosentan is an inducer of CYP2C9 and CYP3A4 and possibly also of CYP2C19 and the P-glycoprotein. *In vitro*, bosentan inhibits the bile salt export pump in hepatocyte cultures.

*In vitro* data demonstrated that bosentan had no relevant inhibitory effect on the CYP isoenzymes tested (CYP1A2, 2A6, 2B6, 2C8, 2C9, 2D6, 2E1, 3A4). Consequently, bosentan is not expected to increase the plasma concentrations of medicinal products metabolised by these isoenzymes.

### Comparison between formulations

In a pharmacokinetic crossover study (AC-052-116), 16 healthy adult subjects received 62.5 mg bosentan using the 62.5 mg film-coated tablet formulation or 64 mg bosentan using the 32 mg dispersible tablet formulation. Following treatment with the dispersible tablet, exposure to bosentan was lower than with the film-coated tablet (ratio of geometric means for  $AUC_{0-\infty}$  0.87 [90% CI: 0.78, 0.97]).  $T_{max}$  and  $t_{1/2}$  of bosentan were not significantly affected by the formulation.

### Pharmacokinetics in special populations

Based on the investigated range of each variable, it is not expected that the pharmacokinetics of bosentan will be influenced by gender, body weight, race, or age in the adult population to any relevant extent.

#### *Children*

Pharmacokinetics were studied in paediatric patients in 4 clinical studies (BREATHE-3, FUTURE 1, FUTURE-3 and FUTURE-4; see section 5.1). Due to limited data in children below 2 years of age, pharmacokinetics remain not well characterised in this age category.

Study AC-052-356 (BREATHE-3) evaluated the pharmacokinetics of single and multiple oral doses of the film-coated tablet formulation of bosentan in 19 children aged from 3 to 15 years with PAH who were dosed on the basis of body weight with 2 mg/kg twice daily. In this study, the exposure to bosentan decreased with time in a manner consistent with the known auto-induction properties of bosentan. The mean  $AUC$  (CV%) values of bosentan in paediatric patients treated with 31.25, 62.5 or 125 mg twice daily were 3,496 (49), 5,428 (79), and 6,124 (27) ng·h/mL, respectively, and were lower than the value of 8,149 (47) ng·h/mL observed in adult patients with PAH receiving 125 mg twice daily. At steady state, the systemic exposures in paediatric patients weighing 10–20 kg, 20–40 kg and > 40 kg were 43%, 67% and 75%, respectively, of the adult systemic exposure.

In study AC-052-365 (FUTURE 1), dispersible tablets were administered in 36 PAH children aged from 2 to 11 years. No dose proportionality was observed, as steady-state bosentan plasma concentrations and  $AUCs$  were similar at oral doses of 2 and 4 mg/kg ( $AUC_{\tau}$ : 3,577 ng·h/mL and 3,371 ng·h/mL for 2 mg/kg twice daily and 4 mg/kg twice daily, respectively). The average exposure to bosentan in these paediatric patients was about half the exposure in adult patients at the 125 mg twice daily maintenance dose but showed a large overlap with the exposures in adults.

In study AC-052-373 (FUTURE 3), using dispersible tablets, the exposure to bosentan in the patients treated with 2 mg/kg twice daily was comparable to that in the FUTURE 1 study. In the overall population ( $n = 31$ ), 2 mg/kg twice daily resulted in a daily exposure of 8,535 ng·h/mL;  $AUC_{\tau}$  was 4,268 ng·h/mL (CV: 61%). In patients between 3 months and 2 years the daily exposure was 7,879 ng·h/mL;  $AUC_{\tau}$  was 3,939 ng·h/mL (CV: 72%). In patients between 3 months and 1 year ( $n = 2$ )  $AUC_{\tau}$  was 5,914 ng·h/mL (CV: 85%), and in patients between 1 and 2 years ( $n = 7$ )  $AUC_{\tau}$  was 3,507 ng·h/mL (CV: 70%). In the patients above 2 years ( $n = 22$ ) the daily exposure was 8,820 ng·h/mL;  $AUC_{\tau}$  was 4,410 ng·h/mL (CV: 58%). Dosing bosentan 2 mg/kg three times daily did not increase exposure; daily exposure was 7,275 ng·h/mL (CV: 83%,  $n = 27$ ).

Based on the findings in studies BREATHE-3, FUTURE 1, and FUTURE-3, it appears that the exposure to bosentan reaches a plateau at lower doses in paediatric patients than in adults, and that doses higher than 2 mg/kg twice daily (4 mg/kg twice daily or 2 mg/kg three times daily) will not result in greater exposure to bosentan in paediatric patients.

In study AC-052-391 (FUTURE 4) conducted in neonates, bosentan concentrations increased slowly and continuously over the first dosing interval, resulting in low exposure ( $AUC_{0-12}$  in whole blood: 164 ng·h/mL, n = 11). At steady state,  $AUC_{\tau}$  was 6,165 ng·h/mL (CV: 133%, n = 7), which is similar to the exposure observed in adult PAH patients receiving 125 mg twice daily and taking into account a blood/plasma distribution ratio of 0.6.

The consequences of these findings regarding hepatotoxicity are unknown. Gender and concomitant use of intravenous epoprostenol had no significant effect on the pharmacokinetics of bosentan.

#### *Hepatic impairment*

In patients with mildly impaired liver function (Child-Pugh class A) no relevant changes in the pharmacokinetics have been observed. The steady-state AUC of bosentan was 9% higher and the AUC of the active metabolite, Ro 48-5033, was 33% higher in patients with mild hepatic impairment than in healthy volunteers.

The impact of moderately impaired liver function (Child-Pugh class B) on the pharmacokinetics of bosentan and its primary metabolite Ro 48-5033 was investigated in a study including 5 patients with pulmonary hypertension associated with portal hypertension and Child-Pugh class B hepatic impairment, and 3 patients with PAH from other causes and normal liver function. In the patients with Child-Pugh class B liver impairment, the mean (95% CI) steady-state AUC of bosentan was 360 (212–613) ng·h/mL, i.e., 4.7 times higher, and the mean (95% CI) AUC of the active metabolite Ro 48-5033 was 106 (58.4–192) ng·h/mL, i.e., 12.4 times higher than in the patients with normal liver function (bosentan: mean [95% CI] AUC: 76.1 [9.07–638] ng·h/mL; Ro 48-5033: mean [95% CI] AUC 8.57 [1.28–57.2] ng·h/ml). Though the number of patients included was limited and with high variability, these data indicate a marked increase in the exposure to bosentan and its primary metabolite Ro 48-5033 in patients with moderate liver function impairment (Child-Pugh class B).

The pharmacokinetics of bosentan have not been studied in patients with Child-Pugh class C hepatic impairment. Tracleer is contraindicated in patients with moderate to severe hepatic impairment, i.e., Child-Pugh class B or C (see section 4.3).

#### *Renal impairment*

In patients with severe renal impairment (creatinine clearance 15–30 mL/min), plasma concentrations of bosentan decreased by approximately 10%. Plasma concentrations of bosentan metabolites increased about 2-fold in these patients as compared with subjects with normal renal function. No dose adjustment is required in patients with renal impairment. There is no specific clinical experience in patients undergoing dialysis. Based on physicochemical properties and the high degree of protein binding, bosentan is not expected to be removed from the circulation by dialysis to any significant extent (see section 4.2).

### **5.3 Preclinical safety data**

A 2-year carcinogenicity study in mice showed an increased combined incidence of hepatocellular adenomas and carcinomas in males, but not in females, at plasma concentrations about 2 to 4 times the plasma concentrations achieved at the therapeutic dose in humans. In rats, oral administration of bosentan for 2 years produced a small, significant increase in the combined incidence of thyroid follicular cell adenomas and carcinomas in males, but not in females, at plasma concentrations about 9 to 14 times the plasma concentrations achieved at the therapeutic dose in humans. Bosentan was negative in tests for genotoxicity. There was evidence of a mild thyroid hormonal imbalance induced by bosentan in rats. However, there was no evidence of bosentan affecting thyroid function (thyroxine, TSH) in humans.

The effect of bosentan on mitochondrial function is unknown.

Bosentan has been shown to be teratogenic in rats at plasma levels higher than 1.5 times the plasma concentrations achieved at the therapeutic dose in humans. Teratogenic effects, including malformations of the head and face and of the major vessels, were dose dependent. The similarities of the pattern of malformations observed with other ET receptor antagonists and in ET knock-out mice indicate a class effect. Appropriate precautions must be taken for women of childbearing potential (see sections 4.3, 4.4 and 4.6).

Development of testicular tubular atrophy and impaired fertility has been linked with chronic administration of endothelin receptor antagonists in rodents.

In fertility studies in male and female rats, no effects on sperm count, motility and viability, or on mating performance or fertility were observed at exposures that were 21 and 43 times the expected therapeutic level in humans, respectively; nor was there any adverse effect on the development of the pre-implantation embryo or on implantation.

Slightly increased incidence of testicular tubular atrophy was observed in rats given bosentan orally at doses as low as 125 mg/kg/day (about 4 times the maximum recommended human dose [MRHD] and the lowest doses tested) for two years but not at doses as high as 1,500 mg/kg/day (about 50 times the MRHD) for 6 months. In a juvenile rat toxicity study, where rats were treated from Day 4 *post partum* up to adulthood, decreased absolute weights of testes and epididymides, and reduced number of sperm in epididymides were observed after weaning. The NOAEL was 21 times (at Day 21 *post partum*) and 2.3 times (Day 69 *post partum*) the human therapeutic exposure, respectively.

However, no effects on general development, growth, sensory, cognitive function and reproductive performance were detected at 7 (males) and 19 (females) times the human therapeutic exposure at Day 21 *post partum*. At adult age (Day 69 *post partum*), no effects of bosentan were detected at 1.3 (males) and 2.6 (females) times the therapeutic exposure in children with PAH.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Cellulose microcrystalline  
Calcium hydrogen phosphate anhydrous  
Croscarmellose sodium  
Silica colloidal anhydrous  
Tartaric acid  
Tutti frutti flavour  
Aspartame (E951)  
Acesulfame potassium  
Magnesium stearate

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

5 years

The remaining parts of a divided dispersible tablet can be stored at room temperature and should be used within 7 days.

#### **6.4 Special precautions for storage**

Do not store above 25 °C.

#### **6.5 Nature and contents of container**

Aluminium/Aluminium peel-push blisters containing 14 dispersible tablets.  
Cartons contain 56 dispersible tablets.

#### **6.6 Special precautions for disposal and other handling**

The dispersible tablet is contained in a child-proof blister.

Each dispersible tablet can be dissolved in water to make a liquid medicine, by adding the tablet to a little water on a spoon, using enough water to cover the whole tablet. When the tablet has fully dissolved, give the liquid to the patient.

If necessary, the dispersible tablet can be divided by breaking along the lines cut into the surface. Hold the tablet between the thumb and index finger on either side of one of the lines, with the line facing upwards, and break the tablet along the line (see figure below).



The remaining parts of a divided dispersible tablet can be stored at room temperature and should be used within 7 days.

### **7. MARKETING AUTHORISATION HOLDER**

Actelion Registration Ltd  
Chiswick Tower, 13<sup>th</sup> Floor  
389 Chiswick High Road  
London W4 4AL  
United Kingdom

### **8. MARKETING AUTHORISATION NUMBERS**

EU/1/02/220/006

### **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 15 May 2002

Date of renewal: 20 April 2012

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>.

## **ANNEX II**

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**



## **A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer responsible for batch release

Actelion Manufacturing GmbH  
Emil-Barell-Strasse 7  
79639 Grenzach-Wyhlen  
Germany

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2)

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic Safety Update Reports**

The requirements for submission of periodic safety update reports including liver reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2. of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency.
  - Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.
- **Additional risk minimisation measures**

### ***Controlled Distribution System:***

The MAH shall agree the details of a controlled distribution system with the National Competent Authorities and must implement such programme nationally to ensure that prior to prescribing all health care professionals who intend to prescribe and/or dispense Tracleer are provided with a Prescriber Kit containing the following:

- Information about Tracleer
- Patient Information Booklet/Patient Alert Card

The information provided about Tracleer shall contain the following key elements:

- That Tracleer is teratogenic in animals.
  - o Use in pregnant women is contraindicated.
  - o The need for effective contraception.
  - o That there is an interaction with hormonal contraceptives.
  - o Monthly pregnancy tests in women of childbearing potential are recommended.
- That Tracleer is hepatotoxic.
  - o Tracleer should not be used in Child Pugh Class B or C, i.e., moderate to severe hepatic impairment.
  - o Need for liver function tests to be measured:
    - Prior to initiation of treatment.
    - At monthly intervals during complete course of the treatment.
    - Two weeks after any dose increase.
  - o Need for close monitoring of liver function and dosage adjustment if levels rise above  $3 \times$  upper limit normal (ULN):
    - $>3$  and  $\leq 5 \times$  ULN: Confirm levels and if confirmed, reduce the daily dose or stop treatment and monitor liver function at least every 2 weeks.
    - $>5$  and  $\leq 8 \times$  ULN: Confirm levels and if confirmed stop treatment and monitor liver function at least every 2 weeks.  
In the above circumstances, if the levels return to pre-treatment values, continuing or re-introducing Tracleer may be considered.
    - $> 8 \times$  ULN or any of the above with associated clinical symptoms of liver injury: Treatment must be stopped and re-introduction of Tracleer is not to be considered.
- That treatment with Tracleer is associated with a decrease in haemoglobin.
  - o Need for blood monitoring:
    - Prior to initiation of treatment.
    - Monthly during the first 4 months.
    - Quarterly thereafter.
- That treatment with Tracleer might be associated with a decrease of sperm count.
- That co-administration of Tracleer with cyclosporine is contraindicated.
- That the safety database of Tracleer in the indication to reduce the number of new digital ulcers in patients with systemic sclerosis and ongoing digital ulcer disease is limited and physicians are encouraged to enrol patients in the surveillance programme/registry to further increase knowledge about the product. The surveillance programme/registry should prompt doctors to report adverse reactions.

The patient information shall contain the following information:

- That Tracleer is teratogenic in animals.
- That pregnant women must not take Tracleer.
- That women of childbearing potential must use effective contraception.
- That hormonal contraceptives on their own are not effective.
- The need for regular pregnancy tests.
- That Tracleer causes a decrease in haemoglobin and the need for regular blood tests.
- That Tracleer is hepatotoxic and the need for regular monitoring of liver function.

The Patient Alert Card:

- Patient Alert Card specifically aimed at facilitating patients' awareness of the need for regular blood tests for liver function.
- Patient Alert Card specifically aimed at informing patients of the need to avoid pregnancy and to ensure effective contraceptive measures are used.

### ***Digital Ulcer Outcome (DUO) Registry***

The MAH shall set up a surveillance programme/registry to collect information on the demographics, safety and outcome data from patients prescribed Tracleer to reduce the number of new digital ulcers in patients with systemic sclerosis and ongoing digital ulcer disease. The data to be collected shall be agreed with the CHMP. Details of the operation of the surveillance programme/registry shall be agreed with the National Competent Authorities in each Member State.

- **Obligation to conduct post-authorisation measures**

<b>Description</b>	<b>Due date</b>
Non-interventional post-authorisation safety study (PASS): Digital Ulcer Outcome (DUO) Registry: a multicentre, prospective, observational, non-interventional programme to document adherence to SmPC requirements for liver function, pregnancy testing	Yearly submission cycle with PSUR
Non-interventional post-authorisation safety study (PASS): Study AC-052-516: Disease characteristics and outcomes of pulmonary arterial hypertension in children and adolescents in real-world clinical settings to collect further data on long-term safety and outcomes in paediatric patients with PAH (Systematic review of prospective observational registries)	3-yearly submission cycle

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING  
CARTON WITH 14, 56 AND 112 TABLETS**

**OUTER CARTON/BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Tracleer 62.5 mg film-coated tablets

bosentan

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 62.5 mg bosentan (as monohydrate)

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

14 film-coated tablets  
56 film-coated tablets  
112 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use  
Read the package leaflet before use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT  
OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP {MM/YYYY}

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30 °C

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Actelion Registration Ltd  
Chiswick Tower, 13<sup>th</sup> Floor  
389 Chiswick High Road  
London W4 4AL  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/02/220/001  
EU/1/02/220/002  
EU/1/02/220/003

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Tracleer 62.5 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING  
CARTON WITH 56 AND 112 TABLETS**

**OUTER CARTON/BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Tracleer 125 mg film-coated tablets

bosentan

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 125 mg bosentan (as monohydrate)

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

56 film-coated tablets

112 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use

Read the package leaflet before use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT  
OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP {MM/YYYY}

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30 °C



**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Actelion Registration Ltd  
Chiswick Tower, 13<sup>th</sup> Floor  
389 Chiswick High Road  
London W4 4AL  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/02/220/004  
EU/1/02/220/005

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Tracleer 125 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING  
CARTON WITH 56 TABLETS**

**OUTER CARTON/BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Tracleer 32 mg dispersible tablets

bosentan

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each dispersible tablet contains 32 mg bosentan (as monohydrate)

**3. LIST OF EXCIPIENTS**

Aspartame (E951), see the package leaflet for further information  
Aspartame (E951) may be harmful for people with phenylketonuria

**4. PHARMACEUTICAL FORM AND CONTENTS**

56 dispersible tablets (14 x 4)

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use  
Read the package leaflet before use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT  
OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP {MM/YYYY}

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 25 °C

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Actelion Registration Ltd  
Chiswick Tower, 13<sup>th</sup> Floor  
389 Chiswick High Road  
London W4 4AL  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/02/220/006

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Tracleer 32 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING  
CARTON WITH 56 TABLETS**

**OUTER CARTON & BOTTLE LABEL/ BOTTLES**

**1. NAME OF THE MEDICINAL PRODUCT**

Tracleer 62.5 mg film-coated tablets

bosentan

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 62.5 mg bosentan (as monohydrate)

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

56 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use

Read the package leaflet before use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

Do not swallow the desiccant

**8. EXPIRY DATE**

EXP {MM/YYYY}

Use within 30 days after the first opening

Open date:

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Actelion Registration Ltd  
Chiswick Tower, 13<sup>th</sup> Floor  
389 Chiswick High Road  
London W4 4AL  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/02/220/007

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Tracleer 62.5 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING  
CARTON WITH 56 TABLETS**

**OUTER CARTON & BOTTLE LABEL/ BOTTLES**

**1. NAME OF THE MEDICINAL PRODUCT**

Tracleer 125 mg film-coated tablets

bosentan

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 125 mg bosentan (as monohydrate)

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

56 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use

Read the package leaflet before use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

Do not swallow the desiccant

**8. EXPIRY DATE**

EXP {MM/YYYY}

Use within 30 days after the first opening

Open date:

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Actelion Registration Ltd  
Chiswick Tower, 13<sup>th</sup> Floor  
389 Chiswick High Road  
London W4 4AL  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/02/220/008

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Tracleer 125 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Tracleer 62.5 mg tablets

bosentan

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Actelion

**3. EXPIRY DATE**

EXP {MM/YYYY}

**4. BATCH NUMBER**

Lot

**5. OTHER**



**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Tracleer 125 mg tablets

bosentan

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Actelion

**3. EXPIRY DATE**

EXP {MM/YYYY}

**4. BATCH NUMBER**

Lot

**5. OTHER**

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Tracleer 32 mg dispersible tablets

bosentan

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Actelion

**3. EXPIRY DATE**

EXP {MM/YYYY}

**4. BATCH NUMBER**

Lot

**5. OTHER**

**PATIENT ALERT CARD**

((Front cover))

**Important Safety Alerts for Patients taking  
Tracleer (bosentan)**

This card contains important information about Tracleer. Please read this card carefully before starting your treatment with Tracleer.

Your name: \_\_\_\_\_

Prescribing doctor: \_\_\_\_\_

If you have questions about Tracleer ask your doctor.

Actelion Registration Ltd

((Back cover))

**Contraception**

Do you currently use or take contraceptives?

Yes  No

If Yes, write the names of these here:

\_\_\_\_\_  
\_\_\_\_\_

Take this card to your doctor or your gynaecologist at your next visit and he/she will be able to advise you on whether you need to use additional or alternative contraceptive methods.

((Inside 1))

**If you are a woman of childbearing age read this page carefully**

**Pregnancy**

Tracleer may harm the development of the foetus. Therefore, you must not take Tracleer if you are pregnant and you must also not become pregnant while taking Tracleer.

Moreover, if you are suffering from pulmonary hypertension disease, the occurrence of a pregnancy can severely deteriorate the symptoms of your disease. If you suspect you may be pregnant, tell your doctor or gynaecologist.

**Contraception**

Birth control based on hormones – such as oral contraceptives or birth control pills, hormone injections, implants, or birth control skin patches don't reliably prevent pregnancy in women who are taking Tracleer. You need to use a barrier form of birth control – like a condom, diaphragm or vaginal sponge – in addition to any of these kinds of hormonal birth control. Be sure to discuss any questions you may have with your doctor or your gynaecologist – complete the details on the back of this card and take it to your doctor or gynaecologist at your next visit.

You should have a pregnancy test before initiation of Tracleer and every month during the treatment even if you think that you are not pregnant.

Date of first monthly test: \_\_\_\_\_

((Inside 2))

**Blood Test for Liver Function**

Some patients taking Tracleer were found to have abnormal liver function tests. During treatment with Tracleer, your doctor will arrange for regular blood tests to check for changes in your liver function.

**Remember to have your liver blood test every month.**

**After an increase in dose, an additional test will be done after 2 weeks.**

Date of first monthly test: \_\_\_\_\_

Your monthly liver blood test schedule:

- |                                    |                                    |                                    |
|------------------------------------|------------------------------------|------------------------------------|
| <input type="checkbox"/> Jan _____ | <input type="checkbox"/> May _____ | <input type="checkbox"/> Sep _____ |
| <input type="checkbox"/> Feb _____ | <input type="checkbox"/> Jun _____ | <input type="checkbox"/> Oct _____ |
| <input type="checkbox"/> Mar _____ | <input type="checkbox"/> Jul _____ | <input type="checkbox"/> Nov _____ |
| <input type="checkbox"/> Apr _____ | <input type="checkbox"/> Aug _____ | <input type="checkbox"/> Dec _____ |

**B. PACKAGE LEAFLET**

## Package leaflet: Information for the user

### Tracleer 62.5 mg film-coated tablets Bosentan

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, please ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

#### What is in this leaflet

1. What Tracleer is and what it is used for
2. What you need to know before you take Tracleer
3. How to take Tracleer
4. Possible side effects
5. How to store Tracleer
6. Contents of the pack and other information

#### 1. What Tracleer is and what it is used for

Tracleer tablets contain bosentan, which blocks a naturally occurring hormone called endothelin-1 (ET-1), which causes blood vessels to narrow. Tracleer therefore causes blood vessels to expand and belongs to the class of medicines called “endothelin receptor antagonists”.

Tracleer is used to treat:

- **Pulmonary arterial hypertension (PAH):** PAH is a disease of severe narrowing of the blood vessels in the lungs resulting in high blood pressure in the blood vessels (the pulmonary arteries) that carry blood from the heart to the lungs. This pressure reduces the amount of oxygen that can get into the blood in the lungs, making physical activity more difficult. Tracleer widens the pulmonary arteries, making it easier for the heart to pump blood through them. This lowers the blood pressure and relieves the symptoms.

Tracleer is used to treat patients with class III PAH to improve exercise capacity (the ability to carry out physical activity) and symptoms. The ‘class’ reflects the seriousness of the disease: ‘class III’ involves marked limitation of physical activity. Some improvements have also been shown in patients with class II PAH. ‘Class II’ involves slight limitation of physical activity. The PAH for which Tracleer is indicated can be:

- primary (with no identified cause or familial);
  - caused by scleroderma (also called systemic sclerosis, a disease where there is abnormal growth of the connective tissue that supports the skin and other organs);
  - caused by congenital (inborn) heart defects with shunts (abnormal passageways) causing abnormal flow of blood through the heart and lungs.
- **Digital ulcers:** (sores on the fingers and toes) in adult patients with a condition called scleroderma. Tracleer reduces the number of new finger and toe ulcers that appear.

## 2. What you need to know before you take Tracleer

### Do not take Tracleer:

- **if you are allergic to bosentan** or any of the other ingredients of this medicine (listed in section 6)
- **if you have liver problems** (ask your doctor)
- **if you are pregnant, or could get pregnant** because you are not using reliable contraceptive methods. Please read the information under “Contraceptives” and “Other medicines and Tracleer”
- **if you are taking cyclosporine A** (a medicine used after a transplant or to treat psoriasis)

If any of these apply to you, tell your doctor.

### Warnings and precautions

#### Tests your doctor will do before treatment

- a blood test to check your liver function
- a blood test to check for anaemia (low haemoglobin)
- a pregnancy test if you are a woman of childbearing potential

Some patients taking Tracleer have been found to have abnormal liver function tests and anaemia (low haemoglobin).

#### Tests your doctor will do during treatment

During treatment with Tracleer, your doctor will arrange for regular blood tests to check for changes in your liver function and haemoglobin level.

For all these tests please refer also to the Patient Alert Card (inside your pack of Tracleer tablets). It is important that you have these regular blood tests as long as you are taking Tracleer. We suggest you write the date of your most recent test and also of your next test (ask your doctor for the date) on the Patient Alert Card, to help you remember when your next test is due.

#### Blood tests for liver function

These will be done every month for the duration of treatment with Tracleer. After an increase in dose an additional test will be done after 2 weeks.

#### **Blood tests for anaemia**

These will be done every month for the first 4 months of treatment, then every 3 months after that, as patients taking Tracleer may get anaemia.

If these results are abnormal, your doctor may decide to reduce your dose or stop treatment with Tracleer and to perform further tests to investigate the cause.

#### **Children and adolescents**

Tracleer is not recommended in paediatric patients with systemic sclerosis and ongoing digital ulcer disease. Please see also section 3. How to take Tracleer.

#### **Other medicines and Tracleer**

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. It is especially important to tell your doctor if you are taking:

- cyclosporine A (a medicine used after transplants and to treat psoriasis), which must not be used together with Tracleer.
- sirolimus or tacrolimus, which are medicines used after transplants, as these are not recommended to be used together with Tracleer.

- glibenclamide (a diabetes medicine), rifampicin (a tuberculosis medicine) or fluconazole (a medicine against fungal infections), nevirapine (an HIV medicine), as these medicines are not recommended to be used together with Tracleer.
- other medicines for the treatment of HIV infection, which may require special monitoring if used together with Tracleer.
- hormonal contraceptives, which are not effective as the sole method of contraception when you take Tracleer. Inside your pack of Tracleer tablets you will find a Patient Alert Card which you should read carefully. Your doctor and/or gynaecologist will establish the contraception which is appropriate for you.

### **Driving and using machines**

Tracleer has no or negligible influence on the ability to drive and use machines. However, Tracleer can induce hypotension (decrease of your blood pressure) which can make you feel dizzy, affect your vision and affect your ability to drive and use machines. Therefore, if you feel dizzy or that your vision is blurred while taking Tracleer, do not drive or operate any tools or machines.

### **Women of childbearing age**

Do NOT take Tracleer if you are pregnant or planning to become pregnant.

### **Pregnancy tests**

Tracleer may harm unborn babies conceived before starting or during treatment. If you are a woman who could become pregnant, your doctor will ask you to take a pregnancy test before you start taking Tracleer, and regularly while you are taking Tracleer.

### **Contraceptives**

If it is possible that you could become pregnant, use a reliable form of birth control (contraception) while you are taking Tracleer. Your doctor or gynaecologist will advise you about reliable contraceptive methods while taking Tracleer. Because Tracleer may make hormonal contraception (e.g., oral, injection, implant, or skin patches) ineffective, this method on its own is not reliable. Therefore, if you use hormonal contraceptives you must also use a barrier method (e.g., female condom, diaphragm, contraceptive sponge, or your partner must also use a condom). Inside your pack of Tracleer tablets you will find a Patient Alert Card. You should complete this card and take it to your doctor at your next visit so that your doctor or gynaecologist can assess whether you need additional or alternative reliable contraceptive methods. Monthly pregnancy tests are recommended while you are taking Tracleer and are of childbearing age.

Tell your doctor immediately if you become pregnant while you are taking Tracleer, or plan to become pregnant in the near future.

### **Breast-feeding**

**Tell your doctor immediately if you are breast-feeding.** You are advised to stop breast-feeding if Tracleer is prescribed for you, because it is not known whether this medicine passes into breast milk.

### **Fertility**

If you are a man taking Tracleer, it is possible that this medicine may lower your sperm count. It cannot be excluded that this may affect your ability to father a child. Talk to your doctor if you have any questions or concerns about this.

## **3. How to take Tracleer**

Treatment with Tracleer should only be started and monitored by a doctor who has experience in the treatment of PAH or systemic sclerosis. Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

### **Tracleer with food and drink**

Tracleer can be taken with or without food.

### **Recommended dose**

#### **Adult**

The treatment in adults is usually started for the first 4 weeks with 62.5 mg twice daily (morning and evening), from then your doctor will usually advise you to take a 125 mg tablet twice daily, depending on how you react to Tracleer.

#### **Children and adolescents**

The dose recommendation in children is only for PAH. For children aged 1 year and older, treatment with Tracleer is usually started with 2 mg per kg bodyweight twice daily (morning and evening). Your doctor will advise you on your dosing.

Please note that Tracleer is also available as a dispersible 32 mg tablet formulation, which may make correct dosing easier for children and patients with low body weight or difficulties to swallow film-coated tablets.

If you have the impression that the effect of Tracleer is too strong or too weak, talk to your doctor in order to find out whether your dose needs to be changed.

#### **How to take Tracleer**

Tablets should be taken (morning and evening), swallowed with water. The tablets can be taken with or without food.

#### **If you take more Tracleer than you should**

If you take more tablets than you have been told to take, contact your doctor immediately.

#### **If you forget to take Tracleer**

If you forget to take Tracleer, take a dose as soon as you remember, then continue to take your tablets at the usual times. Do not take a double dose to make up for forgotten tablets.

#### **If you stop taking Tracleer**

Suddenly stopping your treatment with Tracleer may lead to your symptoms getting worse. Do not stop taking Tracleer unless your doctor tells you to. Your doctor may tell you to reduce the dose over a few days before stopping completely.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The most serious side effects with Tracleer are

- Abnormal liver function which may affect more than 1 in 10 people
- Anaemia (low blood value) which may affect up to 1 in 10 people. Anaemia may occasionally require blood transfusion

Your liver and blood values will be monitored during treatment with Tracleer (see section 2). It is important that you have these tests as ordered by your doctor.



Signs that your liver may not be working properly include:

- nausea (urge to vomit)
- vomiting
- fever (high temperature)
- pain in your stomach (abdomen)
- jaundice (yellowing of your skin or the whites of your eyes)
- dark-coloured urine
- itching of your skin
- lethargy or fatigue (unusual tiredness or exhaustion)
- flu-like syndrome (joint and muscle pain with fever)

If you notice any of these signs **tell your doctor immediately**.

Other side effects:

**Very common** (may affect **more than one in 10** people):

- Headache
- Oedema (swelling of the legs and ankles or other signs of fluid retention)

**Common** (may affect **up to one in 10** people):

- Flushed appearance or redness of skin
- Hypersensitivity reactions (including skin inflammation, itching and rash)
- Gastrooesophageal reflux disease (acid reflux)
- Diarrhoea
- Syncope (fainting)
- Palpitations (fast or irregular heart beats)
- Low blood pressure
- Nasal congestion

**Uncommon** (may affect **up to one in 100** people):

- Thrombocytopenia (low number of blood platelets)
- Neutropenia/leukopenia (low number of white blood cells)
- Elevated liver function tests with hepatitis (inflammation of the liver) including possible exacerbation of underlying hepatitis and/or jaundice (yellowing of the skin or the whites of the eyes)

**Rare** (may affect **up to one in 1000** people):

- Anaphylaxis (general allergic reaction), angioedema (swelling, most commonly around the eyes, lips, tongue or throat)
- Cirrhosis (scarring) of the liver, liver failure (serious disturbance of liver function)

Blurred vision have also been reported at an unknown frequency (frequency cannot be estimated from the available data).

### **Side effects in children and adolescents**

The side effects that have been reported in children treated with Tracleer are the same as those in adults.

### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Tracleer

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and on the blister after “EXP”.

For white high-density polyethylene bottles, use within 30 days after the first opening.

For PVC/PE/PVDC/aluminium-blisters:

Do not store above 30 °C.

For white high-density polyethylene bottles:

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away any medicines you no longer use. These measures will help to protect the environment.

## 6. Contents of the pack and other information

### What Tracleer contains

- **Tracleer 62.5 mg film-coated tablets:** The active substance is bosentan as monohydrate. Each tablet contains 62.5 mg of bosentan (as monohydrate).
- **The other ingredients** in the tablet core are maize starch, pregelatinised starch, sodium starch glycollate, povidone, glycerol dibehenate and magnesium stearate. **The film-coat** contains hypromellose, glycerol triacetate, talc, titanium dioxide (E171), iron oxide yellow (E172), iron oxide red (E172) and ethylcellulose.

### What Tracleer looks like and contents of the pack

Tracleer 62.5 mg film-coated tablets are orange-white, round film-coated tablets with “62,5” on one side.

**PVC/PE/PVDC/aluminium-blisters** containing **14 film-coated tablets**. Cartons contain 14, 56 or 112 film-coated tablets (Tracleer 62.5 mg film-coated tablets).

**White high-density polyethylene bottles with a silica gel desiccant** containing 56 film-coated tablets. Cartons contain 56 film-coated tablets (Tracleer 62.5 mg film-coated tablets).

Do not swallow the desiccant.

Not all pack sizes may be marketed.

**Marketing authorisation holder:**

Actelion Registration Ltd  
Chiswick Tower, 13<sup>th</sup> Floor  
389 Chiswick High Road  
London W4 4AL  
United Kingdom

**Manufacturer:**

Actelion Manufacturing GmbH  
Emil-Barell-Strasse 7  
79639 Grenzach-Wyhlen  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

**België/Belgique/Belgien**

Actelion Pharmaceuticals Belgium N.V.  
Tél/Tel: +32-(0)15 284 777

**Lietuva**

UAB ALGOL PHARMA  
Tel: +370 37 40 86 81

**България**

Аквахим АД  
Тел.: +359 2 807 50 00

**Luxembourg/Luxemburg**

Actelion Pharmaceuticals Belgium N.V.  
Tél/Tel: +32-(0)15 284 777

**Česká republika**

Actelion Pharmaceuticals CZ, s.r.o.  
Tel: +420 221 968 006

**Magyarország**

Actelion Pharmaceuticals Hungaria Kft.  
Tel: +36-1-413-3270

**Danmark**

Actelion Danmark,  
Filial af Actelion Pharmaceuticals Sverige AB,  
Sverige  
Tlf: +45 3694 45 95

**Malta**

Actelion Pharmaceuticals UK Ltd  
Tel: +44 208 987 3333

**Deutschland**

Actelion Pharmaceuticals Deutschland GmbH  
Tel: +49 761 45 64 0

**Nederland**

Actelion Pharmaceuticals Nederland B.V.  
Tel: +31 (0)348 435950

**Eesti**

Algol Pharma OÜ  
Tel: +372 605 6014

**Norge**

Actelion Pharmaceuticals Sverige AB,  
Filial Norge  
Tlf: +47 22480370

**Ελλάδα**

Actelion Pharmaceuticals Ελλάς A.E.  
Τηλ: +30 210 675 25 00

**Österreich**

Actelion Pharmaceuticals Austria GmbH  
Tel: +43 1 505 4527

**España**

Actelion Pharmaceuticals España S.L.  
Tel: +34 93 366 43 99

**Polska**

Actelion Pharma Polska Sp. z o.o.  
Tel: +48 (22) 262 31 00

**France**

Actelion Pharmaceuticals France SAS  
Tél: +33 1 58 62 32 32

**Hrvatska**

Medis Adria d.o.o.  
Tel: +385 (0) 1 2303 446

**Ireland**

Actelion Pharmaceuticals UK Ltd  
Tel: +44 208 987 3333

**Ísland**

Actelion Pharmaceuticals Sverige AB  
Sími: +46 8 544 982 50

**Italia**

Actelion Pharmaceuticals Italia S.r.l.  
Tel: +39 0542 64 87 40

**Κύπρος**

Actelion Pharmaceuticals Ελλάς A.E.  
Τηλ: +30 210 675 25 00

**Latvija**

Algol Pharma SIA  
Tel: +371 6761 9365

**Portugal**

Actelion Pharmaceuticals Portugal Lda.  
Tel: +351 21 358 6120

**România**

Geneva Romfarm International SRL  
Tel: +40 (021) 231 3561

**Slovenija**

Medis d.o.o.  
Tel: +386-(0)1 589 69 00

**Slovenská republika**

Actelion Pharmaceuticals SK, s.r.o.  
Tel: +420 221 968 006

**Suomi/Finland**

Actelion Pharmaceuticals Sverige AB,  
Filial Finland  
Puh/Tel: +358 9 2510 7720

**Sverige**

Actelion Pharmaceuticals Sverige AB  
Tel: +46 8 544 982 50

**United Kingdom**

Actelion Pharmaceuticals UK Ltd  
Tel: +44 208 987 3333

**This leaflet was last revised in**

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu/>.

## Package leaflet: Information for the user

### Tracleer 125 mg film-coated tablets

Bosentan

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, please ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

#### What is in this leaflet

1. What Tracleer is and what it is used for
2. What you need to know before you take Tracleer
3. How to take Tracleer
4. Possible side effects
5. How to store Tracleer
6. Contents of the pack and other information

#### 1. What Tracleer is and what it is used for

Tracleer tablets contain bosentan, which blocks a naturally occurring hormone called endothelin-1 (ET-1), which causes blood vessels to narrow. Tracleer therefore causes blood vessels to expand and belongs to the class of medicines called “endothelin receptor antagonists”.

Tracleer is used to treat:

- **Pulmonary arterial hypertension (PAH):** PAH is a disease of severe narrowing of the blood vessels in the lungs resulting in high blood pressure in the blood vessels (the pulmonary arteries) that carry blood from the heart to the lungs. This pressure reduces the amount of oxygen that can get into the blood in the lungs, making physical activity more difficult. Tracleer widens the pulmonary arteries, making it easier for the heart to pump blood through them. This lowers the blood pressure and relieves the symptoms.

Tracleer is used to treat patients with class III PAH to improve exercise capacity (the ability to carry out physical activity) and symptoms. The ‘class’ reflects the seriousness of the disease: ‘class III’ involves marked limitation of physical activity. Some improvements have also been shown in patients with class II PAH. ‘Class II’ involves slight limitation of physical activity. The PAH for which Tracleer is indicated can be:

- primary (with no identified cause or familial);
  - caused by scleroderma (also called systemic sclerosis, a disease where there is abnormal growth of the connective tissue that supports the skin and other organs);
  - caused by congenital (inborn) heart defects with shunts (abnormal passageways) causing abnormal flow of blood through the heart and lungs.
- **Digital ulcers:** (sores on the fingers and toes) in adult patients with a condition called scleroderma. Tracleer reduces the number of new finger and toe ulcers that appear.

## 2. What you need to know before you take Tracleer

### Do not take Tracleer:

- **if you are allergic to bosentan** or any of the other ingredients of this medicine (listed in section 6)
- **if you have liver problems** (ask your doctor)
- **if you are pregnant, or could get pregnant** because you are not using reliable contraceptive methods. Please read the information under “Contraceptives” and “Other medicines and Tracleer”
- **if you are taking cyclosporine A** (a medicine used after a transplant or to treat psoriasis)

If any of these apply to you, tell your doctor.

### Warnings and precautions

#### Tests your doctor will do before treatment

- a blood test to check your liver function
- a blood test to check for anaemia (low haemoglobin)
- a pregnancy test if you are a woman of childbearing potential

Some patients taking Tracleer have been found to have abnormal liver function tests and anaemia (low haemoglobin).

#### Tests your doctor will do during treatment

During treatment with Tracleer, your doctor will arrange for regular blood tests to check for changes in your liver function and haemoglobin level.

For all these tests please refer also to the Patient Alert Card (inside your pack of Tracleer tablets). It is important that you have these regular blood tests as long as you are taking Tracleer. We suggest you write the date of your most recent test and also of your next test (ask your doctor for the date) on the Patient Alert Card, to help you remember when your next test is due.

#### Blood tests for liver function

These will be done every month for the duration of treatment with Tracleer. After an increase in dose an additional test will be done after 2 weeks.

#### **Blood tests for anaemia**

These will be done every month for the first 4 months of treatment, then every 3 months after that, as patients taking Tracleer may get anaemia.

If these results are abnormal, your doctor may decide to reduce your dose or stop treatment with Tracleer and to perform further tests to investigate the cause.

#### **Children and adolescents**

Tracleer is not recommended in paediatric patients with systemic sclerosis and ongoing digital ulcer disease. Please see also section 3. How to take Tracleer.

#### **Other medicines and Tracleer**

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. It is especially important to tell your doctor if you are taking:

- cyclosporine A (a medicine used after transplants and to treat psoriasis), which must not be used together with Tracleer.
- sirolimus or tacrolimus, which are medicines used after transplants, as these are not recommended to be used together with Tracleer.

- glibenclamide (a diabetes medicine), rifampicin (a tuberculosis medicine) or fluconazole (a medicine against fungal infections), nevirapine (an HIV medicine), as these medicines are not recommended to be used together with Tracleer.
- other medicines for the treatment of HIV infection, which may require special monitoring if used together with Tracleer.
- hormonal contraceptives, which are not effective as the sole method of contraception when you take Tracleer. Inside your pack of Tracleer tablets you will find a Patient Alert Card which you should read carefully. Your doctor and/or gynaecologist will establish the contraception which is appropriate for you.

### **Driving and using machines**

Tracleer has no or negligible influence on the ability to drive and use machines. However, Tracleer can induce hypotension (decrease of your blood pressure) which can make you feel dizzy, affect your vision and affect your ability to drive and use machines. Therefore, if you feel dizzy or that your vision is blurred while taking Tracleer, do not drive or operate any tools or machines.

### **Women of childbearing age**

Do NOT take Tracleer if you are pregnant or planning to become pregnant.

### **Pregnancy tests**

Tracleer may harm unborn babies conceived before starting or during treatment. If you are a woman who could become pregnant, your doctor will ask you to take a pregnancy test before you start taking Tracleer, and regularly while you are taking Tracleer.

### **Contraceptives**

If it is possible that you could become pregnant, use a reliable form of birth control (contraception) while you are taking Tracleer. Your doctor or gynaecologist will advise you about reliable contraceptive methods while taking Tracleer. Because Tracleer may make hormonal contraception (e.g., oral, injection, implant, or skin patches) ineffective, this method on its own is not reliable. Therefore, if you use hormonal contraceptives you must also use a barrier method (e.g., female condom, diaphragm, contraceptive sponge, or your partner must also use a condom). Inside your pack of Tracleer tablets you will find a Patient Alert Card. You should complete this card and take it to your doctor at your next visit so that your doctor or gynaecologist can assess whether you need additional or alternative reliable contraceptive methods. Monthly pregnancy tests are recommended while you are taking Tracleer and are of childbearing age.

Tell your doctor immediately if you become pregnant while you are taking Tracleer, or plan to become pregnant in the near future.

### **Breast-feeding**

**Tell your doctor immediately if you are breast-feeding.** You are advised to stop breast-feeding if Tracleer is prescribed for you, because it is not known whether this medicine passes into breast milk.

### **Fertility**

If you are a man taking Tracleer, it is possible that this medicine may lower your sperm count. It cannot be excluded that this may affect your ability to father a child. Talk to your doctor if you have any questions or concerns about this.

### **3. How to take Tracleer**

Treatment with Tracleer should only be started and monitored by a doctor who has experience in the treatment of PAH or systemic sclerosis. Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

#### **Tracleer with food and drink**

Tracleer can be taken with or without food.

#### **Recommended dose**

##### **Adult**

The treatment in adults is usually started for the first 4 weeks with 62.5 mg twice daily (morning and evening), from then your doctor will usually advise you to take a 125 mg tablet twice daily, depending on how you react to Tracleer.

##### **Children and adolescents**

The dose recommendation in children is only for PAH. For children aged 1 year and older, treatment with Tracleer is usually started with 2 mg per kg bodyweight twice daily (morning and evening). Your doctor will advise you on your dosing.

Please note that Tracleer is also available as a dispersible 32 mg tablet formulation, which may make correct dosing easier for children and patients with low body weight or difficulties to swallow film-coated tablets.

If you have the impression that the effect of Tracleer is too strong or too weak, talk to your doctor in order to find out whether your dose needs to be changed.

#### **How to take Tracleer**

Tablets should be taken (morning and evening), swallowed with water. The tablets can be taken with or without food.

#### **If you take more Tracleer than you should**

If you take more tablets than you have been told to take, contact your doctor immediately.

#### **If you forget to take Tracleer**

If you forget to take Tracleer, take a dose as soon as you remember, then continue to take your tablets at the usual times. Do not take a double dose to make up for forgotten tablets.

#### **If you stop taking Tracleer**

Suddenly stopping your treatment with Tracleer may lead to your symptoms getting worse. Do not stop taking Tracleer unless your doctor tells you to. Your doctor may tell you to reduce the dose over a few days before stopping completely.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The most serious side effects with Tracleer are



- Abnormal liver function which may affect more than 1 in 10 people
- Anaemia (low blood value) which may affect up to 1 in 10 people. Anaemia may occasionally require blood transfusion

Your liver and blood values will be monitored during treatment with Tracleer (see section 2). It is important that you have these tests as ordered by your doctor.

Signs that your liver may not be working properly include:

- nausea (urge to vomit)
- vomiting
- fever (high temperature)
- pain in your stomach (abdomen)
- jaundice (yellowing of your skin or the whites of your eyes)
- dark-coloured urine
- itching of your skin
- lethargy or fatigue (unusual tiredness or exhaustion)
- flu-like syndrome (joint and muscle pain with fever)

If you notice any of these signs **tell your doctor immediately**.

Other side effects:

**Very common** (may affect **more than one in 10** people):

- Headache
- Oedema (swelling of the legs and ankles or other signs of fluid retention)

**Common** (may affect **up to one in 10** people):

- Flushed appearance or redness of skin
- Hypersensitivity reactions (including skin inflammation, itching and rash)
- Gastrooesophageal reflux disease (acid reflux)
- Diarrhoea
- Syncope (fainting)
- Palpitations (fast or irregular heart beats)
- Low blood pressure
- Nasal congestion

**Uncommon** (may affect **up to one in 100** people):

- Thrombocytopenia (low number of blood platelets)
- Neutropenia/leukopenia (low number of white blood cells)
- Elevated liver function tests with hepatitis (inflammation of the liver) including possible exacerbation of underlying hepatitis and/or jaundice (yellowing of the skin or the whites of the eyes)

**Rare** (may affect **up to one in 1000** people):

- Anaphylaxis (general allergic reaction), angioedema (swelling, most commonly around the eyes, lips, tongue or throat)
- Cirrhosis (scarring) of the liver, liver failure (serious disturbance of liver function)

Blurred vision have also been reported at an unknown frequency (frequency cannot be estimated from the available data).

### **Side effects in children and adolescents**

The side effects that have been reported in children treated with Tracleer are the same as those in adults.

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Tracleer**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and on the blister after “EXP”.

For white high-density polyethylene bottles, use within 30 days after the first opening.

For PVC/PE/PVDC/aluminium-blisters:  
Do not store above 30 °C.

For white high-density polyethylene bottles:  
This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away any medicines you no longer use. These measures will help to protect the environment.

## **6. Contents of the pack and other information**

### **What Tracleer contains**

- **Tracleer 125 mg film-coated tablets:** The active substance is bosentan as monohydrate. Each tablet contains 125 mg of bosentan (as monohydrate).
- **The other ingredients** in the tablet core are maize starch, pregelatinised starch, sodium starch glycollate, povidone, glycerol dibehenate and magnesium stearate. **The film-coat** contains hypromellose, glycerol triacetate, talc, titanium dioxide (E171), iron oxide yellow (E172), iron oxide red (E172) and ethylcellulose.

### **What Tracleer looks like and contents of the pack**

Tracleer 125 mg film-coated tablets are orange-white, oval film-coated tablets with “125” on one side.

**PVC/PE/PVDC/aluminium-blisters** containing **14 film-coated tablets**. Cartons contain 56 or 112 film-coated tablets (Tracleer 125 mg film-coated tablets).

**White high-density polyethylene bottles with a silica gel desiccant** containing 56 film-coated tablets. Cartons contain 56 film-coated tablets (Tracleer 125 mg film-coated tablets).

Do not swallow the desiccant.

Not all pack sizes may be marketed.

**Marketing authorisation holder:**

Actelion Registration Ltd  
Chiswick Tower, 13<sup>th</sup> Floor  
389 Chiswick High Road  
London W4 4AL  
United Kingdom

**Manufacturer:**

Actelion Manufacturing GmbH  
Emil-Barell-Strasse 7  
79639 Grenzach-Wyhlen  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

**België/Belgique/Belgien**

Actelion Pharmaceuticals Belgium N.V.  
Tél/Tel: +32-(0)15 284 777

**Lietuva**

UAB ALGOL PHARMA  
Tel: +370 37 40 86 81

**България**

АКВАХИМ АД  
Тел.: +359 2 807 50 00

**Luxembourg/Luxemburg**

Actelion Pharmaceuticals Belgium N.V.  
Tél/Tel: +32-(0)15 284 777

**Česká republika**

Actelion Pharmaceuticals CZ, s.r.o.  
Tel: +420 221 968 006

**Magyarország**

Actelion Pharmaceuticals Hungaria Kft.  
Tel: +36-1-413-3270

**Danmark**

Actelion Danmark,  
Filial af Actelion Pharmaceuticals Sverige AB,  
Sverige  
Tlf: +45 3694 45 95

**Malta**

Actelion Pharmaceuticals UK Ltd  
Tel: +44 208 987 3333

**Deutschland**

Actelion Pharmaceuticals Deutschland GmbH  
Tel: +49 761 45 64 0

**Nederland**

Actelion Pharmaceuticals Nederland B.V.  
Tel: +31 (0)348 435950

**Eesti**

Algol Pharma OÜ  
Tel: +372 605 6014

**Norge**

Actelion Pharmaceuticals Sverige AB,  
Filial Norge  
Tlf: +47 22480370

**Ελλάδα**

Actelion Pharmaceuticals Ελλάς A.E.  
Τηλ: +30 210 675 25 00

**Österreich**

Actelion Pharmaceuticals Austria GmbH  
Tel: +43 1 505 4527

**España**

Actelion Pharmaceuticals España S.L.  
Tel: +34 93 366 43 99

**France**

Actelion Pharmaceuticals France SAS  
Tél: +33 1 58 62 32 32

**Hrvatska**

Medis Adria d.o.o.  
Tel: +385 (0) 1 2303 446

**Ireland**

Actelion Pharmaceuticals UK Ltd  
Tel: +44 208 987 3333

**Ísland**

Actelion Pharmaceuticals Sverige AB  
Sími: +46 8 544 982 50

**Italia**

Actelion Pharmaceuticals Italia S.r.l.  
Tel: +39 0542 64 87 40

**Κύπρος**

Actelion Pharmaceuticals Ελλάς A.E.  
Τηλ: +30 210 675 25 00

**Latvija**

Algol Pharma SIA  
Tel: +371 6761 9365

**Polska**

Actelion Pharma Polska Sp. z o.o.  
Tel: +48 (22) 262 31 00

**Portugal**

Actelion Pharmaceuticals Portugal Lda.  
Tel: +351 21 358 6120

**România**

Geneva Romfarm International SRL  
Tel: +40 (021) 231 3561

**Slovenija**

Medis d.o.o.  
Tel: +386-(0)1 589 69 00

**Slovenská republika**

Actelion Pharmaceuticals SK, s.r.o.  
Tel: +420 221 968 006

**Suomi/Finland**

Actelion Pharmaceuticals Sverige AB,  
Filial Finland  
Puh/Tel: +358 9 2510 7720

**Sverige**

Actelion Pharmaceuticals Sverige AB  
Tel: +46 8 544 982 50

**United Kingdom**

Actelion Pharmaceuticals UK Ltd  
Tel: +44 208 987 3333

**This leaflet was last revised in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu/>.

## Package leaflet: Information for the user

### Tracleer 32 mg dispersible tablets Bosentan

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, please ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

#### What is in this leaflet

1. What Tracleer is and what it is used for
2. What you need to know before you take Tracleer
3. How to take Tracleer
4. Possible side effects
5. How to store Tracleer
6. Contents of the pack and other information

#### 1. What Tracleer is and what it is used for

Tracleer tablets contain bosentan, which blocks a naturally occurring hormone called endothelin-1 (ET-1), which causes blood vessels to narrow. Tracleer therefore causes blood vessels to expand and belongs to the class of medicines called “endothelin receptor antagonists”.

Tracleer is used to treat:

- **Pulmonary arterial hypertension (PAH):** PAH is a disease of severe narrowing of the blood vessels in the lungs resulting in high blood pressure in the blood vessels (the pulmonary arteries) that carry blood from the heart to the lungs. This pressure reduces the amount of oxygen that can get into the blood in the lungs, making physical activity more difficult. Tracleer widens the pulmonary arteries, making it easier for the heart to pump blood through them. This lowers the blood pressure and relieves the symptoms.

Tracleer is used to treat patients with class III PAH to improve exercise capacity (the ability to carry out physical activity) and symptoms. The ‘class’ reflects the seriousness of the disease: ‘class III’ involves marked limitation of physical activity. Some improvements have also been shown in patients with class II PAH. ‘Class II’ involves slight limitation of physical activity. The PAH for which Tracleer is indicated can be:

- primary (with no identified cause or familial);
  - caused by scleroderma (also called systemic sclerosis, a disease where there is abnormal growth of the connective tissue that supports the skin and other organs);
  - caused by congenital (inborn) heart defects with shunts (abnormal passageways) causing abnormal flow of blood through the heart and lungs.
- **Digital ulcers:** (sores on the fingers and toes) in adult patients with a condition called scleroderma. Tracleer reduces the number of new finger and toe ulcers that appear.

## 2. What you need to know before you take Tracleer

### Do not take Tracleer:

- **if you are allergic to bosentan** or any of the other ingredients of this medicine (listed in section 6)
- **if you have liver problems** (ask your doctor)
- **if you are pregnant, or could get pregnant** because you are not using reliable contraceptive methods. Please read the information under “Contraceptives” and “Other medicines and Tracleer”
- **if you are taking cyclosporine A** (a medicine used after a transplant or to treat psoriasis)

If any of these apply to you, tell your doctor.

### Warnings and precautions

#### Tests your doctor will do before treatment

- a blood test to check your liver function
- a blood test to check for anaemia (low haemoglobin)
- a pregnancy test if you are a woman of childbearing potential

Some patients taking Tracleer have been found to have abnormal liver function tests and anaemia (low haemoglobin).

#### Tests your doctor will do during treatment

During treatment with Tracleer, your doctor will arrange for regular blood tests to check for changes in your liver function and haemoglobin level.

For all these tests please refer also to the Patient Alert Card (inside your pack of Tracleer tablets). It is important that you have these regular blood tests as long as you are taking Tracleer. We suggest you write the date of your most recent test and also of your next test (ask your doctor for the date) on the Patient Alert Card, to help you remember when your next test is due.

#### Blood tests for liver function

These will be done every month for the duration of treatment with Tracleer. After an increase in dose an additional test will be done after 2 weeks.

#### **Blood tests for anaemia**

These will be done every month for the first 4 months of treatment, then every 3 months after that, as patients taking Tracleer may get anaemia.

If these results are abnormal, your doctor may decide to reduce your dose or stop treatment with Tracleer and to perform further tests to investigate the cause.

#### **Children and adolescents**

Tracleer is not recommended in paediatric patients with systemic sclerosis and ongoing digital ulcer disease. Please see also section 3. How to take Tracleer.

#### **Other medicines and Tracleer**

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. It is especially important to tell your doctor if you are taking:

- cyclosporine A (a medicine used after transplants and to treat psoriasis), which must not be used together with Tracleer.
- sirolimus or tacrolimus, which are medicines used after transplants, as these are not recommended to be used together with Tracleer.

- glibenclamide (a diabetes medicine), rifampicin (a tuberculosis medicine) or fluconazole (a medicine against fungal infections), nevirapine (an HIV medicine), as these medicines are not recommended to be used together with Tracleer.
- other medicines for the treatment of HIV infection, which may require special monitoring if used together with Tracleer.
- hormonal contraceptives, which are not effective as the sole method of contraception when you take Tracleer. Inside your pack of Tracleer tablets you will find a Patient Alert Card which you should read carefully. Your doctor and/or gynaecologist will establish the contraception which is appropriate for you.

### **Driving and using machines**

Tracleer has no or negligible influence on the ability to drive and use machines. However, Tracleer can induce hypotension (decrease of your blood pressure) which can make you feel dizzy, affect your vision and affect your ability to drive and use machines. Therefore, if you feel dizzy or that your vision is blurred while taking Tracleer, do not drive or operate any tools or machines.

### **Women of childbearing age**

Do NOT take Tracleer if you are pregnant or planning to become pregnant.

### **Pregnancy tests**

Tracleer may harm unborn babies conceived before starting or during treatment. If you are a woman who could become pregnant, your doctor will ask you to take a pregnancy test before you start taking Tracleer, and regularly while you are taking Tracleer.

### **Contraceptives**

If it is possible that you could become pregnant, use a reliable form of birth control (contraception) while you are taking Tracleer. Your doctor or gynaecologist will advise you about reliable contraceptive methods while taking Tracleer. Because Tracleer may make hormonal contraception (e.g., oral, injection, implant, or skin patches) ineffective, this method on its own is not reliable. Therefore, if you use hormonal contraceptives you must also use a barrier method (e.g., female condom, diaphragm, contraceptive sponge, or your partner must also use a condom). Inside your pack of Tracleer tablets you will find a Patient Alert Card. You should complete this card and take it to your doctor at your next visit so that your doctor or gynaecologist can assess whether you need additional or alternative reliable contraceptive methods. Monthly pregnancy tests are recommended while you are taking Tracleer and are of childbearing age.

Tell your doctor immediately if you become pregnant while you are taking Tracleer, or plan to become pregnant in the near future.

### **Breast-feeding**

**Tell your doctor immediately if you are breast-feeding.** You are advised to stop breast-feeding if Tracleer is prescribed for you, because it is not known whether this medicine passes into breast milk.

### **Fertility**

If you are a man taking Tracleer, it is possible that this medicine may lower your sperm count. It cannot be excluded that this may affect your ability to father a child. Talk to your doctor if you have any questions or concerns about this.

### **Important information about some of the ingredients of Tracleer**

Each Tracleer 32 mg dispersible tablets contains 3.7 mg of Aspartame (E951) which is a source of phenylalanine. Aspartame may be harmful for people with phenylketonuria.

### 3. How to take Tracleer

Treatment with Tracleer should only be started and monitored by a doctor who has experience in the treatment of PAH or systemic sclerosis. Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

#### Tracleer with food and drink

Tracleer can be taken with or without food.

#### Recommended dose

##### Adult

The treatment in adults is usually started for the first 4 weeks with 62.5 mg twice daily (morning and evening), from then your doctor will usually advise you to take a 125 mg tablet twice daily, depending on how you react to Tracleer.

##### Children and adolescents

The dose recommendation in children is only for PAH. For children aged 1 year and older, treatment with Tracleer is usually started with 2 mg per kg bodyweight twice daily (morning and evening). Your doctor will advise you on your dosing.

If necessary the dispersible tablet can be divided along the break-marks into four equal parts.


If you have the impression that the effect of Tracleer is too strong or too weak, talk to your doctor in order to find out whether your dose needs to be changed.

#### How to take Tracleer

Tablets should be taken (morning and evening), swallowed with water. The tablets can be taken with or without food.

The dispersible tablet is contained in a child-proof blister.

To remove the dispersible tablet:



1. Separate the individual blister cavity at the perforations.
2. Peel off the top layer.
3. Push the pharmaceutical product through the foil.

Each Tracleer dispersible tablet can be dissolved in water to make a liquid medicine. To make a liquid medicine, add the tablet to a little water on a spoon. Use enough water to cover the whole tablet. Leave for about one minute, until the tablet has fully dissolved, and then swallow all of the liquid. Add a little more water to the spoon and swallow all of the liquid to make sure all of the medicine has been taken. If possible, you should drink a glass of water to ensure that all the medicine has been taken.



If necessary the dispersible tablet can be divided along the break-marks. Hold the tablet between the thumb and the index finger on either side of the score line, with the score line facing upwards. Separate into halves by breaking the tablet along the break-marks (see figure below).



#### **If you take more Tracleer than you should**

If you take more tablets than you have been told to take, contact your doctor immediately.

#### **If you forget to take Tracleer**

If you forget to take Tracleer, take a dose as soon as you remember, then continue to take your tablets at the usual times. Do not take a double dose to make up for forgotten tablets.

#### **If you stop taking Tracleer**

Suddenly stopping your treatment with Tracleer may lead to your symptoms getting worse. Do not stop taking Tracleer unless your doctor tells you to. Your doctor may tell you to reduce the dose over a few days before stopping completely.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The most serious side effects with Tracleer are

- Abnormal liver function which may affect more than 1 in 10 people
- Anaemia (low blood value) which may affect up to 1 in 10 people. Anaemia may occasionally require blood transfusion

Your liver and blood values will be monitored during treatment with Tracleer (see section 2). It is important that you have these tests as ordered by your doctor.

Signs that your liver may not be working properly include:

- nausea (urge to vomit)
- vomiting
- fever (high temperature)
- pain in your stomach (abdomen)
- jaundice (yellowing of your skin or the whites of your eyes)
- dark-coloured urine
- itching of your skin
- lethargy or fatigue (unusual tiredness or exhaustion)
- flu-like syndrome (joint and muscle pain with fever)

If you notice any of these signs **tell your doctor immediately**.

Other side effects:

**Very common** (may affect **more than one in 10** people):

- Headache
- Oedema (swelling of the legs and ankles or other signs of fluid retention)

**Common** (may affect **up to one in 10** people):

- Flushed appearance or redness of skin
- Hypersensitivity reactions (including skin inflammation, itching and rash)
- Gastrooesophageal reflux disease (acid reflux)
- Diarrhoea
- Syncope (fainting)
- Palpitations (fast or irregular heart beats)
- Low blood pressure
- Nasal congestion

**Uncommon** (may affect **up to one in 100** people):

- Thrombocytopenia (low number of blood platelets)
- Neutropenia/leukopenia (low number of white blood cells)
- Elevated liver function tests with hepatitis (inflammation of the liver) including possible exacerbation of underlying hepatitis and/or jaundice (yellowing of the skin or the whites of the eyes)

**Rare** (may affect **up to one in 1000** people):

- Anaphylaxis (general allergic reaction), angioedema (swelling, most commonly around the eyes, lips, tongue or throat)
- Cirrhosis (scarring) of the liver, liver failure (serious disturbance of liver function)

Blurred vision have also been reported at an unknown frequency (frequency cannot be estimated from the available data).

### **Side effects in children and adolescents**

The side effects that have been reported in children treated with Tracleer are the same as those in adults.

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Tracleer**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and on the blister after “EXP”.

Do not store above 25 °C.

Remaining parts of a divided dispersible tablet can be stored at room temperature and should be used within 7 days.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away any medicines you no longer use. These measures will help to protect the environment.

## **6. Contents of the pack and other information**

### **What Tracleer contains**

- The active substance is bosentan as monohydrate. Each dispersible tablet contains 32 mg of bosentan (as monohydrate).
- The other ingredients are cellulose microcrystalline, calcium hydrogen phosphate anhydrous, croscarmellose sodium, silica colloidal anhydrous, tartaric acid, tutti frutti flavour, aspartame (E951, please read further information at the end of section 2), acesulfame potassium, magnesium stearate.

### **What Tracleer looks like and contents of the pack**

Tracleer 32 mg dispersible tablets are pale yellow to off-white, clover-shape dispersible tablets, quadrisectioned on one side and debossed with “32” on the other side.

Peel-push blisters containing 14 dispersible tablets; cartons contain 56 dispersible tablets.

### **Marketing authorisation holder:**

Actelion Registration Ltd  
Chiswick Tower, 13<sup>th</sup> Floor  
389 Chiswick High Road  
London W4 4AL  
United Kingdom

### **Manufacturer:**

Actelion Manufacturing GmbH  
Emil-Barell-Strasse 7  
79639 Grenzach-Wyhlen  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

### **België/Belgique/Belgien**

Actelion Pharmaceuticals Belgium N.V.  
Tél/Tel: +32-(0)15 284 777

### **Lietuva**

UAB ALGOL PHARMA  
Tel: +370 37 40 86 81

### **България**

Аквахим АД  
Тел.: +359 2 807 50 00

### **Luxembourg/Luxemburg**

Actelion Pharmaceuticals Belgium N.V.  
Tél/Tel: +32-(0)15 284 777

### **Česká republika**

Actelion Pharmaceuticals CZ, s.r.o.  
Tel: +420 221 968 006

### **Magyarország**

Actelion Pharmaceuticals Hungaria Kft.  
Tel: +36-1-413-3270

**Danmark**

Actelion Danmark,  
Filial af Actelion Pharmaceuticals Sverige AB,  
Sverige  
Tlf: +45 3694 45 95

**Deutschland**

Actelion Pharmaceuticals Deutschland GmbH  
Tel: +49 761 45 64 0

**Eesti**

Algol Pharma OÜ  
Tel: +372 605 6014

**Ελλάδα**

Actelion Pharmaceuticals Ελλάς A.E.  
Τηλ: +30 210 675 25 00

**España**

Actelion Pharmaceuticals España S.L.  
Tel: +34 93 366 43 99

**France**

Actelion Pharmaceuticals France SAS  
Tél: +33 1 58 62 32 32

**Hrvatska**

Medis Adria d.o.o.  
Tel: +385 (0) 1 2303 446

**Ireland**

Actelion Pharmaceuticals UK Ltd  
Tel: +44 208 987 3333

**Ísland**

Actelion Pharmaceuticals Sverige AB  
Sími: +46 8 544 982 50

**Italia**

Actelion Pharmaceuticals Italia S.r.l.  
Tel: +39 0542 64 87 40

**Κύπρος**

Actelion Pharmaceuticals Ελλάς A.E.  
Τηλ: +30 210 675 25 00

**Latvija**

Algol Pharma SIA  
Tel: +371 6761 9365

**Malta**

Actelion Pharmaceuticals UK Ltd  
Tel: +44 208 987 3333

**Nederland**

Actelion Pharmaceuticals Nederland B.V.  
Tel: +31 (0)348 435950

**Norge**

Actelion Pharmaceuticals Sverige AB,  
Filial Norge  
Tlf: +47 22480370

**Österreich**

Actelion Pharmaceuticals Austria GmbH  
Tel: +43 1 505 4527

**Polska**

Actelion Pharma Polska Sp. z o.o.  
Tel: +48 (22) 262 31 00

**Portugal**

Actelion Pharmaceuticals Portugal Lda.  
Tel: +351 21 358 6120

**România**

Geneva Romfarm International SRL  
Tel: +40 (021) 231 3561

**Slovenija**

Medis d.o.o.  
Tel: +386-(0)1 589 69 00

**Slovenská republika**

Actelion Pharmaceuticals SK, s.r.o.  
Tel: +420 221 968 006

**Suomi/Finland**

Actelion Pharmaceuticals Sverige AB,  
Filial Finland  
Puh/Tel: +358 9 2510 7720

**Sverige**

Actelion Pharmaceuticals Sverige AB  
Tel: +46 8 544 982 50

**United Kingdom**

Actelion Pharmaceuticals UK Ltd  
Tel: +44 208 987 3333

**This leaflet was last revised in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu/>.