

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

ZOSTAVAX powder and solvent for suspension for injection
ZOSTAVAX powder and solvent for suspension for injection in a pre-filled syringe

shingles (herpes zoster) vaccine (live)

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

After reconstitution, 1 dose (0.65 mL) contains:

Varicella-zoster virus¹, Oka/Merck strain, (live, attenuated) not less than 19,400 PFU²

¹produced in human diploid (MRC-5) cells

²PFU = Plaque-forming units

This vaccine may contain traces of neomycin. See sections 4.3 and 4.4.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Powder and solvent for suspension for injection.

The powder is a white to off-white compact crystalline plug.
The solvent is a clear, colourless fluid.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

ZOSTAVAX is indicated for prevention of herpes zoster (“zoster” or shingles) and herpes zoster-related post-herpetic neuralgia (PHN).

ZOSTAVAX is indicated for immunization of individuals 50 years of age or older.

4.2 Posology and method of administration

Posology

Individuals should receive a single dose (0.65 mL).

The need for a booster dose is not known. See sections 4.8 and 5.1.

Paediatric population

The safety and efficacy of ZOSTAVAX in children and adolescents have not been established. No data are available.

There is no relevant use of ZOSTAVAX in children and adolescents for prevention of primary varicella infection (chickenpox).

Method of administration

The vaccine can be injected subcutaneously (SC) or intramuscularly (IM), preferably in the deltoid region (see sections 4.8 and 5.1).

The vaccine should be administered subcutaneously in patients with severe thrombocytopenia or any coagulation disorder (see section 4.4).

The vaccine should under no circumstances be injected intravascularly.

For precautions to be taken before handling or administering the medicinal product see section 6.6.

For instructions on reconstitution of the medicinal product before administration see section 6.6.

4.3 Contraindications

- History of hypersensitivity to the active substance, to any of the excipients or trace residuals (e.g. neomycin) (see sections 4.4 and 6.1).
- Primary and acquired immunodeficiency states due to conditions such as: acute and chronic leukaemias; lymphoma; other conditions affecting the bone marrow or lymphatic system; immunosuppression due to HIV/AIDS (see sections 4.4, 4.8 and 5.1); cellular immune deficiencies.
- Immunosuppressive therapy (including high-dose corticosteroids) (see sections 4.4 and 4.8); however, ZOSTAVAX is not contraindicated for use in individuals who are receiving topical/inhaled corticosteroids or low-dose systemic corticosteroids or in patients who are receiving corticosteroids as replacement therapy, e.g., for adrenal insufficiency (see sections 4.8 and 5.1).
- Active untreated tuberculosis.
- Pregnancy. Furthermore, pregnancy should be avoided for 1 month following vaccination (see section 4.6).

4.4 Special warnings and precautions for use

Appropriate medical treatment and supervision should always be readily available in case of a rare anaphylactic/anaphylactoid reaction following the administration of the vaccine, as there is a possibility of hypersensitivity reactions, not only to the active substances, but also to the excipients and trace residuals (e.g. neomycin) present in the vaccine (see sections 4.3, 4.8 and 6.1).

Neomycin allergy generally manifests as a contact dermatitis. However, a history of contact dermatitis due to neomycin is not a contraindication to receiving live virus vaccines.

ZOSTAVAX is a live, attenuated varicella-zoster vaccine and administration may result in disseminated disease in individuals who are immunosuppressed or immunodeficient. Patients who previously received immune suppressive therapy should be carefully evaluated for the reconstitution of the immune system prior to receiving Zostavax (see section 4.3).

The safety and efficacy of ZOSTAVAX have not been established in adults who are known to be infected with HIV with or without evidence of immunosuppression (see section 4.3) however, a phase II safety and immunogenicity study in HIV-infected adults with conserved immune function (CD 4+T cell count ≥ 200 cells/ μ L) has been completed (see sections 4.8 and 5.1).

This vaccine should be given subcutaneously to individuals with severe thrombocytopenia or any coagulation disorder, because these individuals may bleed following intramuscular injections.

ZOSTAVAX is not indicated for treatment of zoster or PHN.

Immunisation should be postponed in individuals suffering from moderate to severe acute febrile illness or infection.

As for any vaccine, vaccination with ZOSTAVAX may not result in protection in all vaccine recipients. See section 5.1.

Transmission

In clinical trials with ZOSTAVAX, transmission of the vaccine virus has not been reported. However, post-marketing experience with varicella vaccines suggests that transmission of vaccine virus may occur rarely between vaccinees who develop a varicella-like rash and susceptible contacts [for example, varicella-zoster virus (VZV) susceptible infant grandchildren]. Transmission of vaccine virus from varicella vaccine recipients who do not develop a varicella-like rash has also been reported. This is a theoretical risk for vaccination with ZOSTAVAX. The risk of transmitting the attenuated vaccine virus from a vaccinee to a susceptible contact should be weighed against the risk of developing natural zoster and potentially transmitting wild-type VZV to a susceptible contact.

4.5 Interaction with other medicinal products and other forms of interaction

ZOSTAVAX can be administered concomitantly with inactivated influenza vaccine as separate injections and at different body sites (see section 5.1).

ZOSTAVAX and 23-valent pneumococcal polysaccharide vaccine should not be given concomitantly because concomitant use in a clinical trial resulted in reduced immunogenicity of ZOSTAVAX (see section 5.1). Therefore, administration of the two vaccines should be considered to be separated by at least 4 weeks.

No data are currently available regarding concomitant use with other vaccines.

Concurrent administration of ZOSTAVAX and anti-viral medications known to be effective against VZV has not been evaluated.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no data on the use of ZOSTAVAX in pregnant women. Traditional non-clinical studies are insufficient with respect to reproductive toxicity (see section 5.3). However naturally-occurring varicella-zoster virus infection is known to sometimes cause foetal harm. ZOSTAVAX is not recommended to be administered to pregnant women. In any case, pregnancy should be avoided for one month following vaccination (see section 4.3).

Breast-feeding

It is unknown whether VZV is secreted in human milk. A risk to the newborns/infants cannot be excluded. A decision must be made whether to discontinue breast-feeding or to not administer ZOSTAVAX taking into account the benefit of breast-feeding for the child and the benefit of vaccination for the woman.

Fertility

ZOSTAVAX has not been evaluated in fertility studies.

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive or use machines have been performed. However, ZOSTAVAX is expected to have no or negligible influence on ability to drive and use machines.

4.8 Undesirable effects

a. Summary of the safety profile

The most common adverse reactions reported in pivotal clinical trials were injection-site reactions. Headache and pain in the extremity were the most common systemic adverse reactions. Most of these local and systemic adverse reactions were reported as mild in intensity. Vaccine-related serious adverse reactions were reported for 0.01% subjects vaccinated with ZOSTAVAX and subjects who received placebo.

Data from a clinical trial (n=368) demonstrated that the current refrigerated formulation has a safety profile comparable to that of the frozen formulation.

b. Tabulated summary of adverse events

In clinical trials, general safety has been evaluated in more than 57,000 adults vaccinated with ZOSTAVAX.

Table 1 presents vaccine-related injection-site and systemic adverse reactions reported at a significantly greater incidence in the vaccine group versus the placebo group within 42 days postvaccination in the ZOSTAVAX Efficacy and Safety trial (ZEST) study and in the Adverse Event Monitoring Substudy of Shingles Prevention Study (SPS).

Additional adverse reactions, spontaneously reported through post-marketing surveillance, are also included in Table 1. As these events are reported voluntarily from a population of uncertain size, it is not possible to reliably calculate their frequency or establish a causal relationship to vaccine exposure. Consequently, the frequencies of these adverse reactions have been estimated based on the adverse events reported in SPS and ZEST (regardless of vaccine relationship assigned by the investigator).

The adverse reactions are assigned frequency categories using the following convention:

Very Common ($\geq 1/10$);

Common ($\geq 1/100$ to $< 1/10$);

Uncommon ($\geq 1/1,000$ to $< 1/100$);

Rare ($\geq 1/10,000$ to $< 1/1,000$);

Very rare ($< 1/10,000$)

Table 1: Adverse Reactions from Clinical Trial Experience and Post-Marketing Surveillance

| MedDRA System Organ Class | Adverse reaction terms | Frequency |
|--|--|-------------|
| Infections and infestations | Varicella, Herpes zoster (vaccine strain) | Very rare |
| Blood and lymphatic system disorders | Lymphadenopathy (cervical, axillary) | Uncommon |
| Immune system disorders | Hypersensitivity reactions including anaphylactic reactions | Rare |
| Nervous system disorders | Headache ¹ | Common |
| Eye Disorders | Necrotizing retinitis (patients on immunosuppressive therapy) | Very rare |
| Gastrointestinal disorders | Nausea | Uncommon |
| Skin and subcutaneous tissue disorders | Rash | Common |
| Musculoskeletal and connective tissue disorders | Arthralgia, Myalgia, Pain in extremity ¹ | Common |
| General disorders and administration site conditions | Injection site: Erythema ^{1,2} , Pain/tenderness ^{1,2} , Pruritus ^{1,2} , Swelling ^{1,2} | Very common |
| | Injection site: Induration ¹ , Haematoma ¹ , Warmth ¹ , Rash, Pyrexia | Common |
| | Injection site urticaria | Rare |

¹ Clinical trials experience.

² Solicited adverse reaction within 5 days postvaccination.

c. Description of selected adverse reactions

Injection site reactions

Vaccine-related injection-site adverse reactions were significantly greater for subjects vaccinated with ZOSTAVAX versus subjects who received placebo. In SPS, the overall incidence of vaccine-related injection-site adverse reactions were 48% for ZOSTAVAX and 17% for placebo in subjects 60 years of age and older.

In ZEST, the overall incidence of vaccine-related injection site adverse reactions were 63.9% for ZOSTAVAX and 14.4% for placebo in subjects 50 to 59 years of age. Most of these adverse reactions were reported as mild in intensity.

In other clinical trials evaluating ZOSTAVAX in subjects 50 years of age or older, including a study of concomitantly administered inactivated influenza vaccine, a higher rate of injection-site adverse experiences of mild-to-moderate intensity was reported among subjects 50-59 years of age compared with subjects ≥ 60 years of age (see section 5.1).

ZOSTAVAX was administered either subcutaneously (SC) or intramuscularly (IM) in subjects 50 years of age or older (see section 5.1). The general safety profiles of the SC and IM routes were otherwise comparable, but injection-site adverse reactions were significantly less frequent in the IM group (34%) compared with the SC group (64%).

Herpes zoster/herpes zoster-like rashes and Varicella/varicella-like rashes in clinical trials

In clinical trials the number of herpes zoster/herpes zoster-like rashes within the 42-day postvaccination was low in both ZOSTAVAX and placebo groups. The majority of rashes have been rated as mild to moderate; no complications from rash have been observed in the clinical setting. Most of the reporting rashes that were VZV positive by PCR analysis were associated with wild-type VZV.

In SPS and ZEST, the number of subjects who reported herpes zoster/herpes zoster-like rashes was less than 0.2% for ZOSTAVAX and placebo groups, with no significant difference observed between the two groups. The number of subjects who reported varicella/varicella-like rashes was less than 0.7% for ZOSTAVAX and placebo.

The Oka/Merck strain of VZV was not detected from any specimens in SPS or ZEST. VZV was detected in one (0.01%) specimen from a ZOSTAVAX recipient reporting a varicella/varicella-like rash; however, the virus strain (wild type or Oka/Merck strain) could not be determined. Across all other clinical trials, the Oka/Merck strain was identified by PCR analysis from the lesion specimens of only two subjects who reported varicella-like rashes (onset on Day 8 and 17).

d. Special populations

Adults with a history of herpes zoster (HZ) prior to vaccination

ZOSTAVAX was administered to subjects 50 years of age or older with a history of herpes zoster (HZ) prior to vaccination (see section 5.1). The safety profile was generally similar to that seen in the Adverse Event Monitoring Substudy of the SPS.

Adults on chronic/maintenance systemic corticosteroids

In subjects 60 years of age or older who were receiving chronic/maintenance systemic corticosteroid therapy at a daily dose equivalent of 5 to 20 mg of prednisone for at least 2 weeks prior to enrollment, and 6 weeks or more following vaccination, the safety profile was generally comparable to that seen in the Adverse Event Monitoring Substudy of the SPS (see sections 4.3 and 5.1).

HIV-infected adults with conserved immune function

In a clinical trial, ZOSTAVAX was administered to HIV-infected adults (18 years of age or older, CD4+ T cell count \geq 200 cells/ μ L) (see section 5.1). The safety profile was generally similar to the Adverse Event Monitoring Substudy of the SPS. Adverse events were followed up to Day 42 post vaccination and serious adverse events throughout the entire study period (i.e. through Day 180). Of the 295 ZOSTAVAX recipients, one case of serious vaccine related maculo-papular rash was reported on Day 4 following Dose 1 of ZOSTAVAX (see section 4.3).

VZV-seronegative adults

Based on limited data from 2 clinical trials that enrolled VZV-seronegative or low seropositive subjects (30 years of age or older) receiving live attenuated zoster vaccine, injection site and systemic adverse experiences were generally similar to those reported by other subjects who received ZOSTAVAX in clinical trials, with 2 of the 27 subjects reporting fever. No subjects reported varicella-like or herpes zoster-like rashes. No serious vaccine-related adverse experiences were reported.

e. Other studies

Adults receiving additional doses/revaccination

In a clinical study, adults 60 years of age or older received a second dose of ZOSTAVAX 42 days following the initial dose (see section 5.1). The frequency of vaccine-related adverse experiences after the second dose of ZOSTAVAX was generally similar to that seen with the first dose.

In another study, ZOSTAVAX was administered as a booster dose to HZ history-negative subjects 70 years of age or older who had received a first dose approximately 10 years previously, and as a first dose to HZ history-negative subjects 70 years of age or older (see section 5.1). The frequency of vaccine-related adverse experiences after the booster dose of ZOSTAVAX was generally similar to that seen with the first dose.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare

professionals are asked to report any suspected adverse reactions via [the national reporting system listed in Appendix V](#).

4.9 Overdose

Administration of a higher than recommended dose of ZOSTAVAX was reported rarely and the adverse reaction profile was comparable to that observed with the recommended dose of ZOSTAVAX.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Vaccines, Viral Vaccine; ATC code: J07BK02

Mechanism of action

Anyone who has been infected with VZV, including those without a clinical history of varicella, is at risk for developing zoster. This risk appears to be causally related to a decline in VZV-specific immunity. ZOSTAVAX was shown to boost VZV-specific immunity, which is thought to be the mechanism by which it protects against zoster and its complications (See Immunogenicity).

Clinical Efficacy

The protective clinical efficacy of ZOSTAVAX was demonstrated in two large, randomised, placebo controlled clinical trials where subjects received ZOSTAVAX subcutaneously (see Tables 2 and 3).

ZOSTAVAX Efficacy and Safety Trial (ZEST) in subjects 50 to 59 years of age:

The ZEST study was a placebo-controlled, double-blind clinical trial in which 22,439 subjects were randomised to receive a single dose of either ZOSTAVAX or placebo and were followed for the development of zoster for a median of 1.3 years (range 0 to 2 years). Final determination of zoster cases was made by Polymerase Chain Reaction (PCR) [86%], or in the absence of virus detection, as determined by a clinical evaluation committee [14%]. ZOSTAVAX significantly decreased the incidence of zoster compared to placebo (see Table 2).

Table 2: Efficacy of ZOSTAVAX on zoster incidence compared with placebo in the ZEST trial in subjects 50 to 59 years of age*

| ZOSTAVAX | | | Placebo | | | Vaccine efficacy (95% CI) |
|--------------------|------------------------|---|--------------------|------------------------|---|------------------------------|
| Number of subjects | Number of zoster cases | Incidence rate of zoster per 1,000 person years | Number of subjects | Number of zoster cases | Incidence rate of zoster per 1,000 person years | |
| 11,211 | 30 | 2.0 | 11,228 | 99 | 6.6 | 70% (54%, 81%) |

*The analysis was performed on the intent-to-treat (ITT) population that included all subjects randomized in the ZEST study

Shingles Prevention Study (SPS) in Subjects 60 years of age and older:

The SPS study was a placebo-controlled, double-blind clinical trial in which 38,546 subjects were randomised to receive a single dose of either ZOSTAVAX or placebo and were followed for the development of zoster for a median of 3.1 years (range 31 days to 4.9 years).

ZOSTAVAX significantly decreased the incidence of zoster compared with placebo (see Table 3).

Table 3: Efficacy of ZOSTAVAX on zoster incidence compared with placebo in the SPS in subjects 60 years of age and older*

| <i>Age group</i> [†] | ZOSTAVAX | | | Placebo | | | Vaccine efficacy (95% CI) |
|-------------------------------|--------------------|------------------------|---|--------------------|------------------------|---|----------------------------------|
| | Number of subjects | Number of zoster cases | Incidence rate of zoster per 1,000 person years | Number of subjects | Number of zoster cases | Incidence rate of zoster per 1,000 person years | |
| ≥ 60 | 19,254 | 315 | 5.4 | 19,247 | 642 | 11.1 | 51% (44%, 58%) |
| 60-69 | 10,370 | 122 | 3.9 | 10,356 | 334 | 10.8 | 64% (56%, 71%) |
| ≥ 70 | 8,884 | 193 | 7.2 | 8,891 | 308 | 11.5 | 38% (25%, 48%) |
| 70-79 | 7,621 | 156 | 6.7 | 7,559 | 261 | 11.4 | 41% (28%, 52%) |

* The analysis was performed on the Modified Intent-To-Treat (MITT) population that included all subjects randomized in the study who were followed for at least 30 days postvaccination and did not develop an evaluable case of zoster within the first 30 days post vaccination

† Age strata at randomization were 60-69 and ≥ 70 years of age

In the SPS, the reduction in zoster was seen in almost all dermatomes. Ophthalmic zoster occurred in 35 subjects vaccinated with ZOSTAVAX vs. 69 subjects who received placebo. Impaired vision occurred in 2 subjects vaccinated with ZOSTAVAX vs. 9 who received placebo.

ZOSTAVAX significantly decreased the incidence of Post-herpetic Neuralgia (PHN) compared with placebo (see Table 4). In subjects who developed zoster, ZOSTAVAX decreased the risk of subsequently developing PHN. In the vaccine group, the risk of developing PHN after zoster was 9 % (27/315), while in the placebo group it was 13 % (80/642). This effect was more prominent in the group of older subjects (≥70 years of age), where the risk of developing PHN after zoster was reduced to 10% in the vaccine group vs. 19% for the placebo group.

Table 4: Efficacy of ZOSTAVAX on PHN[†] incidence compared with placebo in the SPS in subjects 60 years of age and older*

| Age group [‡] | ZOSTAVAX | | | Placebo | | | Vaccine efficacy (95% CI) |
|------------------------|--------------------|---------------------|--|--------------------|---------------------|--|--------------------------------|
| | Number of subjects | Number of PHN cases | Incidence rate of PHN per 1,000 person years | Number of subjects | Number of PHN cases | Incidence rate of PHN per 1,000 person years | |
| ≥ 60 | 19,254 | 27 | 0.5 | 19,247 | 80 | 1.4 | 67% [§] (48%, 79%) |
| 60-69 | 10,370 | 8 | 0.3 | 10,356 | 23 | 0.7 | 66% (20%, 87%) |
| ≥ 70 | 8,884 | 19 | 0.7 | 8,891 | 57 | 2.1 | 67% (43%, 81%) |
| 70-79 | 7,621 | 12 | 0.5 | 7,559 | 45 | 2.0 | 74% (49%, 87%) |

[†] PHN was defined as zoster-associated pain rated as ≥3 (on a 0-10 scale), persisting or appearing more than 90 days after onset of zoster rash using Zoster Brief Pain Inventory (ZBPI).

* The table is based on the Modified Intent-To-Treat (MITT) population that included all subjects randomized in the study who were followed for at least 30 days postvaccination and did not develop an evaluable case of zoster within the first 30 days postvaccination.

[‡] Age strata at randomization were 60-69 and ≥70 years of age.

[§] Age-adjusted estimate based on the age strata (60-69 and ≥70 years of age) at randomization.

ZOSTAVAX significantly reduced the zoster pain Burden of Illness (BOI) score (see Table 5).

Table 5: Reduction of the zoster-associated pain by the BOI[†] score in the SPS in subjects 60 years of age and older

| Age group [‡] | ZOSTAVAX | | | Placebo | | | Vaccine efficacy (95% CI) |
|------------------------|--------------------|----------------------------------|----------------|--------------------|----------------------------------|----------------|---------------------------|
| | Number of subjects | Number of zoster confirmed cases | Mean BOI score | Number of subjects | Number of zoster confirmed cases | Mean BOI score | |
| ≥ 60 | 19,254 | 315 | 2.21 | 19,247 | 642 | 5.68 | 61% (51%, 69%) |
| 60-69 | 10,370 | 122 | 1.5 | 10,356 | 334 | 4.33 | 66% (52%, 76%) |
| ≥ 70 | 8,884 | 193 | 3.47 | 8,891 | 308 | 7.78 | 55% (40%, 67%) |
| 70-79 | 7,621 | 156 | 3.04 | 7,559 | 261 | 7.43 | 59% (43%, 71%) |

[†] The zoster pain BOI score is a composite score that incorporates the incidence, severity, and duration of acute and chronic zoster-associated pain over a 6 month follow-up period.

[‡] Age strata at randomization were 60-69 and ≥70 years of age.

Prevention of HZ cases with severe pain in the entire SPS study population

ZOSTAVAX reduced the incidence of zoster with severe and long-lasting pain (severity-by-duration score >600) by 73% (95% CI: [46 to 87%]) compared with placebo (11 vs. 40 cases, respectively).

Reduction of zoster pain severity-by-duration in vaccinated individuals who developed zoster

With regard to the acute pain (pain between 0-30 days) there was no statistically significant difference between the vaccine group and the placebo group.

However, among vaccinated individuals who developed PHN, ZOSTAVAX significantly reduced PHN-associated (chronic) pain compared with placebo. In the period from 90 days after rash onset to the end of follow-up, there was a 57% reduction in the severity-by-duration score (average scores of 347 for ZOSTAVAX and 805 for placebo; p=0.016).

Overall, among vaccinated individuals who developed zoster, ZOSTAVAX significantly reduced overall acute and chronic zoster-associated pain compared with placebo. Over the 6-month (acute and chronic) follow-up period, there was a 22% reduction ($p = 0.008$) in the severity-by-duration score and a 52% (95% CI: [7 to 74%]) reduction (from 6.2% to 3.5%) in the risk of having zoster with severe and long-lasting pain (severity-by-duration score of >600).

Zostavax persistence of protection

The persistence of protection following vaccination has been evaluated through longer-term follow-up in Short-term Persistence Substudy (STPS) and Long-term Persistence Substudy (LTPS) and supports the continued benefit of ZOSTAVAX throughout the follow-up periods studied. The STPS was initiated to accrue additional information on the persistence of vaccine efficacy for subjects who received ZOSTAVAX in SPS.

Persistence of ZOSTAVAX efficacy was studied 4 to 7 years postvaccination in the STPS, which included 7,320 subjects previously vaccinated with ZOSTAVAX and 6,950 subjects previously vaccinated with placebo in the SPS (mean age at enrollment was 73.3 years); and 7 to 10 years postvaccination in the Long-term Persistence Substudy (LTPS), which included 6,867 subjects previously vaccinated with ZOSTAVAX (mean age at enrollment into the LTPS was 74.5 years). The median follow-up was ~ 1.2 years (range is one day to 2.2 years) and ~ 3.9 years (range is one week to 4.75 years) in STPS and LTPS, respectively. During the course of the STPS, placebo recipients were offered ZOSTAVAX, at which time they were considered to have completed the STPS. A concurrent placebo control was not available in the LTPS; data from prior placebo recipients were used to estimate vaccine efficacy.

In the STPS, there were 84 evaluable zoster cases [8.4/1,000 person-years] in the ZOSTAVAX group and 95 evaluable cases [14.0/1,000 person-years] in the placebo group. The estimated vaccine efficacy during the STPS follow-up period was 40% (95% CI: [18 to 56%]) for zoster incidence, 60% (95% CI: [-10 to 87%]) for PHN incidence and 50% (95% CI: [14 to 71%]) for zoster BOI.

In the LTPS, there were 263 evaluable zoster cases reported among 261 patients [10.3/1000 person-years]. The estimated vaccine efficacy during the LTPS follow-up period was 21% (95% CI: [11 to 30%]) for zoster incidence, 35% (95% CI: [9 to 56%]) for PHN incidence and 37% (95% CI: [27 to 46%]) for zoster BOI.

Immunogenicity of ZOSTAVAX

Shingles Prevention Study (SPS)

Within SPS, immune responses to vaccination were evaluated in a subset of the enrolled subjects ($N=1395$). ZOSTAVAX elicited significantly higher VZV-specific immune responses at 6 weeks postvaccination compared with placebo.

ZOSTAVAX Efficacy and Safety Trial (ZEST)

Within ZEST, immune responses to vaccination were evaluated in a random 10% subcohort ($n=1,136$ for ZOSTAVAX and $n=1,133$ for placebo) of the subjects enrolled in the ZEST. ZOSTAVAX elicited significantly higher VZV-specific immune responses at 6 weeks postvaccination compared with placebo.

When evaluated at 4 weeks postvaccination, the immunogenicity of the current refrigerator-stable formulation was shown to be similar to the immunogenicity of the earlier frozen formulation of ZOSTAVAX.

Subjects who received ZOSTAVAX by SC (subcutaneous) or IM (intramuscular) route

In an open-label, randomised, controlled clinical trial, ZOSTAVAX was administered either by SC route or by IM route to 353 subjects 50 years of age or older. Subjects with severe thrombocytopenia or any other coagulation disorder were excluded. The VZV specific immune responses to ZOSTAVAX at Week 4 post-vaccination were comparable whether administered by SC or IM route.

Concomitant administration

In a double-blind, controlled clinical trial, 762 adults 50 years of age and older were randomised to receive a single dose of ZOSTAVAX administered either concomitantly (N=382) or nonconcomitantly (N=380) with inactivated split influenza vaccine. The VZV-specific immune responses to both vaccines at 4 weeks postvaccination were similar, whether administered concomitantly or nonconcomitantly.

In a double-blind, controlled clinical trial, 473 adults, 60 years of age or older, were randomised to receive a single dose of ZOSTAVAX either concomitantly (N=237), or nonconcomitantly with 23-valent pneumococcal polysaccharide vaccine (N=236). At four weeks postvaccination, the VZV-specific immune responses following concomitant use were not similar to the VZV-specific immune responses following nonconcomitant administration. Therefore consider administration of the two vaccines separated by at least 4 weeks.

Subjects with a history of herpes zoster (HZ) prior to vaccination

In a double-blind, placebo-controlled, randomised clinical trial, ZOSTAVAX was administered to 100 subjects 50 years of age or older with a history of herpes zoster prior to vaccination to assess immunogenicity and safety (see section 4.8) of ZOSTAVAX. ZOSTAVAX induced a significantly higher VZV-specific immune response at 4 weeks postvaccination, compared with placebo. VZV-specific immune responses were generally similar in subjects 50 to 59 compared to subjects ≥ 60 years of age.

Adults receiving additional doses/revaccination

The need for, or timing of, a booster dose with ZOSTAVAX has not yet been determined. In an open-label study, ZOSTAVAX was administered as: (1) a booster dose to 201 zoster history-negative subjects 70 years of age or older who had received a first dose approximately 10 years previously as participants in the SPS, and (2) a first dose to 199 zoster history-negative subjects 70 years of age or older. The VZV-specific immune responses to vaccine 6 weeks postvaccination was comparable in the booster dose and first dose group.

Subjects on chronic/maintenance systemic corticosteroids

In a double-blind, placebo-controlled, randomised clinical trial, ZOSTAVAX was administered to 206 subjects 60 years of age or older who were receiving chronic/maintenance systemic corticosteroid therapy at a daily dose equivalent of 5 to 20 mg of prednisone for at least 2 weeks prior to enrollment, and 6 weeks or more following vaccination to assess the immunogenicity and safety profile of ZOSTAVAX. Compared with placebo, ZOSTAVAX induced a higher VZV-specific immune response at 6 weeks postvaccination.

HIV-infected adults with conserved immune function

In a double-blind, placebo-controlled randomised clinical trial, ZOSTAVAX was administered to HIV-infected adults (18 years of age or older; median age 49 years) on appropriate antiretroviral therapy with conserved immune function (CD4+ T cell count ≥ 200 cells/ μ L). Although, ZOSTAVAX is indicated as a single dose regimen (see section 4.2), a two-dose regimen was used. 286 subjects received two doses and 9 subjects received only one dose. The VZV-specific immune responses following Doses 1 and 2 were similar (see section 4.3).

Immunocompromised subjects

The vaccine has not been studied in subjects with impaired immunity.

The European Medicines Agency has waived the obligation to submit the results of studies with ZOSTAVAX in all the subsets of paediatric population (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

Not applicable.

5.3 Preclinical safety data

Traditional non-clinical studies were not performed, but there are no non-clinical concerns considered relevant to clinical safety beyond data included in other sections of the Summary of Product Characteristics (SmPC).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Powder:

Sucrose
Hydrolysed gelatin
Sodium chloride
Potassium dihydrogen phosphate
Potassium chloride
Monosodium L-glutamate monohydrate
Disodium phosphate
Sodium hydroxide (to adjust pH)
Urea

Solvent:

Water for injections

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

18 months.

After reconstitution, the vaccine should be used immediately. However, in-use stability has been demonstrated for 30 minutes when stored at 20 °C – 25 °C.

6.4 Special precautions for storage

Store and transport refrigerated (2 °C – 8 °C).
Do not freeze.
Store in the original package in order to protect from light.

For storage conditions after reconstitution of the medicinal product, see section 6.3.

6.5 Nature and contents of container

ZOSTAVAX with solvent for reconstitution supplied in a vial:

Powder in a vial (glass) with a stopper (butyl rubber) and flip off cap (aluminium) and solvent in a vial (glass) with a stopper (chlorobutyl rubber) and flip off cap (aluminium) in a pack size of 1 or 10.

ZOSTAVAX with solvent for reconstitution supplied in a pre-filled syringe:

Powder in a vial (glass) with a stopper (butyl rubber) and flip off cap (aluminium) and solvent in a pre-filled syringe (glass) with plunger stopper (chlorobutyl rubber) and tip cap (styrene-butadiene rubber) with one or two unattached needles in a pack size of 1, 10 or 20.

Powder in a vial (glass) with a stopper (butyl rubber) and flip off cap (aluminium) and solvent in a pre-filled syringe (glass) with plunger stopper (chlorobutyl rubber) and tip cap (styrene-butadiene rubber) without needle in pack size of 1, 10 or 20.

Powder in a vial (glass) with a stopper (butyl rubber) and flip off cap (aluminium) and solvent in a pre-filled syringe (glass) with plunger stopper (chlorobutyl rubber) and needle shield (natural rubber), in a pack size of 1 or 10.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Avoid contact with disinfectants as they may inactivate the vaccine virus.

To reconstitute the vaccine, use the solvent provided. When reconstituted, ZOSTAVAX is a semi-hazy to translucent, off-white to pale yellow liquid.

It is important to use a separate sterile syringe and needle for each patient to prevent transmission of infectious agents from one individual to another.

Reconstitution instructions

ZOSTAVAX with solvent for reconstitution supplied in a vial:

Withdraw the entire contents of the solvent vial into a syringe.

Inject all the solvent into the vial of lyophilised vaccine.

Gently agitate to dissolve completely.

Withdraw the entire content of the reconstituted vaccine using the same syringe. Inject the vaccine.

The reconstituted vaccine should be inspected visually for any foreign particulate matter and/or abnormal physical appearance prior to administration. In the event of either being observed, discard the vaccine.

ZOSTAVAX with solvent for reconstitution supplied in a pre-filled syringe:

If two needles are provided, separate needles should be used for the reconstitution and administration of the vaccine.

To reconstitute the vaccine, inject all the solvent in the pre-filled syringe into the vial of lyophilised vaccine and gently agitate to mix thoroughly.

Withdraw the entire contents of the reconstituted vaccine using the same syringe. Inject the vaccine.

One or 2 separate needles may be available in the secondary packaging of the presentation containing the pre-filled syringe without attached needle.

The needle should be pushed into the extremity of the syringe and rotated a quarter of a turn (90°) to secure the connection.

The reconstituted vaccine should be inspected visually for any foreign particulate matter and/or abnormal physical appearance prior to administration. In the event of either being observed, discard the vaccine.

It is recommended that the vaccine be administered immediately after reconstitution, to minimize loss of potency. Discard reconstituted vaccine if it is not used within 30 minutes.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

MSD VACCINS
162 avenue Jean Jaurès
69007 Lyon
France

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/06/341/001
EU/1/06/341/002
EU/1/06/341/003
EU/1/06/341/004
EU/1/06/341/005
EU/1/06/341/006
EU/1/06/341/007
EU/1/06/341/008
EU/1/06/341/009
EU/1/06/341/010
EU/1/06/341/011
EU/1/06/341/012
EU/1/06/341/013

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 19 May 2006
Date of latest renewal: 24 March 2011

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

ANNEX II

- A. MANUFACTURERS OF THE BIOLOGICAL ACTIVE SUBSTANCES AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURERS OF THE BIOLOGICAL ACTIVE SUBSTANCES AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturers of the biological active substances

Merck Sharp & Dohme Corp.
Sumneytown Pike
West Point
Pennsylvania 19486
U.S.A.

Merck Sharp & Dohme Corp.
5325 Old Oxford Road
Durham
North Carolina 27712
U.S.A

Name and address of the manufacturer responsible for batch release

Merck Sharp & Dohme B.V.
Waarderweg 39
2031 BN Haarlem
The Netherlands

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to medical prescription.

- **Official batch release**

In accordance with Article 114 of Directive 2001/83/EC, the official batch release will be undertaken by a state laboratory or a laboratory designated for that purpose.

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

- **Periodic Safety Update Reports**

The requirements for submission of periodic safety update reports for this medicinal products are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

- **Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;

- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

ZOSTAVAX - Powder in vial and solvent in vial - Pack of 1, 10

1. NAME OF THE MEDICINAL PRODUCT

ZOSTAVAX powder and solvent for suspension for injection
shingles (herpes zoster) vaccine (live)

2. STATEMENT OF ACTIVE SUBSTANCE(S)

After reconstitution, 1 dose (0.65 mL) contains:
Varicella-zoster virus, Oka/Merck strain, (live, attenuated) ≥ 19400 PFU

3. LIST OF EXCIPIENTS

Sucrose, hydrolysed gelatin, urea, sodium chloride, potassium dihydrogen phosphate, potassium chloride, monosodium L-glutamate monohydrate, disodium phosphate, sodium hydroxide.

4. PHARMACEUTICAL FORM AND CONTENTS

Powder and solvent for suspension for injection
1 vial (powder) + 1 vial (solvent)
10 vials (powder) + 10 vials (solvent)

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Subcutaneous or intramuscular use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store and transport refrigerated. Do not freeze. Keep the vial in the outer carton to protect from light.
After reconstitution, use immediately or within 30 minutes if stored at 20°C-25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

MSD VACCINS
162 avenue Jean Jaurès
69007 Lyon
France

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/06/341/001 – pack of 1
EU/1/06/341/002 – pack of 10

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS
VIAL OF POWDER

1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION

ZOSTAVAX powder for injection
SC/IM

2. METHOD OF ADMINISTRATION

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT

1 dose

6. OTHER

MSD VACCINS

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS
VIAL OF SOLVENT

1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION

Solvent for ZOSTAVAX

2. METHOD OF ADMINISTRATION

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT

1 dose

6. OTHER

MSD VACCINS

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

ZOSTAVAX - Powder in vial and solvent in pre-filled syringe with attached needle Pack of 1, 10

1. NAME OF THE MEDICINAL PRODUCT

ZOSTAVAX powder and solvent for suspension for injection in a pre-filled syringe shingles (herpes zoster) vaccine (live)

2. STATEMENT OF ACTIVE SUBSTANCE(S)

After reconstitution, 1 dose (0.65 mL) contains:
Varicella-zoster virus, Oka/Merck strain, (live, attenuated) $\geq 19,400$ PFU

3. LIST OF EXCIPIENTS

Sucrose, hydrolysed gelatin, urea, sodium chloride, potassium dihydrogen phosphate, potassium chloride, monosodium L-glutamate monohydrate, disodium phosphate, sodium hydroxide.

4. PHARMACEUTICAL FORM AND CONTENTS

Powder and solvent for suspension for injection in a pre-filled syringe
1 vial (powder) + 1 pre-filled syringe with needle (solvent)
10 vials (powder) + 10 pre-filled syringes with needle (solvent)

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Subcutaneous or intramuscular use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store and transport refrigerated. Do not freeze. Keep the vial in the outer carton to protect from light.

After reconstitution, use immediately or within 30 minutes if stored at 20°C-25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

MSD VACCINS
162 avenue Jean Jaurès
69007 Lyon
France

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/06/341/003 – pack of 1
EU/1/06/341/004 – pack of 10

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

ZOSTAVAX - Powder in vial and solvent in pre-filled syringe without needle - Pack of 1, 10, 20

1. NAME OF THE MEDICINAL PRODUCT

ZOSTAVAX powder and solvent for suspension for injection in a pre-filled syringe
shingles (herpes zoster) vaccine (live)

2. STATEMENT OF ACTIVE SUBSTANCE(S)

After reconstitution, 1dose (0.65 mL) contains:
Varicella-zoster virus, Oka/Merck strain, (live, attenuated) $\geq 19,400$ PFU

3. LIST OF EXCIPIENTS

Sucrose, hydrolysed gelatin, urea, sodium chloride, potassium dihydrogen phosphate, potassium chloride, monosodium L-glutamate monohydrate, disodium phosphate, sodium hydroxide.

4. PHARMACEUTICAL FORM AND CONTENTS

Powder and solvent for suspension for injection in a pre-filled syringe
1 vial (powder) + 1 pre-filled syringe without needle (solvent)
10 vials (powder) + 10 pre-filled syringes without needle (solvent)
20 vials (powder) + 20 pre-filled syringes without needle (solvent)

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Subcutaneous or intramuscular use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store and transport refrigerated. Do not freeze. Keep the vial in the outer carton to protect from light.

After reconstitution, use immediately or within 30 minutes if stored at 20°C-25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

MSD VACCINS
162 avenue Jean Jaurès
69007 Lyon
France

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/06/341/005 – pack of 1
EU/1/06/341/006 – pack of 10
EU/1/06/341/007 – pack of 20

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

ZOSTAVAX - Powder in vial and solvent in pre-filled syringe with one unattached needle - Pack of 1, 10, 20

1. NAME OF THE MEDICINAL PRODUCT

ZOSTAVAX powder and solvent for suspension for injection in a pre-filled syringe shingles (herpes zoster) vaccine (live)

2. STATEMENT OF ACTIVE SUBSTANCE(S)

After reconstitution, 1dose (0.65 mL) contains:
Varicella-zoster virus Oka/Merck strain, (live, attenuated) $\geq 19,400$ PFU

3. LIST OF EXCIPIENTS

Sucrose, hydrolysed gelatin, urea, sodium chloride, potassium dihydrogen phosphate, potassium chloride, monosodium L-glutamate monohydrate, disodium phosphate, sodium hydroxide.

4. PHARMACEUTICAL FORM AND CONTENTS

Powder and solvent for suspension for injection in a pre-filled syringe
1 vial (powder) + 1 pre-filled syringe (solvent) + 1 needle
10 vials (powder) + 10 pre-filled syringes (solvent) + 10 needles
20 vials (powder) + 20 pre-filled syringes (solvent) + 20 needles

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Subcutaneous or intramuscular use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store and transport refrigerated. Do not freeze. Keep the vial in the outer carton to protect from light.

After reconstitution, use immediately or within 30 minutes if stored at 20°C-25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

MSD VACCINS
162 avenue Jean Jaurès
69007 Lyon
France

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/06/341/008 - pack of 1
EU/1/06/341/009 - pack of 10
EU/1/06/341/010 - pack of 20

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

ZOSTAVAX - Powder in vial and solvent in pre-filled syringe with 2 unattached needles - Pack of 1, 10, 20

1. NAME OF THE MEDICINAL PRODUCT

ZOSTAVAX powder and solvent for suspension for injection in a pre-filled syringe shingles (herpes zoster) vaccine (live)

2. STATEMENT OF ACTIVE SUBSTANCE(S)

After reconstitution, 1 dose (0.65 mL) contains:
Varicella-zoster virus Oka/Merck strain, (live, attenuated) $\geq 19,400$ PFU

3. LIST OF EXCIPIENTS

Sucrose, hydrolysed gelatin, urea, sodium chloride, potassium dihydrogen phosphate, potassium chloride, monosodium L-glutamate monohydrate, disodium phosphate, sodium hydroxide.

4. PHARMACEUTICAL FORM AND CONTENTS

Powder and solvent for suspension for injection in a pre-filled syringe
1 vial (powder) + 1 pre-filled syringe (solvent) + 2 needles
10 vials (powder) + 10 pre-filled syringes (solvent) + 20 needles
20 vials (powder) + 20 pre-filled syringes (solvent) + 40 needles

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Subcutaneous or intramuscular use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store and transport refrigerated. Do not freeze. Keep the vial in the outer carton to protect from light.

After reconstitution, use immediately or within 30 minutes if stored at 20°C-25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

MSD VACCINS
162 avenue Jean Jaurès
69007 Lyon
France

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/06/341/011 – pack of 1
EU/1/06/341/012 – pack of 10
EU/1/06/341/013 – pack of 20

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS
VIAL OF POWDER

1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION

ZOSTAVAX powder for injection
SC/IM

2. METHOD OF ADMINISTRATION

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT

1 dose

6. OTHER

MSD VACCINS

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS
PRE-FILLED SYRINGE OF SOLVENT**

1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION

Solvent for ZOSTAVAX

2. METHOD OF ADMINISTRATION

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT

1 dose

6. OTHER

MSD VACCINS

B. PACKAGE LEAFLET

Package Leaflet: Information for the user

ZOSTAVAX

Powder and solvent for suspension for injection shingles (herpes zoster) vaccine (live)

Read all of this leaflet carefully before you are vaccinated because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This vaccine has been prescribed for you only. Do not pass it on to others.
- If you get any of the side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What ZOSTAVAX is and what it is used for
2. What you need to know before you receive ZOSTAVAX
3. How to use ZOSTAVAX
4. Possible side effects
5. How to store ZOSTAVAX
6. Contents of the pack and other information

1. What ZOSTAVAX is and what it is used for

ZOSTAVAX is a vaccine used to prevent shingles (zoster) and zoster-related post-herpetic neuralgia (PHN), the long-lasting nerve pain that follows shingles.

ZOSTAVAX is used to vaccinate individuals 50 years of age or older.

ZOSTAVAX cannot be used to treat existing shingles or the pain associated with existing shingles.

Disease information on shingles:

What is shingles?

Shingles is a painful, blistering rash. It usually occurs in one part of the body and can last for several weeks. It may lead to severe and long-lasting pain and scarring. Less commonly, bacterial skin infections, weakness, muscle paralysis, loss of hearing or vision can occur. Shingles is caused by the same virus that causes chickenpox. After you have had chickenpox, the virus that caused it stays in your body in nerve cells. Sometimes, after many years, the virus becomes active again and causes shingles.

What is PHN?

After the shingles blisters heal, pain can last for months or years and may be severe. This long-lasting nerve pain is called post-herpetic neuralgia or PHN.

2. What you need to know before you receive ZOSTAVAX

Do not receive ZOSTAVAX

- if you are allergic (hypersensitive) to any of the components of this vaccine (including neomycin or any of the other ingredients listed in section 6)
- if you have a blood disorder or any type of cancer that weakens your immune system

- if you have been told by your doctor that you have a weakened immune system as a result of a disease, medicines, or other treatment
- if you have active untreated tuberculosis
- if you are pregnant (in addition, pregnancy should be avoided for 1 month after vaccination, see **Pregnancy and breast-feeding**).

Warnings and precautions

If you have experienced any of the following, talk to your doctor or pharmacist before receiving ZOSTAVAX:

- if you have or have had any medical problems or any allergies
- if you have a fever
- if you have HIV infection

Tell your doctor if you have ever had an allergic reaction to any of the ingredients (including neomycin or any of the ingredients listed under section 6) before you receive this vaccine.

As with many vaccines, ZOSTAVAX may not completely protect all persons who are vaccinated.

If you have a blood clotting disorder or low levels of platelets, the vaccine should be given under the skin because bleeding may occur following administration into the muscle.

Other medicines and ZOSTAVAX

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines or vaccines.

ZOSTAVAX can be administered at the same time as inactivated influenza vaccine. The two vaccines should be given as separate injections at different body sites.

ZOSTAVAX should not be given at the same time as the 23-valent pneumococcal polysaccharide vaccine. For more information about these vaccines, talk to your doctor or health care provider because it may be better to get these vaccines at least 4 weeks apart.

Pregnancy and breast-feeding

ZOSTAVAX should not be given to pregnant women. Women of child-bearing potential should take the necessary precautions to avoid pregnancy for 1 month following vaccination.

Inform your doctor if you are breast-feeding or intending to breast-feed. Your doctor will decide if ZOSTAVAX should be given.

If you are pregnant or breast-feeding, think you may be pregnant, or are planning to have a baby, ask your doctor or pharmacist for advice before receiving this vaccine.

Driving and using machines

There is no information to suggest that ZOSTAVAX affects the ability to drive or use machines.

ZOSTAVAX contains sodium

ZOSTAVAX contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially 'sodium-free'.

3. How to use ZOSTAVAX

ZOSTAVAX should be injected under the skin or into the muscle, preferably in the upper arm.

If you have a blood clotting disorder or low levels of platelets in your blood, the injection will be given under the skin.

ZOSTAVAX is given as a single dose.

Reconstitution instructions intended for healthcare professionals are included at the end of the leaflet.

4. Possible side effects

Like all vaccines and medicines, this vaccine can cause side effects, although not everybody gets them.

Rarely (may affect up to 1 in 1,000 people), allergic reactions may occur. Some of these reactions may be serious and may include difficulty in breathing or swallowing. If you have an allergic reaction, call your doctor right away.

The following side effects have been observed:

- Very common (may affect more than 1 in 10 people): Redness, pain, swelling and itching at the injection site*
- Common (may affect up to 1 in 10 people): Warmth, bruising, hard lump, and rash at the injection site*; headache*; pain in the arm or leg*; joint pain, muscle pain; fever; rash
- Uncommon (may affect up to 1 in 100 people): Nausea; swollen gland (neck, armpit)
- Rare (may affect up to 1 in 1,000 people): Hives at the injection site
- Very rare (may affect up to 1 in 10,000 people): Varicella (chicken pox); shingles; damage of retina caused by inflammation resulting in changes in sight (in patients under immunosuppressive therapy).

*These adverse reactions have been observed in clinical trials and through postmarketing surveillance; most of those observed in clinical trials were reported as mild in intensity.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this vaccine.

5. How to store ZOSTAVAX

Keep this vaccine out of the sight and reach of children.

Do not use this vaccine after the expiry date which is stated on the outer carton after EXP. The expiry date refers to the last day of that month.

Store and transport refrigerated (2°C - 8°C). Do not freeze. Keep the vial in the outer carton in order to protect from light.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What ZOSTAVAX contains

After reconstitution, 1 dose (0.65 mL) contains:

The active substance is:

Varicella-zoster virus¹, Oka/Merck strain, (live, attenuated) not less than 19,400 PFU (plaque-forming units).

¹ Produced in human diploid (MRC-5) cells

The other ingredients are:

Powder

Sucrose, hydrolysed gelatin, sodium chloride, potassium dihydrogen phosphate, potassium chloride, monosodium L-glutamate monohydrate, disodium phosphate, sodium hydroxide (to adjust pH), and urea.

Solvent

Water for injections

What ZOSTAVAX looks like and contents of the pack

The vaccine is a powder for suspension for injection contained in a single-dose vial, which should be reconstituted with the solvent provided with the vial of powder.

The solvent is a clear and colourless liquid. Before mixing with the solvent, the powder is a white to off-white compact crystalline plug.

ZOSTAVAX is available in packs of 1 or 10. Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder: MSD VACCINS, 162 avenue Jean Jaurès, 69007 Lyon, France

Manufacturer: Merck Sharp and Dohme, B.V., Waarderweg, 39, 2031 BN Haarlem, The Netherlands

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder.

België/Belgique/Belgien

MSD Belgium BVBA/SPRL

Tél/Tel: 0800 38 693 (+32(0)27766211)

dpoc_belux@merck.com

Lietuva

UAB Merck Sharp & Dohme

Tel.: +370.5.2780.247

msd_lietuva@merck.com

България

Мерк Шарп и Доум България ЕООД,

тел.: + 359 2 819 3737

info-msdbg@merck.com

Luxembourg/Luxemburg

MSD Belgium BVBA/SPRL

Tél/Tel: +32(0)27766211

dpoc_belux@merck.com

Česká republika

Merck Sharp & Dohme s.r.o.

Tel.: +420 233 010 111

dpoc_czechslovak@merck.com

Magyarország

MSD Pharma Hungary Kft.

Tel.: + 36.1.888.5300

hungary_msd@merck.com

Danmark

MSD Danmark ApS

Tlf: + 45 4482 4000

Malta

Merck Sharp & Dohme Cyprus Limited.

Tel: 8007 4433 (+356 99917558)

dkmail@merck.com

Deutschland

MSD SHARP & DOHME GMBH
Tel: 0800 673 673 673 (+49 (0) 89 4561 2612)
e-mail@msd.de

Eesti

Merck Sharp & Dohme OÜ,
Tel: +372.614.4200
msdeesti@merck.com

Ελλάδα

MSD A.Φ.B.E.E.
Τηλ: +30 210 98 97 300
dpoc_greece@merck.com

España

Merck Sharp & Dohme de España, S.A.
Tel: +34 91 321 06 00
msd_info@merck.com

France

MSD VACCINS
Tél: +33 (0)1 80 46 40 40

Hrvatska

Merck Sharp & Dohme d.o.o.
Tel: +385 1 66 11 333
croatia_info@merck.com

Ireland

Merck Sharp & Dohme Ireland (Human Health)
Limited
Tel: +353 (0)1 2998700
medinfo_ireland@merck.com

Ísland

Vistor hf.
Sími: + 354 535 7000

Italia

MSD Italia S.r.l.
Tel: +39 06 361911
medicalinformation.it@merck.com

Κύπρος

Merck Sharp & Dohme Cyprus Limited
Τηλ: 800 00 673 (+357 22866700)
cyprus_info@merck.com

Latvija

SIA Merck Sharp & Dohme Latvija
Tel: +371.67364.224
msd_lv@merck.com

malta_info@merck.com

Nederland

Merck Sharp & Dohme BV
Tel: 0800 9999000
(+31 23 5153153)
medicalinfo.nl@merck.com

Norge

MSD (Norge) AS
Tlf: +47 32 20 73 00
msdnorge@msd.no

Österreich

Merck Sharp & Dohme Ges.m.b.H.
Tel: +43 (0) 1 26 044
msd-medizin@merck.com

Polska

MSD Polska Sp. z o.o.
Tel.: +48.22.549.51.00
msdpolska@merck.com

Portugal

Merck Sharp & Dohme, Lda
Tel: +351 21 4465700
clic@merck.com

România

Merck Sharp & Dohme Romania S.R.L
Tel: + 4021 529 29 00
msdromania@merck.com

Slovenija

Merck Sharp & Dohme, inovativna zdravila d.o.o.
Tel: +386.1.520.4201
msd.slovenia@merck.com

Slovenská republika

Merck Sharp & Dohme, s. r. o
Tel: +421 2 58282010
dpoc_czechslovak@merck.com

Suomi/Finland

MSD Finland Oy
Puh/Tel: +358 (0)9 804 650
info@msd.fi

Sverige

Merck Sharp & Dohme (Sweden) AB
Tel: +46 77 5700488
medicinskinfo@merck.com

United Kingdom

Merck Sharp & Dohme Limited
Tel: +44 (0) 1992 467272
medicalinformationuk@merck.com

This leaflet was last revised in:

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency website:
<http://www.ema.europa.eu>.

The following information is intended for healthcare professionals only:

Reconstitution instructions

The solvent is a clear and colourless liquid. Before mixing with the solvent, the powder is a white to off-white compact crystalline plug. When completely reconstituted, the vaccine is a semi-hazy to translucent, off-white to pale yellow liquid.

Withdraw the entire volume of solvent into a syringe. Inject the entire content of the syringe into the vial containing the powder. Gently agitate to dissolve completely. Withdraw the entire content of the reconstituted vaccine vial using the same syringe. Inject the vaccine.

It is recommended that the vaccine be administered immediately after reconstitution to minimise loss of potency. Discard if reconstituted vaccine is not used within 30 minutes.

The reconstituted vaccine should be inspected visually for any foreign particulate matter and/or abnormal physical appearance prior to administration. In the event of either being observed, discard the vaccine.

Any unused product or waste material should be disposed of in accordance with local requirements.

See also section 3. How to use ZOSTAVAX.

Package Leaflet: Information for the user

ZOSTAVAX

Powder and solvent for suspension for injection in a pre-filled syringe shingles (herpes zoster) vaccine (live)

Read all of this leaflet carefully before you are vaccinated because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This vaccine has been prescribed for you only. Do not pass it on to others.
- If you get any of the side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What ZOSTAVAX is and what it is used for
2. What you need to know before you receive ZOSTAVAX
3. How to use ZOSTAVAX
4. Possible side effects
5. How to store ZOSTAVAX
6. Contents of the pack and other information

1. What ZOSTAVAX is and what it is used for

ZOSTAVAX is a vaccine used to prevent shingles (zoster) and zoster-related post-herpetic neuralgia (PHN), the long-lasting nerve pain that follows shingles.

ZOSTAVAX is used to vaccinate individuals 50 years of age or older.

ZOSTAVAX cannot be used to treat existing shingles or the pain associated with existing shingles.

Disease information on shingles:

What is shingles?

Shingles is a painful, blistering rash. It usually occurs in one part of the body and can last for several weeks. It may lead to severe and long-lasting pain and scarring. Less commonly, bacterial skin infections, weakness, muscle paralysis, loss of hearing or vision can occur. Shingles is caused by the same virus that causes chickenpox. After you have had chickenpox, the virus that caused it stays in your body in nerve cells. Sometimes, after many years, the virus becomes active again and causes shingles.

What is PHN?

After the shingles blisters heal, pain can last for months or years and may be severe. This long-lasting nerve pain is called post-herpetic neuralgia or PHN.

2. What you need to know before you receive ZOSTAVAX

Do not receive ZOSTAVAX

- if you are allergic (hypersensitive) to any of the components of this vaccine (including neomycin or any of the other ingredients listed in section 6)
- if you have a blood disorder or any type of cancer that weakens your immune system

- if you have been told by your doctor that you have a weakened immune system as a result of a disease, medicines, or other treatment
- if you have active untreated tuberculosis
- if you are pregnant (in addition, pregnancy should be avoided for 1 month after vaccination, see **Pregnancy and breast-feeding**).

Warnings and precautions

If you have experienced any of the following, talk to your doctor or pharmacist before receiving ZOSTAVAX:

- if you have or have had any medical problems or any allergies
- if you have a fever
- if you have HIV infection

Tell your doctor if you have ever had an allergic reaction to any of the ingredients (including neomycin or any of the ingredients listed under section 6) before you receive this vaccine.

As with many vaccines, ZOSTAVAX may not completely protect all persons who are vaccinated.

If you have a blood clotting disorder or low levels of platelets, the vaccine should be given under the skin because bleeding may occur following administration into the muscle.

Other medicines and ZOSTAVAX

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines or vaccines.

ZOSTAVAX can be administered at the same time as inactivated influenza vaccine. The two vaccines should be given as separate injections at different body sites.

ZOSTAVAX should not be given at the same time as the 23-valent pneumococcal polysaccharide vaccine. For more information about these vaccines, talk to your doctor or health care provider because it may be better to get these vaccines at least 4 weeks apart.

Pregnancy and breast-feeding

ZOSTAVAX should not be given to pregnant women. Women of child-bearing age should take the necessary precautions to avoid pregnancy for 1 month following vaccination.

Inform your doctor if you are breast-feeding or intending to breast-feed. Your doctor will decide if ZOSTAVAX should be given.

If you are pregnant or breast-feeding, think you may be pregnant, or are planning to have a baby, ask your doctor or pharmacist for advice before receiving this vaccine.

Driving and using machines

There is no information to suggest that ZOSTAVAX affects the ability to drive or use machines.

ZOSTAVAX contains sodium

ZOSTAVAX contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially 'sodium-free'.

3. How to use ZOSTAVAX

ZOSTAVAX should be injected under the skin or into the muscle, preferably in the upper arm.

If you have a blood clotting disorder or low levels of platelets in your blood, the injection will be given under the skin.

ZOSTAVAX is given as a single dose.

Reconstitution instructions intended for healthcare professionals are included at the end of the leaflet.

4. Possible side effects

Like all vaccines and medicines, this vaccine can cause side effects, although not everybody gets them.

Rarely (may affect up to 1 in 1,000 people), allergic reactions may occur. Some of these reactions may be serious and may include difficulty in breathing or swallowing. If you have an allergic reaction, call your doctor right away.

The following side effects have been observed:

- Very common (may affect more than 1 in 10 people): Redness, pain, swelling and itching at the injection site*
- Common (may affect up to 1 in 10 people): Warmth, bruising, hard lump, and rash at the injection site*; headache*; pain in the arm or leg*; joint pain, muscle pain; fever; rash
- Uncommon (may affect up to 1 in 100 people): Nausea; swollen gland (neck, armpit)
- Rare (may affect up to 1 in 1,000 people): Hives at the injection site
- Very rare (may affect up to 1 in 10,000 people): Varicella (chicken pox); shingles; damage of retina caused by inflammation resulting in changes in sight (in patients under immunosuppressive therapy).

*These adverse reactions have been observed in clinical trials and through postmarketing surveillance; most of those observed in clinical trials were reported as mild in intensity.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this vaccine.

5. How to store ZOSTAVAX

Keep this vaccine out of the sight and reach of children.

Do not use this vaccine after the expiry date which is stated on the outer carton after EXP. The expiry date refers to the last day of that month.

Store and transport refrigerated (2°C - 8°C). Do not freeze. Keep the vial in the outer carton in order to protect from light.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What ZOSTAVAX contains

After reconstitution, 1 dose (0.65 mL) contains:

The active substance is:

Varicella-zoster virus¹, Oka/Merck strain, (live, attenuated) not less than 19,400 PFU (plaque-forming units).

¹Produced in human diploid (MRC-5) cells

The other ingredients are:

Powder

Sucrose, hydrolysed gelatin, sodium chloride, potassium dihydrogen phosphate, potassium chloride, monosodium L-glutamate monohydrate, disodium phosphate, sodium hydroxide (to adjust pH), and urea.

Solvent

Water for injections

What ZOSTAVAX looks like and contents of the pack

The vaccine is a powder for suspension for injection contained in a single-dose vial, which should be reconstituted with the solvent provided with the vial of powder.

The solvent is a clear and colourless liquid. Before mixing with the solvent, the powder is a white to off-white compact crystalline plug.

One pack of ZOSTAVAX contains a vial and a prefilled syringe with or without attached needles. One or 2 separate needles may be available in the secondary packaging of the presentation containing the pre-filled syringe without the attached needle

ZOSTAVAX is available in packs of 1, 10 or 20 with or without needles. Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder: MSD VACCINS, 162 avenue Jean Jaurès, 69007 Lyon, France

Manufacturer: Merck Sharp and Dohme, B.V., Waarderweg, 39, 2031 BN Haarlem, The Netherlands

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder.

België/Belgique/Belgien

MSD Belgium BVBA/SPRL

Tél/Tel: 0800 38 693 (+32(0)27766211)

dpoc_belux@merck.com

Lietuva

UAB Merck Sharp & Dohme

Tel.: +370.5.2780.247

msd_lietuva@merck.com

България

Мерк Шарп и Доум България ЕООД,

тел.: + 359 2 819 3737

info-msdbg@merck.com

Luxembourg/Luxemburg

MSD Belgium BVBA/SPRL

Tél/Tel: +32(0)27766211

dpoc_belux@merck.com

Česká republika

Merck Sharp & Dohme s.r.o.

Tel.: +420 233 010 111

dpoc_czechslovak@merck.com

Magyarország

MSD Pharma Hungary Kft.

Tel.: + 36.1.888.5300

hungary_msd@merck.com

Danmark

MSD Danmark ApS
Tlf: + 45 4482 4000
dkmail@merck.com

Deutschland

MSD SHARP & DOHME GMBH
Tel: 0800 673 673 673 (+49 (0) 89 4561 2612)
e-mail@msd.de

Eesti

Merck Sharp & Dohme OÜ,
Tel: +372.614.4200
msdeesti@merck.com

Ελλάδα

MSD A.Φ.B.E.E.
Τηλ: +30 210 98 97 300
dpoc_greece@merck.com

España

Merck Sharp & Dohme de España, S.A.
Tel: +34 91 321 06 00
msd_info@merck.com

France

MSD VACCINS
Tél: +33 (0)1 80 46 40 40

Hrvatska

Merck Sharp & Dohme d.o.o.
Tel: +385 1 66 11 333
croatia_info@merck.com

Ireland

Merck Sharp & Dohme Ireland (Human Health)
Limited
Tel: +353 (0)1 2998700
medinfo_ireland@merck.com

Ísland

Vistor hf.
Sími: + 354 535 7000

Italia

MSD Italia S.r.l.
Tel: +39 06 361911
medicalinformation.it@merck.com

Κύπρος

Merck Sharp & Dohme Cyprus Limited
Τηλ: 800 00 673 (+357 22866700)
cyprus_info@merck.com

Malta

Merck Sharp & Dohme Cyprus Limited.
Tel: 8007 4433 (+356 99917558)
malta_info@merck.com

Nederland

Merck Sharp & Dohme BV
Tel: 0800 9999000
(+31 23 5153153)
medicalinfo.nl@merck.com

Norge

MSD (Norge) AS
Tlf: +47 32 20 73 00
msdnorge@msd.no

Österreich

Merck Sharp & Dohme Ges.m.b.H.
Tel: +43 (0) 1 26 044
msd-medizin@merck.com

Polska

MSD Polska Sp. z o.o.
Tel.: +48.22.549.51.00
msdpolska@merck.com

Portugal

Merck Sharp & Dohme, Lda
Tel: +351 21 4465700
clic@merck.com

România

Merck Sharp & Dohme Romania S.R.L.
Tel: + 4021 529 29 00
msdromania@merck.com

Slovenija

Merck Sharp & Dohme, inovativna zdravila d.o.o.
Tel: +386.1.520.4201
msd.slovenia@merck.com

Slovenská republika

Merck Sharp & Dohme, s. r. o
Tel: +421 2 58282010
dpoc_czechslovak@merck.com

Suomi/Finland

MSD Finland Oy
Puh/Tel: +358 (0)9 804 650
info@msd.fi

Sverige

Merck Sharp & Dohme (Sweden) AB
Tel: +46 77 5700488
medicinskinfo@merck.com

Latvija

SIA Merck Sharp & Dohme Latvija
Tel: +371.67364.224
msd_lv@merck.com

United Kingdom

Merck Sharp & Dohme Limited
Tel: +44 (0) 1992 467272
medicalinformationuk@merck.com

This leaflet was last revised in:

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency website:
<http://www.ema.europa.eu>.

The following information is intended for healthcare professionals only:

Reconstitution instructions

The solvent is a clear and colourless liquid. Before mixing with the solvent, the powder is a white to off-white compact crystalline plug. When completely reconstituted, the vaccine is a semi-hazy to translucent, off-white to pale yellow liquid.

Inject the entire content of the pre-filled syringe into the vial containing the powder. Gently agitate to dissolve completely. Withdraw the entire content of the reconstituted vaccine using the same syringe. Inject the vaccine. One or 2 separate needles may be available in the packaging of the presentation containing the pre-filled syringe without attached needle. The needle should be pushed into the extremity of the syringe and rotated a quarter of a turn (90°) to secure the connection.

It is recommended that the vaccine be administered immediately after reconstitution to minimise loss of potency. Discard if reconstituted vaccine is not used within 30 minutes.

The reconstituted vaccine should be inspected visually for any foreign particulate matter and/or abnormal physical appearance prior to administration. In the event of either being observed, discard the vaccine.

Any unused product or waste material should be disposed of in accordance with local requirements.

See also section 3. How to use ZOSTAVAX.