

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Prevenar 13 suspension for injection  
Pneumococcal polysaccharide conjugate vaccine (13-valent, adsorbed)

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

1 dose (0.5 ml) contains:

Pneumococcal polysaccharide serotype 1 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 3 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 4 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 5 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 6A <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 6B <sup>1</sup>	4.4 µg
Pneumococcal polysaccharide serotype 7F <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 9V <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 14 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 18C <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 19A <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 19F <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 23F <sup>1</sup>	2.2 µg

<sup>1</sup>Conjugated to CRM<sub>197</sub> carrier protein, adsorbed on aluminium phosphate.

1 dose (0.5 ml) contains approximately 32 µg CRM<sub>197</sub> carrier protein and 0.125 mg aluminium.

### Excipients with known effect

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Suspension for injection.  
The vaccine is a homogeneous white suspension.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Active immunisation for the prevention of invasive disease, pneumonia and acute otitis media caused by *Streptococcus pneumoniae* in infants, children and adolescents from 6 weeks to 17 years of age.

Active immunisation for the prevention of invasive disease and pneumonia caused by *Streptococcus pneumoniae* in adults ≥18 years of age and the elderly.

See sections 4.4 and 5.1 for information on protection against specific pneumococcal serotypes.

The use of Prevenar 13 should be determined on the basis of official recommendations taking into consideration the risk of invasive disease and pneumonia in different age groups, underlying comorbidities as well as the variability of serotype epidemiology in different geographical areas.

## 4.2 Posology and method of administration

The immunisation schedules for Prevenar 13 should be based on official recommendations.

### Posology

#### Infants and children aged 6 weeks to 5 years

It is recommended that infants who receive a first dose of Prevenar 13 complete the vaccination course with Prevenar 13.

#### Infants aged 6 weeks-6 months

##### *Three-dose primary series*

The recommended immunisation series consists of four doses, each of 0.5 ml. The primary infant series consists of three doses, with the first dose usually given at 2 months of age and with an interval of at least 1 month between doses. The first dose may be given as early as six weeks of age. The fourth (booster) dose is recommended between 11 and 15 months of age.

##### *Two-dose primary series*

Alternatively, when Prevenar 13 is given as part of a routine infant immunisation programme, a series consisting of three doses, each of 0.5 ml, may be given. The first dose may be administered from the age of 2 months, with a second dose 2 months later. The third (booster) dose is recommended between 11 and 15 months of age (see section 5.1).

#### Preterm infants (<37 weeks gestation)

In preterm infants, the recommended immunisation series consists of four doses, each of 0.5 ml. The primary infant series consists of three doses, with the first dose given at 2 months of age and with an interval of at least 1 month between doses. The first dose may be given as early as six weeks of age. The fourth (booster) dose is recommended between 11 and 15 months of age (see sections 4.4 and 5.1).

#### Unvaccinated infants and children $\geq 7$ months of age

##### *Infants aged 7-11 months*

Two doses, each of 0.5 ml, with an interval of at least 1 month between doses. A third dose is recommended in the second year of life.

##### *Children aged 12-23 months*

Two doses, each of 0.5 ml, with an interval of at least 2 months between doses (see section 5.1).

##### *Children and adolescents aged 2-17 years*

One single dose of 0.5 ml.

#### Prevenar 13 vaccine schedule for infants and children previously vaccinated with Prevenar (7-valent) (*Streptococcus pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F, and 23F)

Prevenar 13 contains the same 7 serotypes included in Prevenar, using the same carrier protein CRM<sub>197</sub>.

Infants and children who have begun immunisation with Prevenar may switch to Prevenar 13 at any point in the schedule.

#### *Young children (12-59 months) completely immunised with Prevenar (7-valent)*

Young children who are considered completely immunised with Prevenar (7-valent) should receive one dose of 0.5 ml of Prevenar 13 to elicit immune responses to the 6 additional serotypes. This dose of Prevenar 13 should be administered at least 8 weeks after the final dose of Prevenar (7-valent) (see section 5.1).

#### *Children and adolescents 5–17 years*

Children 5 to 17 years of age may receive a single dose of Prevenar 13 if they have been previously vaccinated with one or more doses of Prevenar. This dose of Prevenar 13 should be administered at least 8 weeks after the final dose of Prevenar (7-valent) (see section 5.1).

#### Adults $\geq$ 18 years of age, and the elderly

One single dose.

The need for revaccination with a subsequent dose of Prevenar 13 has not been established.

Regardless of prior pneumococcal vaccination status, if the use of 23-valent pneumococcal polysaccharide vaccine is considered appropriate, Prevenar 13 should be given first (see sections 4.5 and 5.1).

#### Special Populations

Individuals who have underlying conditions predisposing them to invasive pneumococcal disease (such as sickle cell disease or HIV infection) including those previously vaccinated with one or more doses of 23-valent pneumococcal polysaccharide vaccine may receive at least one dose of Prevenar 13 (see section 5.1).

In individuals with an haematopoietic stem cell transplant (HSCT), the recommended immunisation series consists of four doses of Prevenar 13, each of 0.5 ml. The primary series consists of three doses, with the first dose given at 3 to 6 months after HSCT and with an interval of at least 1 month between doses. A fourth (booster) dose is recommended 6 months after the third dose (see section 5.1).

#### Method of administration

The vaccine should be given by intramuscular injection. The preferred sites are the anterolateral aspect of the thigh (vastus lateralis muscle) in infants or the deltoid muscle of the upper arm in children and adults.

### **4.3 Contraindications**

Hypersensitivity to the active substances, to any of the excipients listed in section 6.1, or to diphtheria toxoid.

As with other vaccines, the administration of Prevenar 13 should be postponed in subjects suffering from acute, severe febrile illness. However, the presence of a minor infection, such as a cold, should not result in the deferral of vaccination.

### **4.4 Special warnings and precautions for use**

Prevenar 13 must not be administered intravascularly.

As with all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of a rare anaphylactic event following the administration of the vaccine.

This vaccine should not be given as an intramuscular injection to individuals with thrombocytopaenia or any coagulation disorder that would contraindicate intramuscular injection, but may be given subcutaneously if the potential benefit clearly outweighs the risks (see section 5.1).

Prevenar 13 will only protect against *Streptococcus pneumoniae* serotypes included in the vaccine, and will not protect against other microorganisms that cause invasive disease, pneumonia, or otitis media. As with any vaccine, Prevenar 13 may not protect all individuals receiving the vaccine from pneumococcal disease. For the most recent epidemiological information in your country you should consult with the relevant national organisation.

Individuals with impaired immune responsiveness, whether due to the use of immuno-suppressive therapy, a genetic defect, human immunodeficiency virus (HIV) infection, or other causes, may have reduced antibody response to active immunisation.

Safety and immunogenicity data are available for a limited number of individuals with sickle cell disease, HIV infection, or with an haematopoietic stem cell transplant (see section 5.1). Safety and immunogenicity data for Prevenar 13 are not available for individuals in other specific immuno-compromised groups (e.g., malignancy or nephrotic syndrome) and vaccination should be considered on an individual basis.

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially 'sodium-free'.

#### Infants and children aged 6 weeks to 5 years

In clinical studies, Prevenar 13 elicited an immune response to all thirteen serotypes included in the vaccine. The immune response for serotype 3 following the booster dose was not increased above the levels seen after the infant vaccination series; the clinical relevance of this observation regarding the induction of serotype 3 immune memory is unknown (see section 5.1).

The proportions of functional antibody responders (OPA titres  $\geq 1:8$ ) to serotypes 1, 3 and 5 were high. However, the OPA geometric mean titres were lower than those against each of the remaining additional vaccine serotypes; the clinical relevance of this observation for protective efficacy is unknown (see section 5.1).

Limited data have demonstrated that Prevenar 7-valent (three-dose primary series) induces an acceptable immune response in infants with sickle cell disease with a safety profile similar to that observed in non-high-risk groups (see section 5.1).

Children younger than 2 years old should receive the appropriate-for-age Prevenar 13 vaccination series (see section 4.2). The use of pneumococcal conjugate vaccine does not replace the use of 23-valent pneumococcal polysaccharide vaccines in children  $\geq 2$  years of age with conditions (such as sickle cell disease, asplenia, HIV infection, chronic illness, or those who are immuno-compromised) placing them at higher risk for invasive disease due to *Streptococcus pneumoniae*. Whenever recommended, children at risk who are  $\geq 24$  months of age and already primed with Prevenar 13 should receive 23-valent pneumococcal polysaccharide vaccine. The interval between the 13-valent pneumococcal conjugate vaccine (Prevenar 13) and the 23-valent pneumococcal polysaccharide vaccine should not be less than 8 weeks. There are no data available to indicate whether the administration of 23-valent pneumococcal polysaccharide vaccine to unprimed children or to children primed with Prevenar 13 might result in hyporesponsiveness to further doses of Prevenar 13.

The potential risk of apnoea and the need for respiratory monitoring for 48-72 h should be considered when administering the primary immunisation series to very premature infants (born  $\leq 28$  weeks of gestation), and particularly for those with a previous history of respiratory immaturity. As the benefit of vaccination is high in this group of infants, vaccination should not be withheld or delayed.

For vaccine serotypes, protection against otitis media is expected to be lower than protection against invasive disease. As otitis media is caused by many organisms other than pneumococcal serotypes represented in the vaccine, protection against all otitis media is expected to be low (see section 5.1).

When Prevenar 13 is administered concomitantly with Infanrix hexa (DTPa-HBV-IPV/Hib), the rates of febrile reactions are similar to those seen with concomitant administration of Prevenar (7-valent) and Infanrix hexa (see section 4.8). Increased reporting rates of convulsions (with or without fever) and hypotonic hyporesponsive episode (HHE) were observed with concomitant administration of Prevenar 13 and Infanrix hexa (see section 4.8).

Antipyretic treatment should be initiated according to local treatment guidelines for children with seizure disorders or with a prior history of febrile seizures and for all children receiving Prevenar 13 simultaneously with vaccines containing whole cell pertussis.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

##### Infants and children aged 6 weeks to 5 years

Prevenar 13 can be given concomitantly with any of the following vaccine antigens, either as monovalent or combination vaccines: diphtheria, tetanus, acellular or whole cell pertussis, *Haemophilus influenzae* type b, inactivated poliomyelitis, hepatitis B (see section 4.4 for guidance on Infanrix hexa), meningococcal serogroup C, measles, mumps, rubella, varicella and rotavirus vaccine.

Prevenar 13 can also be given concomitantly between 12-23 months with the tetanus toxoid conjugated meningococcal polysaccharide serogroups A, C, W and Y vaccine to children who have been adequately primed with Prevenar 13 (as per local recommendations).

Data from a postmarketing clinical study evaluating the impact of prophylactic use of antipyretics (ibuprofen and paracetamol) on the immune response to Prevenar 13 suggest that administration of paracetamol concomitantly or within the same day of vaccination may reduce the immune response to Prevenar 13 after the infant series. Responses to the booster dose administered at 12 months were unaffected. The clinical significance of this observation is unknown.

##### Children and adolescents 6 to 17 years of age

No data are currently available regarding concomitant use with other vaccines.

##### Adults 18 to 49 years of age

No data are available regarding concomitant use with other vaccines.

##### Adults aged 50 years and older

Prevenar 13 may be administered concomitantly with the seasonal trivalent inactivated influenza vaccine (TIV).

In two studies conducted in adults aged 50-59 and 65 years and older, it was demonstrated that Prevenar 13 may be given concomitantly with trivalent inactivated influenza vaccine (TIV). The responses to all three TIV antigens were comparable when TIV was given alone or concomitantly with Prevenar 13.

When Prevenar 13 was given concomitantly with TIV, the immune responses to Prevenar 13 were lower compared to when Prevenar 13 was given alone, however, there was no long-term impact on circulating antibody levels.

In a third study in adults aged 50-93 years, it was demonstrated that Prevenar 13 may be given concomitantly with the seasonal quadrivalent inactivated influenza vaccine (QIV). The immune responses to all four QIV strains were noninferior when Prevenar 13 was given concomitantly with QIV compared to when QIV was given alone.

The immune responses to Prevenar 13 were noninferior when Prevenar 13 was given concomitantly with QIV compared to when Prevenar 13 was given alone. As with concomitant administration with trivalent vaccines, immune responses to some pneumococcal serotypes were lower when both vaccines were given concomitantly.

Concomitant use with other vaccines has not been investigated.

Different injectable vaccines should always be given at different vaccination sites.

Concomitant administration of Prevenar 13 and 23-valent pneumococcal polysaccharide vaccine has not been studied. In clinical studies when Prevenar 13 was given 1 year after 23-valent pneumococcal polysaccharide vaccine the immune responses were lower for all serotypes compared to when Prevenar 13 was given to subjects not previously immunised with 23-valent pneumococcal polysaccharide vaccine. The clinical significance of this is unknown.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

There are no data from the use of pneumococcal 13-valent conjugate vaccine in pregnant women. Therefore the use of Prevenar 13 should be avoided during pregnancy.

##### Breast-feeding

It is unknown whether pneumococcal 13-valent conjugate vaccine is excreted in human milk.

##### Fertility

Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3).

#### **4.7 Effects on ability to drive and use machines**

Prevenar 13 has no or negligible influence on the ability to drive and use machines. However, some of the effects mentioned under section 4.8 “Undesirable effects” may temporarily affect the ability to drive or use machines.

#### **4.8 Undesirable effects**

Analysis of postmarketing reporting rates suggests a potential increased risk of convulsions, with or without fever, and HHE when comparing groups which reported use of Prevenar 13 with Infanrix hexa to those which reported use of Prevenar 13 alone.

Adverse reactions reported in clinical studies or from the postmarketing experience for all age groups are listed in this section per system organ class, in decreasing order of frequency and seriousness. The frequency is defined as follows: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ), very rare ( $< 1/10,000$ ), not known (cannot be estimated from available data).

##### Infants and children aged 6 weeks to 5 years

The safety of the vaccine was assessed in controlled clinical studies where 14,267 doses were given to 4,429 healthy infants from 6 weeks of age at first vaccination and 11-16 months of age at booster dose. In all infant studies, Prevenar 13 was co-administered with routine paediatric vaccines (see section 4.5).

Safety in 354 previously unvaccinated children (7 months to 5 years of age) was also assessed.

The most commonly reported adverse reactions in children 6 weeks to 5 years of age were vaccination-site reactions, fever, irritability, decreased appetite, and increased and/or decreased sleep.

In a clinical study in infants vaccinated at 2, 3, and 4 months of age, fever  $\geq 38^{\circ}\text{C}$  was reported at higher rates among infants who received Prevenar (7-valent) concomitantly with Infanrix hexa (28.3% to 42.3%) than in infants receiving Infanrix hexa alone (15.6% to 23.1%). After a booster dose at 12 to 15 months of age, fever  $\geq 38^{\circ}\text{C}$  was reported in 50.0% of infants who received Prevenar (7-valent) and Infanrix hexa at the same time as compared to 33.6% of infants receiving Infanrix hexa alone. These reactions were mostly moderate (less than or equal to  $39^{\circ}\text{C}$ ) and transient.

An increase in vaccination-site reactions was reported in children older than 12 months compared to rates observed in infants during the primary series with Prevenar 13.

#### *Adverse reactions from clinical studies*

In clinical studies, the safety profile of Prevenar 13 was similar to Prevenar. The following frequencies are based on adverse reactions assessed in Prevenar 13 clinical studies:

#### Immune system disorders:

Rare: Hypersensitivity reaction including face oedema, dyspnoea, bronchospasm

#### Nervous system disorders:

Uncommon: Convulsions (including febrile convulsions)

Rare: Hypotonic-hyporesponsive episode

#### Gastrointestinal disorders:

Very common: Decreased appetite

Common: Vomiting; diarrhoea

#### Skin and subcutaneous tissue disorders:

Common: Rash

Uncommon: Urticaria or urticaria-like rash

#### General disorders and administration site conditions:

Very common: Pyrexia; irritability; any vaccination-site erythema, induration/swelling or pain/tenderness; somnolence; poor quality sleep

Vaccination-site erythema or induration/swelling 2.5 cm–7.0 cm (after the booster dose and in older children [age 2 to 5 years])

Common: Pyrexia  $> 39^{\circ}\text{C}$ ; vaccination-site movement impairment (due to pain); vaccination-site erythema or induration/swelling 2.5 cm–7.0 cm (after infant series)

Uncommon: Vaccination-site erythema, induration/swelling  $> 7.0$  cm; crying

#### *Adverse reactions from Prevenar 13 postmarketing experience*

Although the following adverse drug reactions were not observed in the Prevenar 13 clinical studies in infants and children, the following are considered adverse drug reactions for Prevenar 13 as they were reported in the postmarketing experience. Because these reactions were derived from spontaneous reports, the frequencies could not be determined and are thus considered as not known.

#### Blood and lymphatic system disorders:

Lymphadenopathy (localised to the region of the vaccination-site)

#### Immune system disorders:

Anaphylactic/anaphylactoid reaction including shock; angioedema

#### Skin and subcutaneous tissue disorders:

Erythema multiforme

General disorders and administration site conditions:

Vaccination-site urticaria; vaccination-site dermatitis; vaccination-site pruritus; flushing

Additional information in special populations:

Apnoea in very premature infants ( $\leq 28$  weeks of gestation) (see section 4.4).

Children and adolescents aged 6 to 17 years of age

Safety was evaluated in 592 children (294 children aged 5 to 10 years previously immunised with at least one dose of Prevenar and 298 children aged 10 to 17 years who had not received a pneumococcal vaccine).

The most common adverse events in children and adolescents 6 to 17 years of age were:

Nervous system disorders:

Common: Headaches

Gastrointestinal disorders:

Very common: Decreased appetite

Common: Vomiting; diarrhoea

Skin and subcutaneous tissue disorders:

Common: Rash; urticaria or urticaria-like rash

General disorders and administration site conditions:

Very common: Irritability; any vaccination-site erythema; induration/swelling or pain/tenderness; somnolence; poor quality sleep; vaccination-site tenderness (including impaired movement)

Common: Pyrexia

Other adverse events previously observed in infants and children 6 weeks to 5 years of age may also be applicable to this age group but were not seen in this study possibly due to the small sample size.

Additional information in special populations

Children and adolescents with sickle cell disease, HIV infection, or an haematopoietic stem cell transplant have similar frequencies of adverse reactions, except that headaches, vomiting, diarrhoea, pyrexia, fatigue, arthralgia, and myalgia were very common.

Adults  $\geq 18$  years and the elderly

Safety was assessed in 7 clinical studies including 91,593 adults ranging in age from 18 to 101 years. Prevenar 13 was administered to 48,806 adults; 2,616 (5.4%) aged 50 to 64 years, and 45,291 (92.8%) aged 65 years and older. One of the 7 studies included a group of adults ( $n=899$ ) ranging from 18 to 49 years who received Prevenar 13 and who were not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine. Of the Prevenar 13 recipients 1,916 adults were previously vaccinated with the 23-valent pneumococcal polysaccharide vaccine at least 3 years prior to study vaccination, and 46,890 were 23-valent pneumococcal polysaccharide vaccine unvaccinated.

A trend to lower frequency of adverse reactions was associated with greater age; adults  $> 65$  years of age (regardless of prior pneumococcal vaccination status) reported fewer adverse reactions than younger adults, with adverse reactions generally most common in the youngest adults, 18 to 29 years of age.

Overall, the frequency categories were similar for all age groups, with the exception of vomiting which was very common ( $\geq 1/10$ ) in adults aged 18 to 49 years and common ( $\geq 1/100$  to  $< 1/10$ ) in all other age groups, and pyrexia was very common in adults aged 18 to 29 years and common in all other age groups. Severe vaccination-site pain/tenderness and severe limitation of arm movement was very common in adults 18 to 39 years and common in all other age groups.

*Adverse reactions from clinical studies*

Local reactions and systemic events were solicited daily after each vaccination for 14 days in 6 studies and 7 days in the remaining study. The following frequencies are based on adverse reactions assessed in Prevenar 13 clinical studies in adults:

Metabolism and nutrition disorders:

Very common: Decreased appetite

Nervous system disorders:

Very common: Headaches

Gastrointestinal disorders:

Very common: Diarrhoea; vomiting (in adults aged 18 to 49 years)

Common: Vomiting (in adults aged 50 years and over)

Uncommon: Nausea

Immune system disorders:

Uncommon: Hypersensitivity reaction including face oedema, dyspnoea, bronchospasm

Skin and subcutaneous tissue disorders:

Very common: Rash

General disorders and administration site conditions:

Very common: Chills; fatigue; vaccination-site erythema; vaccination-site induration/swelling; vaccination-site pain/tenderness (severe vaccination-site pain/tenderness very common in adults aged 18 to 39 years); limitation of arm movement (severe limitation of arm movements very common in adults aged 18 to 39 years)

Common: Pyrexia (very common in adults aged 18 to 29 years)

Uncommon: Lymphadenopathy localized to the region of the vaccination-site

Musculoskeletal and connective tissue disorders:

Very common: Arthralgia; myalgia

Overall, no significant differences in frequencies of adverse reactions were seen when Prevenar 13 was given to adults previously vaccinated with the pneumococcal polysaccharide vaccine.

*Additional information in special populations*

Adults with HIV infection have similar frequencies of adverse reactions, except that pyrexia and vomiting were very common and nausea common.

Adults with an haematopoietic stem cell transplant have similar frequencies of adverse reactions, except that pyrexia and vomiting were very common.

Higher frequency in some solicited systemic reactions was observed when Prevenar 13 was administered concomitantly with trivalent inactivated influenza vaccine (TIV) compared to TIV given alone (headache, chills, rash, decreased appetite, arthralgia, and myalgia) or Prevenar 13 given alone (headache, fatigue, chills, decreased appetite, and arthralgia).

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

## **4.9 Overdose**

Overdose with Prevenar 13 is unlikely due to its presentation as a pre-filled syringe. However, in infants and children there have been reports of overdose with Prevenar 13 defined as subsequent doses administered closer than recommended to the previous dose. In general, adverse events reported with overdose are consistent with those that have been reported with doses given in the recommended paediatric schedules of Prevenar 13.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: vaccines, pneumococcal vaccines; ATC code: J07AL02

Prevenar 13 contains the 7 pneumococcal capsular polysaccharides that are in Prevenar (4, 6B, 9V, 14, 18C, 19F, 23F) plus 6 additional polysaccharides (1, 3, 5, 6A, 7F, 19A) all conjugated to CRM<sub>197</sub> carrier protein.

#### Burden of disease

##### *Infants and children aged 6 weeks to 5 years*

Based on serotype surveillance in Europe performed before the introduction of Prevenar, Prevenar 13 is estimated to cover 73-100% (depending on the country) of serotypes causing invasive pneumococcal disease (IPD) in children less than 5 years of age. In this age group, serotypes 1, 3, 5, 6A, 7F, and 19A account for 15.6% to 59.7% of invasive disease, depending on the country, the time period studied, and the use of Prevenar.

Acute otitis media (AOM) is a common childhood disease with different aetiologies. Bacteria can be responsible for 60-70% of clinical episodes of AOM. *S. pneumoniae* is one of the most common causes of bacterial AOM worldwide.

Prevenar 13 is estimated to cover over 90% of serotypes causing antimicrobial-resistant IPD.

##### *Children and adolescents aged 6 to 17 years*

In children and adolescents aged 6 to 17 years, the incidence of pneumococcal disease is low, however, there is an increased risk of morbidity and mortality in those with underlying comorbidities.

##### *Adults ≥18 years and the elderly*

Pneumonia is the most common clinical presentation of pneumococcal disease in adults.

The reported incidence of community-acquired pneumonia (CAP) and IPD in Europe varies by country, increases with age from 50 years and is highest in individuals aged ≥ 65 years. *S. pneumoniae* is the most frequent cause of CAP, and is estimated to be responsible for approximately 30% of all CAP cases requiring hospitalisation in adults in developed countries.

Bacteraemic pneumonia (approximately 80% of IPD in adults), bacteraemia without a focus, and meningitis are the most common manifestations of IPD in adults. Based on surveillance data following

the introduction of Prevenar but before the introduction of Prevenar 13 in childhood vaccination programmes, the pneumococcal serotypes in Prevenar 13 may be responsible for at least 50 – 76% (depending on country) of IPD in adults.

The risk for CAP and IPD in adults also increases with chronic underlying medical conditions, specifically, anatomical or functional asplenia, diabetes mellitus, asthma, chronic cardiovascular, pulmonary, kidney or liver disease, and it is highest in those who are immune-suppressed such as those with malignant haematological diseases or HIV infection.

#### Prevenar 13 immunogenicity clinical studies in infants, children and adolescents

The protective efficacy of Prevenar 13 against IPD has not been studied. As recommended by the World Health Organization (WHO) the assessment of potential efficacy against IPD in infants and young children has been based on a comparison of immune responses to the seven common serotypes shared between Prevenar 13 and Prevenar, for which protective efficacy has been proven (for Prevenar (7-valent) efficacy in infants and children, see below). Immune responses to the additional 6 serotypes were also measured.

#### Immune responses following a three-dose primary infant series

Clinical studies have been conducted in a number of European countries and the US using a range of vaccination schedules, including two randomised non-inferiority studies (Germany using a 2, 3, 4 month primary series [006] and US using a 2, 4, 6 month primary series [004]). In these two studies pneumococcal immune responses were compared using a set of non-inferiority criteria including the percentage of subjects with serum anti-polysaccharide serotype-specific IgG  $\geq 0.35$   $\mu\text{g/ml}$  one month after the primary series and the comparison of IgG geometric mean concentrations (ELISA GMCs); in addition, functional antibody titres (OPA) between subjects receiving Prevenar 13 and Prevenar were compared. For the six additional serotypes, these values were compared with the lowest response among all of the seven common serotypes in the Prevenar recipients.

The non-inferiority immune response comparisons for study 006, based on the proportion of infants achieving anti-polysaccharide IgG concentrations  $\geq 0.35$   $\mu\text{g/ml}$ , are shown in Table 1. Results for study 004 were similar. Prevenar 13 non-inferiority (lower bound of the 95% CI for the difference in percentage of responders at 0.35  $\mu\text{g/ml}$  between groups was  $>-10\%$ ) was demonstrated for all 7 common serotypes, except for serotype 6B in study 006 and serotypes 6B and 9V in study 004, which missed by a small margin. All seven common serotypes met pre-defined non-inferiority criteria for IgG ELISA GMCs. Prevenar 13 elicited comparable, although slightly lower, antibody levels than Prevenar for the 7 common serotypes. The clinical relevance of these differences is not known.

Non-inferiority was met for the 6 additional serotypes based on the proportion of infants achieving antibody concentrations  $\geq 0.35$   $\mu\text{g/ml}$  and comparison of IgG ELISA GMCs in study 006 and was met for 5 out of the 6 serotypes, with the exception of serotype 3 for study 004. For serotype 3, the percentages of Prevenar 13 recipients with serum IgG  $\geq 0.35$   $\mu\text{g/ml}$  were 98.2% (study 006) and 63.5% (study 004).

<b>Table 1: Comparison of the proportion of subjects achieving a pneumococcal anti-polysaccharide IgG antibody concentration <math>\geq 0.35</math> <math>\mu\text{g/ml}</math> after dose 3 of the infant series – study 006</b>			
<b>Serotypes</b>	<b>Prevenar 13 % (N=282-285)</b>	<b>7-valent Prevenar % (N=277-279)</b>	<b>Difference (95 % CI)</b>
<b>7-valent Prevenar serotypes</b>			
4	98.2	98.2	0.0 (-2.5, 2.6)
6B	77.5	87.1	-9.6 (-16.0, -3.3)
9V	98.6	96.4	2.2 (-0.4, 5.2)
14	98.9	97.5	1.5 (-0.9, 4.1)
18C	97.2	98.6	-1.4 (-4.2, 1.2)
19F	95.8	96.0	-0.3 (-3.8, 3.3)
23F	88.7	89.5	-0.8 (-6.0, 4.5)
<b>Additional serotypes in Prevenar 13</b>			
1	96.1	87.1*	9.1 (4.5, 13.9)
3	98.2	87.1	11.2 (7.0, 15.8)
5	93.0	87.1	5.9 (0.8, 11.1)
6A	91.9	87.1	4.8 (-0.3, 10.1)
7F	98.6	87.1	11.5 (7.4, 16.1)
19A	99.3	87.1	12.2 (8.3, 16.8)

\*The serotype in Prevenar with the lowest percent response rate was 6B in study 006 (87.1 %).

Prevenar 13 elicited functional antibodies to all 13 vaccine serotypes in studies 004 and 006. For the 7 common serotypes there were no differences between groups in the proportion of subjects with OPA titres  $\geq 1:8$ . For each of the seven common serotypes,  $>96\%$  and  $>90\%$  of the Prevenar 13 recipients reached an OPA titre  $\geq 1:8$  one month after the primary series in studies 006 and 004, respectively.

For each of the 6 additional serotypes, Prevenar 13 elicited OPA titres  $\geq 1:8$  in 91.4% to 100% of vaccinees one month after the primary series in studies 004/006. The functional antibody (OPA) geometric mean titres for serotypes 1, 3 and 5 were lower than the titres for each of the other additional serotypes; the clinical relevance of this observation for protective efficacy is unknown.

#### Immune responses following a two-dose primary infant series

The immunogenicity after two doses in infants has been documented in four studies. The proportion of infants achieving a pneumococcal anti-capsular polysaccharide IgG concentration  $\geq 0.35$   $\mu\text{g/ml}$  one month after the second dose ranged from 79.6% to 98.5% across 11 of the 13 vaccine serotypes. Smaller proportions of infants achieved this antibody concentration threshold for serotype 6B (27.9% to 57.3%) and 23F (55.8% to 68.1%) for all studies using a 2, 4 month regimen, compared to 58.4% for serotype 6B and 68.6% for 23F for a study using a 3, 5 month regimen. After the booster dose, all vaccine serotypes including 6B and 23F had immune responses consistent with adequate priming with a two-dose primary series. In a UK study, the functional antibody (OPA) responses were comparable for all serotypes including 6B and 23F in the Prevenar and Prevenar 13 arms after the primary series at two and four months of age and after the booster dose at 12 months of age. For Prevenar 13 recipients, the proportion of responders with an OPA titre  $\geq 1:8$  was at least 87 % following the infant series, and at least 93% following the booster dose. The OPA geometric mean titres for serotypes 1, 3 and 5 were lower than the titres for each of the other additional serotypes; the clinical relevance of this observation is unknown.

#### Booster responses following two-dose and three-dose primary infant series

Following the booster dose, antibody concentrations increased from the pre-booster level for all 13 serotypes. Post-booster antibody concentrations were higher for 12 serotypes than those achieved after the infant primary series. These observations are consistent with adequate priming (the induction of immunologic memory). The immune response for serotype 3 following the booster dose was not increased above the levels seen after the infant vaccination series; the clinical relevance of this observation regarding the induction of serotype 3 immune memory is unknown.

Antibody responses to booster doses following two-dose or three-dose infant primary series were comparable for all 13 vaccine serotypes.

For children aged from 7 months to 5 years, age appropriate catch-up immunisation schedules (as described in section 4.2) result in levels of anti-capsular polysaccharide IgG antibody responses to each of the 13 serotypes that are at least comparable to those of a three-dose primary series in infants.

Antibody persistence and immunological memory were evaluated in a study in healthy children who received a single dose of Prevenar 13 at least 2 years after they had been previously immunised with either 4 doses of Prevenar, a 3-dose infant series of Prevenar followed by Prevenar 13 at 12 months of age, or 4 doses of Prevenar 13.

The single dose of Prevenar 13, in children approximately 3.4 years of age regardless of previous vaccination history with Prevenar or Prevenar 13, induced a robust antibody response for both the 7 common serotypes and the 6 additional serotypes in Prevenar 13.

Since the introduction of 7-valent Prevenar in 2000, pneumococcal disease surveillance data have not shown that the immunity elicited by Prevenar in infancy has waned over time.

### Preterm Infants

Safety and immunogenicity of Prevenar 13 given at 2, 3, 4 and 12 months was assessed in approximately 100 prematurely born infants (mean Estimated Gestational Age [EGA], 31 weeks; range, 26 to 36 weeks) and compared with approximately 100 infants born at term (mean EGA, 39 weeks; range, 37 to 42 weeks).

Immune responses in preterm and term infants were compared using the proportion of subjects achieving a pneumococcal polysaccharide IgG binding antibody concentration  $\geq 0.35 \mu\text{g/mL}$  1 month after the infant series, the approach used for immunogenicity comparisons of Prevenar 13 to Prevenar based on WHO guidelines.

More than 85% achieved a pneumococcal polysaccharide IgG binding antibody concentration  $\geq 0.35 \mu\text{g/mL}$  1 month after the infant series, except for serotypes 5 (71.7%), 6A (82.7%), and 6B (72.7%) in the preterm group. For these 3 serotypes, the proportion of responders among preterm infants was significantly lower than among term infants. Approximately one month after the toddler dose, the proportion of subjects in each group achieving this same antibody concentration threshold was  $>97\%$ , except for serotype 3 (71% in preterm infants and 79% in term infants). It is unknown whether immunological memory to all serotypes is induced in pre-term infants. In general, serotype-specific IgG GMCs were lower for preterm infants than term infants.

After the infant series, OPA GMTs were similar in preterm infants compared to term infants except for serotype 5, which was lower in preterm infants. OPA GMTs after the toddler dose relative to those after the infant series were similar or lower for 4 serotypes (4, 14, 18C, and 19F) and were statistically significantly higher for 6 of 13 serotypes (1, 3, 5, 7F, 9V, and 19A) in preterm infants compared to 10 of 13 serotypes (1, 3, 4, 5, 6A, 7F, 9V, 18C, 19A, and 23F) in term infants.

### Children (12-59 months) completely immunised with Prevenar (7-valent)

Following administration of a single dose of Prevenar 13 to children (12-59 months) who are considered completely immunised with Prevenar (7-valent) (either 2 or 3 dose primary series plus booster), the proportion achieving serum IgG levels  $\geq 0.35 \mu\text{g/ml}$  and OPA titres  $\geq 1:8$  was at least 90%. However, 3 (serotypes 1, 5 and 6A) of the 6 additional serotypes showed lower IgG GMC and OPA GMT when compared with children who had received at least one previous vaccination with Prevenar 13. The clinical relevance of the lower GMCs and GMTs is currently unknown.

### Unvaccinated Children (12-23 months)

Studies in unvaccinated children (12-23 months) with Prevenar (7-valent) demonstrated that 2 doses were required to achieve serum IgG concentrations for 6B and 23F similar to those induced by a 3-dose infant series.

### Children and Adolescents 5 to 17 years of age

In an open-label study in 592 healthy children and adolescents including those with asthma (17.4%) who may be predisposed to pneumococcal infection, Prevenar 13 elicited immune responses to all 13 serotypes. A single dose of Prevenar 13 was given to children 5 to 10 years of age previously vaccinated with at least 1 dose of Prevenar, and children and adolescents 10 to 17 years of age who had never received a pneumococcal vaccine.

In both the children 5 to 10 years of age and children and adolescents aged 10 to 17 years, the immune response to Prevenar 13 was non inferior to Prevenar for the 7 common serotypes and to Prevenar 13 for the 6 additional serotypes compared to the immune response after the fourth dose in infants vaccinated at 2, 4, 6 and 12-15 months of age as measured by serum IgG.

In children and adolescents aged 10 to 17 years of age OPA GMTs 1 month after vaccination were noninferior to OPA GMTs in the 5 to 10 year old age group for 12 of the 13 serotypes (except serotype 3).

### Immune responses after subcutaneous administration

Subcutaneous administration of Prevenar 13 was evaluated in a non-comparative study in 185 healthy Japanese infants and children who received 4 doses at 2, 4, 6 and 12-15 months of age. The study demonstrated that safety and immunogenicity were generally comparable with observations made in studies of intramuscular administration.

### Prevenar 13 Effectiveness

#### *Invasive Pneumococcal Disease*

Data published by Public Health England showed that, four years after the introduction of Prevenar as a two dose primary infant series with booster dose in the second year of life and with a 94% vaccine uptake, there was a 98% (95% CI 95; 99) reduction in disease caused by the 7 vaccine serotypes in England and Wales. Subsequently, four years following the switch to Prevenar 13, the additional reduction in incidence of IPD due to the 7 serotypes in Prevenar ranged from 76% in children less than 2 years of age to 91% in children 5-14 years of age. The serotype specific reductions for each of the 5 additional serotypes in Prevenar 13 (no cases of serotype 5 IPD were observed) by age group are shown in Table 2 and ranged from 68% (serotype 3) to 100% (serotype 6A) for children less than 5 years of age. Significant incidence reductions were also observed in older age groups who had not been vaccinated with Prevenar 13 (indirect effect).

<b>Table 2: Serotype specific number of cases and incidence reductions of IPD in 2013/14 compared to 2008/09-2009/10 (2008/10) by age in England and Wales</b>									
	<b>&lt;5 years of age</b>			<b>5 to 64 years of age</b>			<b>≥65 years of age</b>		
	2008-10 <sup>§</sup>	2013/14 <sup>§</sup>	% Incidence reduction (95% CI*)	2008-10 <sup>§</sup>	2013/14 <sup>§</sup>	% Incidence reduction (95% CI*)	2008-10 <sup>§</sup>	2013/14 <sup>§</sup>	% Incidence reduction (95% CI*)
<b>Additional serotypes covered by Prevenar 13</b>									
<b>1</b>	59 (54)	5 (5)	91% (98%; 68%)**	458 (382)	77 (71)	83% (88%; 74%)**	102 (89)	13 (13)	87% (94%; 72%)**
<b>3</b>	26 (24)	8 (8)	68% (89%; 6%)	178 (148)	73 (68)	59% (72%; 38%)**	256 (224)	143 (146)	44% (57%; 27%)**
<b>6A</b>	10 (9)	0 (0)	100% (100%; 62%)**	53 (44)	5 (5)	90% (97%; 56%)**	94 (82)	5 (5)	95% (99%; 81%)**
<b>7F</b>	90 (82)	8 (8)	91% (97%; 74%)**	430 (361)	160 (148)	63% (71%; 50%)**	173 (152)	75 (77)	56% (70%; 37%)**
<b>19A</b>	85 (77)	7 (7)	91% (97%; 75%)**	225 (191)	104 (97)	54% (65%; 32%)**	279 (246)	97 (99)	65% (75%; 53%)**
<sup>§</sup> Corrected for proportion of samples serotyped, missing age, denominator compared with 2009/10, and for the trend in total invasive pneumococcal disease up to 2009/10 (after which no trend correction was applied). * 95% CI inflated from a Poisson interval based on over-dispersion of 2.1 seen from modelling of 2000-06 pre-Prevenar all IPD data. ** p<0.005 to cover 6A where p=0.002									

### *Otitis media (OM)*

In a published study performed in Israel, using a 2-dose primary series plus booster dose in the second year of life, the impact of Prevenar 13 on OM was documented in a population-based active-surveillance system with tympanocentesis culturing of middle ear fluid in Israeli children less than 2 years of age with OM.

Following the introduction of Prevenar and subsequently Prevenar 13 there was a decline in incidence from 2.1 to 0.1 cases per 1000 children (95%) for the Prevenar serotypes plus serotype 6A and a decline in incidence from 0.9 to 0.1 cases per 1000 children (89%) for the additional serotypes 1, 3, 5, 7F, and 19A in Prevenar 13. The annual overall pneumococcal incidence of OM declined from 9.6 to 2.1 cases per 1000 children (78%) between July 2004 (prior to the introduction of Prevenar) and June 2013 (post Prevenar 13 introduction).

### *Pneumonia*

In a multicenter observational study in France comparing the periods before and after the switch from Prevenar to Prevenar 13, there was 16% (2060 to 1725 cases) reduction in all community acquired pneumonia (CAP) cases in emergency departments in children 1 month to 15 years of age. Reductions were 53% (167 to 79 cases) ( $p < 0.001$ ) for CAP cases with pleural effusion and 63% (64 to 24 cases) ( $p < 0.001$ ) for microbiologically confirmed pneumococcal CAP cases. In the second year after the introduction of Prevenar 13 the total number of CAP cases due to the 6 additional vaccine serotypes in Prevenar 13 was reduced from 27 to 7 isolates (74%).

The decrease in all cause pneumonia cases was most pronounced in the younger vaccinated age groups with a decrease of 31.8% (757 to 516 cases) and 16.6% (833 to 695 cases) in the age groups <2 years and 2 to 5 years, respectively. The incidence in older, predominantly non-vaccinated children (>5 years) did not change over the duration of the study.

In an ongoing surveillance system (2004 to 2013) to document the impact of Prevenar and subsequently Prevenar 13 on CAP in children less than 5 years in Southern Israel using a 2 dose primary series with a booster dose in the second year of life, there was a reduction of 68% (95% CI 73; 61) in outpatient visits and 32% (95% CI 39; 22) in hospitalizations for alveolar CAP following the introduction of Prevenar 13 when compared to the period before the introduction of Prevenar.

### Effect on nasopharyngeal carriage

In a surveillance study in France in children presenting with acute otitis media, changes in nasopharyngeal (NP) carriage of pneumococcal serotypes were evaluated following the introduction of Prevenar (7-valent) and subsequently Prevenar 13. Prevenar 13 significantly reduced NP carriage of the 6 additional serotypes (and serotype 6C) combined and individual serotypes 6C, 7F, 19A when compared with Prevenar. A reduction in carriage was also seen for serotype 3 (2.5% vs 1.1%;  $p = 0.1$ ). There was no carriage of serotypes 1 and 5 observed.

The effect of pneumococcal conjugate vaccination on nasopharyngeal carriage was studied in a randomised double-blind study in which infants received either Prevenar 13 or Prevenar (7-valent) at 2, 4, 6 and 12 months of age in Israel. Prevenar 13 significantly reduced newly identified NP acquisition of the 6 additional serotypes (and serotype 6C) combined and of individual serotypes 1, 6A, 6C, 7F, 19A when compared with Prevenar. There was no reduction seen in serotype 3 and for serotype 5 the colonization was too infrequent to assess impact. For 6 of the remaining 7 common serotypes, similar rates of NP acquisition were observed in both vaccine groups; for serotype 19F a significant reduction was observed.

In this study, reductions of *S. pneumoniae* serotypes 19A, 19F, and 6A not susceptible to a number of antibiotics were documented. The reductions ranged between 34% and 62% depending on serotype and antibiotic.

### Prevenar (7-valent vaccine) protective efficacy in infants and children

The efficacy of 7-valent Prevenar was evaluated in two major studies – the Northern California Kaiser Permanente (NCKP) study and the Finnish Otitis Media (FinOM) study. Both studies were randomised, double-blind, active-control studies in which infants were randomised to receive either Prevenar or control vaccine (NCKP, meningococcal serogroup C CRM-conjugate [MnCC] vaccine; FinOM, hepatitis B vaccine) in a four-dose series at 2, 4, 6, and 12-15 months of age. The efficacy results from these studies (for invasive pneumococcal disease, pneumonia, and acute otitis media) are presented below (Table 3).

Test	N	VE <sup>2</sup>	95% CI
NCKP: Vaccine-serotype IPD <sup>3</sup>	30,258	97%	85, 100
NCKP: Clinical pneumonia with abnormal chest X-ray	23,746	35%	4, 56
NCKP: Acute Otitis Media (AOM) <sup>4</sup>	23,746		
Total episodes		7%	4, 10
Recurrent AOM (3 episodes in 6 months, or 4 episodes in 1 year)		9%	3, 15
Recurrent AOM (5 episodes in 6 months, or 6 episodes in 1 year)		23%	7, 36
Tympanostomy tube placement		20%	2, 35
FinOM: AOM	1,662		
Total episodes		6%	-4, 16
All pneumococcal AOM		34%	21, 45
Vaccine-serotype AOM		57%	44, 67
<sup>1</sup> Per protocol			
<sup>2</sup> Vaccine efficacy			
<sup>3</sup> October 1995 to April 20, 1999			
<sup>4</sup> October 1995 to April 30, 1998			

### Prevenar (7-valent) effectiveness

The effectiveness (both direct and indirect effect) of 7-valent Prevenar against pneumococcal disease has been evaluated in both three-dose and two-dose primary infant series immunisation programmes, each with booster doses (Table 4). Following the widespread use of Prevenar, the incidence of IPD has been consistently and substantially reduced.

Using the screening method, serotype-specific effectiveness estimates for 2 doses under the age of 1 year in the UK were 66 % (-29, 91 %) and 100 % (25, 100 %) for serotype 6B and 23F, respectively.

Country (year of introduction)	Recommended schedule	Disease reduction, %	95% CI
UK (England & Wales) <sup>1</sup> (2006)	2, 4, + 13 months	<u>Vaccine serotypes:</u> Two doses under age 1: 85%	49, 95%
USA (2000)	2, 4, 6, + 12-15 months		
Children < 5 <sup>2</sup>		Vaccine serotypes: 98% All serotypes: 77%	97, 99% 73, 79%
Persons ≥ 65 <sup>3</sup>		Vaccine serotypes: 76% All serotypes: 38%	NA NA
Canada (Quebec) <sup>4</sup> (2004)	2, 4, + 12 months	All serotypes: 73% <u>Vaccine serotypes:</u> 2-dose infant series: 99% Completed schedule: 100%	NA 92, 100% 82, 100%
<sup>1</sup> Children < 2 years of age. Calculated vaccine effectiveness as of June 2008 (Broome method).			
<sup>2</sup> 2005 data.			
<sup>3</sup> 2004 data.			
<sup>4</sup> Children < 5 years of age. January 2005 to December 2007. Complete effectiveness for routine 2+1 schedule not yet available.			

### *Acute Otitis Media*

Effectiveness of Prevenar in a 3+1 schedule has also been observed against acute otitis media and pneumonia since its introduction in a national immunisation programme. In a retrospective evaluation of a large US insurance database, AOM visits were reduced by 42.7 % (95 % CI, 42.4-43.1 %), and prescriptions for AOM by 41.9 % in children younger than 2 years of age, compared with a pre-licensure baseline (2004 vs. 1997-99). In a similar analysis, hospitalisations and ambulatory visits for all-cause pneumonia were reduced by 52.4 % and 41.1 %, respectively. For those events specifically identified as pneumococcal pneumonia, the observed reductions in hospitalisations and

ambulatory visits were 57.6 % and 46.9 %, respectively, in children younger than 2 years of age, compared with a pre-licensure baseline (2004 vs. 1997-99). While direct cause-and-effect cannot be inferred from observational analyses of this type, these findings suggest that Prevenar plays an important role in reducing the burden of mucosal disease (AOM and pneumonia) in the target population.

#### Efficacy study in adults 65 years and older

Efficacy against vaccine-type (VT) pneumococcal CAP and IPD was assessed in a large-scale randomised double-blind, placebo-controlled study (Community-Acquired Pneumonia Immunization Trial in Adults–CAPiTA) in the Netherlands. 84,496 subjects, 65 years and older received a single vaccination of either Prevenar 13 or placebo in a 1:1 randomisation.

The CAPiTA study enrolled volunteers  $\geq$  65 years of age whose demographic and health characteristics may differ from those seeking vaccination.

A first episode of hospitalised, chest X-ray confirmed pneumonia was identified in about 2% of this population (n=1,814 subjects) of which 329 cases were confirmed pneumococcal CAP and 182 cases were VT pneumococcal CAP in the per protocol and modified intent to treat (mITT) populations. For the primary endpoint (per protocol population), there were 139 (49 Prevenar 13: 90 Placebo) first episodes of VT-CAP resulting in an efficacy of 45.56% (95.2% CI, 21.82-62.49; p=0.0006).

Efficacy was also demonstrated for the two secondary endpoints in the per protocol population. For the non-bacteraemic/non-invasive (NB/NI) pneumococcal CAP secondary endpoint, there were 93 (33 Prevenar 13: 60 Placebo) first episodes of NB/NI VT pneumococcal CAP resulting in an efficacy of 45.00% (95.2% CI, 14.21-65.31; p=0.0067). For the IPD secondary endpoint, there were 35 (7 Prevenar 13: 28 Placebo) first episodes of VT-IPD, resulting in an efficacy of 75.00% (95.2% CI, 41.06-90.87; p=0.0005).

The duration of protective efficacy against a first episode of VT pneumococcal CAP, NB/NI VT pneumococcal CAP, and VT-IPD extended throughout the 4-year study.

The study was not designed to demonstrate efficacy in subgroups, and the number of subjects  $\geq$  85 years of age was not sufficient to demonstrate efficacy in this age group.

#### Immunogenicity studies in adults $\geq$ 18 years and the elderly

In adults, an antibody threshold of serotype-specific pneumococcal polysaccharide IgG binding antibody concentration associated with protection has not been defined. For all pivotal clinical trials, a serotype-specific opsonophagocytosis assay (OPA) was used as a surrogate to assess potential efficacy against invasive pneumococcal disease and pneumonia. OPA geometric mean titers (GMTs) measured 1-month after each vaccination were calculated. OPA titres are expressed as the reciprocal of the highest serum dilution that reduces survival of the pneumococci by at least 50 %.

Pivotal trials for Prevenar 13 were designed to show that functional OPA antibody responses for the 13 serotypes are non-inferior, and for some serotypes superior, to the 12 serotypes in common with the licensed 23-valent pneumococcal polysaccharide vaccine [1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F] one month after vaccine administration. The response to serotype 6A, which is unique to Prevenar 13, was assessed by demonstration of a 4-fold increase in the specific OPA titer above pre-immunised levels.

Five clinical studies were conducted in Europe and the USA evaluating the immunogenicity of Prevenar 13 in different age groups ranging from 18-95 years of age. Clinical studies with Prevenar 13 currently provide immunogenicity data in adults aged 18 years and older, including adults aged 65 and older previously vaccinated with one or more doses of 23-valent pneumococcal polysaccharide vaccine, 5 years prior to enrollment. Each study included healthy adults and immuno-competent adults with stable underlying conditions known to predispose individuals to pneumococcal infection (i.e.,

chronic cardiovascular disease, chronic pulmonary disease including asthma, renal disorders and diabetes mellitus, chronic liver disease including alcoholic liver disease), and adults with risk factors such as smoking and alcohol abuse.

Immunogenicity and safety of Prevenar 13 has been demonstrated in adults aged 18 years and older including those previously vaccinated with a pneumococcal polysaccharide vaccine.

*Adults not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine*

In a head-to-head, comparative trial conducted in adults aged 60-64 years, subjects received a single dose of either Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine. In the same study another group of adults aged 50-59 years and another group of adults aged 18-49 years received a single dose of Prevenar 13.

Table 5 compares the OPA GMTs, 1-month post-dose, in 60-64 year olds given either a single dose of Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine, and in 50-59 year olds given a single dose of Prevenar 13.

<b>Table 5: OPA GMTs in adults aged 60-64 years given Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine (PPSV23) and in adults aged 50-59 years given Prevenar 13<sup>a,b,c</sup></b>							
	<b>Prevenar 13</b>	<b>Prevenar 13</b>	<b>PPSV23</b>	<b>Prevenar 13</b>		<b>Prevenar 13 Relative</b>	
	<b>50-59 Years</b>	<b>60-64 Years</b>	<b>60-64 Years</b>	<b>50-59 Relative to</b>		<b>to PPSV23,</b>	
	<b>N=350-384</b>	<b>N=359-404</b>	<b>N=367-402</b>	<b>60-64 Years</b>		<b>60-64 Years</b>	
<b>Serotype</b>	<b>GMT</b>	<b>GMT</b>	<b>GMT</b>	<b>GMR</b>	<b>(95% CI)</b>	<b>GMR</b>	<b>(95% CI)</b>
1	200	146	104	1.4	(1.08, 1.73)	1.4	(1.10, 1.78)
3	91	93	85	1.0	(0.81, 1.19)	1.1	(0.90, 1.32)
4	2833	2062	1295	1.4	(1.07, 1.77)	1.6	(1.19, 2.13)
5	269	199	162	1.4	(1.01, 1.80)	1.2	(0.93, 1.62)
6A <sup>†</sup>	4328	2593	213	1.7	(1.30, 2.15)	12.1	(8.63, 17.08)
6B	3212	1984	788	1.6	(1.24, 2.12)	2.5	(1.82, 3.48)
7F	1520	1120	405	1.4	(1.03, 1.79)	2.8	(1.98, 3.87)
9V	1726	1164	407	1.5	(1.11, 1.98)	2.9	(2.00, 4.08)
14	957	612	692	1.6	(1.16, 2.12)	0.9	(0.64, 1.21)
18C	1939	1726	925	1.1	(0.86, 1.47)	1.9	(1.39, 2.51)
19A	956	682	352	1.4	(1.16, 1.69)	1.9	(1.56, 2.41)
19F	599	517	539	1.2	(0.87, 1.54)	1.0	(0.72, 1.28)
23F	494	375	72	1.3	(0.94, 1.84)	5.2	(3.67, 7.33)

<sup>a</sup> Non-inferiority was defined as the lower limit of the 2-sided 95% CI for GMR was greater than 0.5.  
<sup>b</sup> Statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR was greater than 1.  
<sup>c</sup> For serotype 6A<sup>†</sup>, which is unique to Prevenar 13, a statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR being greater than 2.

In adults aged 60-64 years, OPA GMTs to Prevenar 13 were non-inferior to the OPA GMTs elicited to the 23-valent pneumococcal polysaccharide vaccine for the twelve serotypes common to both vaccines. For 9 serotypes, the OPA titers were shown to be statistically significantly greater in Prevenar 13 recipients.

In adults aged 50-59 years, OPA GMTs to all 13 serotypes in Prevenar 13 were non-inferior to the Prevenar 13 responses in adults aged 60-64 years. For 9 serotypes, immune responses were related to age, with adults in the 50-59 years group showing statistically significantly greater responses than adults aged 60-64 years.

In all adults ≥ 50 years who received a single dose of Prevenar 13, the OPA titers to serotype 6A were significantly greater than in adults ≥ 60 years who received a single dose of 23-valent pneumococcal polysaccharide vaccine.

One year after vaccination with Prevenar 13 OPA titers had declined compared to one month after vaccination, however, OPA titers for all serotypes remained higher than levels at baseline:

	OPA GMT levels at baseline	OPA GMT levels one year after Prevenar 13
Adults 50-59 years not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine	5 to 45	20 to 1234
Adults 60-64 years not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine	5 to 37	19 to 733

Table 6 shows OPA GMTs 1-month after a single dose of Prevenar 13 in 18-49 year olds compared to 60-64 year olds.

<b>Table 6: OPA GMTs in adults aged 18-49 years and 60-64 years given Prevenar 13<sup>a,b</sup></b>				
	<b>18-49 Years N=836-866</b>	<b>60-64 Years N=359-404</b>	<b>18-49 Years Relative to 60-64 Years</b>	
<b>Serotype</b>	GMT <sup>b</sup>	GMT <sup>b</sup>	GMR	(95% CI <sup>c</sup> )
1	353	146	2.4	(2.03, 2.87)
3	91	93	1.0	(0.84, 1.13)
4	4747	2062	2.3	(1.92, 2.76)
5	386	199	1.9	(1.55, 2.42)
6A	5746	2593	2.2	(1.84, 2.67)
6B	9813	1984	4.9	(4.13, 5.93)
7F	3249	1120	2.9	(2.41, 3.49)
9V	3339	1164	2.9	(2.34, 3.52)
14	2983	612	4.9	(4.01, 5.93)
18C	3989	1726	2.3	(1.91, 2.79)
19A	1580	682	2.3	(2.02, 2.66)
19F	1533	517	3.0	(2.44, 3.60)
23F	1570	375	4.2	(3.31, 5.31)

<sup>a</sup> Non-inferiority was defined as the lower limit of the 2-sided 95% CI for GMR was greater than 0.5.  
<sup>b</sup> Statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR was greater than 1.  
<sup>c</sup> Confidence intervals (CIs) for the ratio are back transformations of a confidence interval based on the Student t distribution for the mean difference of the logarithms of the measures.

In adults aged 18-49 years, OPA GMTs to all 13 serotypes in Prevenar 13 were non-inferior to the Prevenar 13 responses in adults aged 60-64 years.

One year after vaccination with Prevenar 13 OPA titers had declined compared to one month after vaccination, however OPA titers for all serotypes remained higher than levels at baseline.

	OPA GMT levels at baseline	OPA GMT levels one year after Prevenar 13
Adults 18-49 years not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine	5 to 186	23 to 2948

*Adults previously vaccinated with 23-valent pneumococcal polysaccharide vaccine*

Immune responses to Prevenar 13 and 23-valent pneumococcal polysaccharide vaccine were compared in a head to head trial in adults aged  $\geq 70$  years, who had received a single dose of pneumococcal polysaccharide vaccine at least 5 years before study vaccination.

Table 7 compares the OPA GMTs, 1-month post-dose, in pneumococcal polysaccharide vaccinated adults aged  $\geq 70$  years given a single dose of either Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine.

<b>Table 7: OPA GMTs in pneumococcal polysaccharide vaccinated adults aged <math>\geq 70</math> years given either Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine (PPSV23)<sup>a,b,c</sup></b>				
	<b>Prevenar 13 N=400-426</b>	<b>PPSV23 N=395-445</b>	<b>Prevenar OPA GMT Relative to PPSV23</b>	
<b>Serotype</b>	<b>OPA GMT</b>	<b>OPA GMT</b>	<b>GMR</b>	<b>(95% CI)</b>
1	81	55	1.5	(1.17, 1.88)
3	55	49	1.1	(0.91, 1.35)
4	545	203	2.7	(1.93, 3.74)
5	72	36	2.0	(1.55, 2.63)
6A <sup>†</sup>	903	94	9.6	(7.00, 13.26)
6B	1261	417	3.0	(2.21, 4.13)
7F	245	160	1.5	(1.07, 2.18)
9V	181	90	2.0	(1.36, 2.97)
14	280	285	1.0	(0.73, 1.33)
18C	907	481	1.9	(1.42, 2.50)
19A	354	200	1.8	(1.43, 2.20)
19F	333	214	1.6	(1.17, 2.06)
23F	158	43	3.7	(2.69, 5.09)

<sup>a</sup> Non-inferiority was defined as the lower limit of the 2-sided 95% CI for GMR was greater than 0.5.  
<sup>b</sup> Statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR was greater than 1.  
<sup>c</sup> For serotype 6A<sup>†</sup>, which is unique to Prevenar 13, a statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR greater than 2.

In adults vaccinated with pneumococcal polysaccharide vaccine at least 5 years prior to the clinical study, OPA GMTs to Prevenar 13 were non-inferior to the 23-valent pneumococcal polysaccharide vaccine responses for the 12 serotypes in common. Furthermore, in this study statistically significantly greater OPA GMTs were demonstrated for 10 of the 12 serotypes in common. Immune responses to serotype 6A were statistically significantly greater following vaccination with Prevenar 13 than after 23-valent pneumococcal polysaccharide vaccine.

One year after vaccination with Prevenar 13 in adults aged 70 years and over who were vaccinated with 23-valent pneumococcal polysaccharide vaccine, at least 5 years prior to study entry, OPA titers had declined compared to one month after vaccination, however, OPA titers for all serotypes remained higher than levels at baseline:

	<b>OPA GMT levels at baseline</b>	<b>OPA GMT levels one year after Prevenar 13</b>
Adults $\geq 70$ years vaccinated with 23-valent pneumococcal polysaccharide vaccine at least 5 years prior	9 to 122	18 to 381

### Immune responses in Special Populations

Individuals with the conditions described below have an increased risk of pneumococcal disease. The clinical relevance of the antibody levels elicited by Prevenar 13 in these special populations is unknown.

#### *Sickle cell disease*

An open label single arm study in France, Italy, UK, US, Lebanon, Egypt and Saudi Arabia with 2 doses of Prevenar 13 given 6 months apart was conducted in 158 children and adolescents  $\geq 6$  to  $< 18$  years of age with sickle cell disease who were previously vaccinated with one or more doses of

23-valent pneumococcal polysaccharide vaccine at least 6 months prior to enrollment. After the first vaccination, Prevenar 13 elicited antibody levels measured by both IgG GMCs and OPA GMTs that were statistically significantly higher when compared to levels prior to vaccination. After the second dose immune responses were comparable to those after the first dose. One year after the second dose, antibody levels measured by both IgG GMCs and OPA GMTs were higher than levels prior to the first dose of Prevenar 13, except for the IgG GMCs for serotypes 3 and 5 that were numerically similar.

*Additional Prevenar (7-valent) immunogenicity data: children with sickle cell disease*

The immunogenicity of Prevenar has been investigated in an open-label, multicentre study in 49 infants with sickle cell disease. Children were vaccinated with Prevenar (3 doses one month apart from the age of 2 months), 46 of these children also received a 23-valent pneumococcal polysaccharide vaccine at the age of 15-18 months. After primary immunisation, 95.6 % of the subjects had antibody levels of at least 0.35 µg/ml for all seven serotypes found in Prevenar. A significant increase was seen in the concentrations of antibodies against the seven serotypes after the polysaccharide vaccination, suggesting that immunological memory was well established.

*HIV infection*

*Children and adults not previously vaccinated with a pneumococcal vaccine*

HIV-infected children and adults with CD4  $\geq$  200 cells/ $\mu$ L (mean 717.0 cells/ $\mu$ L), viral load  $<$  50,000 copies/ml (mean 2090.0 copies/ml), free of active AIDS-related illness and not previously vaccinated with a pneumococcal vaccine received 3 doses of Prevenar 13. As per general recommendations, a single dose of 23-valent pneumococcal polysaccharide vaccine was subsequently administered. Vaccines were administered at 1 month intervals. Immune responses were assessed in 259-270 evaluable subjects approximately 1 month after each dose of vaccine. After the first dose, Prevenar 13 elicited antibody levels, measured by both IgG GMCs and OPA GMTs that were statistically significantly higher when compared to levels prior to vaccination. After the second and third dose of Prevenar 13, immune responses were similar or higher than those after the first dose.

*Adults previously vaccinated with 23-valent pneumococcal polysaccharide vaccine*

HIV-infected adults  $\geq$  18 years of age with CD4  $\geq$  200 cells/ $\mu$ L (mean 609.1 cells/ $\mu$ L) and viral load  $<$  50,000 copies/ml (mean 330.6 copies/ml), who were free of active AIDS-related illness and were previously vaccinated with 23-valent pneumococcal polysaccharide vaccine administered at least 6 months prior to enrollment, received 3 doses of Prevenar 13, at enrollment, 6 months, and 12 months after the first dose of Prevenar 13. Immune responses were assessed in 231-255 evaluable subjects approximately 1 month after each dose of Prevenar 13. After the first dose, Prevenar 13 elicited antibody levels measured by both IgG GMCs and OPA GMTs that were statistically significantly higher when compared to levels prior to vaccination. After the second and third dose of Prevenar 13, immune responses were comparable or higher than those after the first dose. In the study 162 subjects had received one prior dose of 23-valent pneumococcal polysaccharide vaccine, 143 subjects 2 prior doses and 26 subjects more than 2 prior doses of 23-valent polysaccharide vaccine. Subjects who received two or more previous doses of 23-valent pneumococcal polysaccharide vaccine showed a similar immune response compared with subjects who received a single previous dose.

*Haematopoietic stem cell transplant*

Children and adults with an allogeneic haematopoietic stem cell transplant (HSCT) at  $\geq$  2 years of age with complete haematologic remission of underlying disease or with very good partial remission in the case of lymphoma and myeloma received three doses of Prevenar 13 with an interval of at least 1 month between doses. The first dose was administered at 3 to 6 months after HSCT. A fourth (booster) dose of Prevenar 13 was administered 6 months after the third dose. As per general recommendations, a single dose of 23-valent pneumococcal polysaccharide vaccine was administered 1 month after the fourth dose of Prevenar 13. Immune responses as measured by IgG GMCs were assessed in 168-211 evaluable subjects approximately 1 month after vaccination. Prevenar 13 elicited increased antibody levels after each dose of Prevenar 13. Immune responses after the fourth dose of Prevenar 13 were significantly increased for all serotypes compared with after the third dose. Functional antibody titers (OPA titers) were not measured in this study.

## **5.2 Pharmacokinetic properties**

Not applicable.

## **5.3 Preclinical safety data**

Non-clinical data revealed no special hazard for humans based on conventional studies of safety pharmacology, single and repeated dose toxicity, local tolerance, and reproduction and developmental toxicity.

# **6. PHARMACEUTICAL PARTICULARS**

## **6.1 List of excipients**

Sodium chloride  
Succinic acid  
Polysorbate 80  
Water for injections

For adjuvant, see section 2.

## **6.2 Incompatibilities**

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

## **6.3 Shelf life**

3 years

## **6.4 Special precautions for storage**

Store in a refrigerator (2°C – 8°C). Do not freeze.

Prevenar 13 is stable at temperatures up to 25°C for four days. At the end of this period Prevenar 13 should be used or discarded. These data are intended to guide health care professionals in case of temporary temperature excursions.

## **6.5 Nature and contents of container**

0.5 ml suspension for injection in pre-filled syringe (Type I glass) with a plunger stopper (latex-free chlorobutyl rubber) and protective-tip cap (latex-free isoprene bromobutyl rubber).

Pack sizes of 1 and 10, with or without needle, and a multipack of 5 packs, each containing 10 pre-filled syringes, with or without needle.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal and other handling**

During storage, a white deposit and clear supernatant can be observed. This does not constitute a sign of deterioration.

The vaccine should be shaken well to obtain a homogeneous white suspension prior to expelling air from the syringe, and should be inspected visually for any particulate matter and/or variation of physical aspect prior to administration. Do not use if the content appears otherwise.

No special requirements for disposal.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Pfizer Limited  
Ramsgate Road  
Sandwich  
Kent CT13 9NJ  
United Kingdom

## **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/09/590/001  
EU/1/09/590/002  
EU/1/09/590/003  
EU/1/09/590/004  
EU/1/09/590/005  
EU/1/09/590/006

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 09 December 2009  
Date of latest renewal: 18 September 2014

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## 1. NAME OF THE MEDICINAL PRODUCT

Prevenar 13 suspension for injection in single dose vial  
Pneumococcal polysaccharide conjugate vaccine (13-valent, adsorbed)

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

1 dose (0.5 ml) contains:

Pneumococcal polysaccharide serotype 1 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 3 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 4 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 5 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 6A <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 6B <sup>1</sup>	4.4 µg
Pneumococcal polysaccharide serotype 7F <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 9V <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 14 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 18C <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 19A <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 19F <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 23F <sup>1</sup>	2.2 µg

<sup>1</sup>Conjugated to CRM<sub>197</sub> carrier protein, adsorbed on aluminium phosphate.

1 dose (0.5 ml) contains approximately 32 µg CRM<sub>197</sub> carrier protein and 0.125 mg aluminium.

### Excipients with known effect

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Suspension for injection in single dose vial.  
The vaccine is a homogeneous white suspension.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Active immunisation for the prevention of invasive disease, pneumonia and acute otitis media caused by *Streptococcus pneumoniae* in infants, children and adolescents from 6 weeks to 17 years of age.

Active immunisation for the prevention of invasive disease and pneumonia caused by *Streptococcus pneumoniae* in adults  $\geq$  18 years of age and the elderly.

See sections 4.4 and 5.1 for information on protection against specific pneumococcal serotypes.

The use of Prevenar 13 should be determined on the basis of official recommendations taking into consideration the risk of invasive disease and pneumonia in different age groups, underlying comorbidities as well as the variability of serotype epidemiology in different geographical areas.

## 4.2 Posology and method of administration

The immunisation schedules for Prevenar 13 should be based on official recommendations.

### Posology

#### Infants and children aged 6 weeks to 5 years

It is recommended that infants who receive a first dose of Prevenar 13 complete the vaccination course with Prevenar 13.

#### Infants aged 6 weeks-6 months

##### *Three-dose primary series*

The recommended immunisation series consists of four doses, each of 0.5 ml. The primary infant series consists of three doses, with the first dose usually given at 2 months of age and with an interval of at least 1 month between doses. The first dose may be given as early as six weeks of age. The fourth (booster) dose is recommended between 11 and 15 months of age.

##### *Two-dose primary series*

Alternatively, when Prevenar 13 is given as part of a routine infant immunisation programme, a series consisting of three doses, each of 0.5 ml, may be given. The first dose may be administered from the age of 2 months, with a second dose 2 months later. The third (booster) dose is recommended between 11 and 15 months of age (see section 5.1).

#### Preterm infants (<37 weeks gestation)

In preterm infants, the recommended immunisation series consists of four doses, each of 0.5 ml. The primary infant series consists of three doses, with the first dose given at 2 months of age and with an interval of at least 1 month between doses. The first dose may be given as early as six weeks of age. The fourth (booster) dose is recommended between 11 and 15 months of age (see sections 4.4 and 5.1).

#### Unvaccinated infants and children $\geq$ 7 months of age

##### *Infants aged 7-11 months*

Two doses, each of 0.5 ml, with an interval of at least 1 month between doses. A third dose is recommended in the second year of life.

##### *Children aged 12-23 months*

Two doses, each of 0.5 ml, with an interval of at least 2 months between doses (see section 5.1).

##### *Children and adolescents aged 2-17 years*

One single dose of 0.5 ml.

#### Prevenar 13 vaccine schedule for infants and children previously vaccinated with Prevenar (7-valent) (*Streptococcus pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F, and 23F)

Prevenar 13 contains the same 7 serotypes included in Prevenar, using the same carrier protein CRM<sub>197</sub>.

Infants and children who have begun immunisation with Prevenar may switch to Prevenar 13 at any point in the schedule.

##### *Young children (12-59 months) completely immunised with Prevenar (7-valent)*

Young children who are considered completely immunised with Prevenar (7-valent) should receive one dose of 0.5 ml of Prevenar 13 to elicit immune responses to the 6 additional serotypes. This dose

of Prevenar 13 should be administered at least 8 weeks after the final dose of Prevenar (7-valent) (see section 5.1).

#### *Children and adolescents 5–17 years*

Children 5 to 17 years of age may receive a single dose of Prevenar 13 if they have been previously vaccinated with one or more doses of Prevenar. This dose of Prevenar 13 should be administered at least 8 weeks after the final dose of Prevenar (7-valent) (see section 5.1).

#### Adults $\geq$ 18 years of age, and the elderly

One single dose.

The need for revaccination with a subsequent dose of Prevenar 13 has not been established.

Regardless of prior pneumococcal vaccination status, if the use of 23-valent pneumococcal polysaccharide vaccine is considered appropriate, Prevenar 13 should be given first (see sections 4.5 and 5.1).

#### Special Populations

Individuals who have underlying conditions predisposing them to invasive pneumococcal disease (such as sickle cell disease or HIV infection) including those previously vaccinated with one or more doses of 23-valent pneumococcal polysaccharide vaccine may receive at least one dose of Prevenar 13 (see section 5.1).

In individuals with an haematopoietic stem cell transplant (HSCT), the recommended immunisation series consists of four doses of Prevenar 13, each of 0.5 ml. The primary series consists of three doses, with the first dose given at 3 to 6 months after HSCT and with an interval of at least 1 month between doses. A fourth (booster) dose is recommended 6 months after the third dose (see section 5.1).

#### Method of administration

The vaccine should be given by intramuscular injection. The preferred sites are the anterolateral aspect of the thigh (vastus lateralis muscle) in infants or the deltoid muscle of the upper arm in children and adults.

### **4.3 Contraindications**

Hypersensitivity to the active substances, to any of the excipients listed in section 6.1, or to diphtheria toxoid.

As with other vaccines, the administration of Prevenar 13 should be postponed in subjects suffering from acute, severe febrile illness. However, the presence of a minor infection, such as a cold, should not result in the deferral of vaccination.

### **4.4 Special warnings and precautions for use**

Prevenar 13 must not be administered intravascularly.

As with all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of a rare anaphylactic event following the administration of the vaccine.

This vaccine should not be given as an intramuscular injection to individuals with thrombocytopaenia or any coagulation disorder that would contraindicate intramuscular injection, but may be given subcutaneously if the potential benefit clearly outweighs the risks (see section 5.1).

Prevenar 13 will only protect against *Streptococcus pneumoniae* serotypes included in the vaccine, and will not protect against other microorganisms that cause invasive disease, pneumonia, or otitis media.

As with any vaccine, Prevenar 13 may not protect all individuals receiving the vaccine from pneumococcal disease. For the most recent epidemiological information in your country you should consult with the relevant national organisation.

Individuals with impaired immune responsiveness, whether due to the use of immuno-suppressive therapy, a genetic defect, human immunodeficiency virus (HIV) infection, or other causes, may have reduced antibody response to active immunisation.

Safety and immunogenicity data are available for a limited number of individuals with sickle cell disease, HIV infection, or with an haematopoietic stem cell transplant (see section 5.1). Safety and immunogenicity data for Prevenar 13 are not available for individuals in other specific immuno-compromised groups (e.g., malignancy or nephrotic syndrome) and vaccination should be considered on an individual basis.

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially 'sodium-free'.

#### Infants and children aged 6 weeks to 5 years

In clinical studies, Prevenar 13 elicited an immune response to all thirteen serotypes included in the vaccine. The immune response for serotype 3 following the booster dose was not increased above the levels seen after the infant vaccination series; the clinical relevance of this observation regarding the induction of serotype 3 immune memory is unknown (see section 5.1).

The proportions of functional antibody responders (OPA titres  $\geq 1:8$ ) to serotypes 1, 3 and 5 were high. However, the OPA geometric mean titres were lower than those against each of the remaining additional vaccine serotypes; the clinical relevance of this observation for protective efficacy is unknown (see section 5.1).

Limited data have demonstrated that Prevenar 7-valent (three-dose primary series) induces an acceptable immune response in infants with sickle cell disease with a safety profile similar to that observed in non-high-risk groups (see section 5.1).

Children younger than 2 years old should receive the appropriate-for-age Prevenar 13 vaccination series (see section 4.2). The use of pneumococcal conjugate vaccine does not replace the use of 23-valent pneumococcal polysaccharide vaccines in children  $\geq 2$  years of age with conditions (such as sickle cell disease, asplenia, HIV infection, chronic illness, or those who are immuno-compromised) placing them at higher risk for invasive disease due to *Streptococcus pneumoniae*. Whenever recommended, children at risk who are  $\geq 24$  months of age and already primed with Prevenar 13 should receive 23-valent pneumococcal polysaccharide vaccine. The interval between the 13-valent pneumococcal conjugate vaccine (Prevenar 13) and the 23-valent pneumococcal polysaccharide vaccine should not be less than 8 weeks. There are no data available to indicate whether the administration of 23-valent pneumococcal polysaccharide vaccine to unprimed children or to children primed with Prevenar 13 might result in hyporesponsiveness to further doses of Prevenar 13.

The potential risk of apnoea and the need for respiratory monitoring for 48-72 h should be considered when administering the primary immunisation series to very premature infants (born  $\leq 28$  weeks of gestation), and particularly for those with a previous history of respiratory immaturity. As the benefit of vaccination is high in this group of infants, vaccination should not be withheld or delayed.

For vaccine serotypes, protection against otitis media is expected to be lower than protection against invasive disease. As otitis media is caused by many organisms other than pneumococcal serotypes represented in the vaccine, protection against all otitis media is expected to be low (see section 5.1).

When Prevenar 13 is administered concomitantly with Infanrix hexa (DTPa-HBV-IPV/Hib), the rates of febrile reactions are similar to those seen with concomitant administration of Prevenar (7-valent) and Infanrix hexa (see section 4.8). Increased reporting rates of convulsions (with or without fever)

and hypotonic hyporesponsive episode (HHE) were observed with concomitant administration of Prevenar 13 and Infanrix hexa (see section 4.8).

Antipyretic treatment should be initiated according to local treatment guidelines for children with seizure disorders or with a prior history of febrile seizures and for all children receiving Prevenar 13 simultaneously with vaccines containing whole cell pertussis.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

##### Infants and children aged 6 weeks to 5 years

Prevenar 13 can be given concomitantly with any of the following vaccine antigens, either as monovalent or combination vaccines: diphtheria, tetanus, acellular or whole cell pertussis, *Haemophilus influenzae* type b, inactivated poliomyelitis, hepatitis B (see section 4.4 for guidance on Infanrix hexa), meningococcal serogroup C, measles, mumps, rubella, varicella and rotavirus vaccine.

Prevenar 13 can also be given concomitantly between 12-23 months with the tetanus toxoid conjugated meningococcal polysaccharide serogroups A, C, W and Y vaccine to children who have been adequately primed with Prevenar 13 (as per local recommendations).

Data from a postmarketing clinical study evaluating the impact of prophylactic use of antipyretics (ibuprofen and paracetamol) on the immune response to Prevenar 13 suggest that administration of paracetamol concomitantly or within the same day of vaccination may reduce the immune response to Prevenar 13 after the infant series. Responses to the booster dose administered at 12 months were unaffected. The clinical significance of this observation is unknown.

##### Children and adolescents 6 to 17 years of age

No data are currently available regarding concomitant use with other vaccines.

##### Adults 18 to 49 years of age

No data are available regarding concomitant use with other vaccines.

##### Adults aged 50 years and older

Prevenar 13 may be administered concomitantly with the seasonal trivalent inactivated influenza vaccine (TIV).

In two studies conducted in adults aged 50-59 and 65 years and older, it was demonstrated that Prevenar 13 may be given concomitantly with trivalent inactivated influenza vaccine (TIV). The responses to all three TIV antigens were comparable when TIV was given alone or concomitantly with Prevenar 13.

When Prevenar 13 was given concomitantly with TIV, the immune responses to Prevenar 13 were lower compared to when Prevenar 13 was given alone, however, there was no long-term impact on circulating antibody levels.

In a third study in adults aged 50-93 years, it was demonstrated that Prevenar 13 may be given concomitantly with the seasonal quadrivalent inactivated influenza vaccine (QIV). The immune responses to all four QIV strains were noninferior when Prevenar 13 was given concomitantly with QIV compared to when QIV was given alone.

The immune responses to Prevenar 13 were noninferior when Prevenar 13 was given concomitantly with QIV compared to when Prevenar 13 was given alone. As with concomitant administration with

trivalent vaccines, immune responses to some pneumococcal serotypes were lower when both vaccines were given concomitantly.

Concomitant use with other vaccines has not been investigated.

Different injectable vaccines should always be given at different vaccination sites.

Concomitant administration of Prevenar 13 and 23-valent pneumococcal polysaccharide vaccine has not been studied. In clinical studies when Prevenar 13 was given 1 year after 23-valent pneumococcal polysaccharide vaccine the immune responses were lower for all serotypes compared to when Prevenar 13 was given to subjects not previously immunised with 23-valent pneumococcal polysaccharide vaccine. The clinical significance of this is unknown.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

There are no data from the use of pneumococcal 13-valent conjugate vaccine in pregnant women. Therefore the use of Prevenar 13 should be avoided during pregnancy.

##### Breast-feeding

It is unknown whether pneumococcal 13-valent conjugate vaccine is excreted in human milk.

##### Fertility

Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3).

#### **4.7 Effects on ability to drive and use machines**

Prevenar 13 has no or negligible influence on the ability to drive and use machines. However, some of the effects mentioned under section 4.8 “Undesirable effects” may temporarily affect the ability to drive or use machines.

#### **4.8 Undesirable effects**

Analysis of postmarketing reporting rates suggests a potential increased risk of convulsions, with or without fever, and HHE when comparing groups which reported use of Prevenar 13 with Infanrix hexa to those which reported use of Prevenar 13 alone.

Adverse reactions reported in clinical studies or from the postmarketing experience for all age groups are listed in this section per system organ class, in decreasing order of frequency and seriousness. The frequency is defined as follows: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ), very rare ( $< 1/10,000$ ), not known (cannot be estimated from available data).

##### Infants and children aged 6 weeks to 5 years

The safety of the vaccine was assessed in controlled clinical studies where 14,267 doses were given to 4,429 healthy infants from 6 weeks of age at first vaccination and 11-16 months of age at booster dose. In all infant studies, Prevenar 13 was co-administered with routine paediatric vaccines (see section 4.5).

Safety in 354 previously unvaccinated children (7 months to 5 years of age) was also assessed.

The most commonly reported adverse reactions in children 6 weeks to 5 years of age were vaccination-site reactions, fever, irritability, decreased appetite, and increased and/or decreased sleep.

In a clinical study in infants vaccinated at 2, 3, and 4 months of age, fever  $\geq 38^{\circ}\text{C}$  was reported at higher rates among infants who received Prevenar (7-valent) concomitantly with Infanrix hexa (28.3% to 42.3%) than in infants receiving Infanrix hexa alone (15.6% to 23.1%). After a booster dose at 12 to 15 months of age, fever  $\geq 38^{\circ}\text{C}$  was reported in 50.0% of infants who received Prevenar (7-valent) and Infanrix hexa at the same time as compared to 33.6% of infants receiving Infanrix hexa alone. These reactions were mostly moderate (less than or equal to  $39^{\circ}\text{C}$ ) and transient.

An increase in vaccination-site reactions was reported in children older than 12 months compared to rates observed in infants during the primary series with Prevenar 13.

#### *Adverse reactions from clinical studies*

In clinical studies, the safety profile of Prevenar 13 was similar to Prevenar. The following frequencies are based on adverse reactions assessed in Prevenar 13 clinical studies:

#### Immune system disorders:

Rare: Hypersensitivity reaction including face oedema, dyspnoea, bronchospasm

#### Nervous system disorders:

Uncommon: Convulsions (including febrile convulsions)

Rare: Hypotonic-hyporesponsive episode

#### Gastrointestinal disorders:

Very common: Decreased appetite

Common: Vomiting; diarrhoea

#### Skin and subcutaneous tissue disorders:

Common: Rash

Uncommon: Urticaria or urticaria-like rash

#### General disorders and administration site conditions:

Very common: Pyrexia; irritability; any vaccination-site erythema, induration/swelling or pain/tenderness; somnolence; poor quality sleep

Vaccination-site erythema or induration/swelling 2.5 cm–7.0 cm (after the booster dose and in older children [age 2 to 5 years])

Common: Pyrexia  $> 39^{\circ}\text{C}$ ; vaccination-site movement impairment (due to pain); vaccination-site erythema or induration/swelling 2.5 cm–7.0 cm (after infant series)

Uncommon: Vaccination-site erythema, induration/swelling  $> 7.0$  cm; crying

#### *Adverse reactions from Prevenar 13 postmarketing experience*

Although the following adverse drug reactions were not observed in the Prevenar 13 clinical studies in infants and children, the following are considered adverse drug reactions for Prevenar 13 as they were reported in the postmarketing experience. Because these reactions were derived from spontaneous reports, the frequencies could not be determined and are thus considered as not known.

#### Blood and lymphatic system disorders:

Lymphadenopathy (localised to the region of the vaccination-site)

#### Immune system disorders:

Anaphylactic/anaphylactoid reaction including shock; angioedema

#### Skin and subcutaneous tissue disorders:

Erythema multiforme

General disorders and administration site conditions:

Vaccination-site urticaria; vaccination-site dermatitis; vaccination-site pruritus; flushing

Additional information in special populations:

Apnoea in very premature infants ( $\leq 28$  weeks of gestation) (see section 4.4).

Children and adolescents aged 6 to 17 years of age

Safety was evaluated in 592 children (294 children aged 5 to 10 years previously immunised with at least one dose of Prevenar and 298 children aged 10 to 17 years who had not received a pneumococcal vaccine).

The most common adverse events in children and adolescents 6 to 17 years of age were:

Nervous system disorders:

Common: Headaches

Gastrointestinal disorders:

Very common: Decreased appetite

Common: Vomiting; diarrhoea

Skin and subcutaneous tissue disorders:

Common: Rash; urticaria or urticaria-like rash

General disorders and administration site conditions:

Very common: Irritability; any vaccination-site erythema; induration/swelling or pain/tenderness; somnolence; poor quality sleep; vaccination-site tenderness (including impaired movement)

Common: Pyrexia

Other adverse events previously observed in infants and children 6 weeks to 5 years of age may also be applicable to this age group but were not seen in this study possibly due to the small sample size.

Additional information in special populations

Children and adolescents with sickle cell disease, HIV infection, or an haematopoietic stem cell transplant have similar frequencies of adverse reactions, except that headaches, vomiting, diarrhoea, pyrexia, fatigue, arthralgia, and myalgia were very common.

Adults  $\geq 18$  years and elderly

Safety was assessed in 7 clinical studies including 91,593 adults ranging in age from 18 to 101 years. Prevenar 13 was administered to 48,806 adults; 2,616 (5.4%) aged 50 to 64 years, and 45,291 (92.8%) aged 65 years and older. One of the 7 studies included a group of adults ( $n=899$ ) ranging from 18 to 49 years who received Prevenar 13 and who were not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine. Of the Prevenar 13 recipients 1,916 adults were previously vaccinated with the 23-valent pneumococcal polysaccharide vaccine at least 3 years prior to study vaccination, and 46,890 were 23-valent pneumococcal polysaccharide vaccine unvaccinated.

A trend to lower frequency of adverse reactions was associated with greater age; adults  $> 65$  years of age (regardless of prior pneumococcal vaccination status) reported fewer adverse reactions than younger adults, with adverse reactions generally most common in the youngest adults, 18 to 29 years of age.

Overall, the frequency categories were similar for all age groups, with the exception of vomiting which was very common ( $\geq 1/10$ ) in adults aged 18 to 49 years and common ( $\geq 1/100$  to  $< 1/10$ ) in all other age groups, and pyrexia was very common in adults aged 18 to 29 years and common in all other age groups. Severe vaccination-site pain/tenderness and severe limitation of arm movement was very common in adults 18 to 39 years and common in all other age groups.

Adverse reactions from clinical studies

Local reactions and systemic events were solicited daily after each vaccination for 14 days in 6 studies and 7 days in the remaining study. The following frequencies are based on adverse reactions assessed in Prevenar 13 clinical studies in adults:

Metabolism and nutrition disorders:

Very common: Decreased appetite

Nervous system disorders:

Very common: Headaches

Gastrointestinal disorders:

Very common: Diarrhoea; vomiting (in adults aged 18 to 49 years)

Common: Vomiting (in adults aged 50 years and over)

Uncommon: Nausea

Immune system disorders:

Uncommon: Hypersensitivity reaction including face oedema, dyspnoea, bronchospasm

Skin and subcutaneous tissue disorders:

Very common: Rash

General disorders and administration site conditions:

Very common: Chills; fatigue; vaccination-site erythema; vaccination-site induration/swelling; vaccination-site pain/tenderness (severe vaccination-site pain/tenderness very common in adults aged 18 to 39 years); limitation of arm movement (severe limitation of arm movements very common in adults aged 18 to 39 years)

Common: Pyrexia (very common in adults aged 18 to 29 years)

Uncommon: Lymphadenopathy localized to the region of the vaccination-site

Musculoskeletal and connective tissue disorders:

Very common: Arthralgia; myalgia

Overall, no significant differences in frequencies of adverse reactions were seen when Prevenar 13 was given to adults previously vaccinated with the pneumococcal polysaccharide vaccine.

Additional information in special populations

Adults with HIV infection have similar frequencies of adverse reactions, except that pyrexia and vomiting were very common and nausea common.

Adults with an haematopoietic stem cell transplant have similar frequencies of adverse reactions, except that pyrexia and vomiting were very common.

Higher frequency in some solicited systemic reactions was observed when Prevenar 13 was administered concomitantly with trivalent inactivated influenza vaccine (TIV) compared to TIV given alone (headache, chills, rash, decreased appetite, arthralgia, and myalgia) or Prevenar 13 given alone (headache, fatigue, chills, decreased appetite, and arthralgia).

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

## **4.9 Overdose**

Overdose with Prevenar 13 is unlikely due to its presentation as a single dose vial. However, in infants and children there have been reports of overdose with Prevenar 13 defined as subsequent doses administered closer than recommended to the previous dose. In general, adverse events reported with overdose are consistent with those that have been reported with doses given in the recommended paediatric schedules of Prevenar 13.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: vaccines, pneumococcal vaccines; ATC code: J07AL02

Prevenar 13 contains the 7 pneumococcal capsular polysaccharides that are in Prevenar (4, 6B, 9V, 14, 18C, 19F, 23F) plus 6 additional polysaccharides (1, 3, 5, 6A, 7F, 19A) all conjugated to CRM<sub>197</sub> carrier protein.

#### Burden of disease

##### *Infants and children aged 6 weeks to 5 years*

Based on serotype surveillance in Europe performed before the introduction of Prevenar, Prevenar 13 is estimated to cover 73-100% (depending on the country) of serotypes causing invasive pneumococcal disease (IPD) in children less than 5 years of age. In this age group, serotypes 1, 3, 5, 6A, 7F, and 19A account for 15.6% to 59.7% of invasive disease, depending on the country, the time period studied, and the use of Prevenar.

Acute otitis media (AOM) is a common childhood disease with different aetiologies. Bacteria can be responsible for 60-70% of clinical episodes of AOM. *S. pneumoniae* is one of the most common causes of bacterial AOM worldwide.

Prevenar 13 is estimated to cover over 90% of serotypes causing antimicrobial-resistant IPD.

##### *Children and adolescents aged 6 to 17 years*

In children and adolescents aged 6 to 17 years, the incidence of pneumococcal disease is low, however, there is an increased risk of morbidity and mortality in those with underlying comorbidities.

##### *Adults ≥ 18 years and the elderly*

Pneumonia is the most common clinical presentation of pneumococcal disease in adults.

The reported incidence of community-acquired pneumonia (CAP) and IPD in Europe varies by country, increases with age from 50 years and is highest in individuals aged ≥ 65 years. *S. pneumoniae* is the most frequent cause of CAP, and is estimated to be responsible for approximately 30% of all CAP cases requiring hospitalisation in adults in developed countries.

Bacteraemic pneumonia (approximately 80% of IPD in adults), bacteraemia without a focus, and meningitis are the most common manifestations of IPD in adults. Based on surveillance data following

the introduction of Prevenar but before the introduction of Prevenar 13 in childhood vaccination programmes, the pneumococcal serotypes in Prevenar 13 may be responsible for at least 50 – 76% (depending on country) of IPD in adults.

The risk for CAP and IPD in adults also increases with chronic underlying medical conditions, specifically, anatomical or functional asplenia, diabetes mellitus, asthma, chronic cardiovascular, pulmonary, kidney or liver disease, and it is highest in those who are immune-suppressed such as those with malignant haematological diseases or HIV infection.

#### Prevenar 13 immunogenicity clinical studies in infants, children and adolescents

The protective efficacy of Prevenar 13 against IPD has not been studied. As recommended by the World Health Organization (WHO) the assessment of potential efficacy against IPD in infants and young children has been based on a comparison of immune responses to the seven common serotypes shared between Prevenar 13 and Prevenar, for which protective efficacy has been proven (for Prevenar (7-valent) efficacy in infants and children, see below). Immune responses to the additional 6 serotypes were also measured.

#### Immune responses following a three-dose primary infant series

Clinical studies have been conducted in a number of European countries and the US using a range of vaccination schedules, including two randomised non-inferiority studies (Germany using a 2, 3, 4 month primary series [006] and US using a 2, 4, 6 month primary series [004]). In these two studies pneumococcal immune responses were compared using a set of non-inferiority criteria including the percentage of subjects with serum anti-polysaccharide serotype-specific IgG  $\geq 0.35$   $\mu\text{g/ml}$  one month after the primary series and the comparison of IgG geometric mean concentrations (ELISA GMCs); in addition, functional antibody titres (OPA) between subjects receiving Prevenar 13 and Prevenar were compared. For the six additional serotypes, these values were compared with the lowest response among all of the seven common serotypes in the Prevenar recipients.

The non-inferiority immune response comparisons for study 006, based on the proportion of infants achieving anti-polysaccharide IgG concentrations  $\geq 0.35$   $\mu\text{g/ml}$ , are shown in Table 1. Results for study 004 were similar. Prevenar 13 non-inferiority (lower bound of the 95% CI for the difference in percentage of responders at 0.35  $\mu\text{g/ml}$  between groups was  $>-10\%$ ) was demonstrated for all 7 common serotypes, except for serotype 6B in study 006 and serotypes 6B and 9V in study 004, which missed by a small margin. All seven common serotypes met pre-defined non-inferiority criteria for IgG ELISA GMCs. Prevenar 13 elicited comparable, although slightly lower, antibody levels than Prevenar for the 7 common serotypes. The clinical relevance of these differences is not known.

Non-inferiority was met for the 6 additional serotypes based on the proportion of infants achieving antibody concentrations  $\geq 0.35$   $\mu\text{g/ml}$  and comparison of IgG ELISA GMCs in study 006 and was met for 5 out of the 6 serotypes, with the exception of serotype 3 for study 004. For serotype 3, the percentages of Prevenar 13 recipients with serum IgG  $\geq 0.35$   $\mu\text{g/ml}$  were 98.2% (study 006) and 63.5% (study 004).

<b>Table 1: Comparison of the proportion of subjects achieving a pneumococcal anti-polysaccharide IgG antibody concentration <math>\geq 0.35</math> <math>\mu\text{g/ml}</math> after dose 3 of the infant series – study 006</b>			
<b>Serotypes</b>	<b>Prevenar 13 % (N=282-285)</b>	<b>7-valent Prevenar % (N=277-279)</b>	<b>Difference (95 % CI)</b>
<b>7-valent Prevenar serotypes</b>			
4	98.2	98.2	0.0 (-2.5, 2.6)
6B	77.5	87.1	-9.6 (-16.0, -3.3)
9V	98.6	96.4	2.2 (-0.4, 5.2)
14	98.9	97.5	1.5 (-0.9, 4.1)
18C	97.2	98.6	-1.4 (-4.2, 1.2)
19F	95.8	96.0	-0.3 (-3.8, 3.3)
23F	88.7	89.5	-0.8 (-6.0, 4.5)
<b>Additional serotypes in Prevenar 13</b>			
1	96.1	87.1*	9.1 (4.5, 13.9)
3	98.2	87.1	11.2 (7.0, 15.8)
5	93.0	87.1	5.9 (0.8, 11.1)
6A	91.9	87.1	4.8 (-0.3, 10.1)
7F	98.6	87.1	11.5 (7.4, 16.1)
19A	99.3	87.1	12.2 (8.3, 16.8)

\*The serotype in Prevenar with the lowest percent response rate was 6B in study 006 (87.1 %).

Prevenar 13 elicited functional antibodies to all 13 vaccine serotypes in studies 004 and 006. For the 7 common serotypes there were no differences between groups in the proportion of subjects with OPA titres  $\geq 1:8$ . For each of the seven common serotypes,  $> 96\%$  and  $> 90\%$  of the Prevenar 13 recipients reached an OPA titre  $\geq 1:8$  one month after the primary series in studies 006 and 004, respectively.

For each of the 6 additional serotypes, Prevenar 13 elicited OPA titres  $\geq 1:8$  in 91.4% to 100% of vaccinees one month after the primary series in studies 004/006. The functional antibody (OPA) geometric mean titres for serotypes 1, 3 and 5 were lower than the titres for each of the other additional serotypes; the clinical relevance of this observation for protective efficacy is unknown.

#### Immune responses following a two-dose primary infant series

The immunogenicity after two doses in infants has been documented in four studies. The proportion of infants achieving a pneumococcal anti-capsular polysaccharide IgG concentration  $\geq 0.35$   $\mu\text{g/ml}$  one month after the second dose ranged from 79.6% to 98.5% across 11 of the 13 vaccine serotypes. Smaller proportions of infants achieved this antibody concentration threshold for serotype 6B (27.9% to 57.3%) and 23F (55.8% to 68.1%) for all studies using a 2, 4 month regimen, compared to 58.4% for serotype 6B and 68.6% for 23F for a study using a 3, 5 month regimen. After the booster dose, all vaccine serotypes including 6B and 23F had immune responses consistent with adequate priming with a two-dose primary series. In a UK study, the functional antibody (OPA) responses were comparable for all serotypes including 6B and 23F in the Prevenar and Prevenar 13 arms after the primary series at two and four months of age and after the booster dose at 12 months of age. For Prevenar 13 recipients, the proportion of responders with an OPA titre  $\geq 1:8$  was at least 87% following the infant series, and at least 93% following the booster dose. The OPA geometric mean titres for serotypes 1, 3 and 5 were lower than the titres for each of the other additional serotypes; the clinical relevance of this observation is unknown.

#### Booster responses following two-dose and three-dose primary infant series

Following the booster dose, antibody concentrations increased from the pre-booster level for all 13 serotypes. Post-booster antibody concentrations were higher for 12 serotypes than those achieved after the infant primary series. These observations are consistent with adequate priming (the induction of immunologic memory). The immune response for serotype 3 following the booster dose was not increased above the levels seen after the infant vaccination series; the clinical relevance of this observation regarding the induction of serotype 3 immune memory is unknown.

Antibody responses to booster doses following two-dose or three-dose infant primary series were comparable for all 13 vaccine serotypes.

For children aged from 7 months to 5 years, age appropriate catch-up immunisation schedules (as described in section 4.2) result in levels of anti-capsular polysaccharide IgG antibody responses to each of the 13 serotypes that are at least comparable to those of a three-dose primary series in infants.

Antibody persistence and immunological memory were evaluated in a study in healthy children who received a single dose of Prevenar 13 at least 2 years after they had been previously immunised with either 4 doses of Prevenar, a 3-dose infant series of Prevenar followed by Prevenar 13 at 12 months of age, or 4 doses of Prevenar 13.

The single dose of Prevenar 13, in children approximately 3.4 years of age regardless of previous vaccination history with Prevenar or Prevenar 13, induced a robust antibody response for both the 7 common serotypes and the 6 additional serotypes in Prevenar 13.

Since the introduction of 7-valent Prevenar in 2000, pneumococcal disease surveillance data have not shown that the immunity elicited by Prevenar in infancy has waned over time.

### Preterm Infants

Safety and immunogenicity of Prevenar 13 given at 2, 3, 4 and 12 months was assessed in approximately 100 prematurely born infants (mean Estimated Gestational Age [EGA], 31 weeks; range, 26 to 36 weeks) and compared with approximately 100 infants born at term (mean EGA, 39 weeks; range, 37 to 42 weeks).

Immune responses in preterm and term infants were compared using the proportion of subjects achieving a pneumococcal polysaccharide IgG binding antibody concentration  $\geq 0.35 \mu\text{g/ml}$  1 month after the infant series, the approach used for immunogenicity comparisons of Prevenar 13 to Prevenar based on WHO guidelines.

More than 85% achieved a pneumococcal polysaccharide IgG binding antibody concentration  $\geq 0.35 \mu\text{g/ml}$  1 month after the infant series, except for serotypes 5 (71.7%), 6A (82.7%), and 6B (72.7%) in the preterm group. For these 3 serotypes, the proportion of responders among preterm infants was significantly lower than among term infants. Approximately one month after the toddler dose, the proportion of subjects in each group achieving this same antibody concentration threshold was  $>97\%$ , except for serotype 3 (71% in preterm infants and 79% in term infants). It is unknown whether immunological memory to all serotypes is induced in pre-term infants. In general, serotype-specific IgG GMCs were lower for preterm infants than term infants.

After the infant series, OPA GMTs were similar in preterm infants compared to term infants except for serotype 5, which was lower in preterm infants. OPA GMTs after the toddler dose relative to those after the infant series were similar or lower for 4 serotypes (4, 14, 18C, and 19F) and were statistically significantly higher for 6 of 13 serotypes (1, 3, 5, 7F, 9V, and 19A) in preterm infants compared to 10 of 13 serotypes (1, 3, 4, 5, 6A, 7F, 9V, 18C, 19A, and 23F) in term infants.

### Children (12-59 months) completely immunised with Prevenar (7-valent)

Following administration of a single dose of Prevenar 13 to children (12-59 months) who are considered completely immunised with Prevenar (7-valent) (either 2 or 3 dose primary series plus booster), the proportion achieving serum IgG levels  $\geq 0.35 \mu\text{g/ml}$  and OPA titres  $\geq 1:8$  was at least 90%. However, 3 (serotypes 1, 5 and 6A) of the 6 additional serotypes showed lower IgG GMC and OPA GMT when compared with children who had received at least one previous vaccination with Prevenar 13. The clinical relevance of the lower GMCs and GMTs is currently unknown.

### Unvaccinated Children (12-23 months)

Studies in unvaccinated children (12 -23 months) with Prevenar (7-valent) demonstrated that 2 doses were required to achieve serum IgG concentrations for 6B and 23F similar to those induced by a 3-dose infant series.

### Children and Adolescents 5 to 17 years of age

In an open-label study in 592 healthy children and adolescents including those with asthma (17.4%) who may be predisposed to pneumococcal infection, Prevenar 13 elicited immune responses to all 13 serotypes. A single dose of Prevenar 13 was given to children 5 to 10 years of age previously vaccinated with at least 1 dose of Prevenar, and children and adolescents 10 to 17 years of age who had never received a pneumococcal vaccine.

In both the children 5 to 10 years of age and children and adolescents aged 10 to 17 years, the immune response to Prevenar 13 was non inferior to Prevenar for the 7 common serotypes and to Prevenar 13 for the 6 additional serotypes compared to the immune response after the fourth dose in infants vaccinated at 2, 4, 6 and 12-15 months of age as measured by serum IgG.

In children and adolescents aged 10 to 17 years of age OPA GMTs 1 month after vaccination were noninferior to OPA GMTs in the 5 to 10 year old age group for 12 of the 13 serotypes (except serotype 3).

### Immune responses after subcutaneous administration

Subcutaneous administration of Prevenar 13 was evaluated in a non-comparative study in 185 healthy Japanese infants and children who received 4 doses at 2, 4, 6 and 12-15 months of age. The study demonstrated that safety and immunogenicity were generally comparable with observations made in studies of intramuscular administration.

### Prevenar 13 Effectiveness

#### *Invasive Pneumococcal Disease*

Data published by Public Health England showed that, four years after the introduction of Prevenar as a two dose primary infant series with booster dose in the second year of life and with a 94% vaccine uptake, there was a 98% (95% CI 95; 99) reduction in disease caused by the 7 vaccine serotypes in England and Wales. Subsequently, four years following the switch to Prevenar 13, the additional reduction in incidence of IPD due to the 7 serotypes in Prevenar ranged from 76% in children less than 2 years of age to 91% in children 5-14 years of age. The serotype specific reductions for each of the 5 additional serotypes in Prevenar 13 (no cases of serotype 5 IPD were observed) by age group are shown in Table 2 and ranged from 68% (serotype 3) to 100% (serotype 6A) for children less than 5 years of age. Significant incidence reductions were also observed in older age groups who had not been vaccinated with Prevenar 13 (indirect effect).

<b>Table 2: Serotype specific number of cases and incidence reductions of IPD in 2013/14 compared to 2008/09-2009/10 (2008/10) by age in England and Wales</b>									
	<b>&lt;5 years of age</b>			<b>5 to 64 years of age</b>			<b>≥65 years of age</b>		
	2008-10 <sup>§</sup>	2013/14 <sup>§</sup>	% Incidence reduction (95% CI*)	2008-10 <sup>§</sup>	2013/14 <sup>§</sup>	% Incidence reduction (95% CI*)	2008-10 <sup>§</sup>	2013/14 <sup>§</sup>	% Incidence reduction (95% CI*)
<b>Additional serotypes covered by Prevenar 13</b>									
<b>1</b>	59 (54)	5 (5)	91% (98%; 68%)**	458 (382)	77 (71)	83% (88%; 74%)**	102 (89)	13 (13)	87% (94%; 72%)**
<b>3</b>	26 (24)	8 (8)	68% (89%; 6%)	178 (148)	73 (68)	59% (72%; 38%)**	256 (224)	143 (146)	44% (57%; 27%)**
<b>6A</b>	10 (9)	0 (0)	100% (100%; 62%)**	53 (44)	5 (5)	90% (97%; 56%)**	94 (82)	5 (5)	95% (99%; 81%)**
<b>7F</b>	90 (82)	8 (8)	91% (97%; 74%)**	430 (361)	160 (148)	63% (71%; 50%)**	173 (152)	75 (77)	56% (70%; 37%)**
<b>19A</b>	85 (77)	7 (7)	91% (97%; 75%)**	225 (191)	104 (97)	54% (65%; 32%)**	279 (246)	97 (99)	65% (75%; 53%)**
<sup>§</sup> Corrected for proportion of samples serotyped, missing age, denominator compared with 2009/10, and for the trend in total invasive pneumococcal disease up to 2009/10 (after which no trend correction was applied). * 95% CI inflated from a Poisson interval based on over-dispersion of 2.1 seen from modelling of 2000-06 pre-Prevenar all IPD data. ** p<0.005 to cover 6A where p=0.002									

### *Otitis media (OM)*

In a published study performed in Israel, using a 2-dose primary series plus booster dose in the second year of life, the impact of Prevenar 13 on OM was documented in a population-based active-surveillance system with tympanocentesis culturing of middle ear fluid in Israeli children less than 2 years of age with OM.

Following the introduction of Prevenar and subsequently Prevenar 13 there was a decline in incidence from 2.1 to 0.1 cases per 1000 children (95%) for the Prevenar serotypes plus serotype 6A and a decline in incidence from 0.9 to 0.1 cases per 1000 children (89%) for the additional serotypes 1, 3, 5, 7F, and 19A in Prevenar 13. The annual overall pneumococcal incidence of OM declined from 9.6 to 2.1 cases per 1000 children (78%) between July 2004 (prior to the introduction of Prevenar) and June 2013 (post Prevenar 13 introduction).

### *Pneumonia*

In a multicenter observational study in France comparing the periods before and after the switch from Prevenar to Prevenar 13, there was 16% (2060 to 1725 cases) reduction in all community acquired pneumonia (CAP) cases in emergency departments in children 1 month to 15 years of age. Reductions were 53% (167 to 79 cases) ( $p<0.001$ ) for CAP cases with pleural effusion and 63% (64 to 24 cases) ( $p<0.001$ ) for microbiologically confirmed pneumococcal CAP cases. In the second year after the introduction of Prevenar 13 the total number of CAP cases due to the 6 additional vaccine serotypes in Prevenar 13 was reduced from 27 to 7 isolates (74%).

The decrease in all cause pneumonia cases was most pronounced in the younger vaccinated age groups with a decrease of 31.8% (757 to 516 cases) and 16.6% (833 to 695 cases) in the age groups <2 years and 2 to 5 years, respectively. The incidence in older, predominantly non-vaccinated children (>5 years) did not change over the duration of the study.

In an ongoing surveillance system (2004 to 2013) to document the impact of Prevenar and subsequently Prevenar 13 on CAP in children less than 5 years in Southern Israel using a 2 dose primary series with a booster dose in the second year of life, there was a reduction of 68% (95% CI 73; 61) in outpatient visits and 32% (95% CI 39; 22) in hospitalizations for alveolar CAP following the introduction of Prevenar 13 when compared to the period before the introduction of Prevenar.

### Effect on nasopharyngeal carriage

In a surveillance study in France in children presenting with acute otitis media, changes in nasopharyngeal (NP) carriage of pneumococcal serotypes were evaluated following the introduction of Prevenar (7-valent) and subsequently Prevenar 13. Prevenar 13 significantly reduced NP carriage of the 6 additional serotypes (and serotype 6C) combined and individual serotypes 6C, 7F, 19A when compared with Prevenar. A reduction in carriage was also seen for serotype 3 (2.5% vs 1.1%;  $p=0.1$ ). There was no carriage of serotypes 1 and 5 observed.

The effect of pneumococcal conjugate vaccination on nasopharyngeal carriage was studied in a randomised double-blind study in which infants received either Prevenar 13 or Prevenar (7-valent) at 2, 4, 6 and 12 months of age in Israel. Prevenar 13 significantly reduced newly identified NP acquisition of the 6 additional serotypes (and serotype 6C) combined and of individual serotypes 1, 6A, 6C, 7F, 19A when compared with Prevenar. There was no reduction seen in serotype 3 and for serotype 5 the colonization was too infrequent to assess impact. For 6 of the remaining 7 common serotypes, similar rates of NP acquisition were observed in both vaccine groups; for serotype 19F a significant reduction was observed.

In this study, reductions of *S. pneumoniae* serotypes 19A, 19F, and 6A not susceptible to a number of antibiotics were documented. The reductions ranged between 34% and 62% depending on serotype and antibiotic.

### Prevenar (7-valent vaccine) protective efficacy in infants and children

The efficacy of 7-valent Prevenar was evaluated in two major studies – the Northern California Kaiser Permanente (NCKP) study and the Finnish Otitis Media (FinOM) study. Both studies were randomised, double-blind, active-control studies in which infants were randomised to receive either Prevenar or control vaccine (NCKP, meningococcal serogroup C CRM-conjugate [MnCC] vaccine; FinOM, hepatitis B vaccine) in a four-dose series at 2, 4, 6, and 12-15 months of age. The efficacy results from these studies (for invasive pneumococcal disease, pneumonia, and acute otitis media) are presented below (Table 3).

Test	N	VE <sup>2</sup>	95% CI
NCKP: Vaccine-serotype IPD <sup>3</sup>	30,258	97%	85, 100
NCKP: Clinical pneumonia with abnormal chest X-ray	23,746	35%	4, 56
NCKP: Acute Otitis Media (AOM) <sup>4</sup>	23,746		
Total episodes		7%	4, 10
Recurrent AOM (3 episodes in 6 months, or 4 episodes in 1 year)		9%	3, 15
Recurrent AOM (5 episodes in 6 months, or 6 episodes in 1 year)		23%	7, 36
Tympanostomy tube placement		20%	2, 35
FinOM: AOM	1,662		
Total episodes		6%	-4, 16
All pneumococcal AOM		34%	21, 45
Vaccine-serotype AOM		57%	44, 67
<sup>1</sup> Per protocol			
<sup>2</sup> Vaccine efficacy			
<sup>3</sup> October 1995 to April 20, 1999			
<sup>4</sup> October 1995 to April 30, 1998			

### Prevenar (7-valent) effectiveness

The effectiveness (both direct and indirect effect) of 7-valent Prevenar against pneumococcal disease has been evaluated in both three-dose and two-dose primary infant series immunisation programmes, each with booster doses (Table 4). Following the widespread use of Prevenar, the incidence of IPD has been consistently and substantially reduced.

Using the screening method, serotype-specific effectiveness estimates for 2 doses under the age of 1 year in the UK were 66 % (-29, 91 %) and 100 % (25, 100 %) for serotype 6B and 23F, respectively.

Country (year of introduction)	Recommended schedule	Disease reduction, %	95% CI
UK (England & Wales) <sup>1</sup> (2006)	2, 4, + 13 months	<u>Vaccine serotypes:</u> Two doses under age 1: 85%	49, 95%
USA (2000)	2, 4, 6, + 12-15 months		
Children < 5 <sup>2</sup>		Vaccine serotypes: 98% All serotypes: 77%	97, 99% 73, 79%
Persons ≥ 65 <sup>3</sup>		Vaccine serotypes: 76% All serotypes: 38%	NA NA
Canada (Quebec) <sup>4</sup> (2004)	2, 4, + 12 months	All serotypes: 73% <u>Vaccine serotypes:</u> 2-dose infant series: 99% Completed schedule: 100%	NA 92, 100% 82, 100%
<sup>1</sup> Children < 2 years of age. Calculated vaccine effectiveness as of June 2008 (Broome method).			
<sup>2</sup> 2005 data.			
<sup>3</sup> 2004 data.			
<sup>4</sup> Children < 5 years of age. January 2005 to December 2007. Complete effectiveness for routine 2+1 schedule not yet available.			

### *Acute Otitis Media*

Effectiveness of Prevenar in a 3+1 schedule has also been observed against acute otitis media and pneumonia since its introduction in a national immunisation programme. In a retrospective evaluation of a large US insurance database, AOM visits were reduced by 42.7 % (95 % CI, 42.4-43.1 %), and prescriptions for AOM by 41.9 % in children younger than 2 years of age, compared with a pre-licensure baseline (2004 vs. 1997-99). In a similar analysis, hospitalisations and ambulatory visits for all-cause pneumonia were reduced by 52.4 % and 41.1 %, respectively. For those events specifically identified as pneumococcal pneumonia, the observed reductions in hospitalisations and

ambulatory visits were 57.6 % and 46.9 %, respectively, in children younger than 2 years of age, compared with a pre-licensure baseline (2004 vs. 1997-99). While direct cause-and-effect cannot be inferred from observational analyses of this type, these findings suggest that Prevenar plays an important role in reducing the burden of mucosal disease (AOM and pneumonia) in the target population.

#### Efficacy study in adults 65 years and older

Efficacy against vaccine-type (VT) pneumococcal CAP and IPD was assessed in a large-scale randomised double-blind, placebo-controlled study (Community-Acquired Pneumonia Immunization Trial in Adults–CAPiTA) in the Netherlands. 84,496 subjects, 65 years and older received a single vaccination of either Prevenar 13 or placebo in a 1:1 randomisation.

The CAPiTA study enrolled volunteers  $\geq 65$  years of age whose demographic and health characteristics may differ from those seeking vaccination.

A first episode of hospitalised, chest X-ray confirmed pneumonia was identified in about 2% of this population (n=1,814 subjects) of which 329 cases were confirmed pneumococcal CAP and 182 cases were VT pneumococcal CAP in the per protocol and modified intent to treat (mITT) populations. For the primary endpoint (per protocol population), there were 139 (49 Prevenar 13: 90 Placebo) first episodes of VT-CAP resulting in an efficacy of 45.56% (95.2% CI, 21.82-62.49; p=0.0006).

Efficacy was also demonstrated for the two secondary endpoints in the per protocol population. For the non-bacteraemic/non-invasive (NB/NI) pneumococcal CAP secondary endpoint, there were 93 (33 Prevenar 13: 60 Placebo) first episodes of NB/NI VT pneumococcal CAP resulting in an efficacy of 45.00% (95.2% CI, 14.21-65.31; p=0.0067). For the IPD secondary endpoint, there were 35 (7 Prevenar 13: 28 Placebo) first episodes of VT-IPD, resulting in an efficacy of 75.00% (95.2% CI, 41.06-90.87; p=0.0005).

The duration of protective efficacy against a first episode of VT pneumococcal CAP, NB/NI VT pneumococcal CAP, and VT-IPD extended throughout the 4-year study.

The study was not designed to demonstrate efficacy in subgroups, and the number of subjects  $\geq 85$  years of age was not sufficient to demonstrate efficacy in this age group.

#### Immunogenicity studies in adults $\geq 18$ years and the elderly

In adults, an antibody threshold of serotype-specific pneumococcal polysaccharide IgG binding antibody concentration associated with protection has not been defined. For all pivotal clinical trials, a serotype-specific opsonophagocytosis assay (OPA) was used as a surrogate to assess potential efficacy against invasive pneumococcal disease and pneumonia. OPA geometric mean titers (GMTs) measured 1-month after each vaccination were calculated. OPA titres are expressed as the reciprocal of the highest serum dilution that reduces survival of the pneumococci by at least 50 %.

Pivotal trials for Prevenar 13 were designed to show that functional OPA antibody responses for the 13 serotypes are non-inferior, and for some serotypes superior, to the 12 serotypes in common with the licensed 23-valent pneumococcal polysaccharide vaccine [1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F] one month after vaccine administration. The response to serotype 6A, which is unique to Prevenar 13, was assessed by demonstration of a 4-fold increase in the specific OPA titer above pre-immunised levels.

Five clinical studies were conducted in Europe and the USA evaluating the immunogenicity of Prevenar 13 in different age groups ranging from 18-95 years of age. Clinical studies with Prevenar 13 currently provide immunogenicity data in adults aged 18 years and older, including adults aged 65 and older previously vaccinated with one or more doses of 23-valent pneumococcal polysaccharide vaccine, 5 years prior to enrollment. Each study included healthy adults and immuno-competent adults with stable underlying conditions known to predispose individuals to pneumococcal infection (i.e.,

chronic cardiovascular disease, chronic pulmonary disease including asthma, renal disorders and diabetes mellitus, chronic liver disease including alcoholic liver disease), and adults with risk factors such as smoking and alcohol abuse.

Immunogenicity and safety of Prevenar 13 has been demonstrated in adults aged 18 years and older including those previously vaccinated with a pneumococcal polysaccharide vaccine.

*Adults not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine*

In a head-to-head, comparative trial conducted in adults aged 60-64 years, subjects received a single dose of either Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine. In the same study another group of adults aged 50-59 years and another group of adults aged 18-49 years received a single dose of Prevenar 13.

Table 5 compares the OPA GMTs, 1-month post-dose, in 60-64 year olds given either a single dose of Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine, and in 50-59 year olds given a single dose of Prevenar 13.

Table 5: OPA GMTs in adults aged 60-64 years given Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine (PPSV23) and in adults aged 50-59 years given Prevenar 13 <sup>a,b,c</sup>							
	Prevenar 13 50-59 Years N=350-384	Prevenar 13 60-64 Years N=359-404	PPSV23 60-64 Years N=367-402	Prevenar 13 50-59 Relative to 60-64 Years		Prevenar 13 Relative to PPSV23, 60-64 Years	
Serotype	GMT	GMT	GMT	GMR	(95% CI)	GMR	(95% CI)
1	200	146	104	1.4	(1.08, 1.73)	1.4	(1.10, 1.78)
3	91	93	85	1.0	(0.81, 1.19)	1.1	(0.90, 1.32)
4	2833	2062	1295	1.4	(1.07, 1.77)	1.6	(1.19, 2.13)
5	269	199	162	1.4	(1.01, 1.80)	1.2	(0.93, 1.62)
6A <sup>†</sup>	4328	2593	213	1.7	(1.30, 2.15)	12.1	(8.63, 17.08)
6B	3212	1984	788	1.6	(1.24, 2.12)	2.5	(1.82, 3.48)
7F	1520	1120	405	1.4	(1.03, 1.79)	2.8	(1.98, 3.87)
9V	1726	1164	407	1.5	(1.11, 1.98)	2.9	(2.00, 4.08)
14	957	612	692	1.6	(1.16, 2.12)	0.9	(0.64, 1.21)
18C	1939	1726	925	1.1	(0.86, 1.47)	1.9	(1.39, 2.51)
19A	956	682	352	1.4	(1.16, 1.69)	1.9	(1.56, 2.41)
19F	599	517	539	1.2	(0.87, 1.54)	1.0	(0.72, 1.28)
23F	494	375	72	1.3	(0.94, 1.84)	5.2	(3.67, 7.33)

<sup>a</sup> Non-inferiority was defined as the lower limit of the 2-sided 95% CI for GMR was greater than 0.5.  
<sup>b</sup> Statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR was greater than 1.  
<sup>c</sup> For serotype 6A<sup>†</sup>, which is unique to Prevenar 13, a statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR being greater than 2.

In adults aged 60-64 years, OPA GMTs to Prevenar 13 were non-inferior to the OPA GMTs elicited to the 23-valent pneumococcal polysaccharide vaccine for the twelve serotypes common to both vaccines. For 9 serotypes, the OPA titers were shown to be statistically significantly greater in Prevenar 13 recipients.

In adults aged 50-59 years, OPA GMTs to all 13 serotypes in Prevenar 13 were non-inferior to the Prevenar 13 responses in adults aged 60-64 years. For 9 serotypes, immune responses were related to age, with adults in the 50-59 years group showing statistically significantly greater responses than adults aged 60-64 years.

In all adults ≥ 50 years who received a single dose of Prevenar 13, the OPA titers to serotype 6A were significantly greater than in adults ≥ 60 years who received a single dose of 23-valent pneumococcal polysaccharide vaccine.

One year after vaccination with Prevenar 13 OPA titers had declined compared to one month after vaccination, however, OPA titers for all serotypes remained higher than levels at baseline:

	OPA GMT levels at baseline	OPA GMT levels one year after Prevenar 13
Adults 50-59 years not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine	5 to 45	20 to 1234
Adults 60-64 years not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine	5 to 37	19 to 733

Table 6 shows OPA GMTs 1-month after a single dose of Prevenar 13 in 18-49 year olds compared to 60-64 year olds.

<b>Table 6: OPA GMTs in adults aged 18-49 years and 60-64 years given Prevenar 13<sup>a,b</sup></b>				
	<b>18-49 Years N=836-866</b>	<b>60-64 Years N=359-404</b>	<b>18-49 Years Relative to 60-64 Years</b>	
<b>Serotype</b>	GMT <sup>b</sup>	GMT <sup>b</sup>	GMR	(95% CI <sup>c</sup> )
1	353	146	2.4	(2.03, 2.87)
3	91	93	1.0	(0.84, 1.13)
4	4747	2062	2.3	(1.92, 2.76)
5	386	199	1.9	(1.55, 2.42)
6A	5746	2593	2.2	(1.84, 2.67)
6B	9813	1984	4.9	(4.13, 5.93)
7F	3249	1120	2.9	(2.41, 3.49)
9V	3339	1164	2.9	(2.34, 3.52)
14	2983	612	4.9	(4.01, 5.93)
18C	3989	1726	2.3	(1.91, 2.79)
19A	1580	682	2.3	(2.02, 2.66)
19F	1533	517	3.0	(2.44, 3.60)
23F	1570	375	4.2	(3.31, 5.31)

<sup>a</sup> Non-inferiority was defined as the lower limit of the 2-sided 95% CI for GMR was greater than 0.5.  
<sup>b</sup> Statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR was greater than 1.  
<sup>c</sup> Confidence intervals (CIs) for the ratio are back transformations of a confidence interval based on the Student t distribution for the mean difference of the logarithms of the measures.

In adults aged 18-49 years, OPA GMTs to all 13 serotypes in Prevenar 13 were non-inferior to the Prevenar 13 responses in adults aged 60-64 years.

One year after vaccination with Prevenar 13 OPA titers had declined compared to one month after vaccination, however OPA titers for all serotypes remained higher than levels at baseline.

	OPA GMT levels at baseline	OPA GMT levels one year after Prevenar 13
Adults 18-49 years not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine	5 to 186	23 to 2948

*Adults previously vaccinated with 23-valent pneumococcal polysaccharide vaccine*

Immune responses to Prevenar 13 and 23-valent pneumococcal polysaccharide vaccine were compared in a head to head trial in adults aged  $\geq 70$  years, who had received a single dose of pneumococcal polysaccharide vaccine at least 5 years before study vaccination.

Table 7 compares the OPA GMTs, 1-month post-dose, in pneumococcal polysaccharide vaccinated adults aged  $\geq 70$  years given a single dose of either Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine.

<b>Table 7 : OPA GMTs in pneumococcal polysaccharide vaccinated adults aged <math>\geq 70</math> years given either Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine (PPSV23)<sup>a,b,c</sup></b>				
	<b>Prevenar 13 N=400-426</b>	<b>PPSV23 N=395-445</b>	<b>Prevenar OPA GMT Relative to PPSV23</b>	
<b>Serotype</b>	<b>OPA GMT</b>	<b>OPA GMT</b>	<b>GMR</b>	<b>(95% CI)</b>
1	81	55	1.5	(1.17, 1.88)
3	55	49	1.1	(0.91, 1.35)
4	545	203	2.7	(1.93, 3.74)
5	72	36	2.0	(1.55, 2.63)
6A <sup>†</sup>	903	94	9.6	(7.00, 13.26)
6B	1261	417	3.0	(2.21, 4.13)
7F	245	160	1.5	(1.07, 2.18)
9V	181	90	2.0	(1.36, 2.97)
14	280	285	1.0	(0.73, 1.33)
18C	907	481	1.9	(1.42, 2.50)
19A	354	200	1.8	(1.43, 2.20)
19F	333	214	1.6	(1.17, 2.06)
23F	158	43	3.7	(2.69, 5.09)

<sup>a</sup> Non-inferiority was defined as the lower limit of the 2-sided 95% CI for GMR was greater than 0.5.  
<sup>b</sup> Statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR was greater than 1.  
<sup>c</sup> For serotype 6A<sup>†</sup>, which is unique to Prevenar 13, a statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR greater than 2.

In adults vaccinated with pneumococcal polysaccharide vaccine at least 5 years prior to the clinical study, OPA GMTs to Prevenar 13 were non-inferior to the 23-valent pneumococcal polysaccharide vaccine responses for the 12 serotypes in common. Furthermore, in this study statistically significantly greater OPA GMTs were demonstrated for 10 of the 12 serotypes in common. Immune responses to serotype 6A were statistically significantly greater following vaccination with Prevenar 13 than after 23-valent pneumococcal polysaccharide vaccine.

One year after vaccination with Prevenar 13 in adults aged 70 years and over who were vaccinated with 23-valent pneumococcal polysaccharide vaccine, at least 5 years prior to study entry, OPA titers had declined compared to one month after vaccination, however, OPA titers for all serotypes remained higher than levels at baseline:

	<b>OPA GMT levels at baseline</b>	<b>OPA GMT levels one year after Prevenar 13</b>
Adults $\geq 70$ years vaccinated with 23-valent pneumococcal polysaccharide vaccine at least 5 years prior	9 to 122	18 to 381

### Immune responses in Special Populations

Individuals with the conditions described below have an increased risk of pneumococcal disease. The clinical relevance of the antibody levels elicited by Prevenar 13 in these special populations is unknown.

#### *Sickle cell disease*

An open label single arm study in France, Italy, UK, US, Lebanon, Egypt and Saudi Arabia with 2 doses of Prevenar 13 given 6 months apart was conducted in 158 children and adolescents  $\geq 6$  to  $< 18$  years of age with sickle cell disease who were previously vaccinated with one or more doses of

23-valent pneumococcal polysaccharide vaccine at least 6 months prior to enrollment. After the first vaccination, Prevenar 13 elicited antibody levels measured by both IgG GMCs and OPA GMTs that were statistically significantly higher when compared to levels prior to vaccination. After the second dose immune responses were comparable to those after the first dose. One year after the second dose, antibody levels measured by both IgG GMCs and OPA GMTs were higher than levels prior to the first dose of Prevenar 13, except for the IgG GMCs for serotypes 3 and 5 that were numerically similar.

*Additional Prevenar (7-valent) immunogenicity data: children with sickle cell disease*

The immunogenicity of Prevenar has been investigated in an open-label, multicentre study in 49 infants with sickle cell disease. Children were vaccinated with Prevenar (3 doses one month apart from the age of 2 months), 46 of these children also received a 23-valent pneumococcal polysaccharide vaccine at the age of 15-18 months. After primary immunisation, 95.6 % of the subjects had antibody levels of at least 0.35 µg/ml for all seven serotypes found in Prevenar. A significant increase was seen in the concentrations of antibodies against the seven serotypes after the polysaccharide vaccination, suggesting that immunological memory was well established.

*HIV infection*

*Children and adults not previously vaccinated with a pneumococcal vaccine*

HIV-infected children and adults with CD4  $\geq$  200 cells/ $\mu$ L (mean 717.0 cells/ $\mu$ L), viral load  $<$  50,000 copies/ml (mean 2090.0 copies/ml), free of active AIDS-related illness and not previously vaccinated with a pneumococcal vaccine received 3 doses of Prevenar 13. As per general recommendations, a single dose of 23-valent pneumococcal polysaccharide vaccine was subsequently administered. Vaccines were administered at 1 month intervals. Immune responses were assessed in 259-270 evaluable subjects approximately 1 month after each dose of vaccine. After the first dose, Prevenar 13 elicited antibody levels, measured by both IgG GMCs and OPA GMTs that were statistically significantly higher when compared to levels prior to vaccination. After the second and third dose of Prevenar 13, immune responses were similar or higher than those after the first dose.

*Adults previously vaccinated with 23-valent pneumococcal polysaccharide vaccine*

HIV-infected adults  $\geq$  18 years of age with CD4  $\geq$  200 cells/ $\mu$ L (mean 609.1 cells/ $\mu$ L) and viral load  $<$  50,000 copies/ml (mean 330.6 copies/ml), who were free of active AIDS-related illness and were previously vaccinated with 23-valent pneumococcal polysaccharide vaccine administered at least 6 months prior to enrollment, received 3 doses of Prevenar 13, at enrollment, 6 months, and 12 months after the first dose of Prevenar 13. Immune responses were assessed in 231-255 evaluable subjects approximately 1 month after each dose of Prevenar 13. After the first dose, Prevenar 13 elicited antibody levels measured by both IgG GMCs and OPA GMTs that were statistically significantly higher when compared to levels prior to vaccination. After the second and third dose of Prevenar 13, immune responses were comparable or higher than those after the first dose. In the study 162 subjects had received one prior dose of 23-valent pneumococcal polysaccharide vaccine, 143 subjects 2 prior doses and 26 subjects more than 2 prior doses of 23-valent polysaccharide vaccine. Subjects who received two or more previous doses of 23-valent pneumococcal polysaccharide vaccine showed a similar immune response compared with subjects who received a single previous dose.

*Haematopoietic stem cell transplant*

Children and adults with an allogeneic haematopoietic stem cell transplant (HSCT) at  $\geq$  2 years of age with complete haematologic remission of underlying disease or with very good partial remission in the case of lymphoma and myeloma received three doses of Prevenar 13 with an interval of at least 1 month between doses. The first dose was administered at 3 to 6 months after HSCT. A fourth (booster) dose of Prevenar 13 was administered 6 months after the third dose. As per general recommendations, a single dose of 23-valent pneumococcal polysaccharide vaccine was administered 1 month after the fourth dose of Prevenar 13. Immune responses as measured by IgG GMCs were assessed in 168-211 evaluable subjects approximately 1 month after vaccination. Prevenar 13 elicited increased antibody levels after each dose of Prevenar 13. Immune responses after the fourth dose of Prevenar 13 were significantly increased for all serotypes compared with after the third dose. Functional antibody titers (OPA titers) were not measured in this study.

## **5.2 Pharmacokinetic properties**

Not applicable.

## **5.3 Preclinical safety data**

Non-clinical data revealed no special hazard for humans based on conventional studies of safety pharmacology, single and repeated dose toxicity, local tolerance, and reproduction and developmental toxicity.

# **6. PHARMACEUTICAL PARTICULARS**

## **6.1 List of excipients**

Sodium chloride  
Succinic acid  
Polysorbate 80  
Water for injections

For adjuvant, see section 2.

## **6.2 Incompatibilities**

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

## **6.3 Shelf life**

3 years

## **6.4 Special precautions for storage**

Store in a refrigerator (2°C–8°C). Do not freeze.

## **6.5 Nature and contents of container**

0.5 ml suspension for injection in a vial (Type I glass) with a latex-free grey chlorobutyl rubber stopper and sealed with an aluminum flip-off seal and a polypropylene flip-off cap.

Pack sizes of 1, 5, 10, 25 and 50.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal and other handling**

During storage, a white deposit and clear supernatant can be observed. This does not constitute a sign of deterioration.

The vaccine should be shaken well to obtain a homogeneous white suspension and should be inspected visually for any particulate matter and/or variation of physical aspect prior to administration. Do not use if the content appears otherwise.

No special requirements for disposal.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

**7.       MARKETING AUTHORISATION HOLDER**

Pfizer Limited  
Ramsgate Road  
Sandwich  
Kent CT13 9NJ  
United Kingdom

**8.       MARKETING AUTHORISATION NUMBER(S)**

EU/1/09/590/007  
EU/1/09/590/008  
EU/1/09/590/009  
EU/1/09/590/010  
EU/1/09/590/011

**9.       DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 09 December 2009  
Date of latest renewal: 18 September 2014

**10.      DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## 1. NAME OF THE MEDICINAL PRODUCT

Prevenar 13 suspension for injection in multidose container  
Pneumococcal polysaccharide conjugate vaccine (13-valent, adsorbed)

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

1 dose (0.5 ml) contains:

Pneumococcal polysaccharide serotype 1 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 3 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 4 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 5 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 6A <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 6B <sup>1</sup>	4.4 µg
Pneumococcal polysaccharide serotype 7F <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 9V <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 14 <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 18C <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 19A <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 19F <sup>1</sup>	2.2 µg
Pneumococcal polysaccharide serotype 23F <sup>1</sup>	2.2 µg

<sup>1</sup>Conjugated to CRM<sub>197</sub> carrier protein, adsorbed on aluminium phosphate.

1 dose (0.5 ml) contains approximately 32 µg CRM<sub>197</sub> carrier protein and 0.125 mg aluminium.

### Excipients with known effect

For the full list of excipients, see section 6.1.

This is a multidose presentation. See section 6.5 for the number of doses per container.

## 3. PHARMACEUTICAL FORM

Suspension for injection in multidose container (4 doses).  
The vaccine is a homogeneous white suspension.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Active immunisation for the prevention of invasive disease, pneumonia and acute otitis media caused by *Streptococcus pneumoniae* in infants, children and adolescents from 6 weeks to 17 years of age.

Active immunisation for the prevention of invasive disease and pneumonia caused by *Streptococcus pneumoniae* in adults ≥ 18 years of age and the elderly.

See sections 4.4 and 5.1 for information on protection against specific pneumococcal serotypes.

The use of Prevenar 13 should be determined on the basis of official recommendations taking into consideration the risk of invasive disease and pneumonia in different age groups, underlying comorbidities as well as the variability of serotype epidemiology in different geographical areas.

## 4.2 Posology and method of administration

The immunisation schedules for Prevenar 13 should be based on official recommendations.

### Posology

#### Infants and children aged 6 weeks to 5 years

It is recommended that infants who receive a first dose of Prevenar 13 complete the vaccination course with Prevenar 13.

#### Infants aged 6 weeks-6 months

##### *Three-dose primary series*

The recommended immunisation series consists of four doses, each of 0.5 ml. The primary infant series consists of three doses, with the first dose usually given at 2 months of age and with an interval of at least 1 month between doses. The first dose may be given as early as six weeks of age. The fourth (booster) dose is recommended between 11 and 15 months of age.

##### *Two-dose primary series*

Alternatively, when Prevenar 13 is given as part of a routine infant immunisation programme, a series consisting of three doses, each of 0.5 ml, may be given. The first dose may be administered from the age of 2 months, with a second dose 2 months later. The third (booster) dose is recommended between 11 and 15 months of age (see section 5.1).

#### Preterm infants (<37 weeks gestation)

In preterm infants, the recommended immunisation series consists of four doses, each of 0.5 ml. The primary infant series consists of three doses, with the first dose given at 2 months of age and with an interval of at least 1 month between doses. The first dose may be given as early as six weeks of age. The fourth (booster) dose is recommended between 11 and 15 months of age (see sections 4.4 and 5.1).

#### Unvaccinated infants and children $\geq$ 7 months of age

##### *Infants aged 7-11 months*

Two doses, each of 0.5 ml, with an interval of at least 1 month between doses. A third dose is recommended in the second year of life.

##### *Children aged 12-23 months*

Two doses, each of 0.5 ml, with an interval of at least 2 months between doses (see section 5.1).

##### *Children and adolescents aged 2-17 years*

One single dose of 0.5 ml.

#### Prevenar 13 vaccine schedule for infants and children previously vaccinated with Prevenar (7-valent) (*Streptococcus pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F, and 23F)

Prevenar 13 contains the same 7 serotypes included in Prevenar, using the same carrier protein CRM<sub>197</sub>.

Infants and children who have begun immunisation with Prevenar may switch to Prevenar 13 at any point in the schedule.

##### *Young children (12-59 months) completely immunised with Prevenar (7-valent)*

Young children who are considered completely immunised with Prevenar (7-valent) should receive one dose of 0.5 ml of Prevenar 13 to elicit immune responses to the 6 additional serotypes. This dose

of Prevenar 13 should be administered at least 8 weeks after the final dose of Prevenar (7-valent) (see section 5.1).

#### *Children and adolescents 5–17 years*

Children 5 to 17 years of age may receive a single dose of Prevenar 13 if they have been previously vaccinated with one or more doses of Prevenar. This dose of Prevenar 13 should be administered at least 8 weeks after the final dose of Prevenar (7-valent) (see section 5.1).

#### Adults $\geq$ 18 years of age, and the elderly

One single dose.

The need for revaccination with a subsequent dose of Prevenar 13 has not been established.

Regardless of prior pneumococcal vaccination status, if the use of 23-valent pneumococcal polysaccharide vaccine is considered appropriate, Prevenar 13 should be given first (see sections 4.5 and 5.1).

#### Special Populations

Individuals who have underlying conditions predisposing them to invasive pneumococcal disease (such as sickle cell disease or HIV infection) including those previously vaccinated with one or more doses of 23-valent pneumococcal polysaccharide vaccine may receive at least one dose of Prevenar 13 (see section 5.1).

In individuals with an haematopoietic stem cell transplant (HSCT), the recommended immunisation series consists of four doses of Prevenar 13, each of 0.5 ml. The primary series consists of three doses, with the first dose given at 3 to 6 months after HSCT and with an interval of at least 1 month between doses. A fourth (booster) dose is recommended 6 months after the third dose (see section 5.1).

#### Method of administration

The vaccine should be given by intramuscular injection. The preferred sites are the anterolateral aspect of the thigh (vastus lateralis muscle) in infants or the deltoid muscle of the upper arm in children and adults.

### **4.3 Contraindications**

Hypersensitivity to the active substances, to any of the excipients listed in section 6.1, or to diphtheria toxoid.

As with other vaccines, the administration of Prevenar 13 should be postponed in subjects suffering from acute, severe febrile illness. However, the presence of a minor infection, such as a cold, should not result in the deferral of vaccination.

### **4.4 Special warnings and precautions for use**

Prevenar 13 must not be administered intravascularly.

As with all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of a rare anaphylactic event following the administration of the vaccine.

This vaccine should not be given as an intramuscular injection to individuals with thrombocytopaenia or any coagulation disorder that would contraindicate intramuscular injection, but may be given subcutaneously if the potential benefit clearly outweighs the risks (see section 5.1).

Prevenar 13 will only protect against *Streptococcus pneumoniae* serotypes included in the vaccine, and will not protect against other microorganisms that cause invasive disease, pneumonia, or otitis media.

As with any vaccine, Prevenar 13 may not protect all individuals receiving the vaccine from pneumococcal disease. For the most recent epidemiological information in your country you should consult with the relevant national organisation.

Individuals with impaired immune responsiveness, whether due to the use of immuno-suppressive therapy, a genetic defect, human immunodeficiency virus (HIV) infection, or other causes, may have reduced antibody response to active immunisation.

Safety and immunogenicity data are available for a limited number of individuals with sickle cell disease, HIV infection, or with an haematopoietic stem cell transplant (see section 5.1). Safety and immunogenicity data for Prevenar 13 are not available for individuals in other specific immuno-compromised groups (e.g., malignancy or nephrotic syndrome) and vaccination should be considered on an individual basis.

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially 'sodium-free'.

#### Infants and children aged 6 weeks to 5 years

In clinical studies, Prevenar 13 elicited an immune response to all thirteen serotypes included in the vaccine. The immune response for serotype 3 following the booster dose was not increased above the levels seen after the infant vaccination series; the clinical relevance of this observation regarding the induction of serotype 3 immune memory is unknown (see section 5.1).

The proportions of functional antibody responders (OPA titres  $\geq 1:8$ ) to serotypes 1, 3 and 5 were high. However, the OPA geometric mean titres were lower than those against each of the remaining additional vaccine serotypes; the clinical relevance of this observation for protective efficacy is unknown (see section 5.1).

Limited data have demonstrated that Prevenar 7-valent (three-dose primary series) induces an acceptable immune response in infants with sickle cell disease with a safety profile similar to that observed in non-high-risk groups (see section 5.1).

Children younger than 2 years old should receive the appropriate-for-age Prevenar 13 vaccination series (see section 4.2). The use of pneumococcal conjugate vaccine does not replace the use of 23-valent pneumococcal polysaccharide vaccines in children  $\geq 2$  years of age with conditions (such as sickle cell disease, asplenia, HIV infection, chronic illness, or those who are immuno-compromised) placing them at higher risk for invasive disease due to *Streptococcus pneumoniae*. Whenever recommended, children at risk who are  $\geq 24$  months of age and already primed with Prevenar 13 should receive 23-valent pneumococcal polysaccharide vaccine. The interval between the 13-valent pneumococcal conjugate vaccine (Prevenar 13) and the 23-valent pneumococcal polysaccharide vaccine should not be less than 8 weeks. There are no data available to indicate whether the administration of 23-valent pneumococcal polysaccharide vaccine to unprimed children or to children primed with Prevenar 13 might result in hyporesponsiveness to further doses of Prevenar 13.

The potential risk of apnoea and the need for respiratory monitoring for 48-72 h should be considered when administering the primary immunisation series to very premature infants (born  $\leq 28$  weeks of gestation), and particularly for those with a previous history of respiratory immaturity. As the benefit of vaccination is high in this group of infants, vaccination should not be withheld or delayed.

For vaccine serotypes, protection against otitis media is expected to be lower than protection against invasive disease. As otitis media is caused by many organisms other than pneumococcal serotypes represented in the vaccine, protection against all otitis media is expected to be low (see section 5.1).

When Prevenar 13 is administered concomitantly with Infanrix hexa (DTPa-HBV-IPV/Hib), the rates of febrile reactions are similar to those seen with concomitant administration of Prevenar (7-valent) and Infanrix hexa (see section 4.8). Increased reporting rates of convulsions (with or without fever)

and hypotonic hyporesponsive episode (HHE) were observed with concomitant administration of Prevenar 13 and Infanrix hexa (see section 4.8).

Antipyretic treatment should be initiated according to local treatment guidelines for children with seizure disorders or with a prior history of febrile seizures and for all children receiving Prevenar 13 simultaneously with vaccines containing whole cell pertussis.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

##### Infants and children aged 6 weeks to 5 years

Prevenar 13 can be given concomitantly with any of the following vaccine antigens, either as monovalent or combination vaccines: diphtheria, tetanus, acellular or whole cell pertussis, *Haemophilus influenzae* type b, inactivated poliomyelitis, hepatitis B (see section 4.4 for guidance on Infanrix hexa), meningococcal serogroup C, measles, mumps, rubella, varicella and rotavirus vaccine.

Prevenar 13 can also be given concomitantly between 12-23 months with the tetanus toxoid conjugated meningococcal polysaccharide serogroups A, C, W and Y vaccine to children who have been adequately primed with Prevenar 13 (as per local recommendations).

Data from a postmarketing clinical study evaluating the impact of prophylactic use of antipyretics (ibuprofen and paracetamol) on the immune response to Prevenar 13 suggest that administration of paracetamol concomitantly or within the same day of vaccination may reduce the immune response to Prevenar 13 after the infant series. Responses to the booster dose administered at 12 months were unaffected. The clinical significance of this observation is unknown.

##### Children and adolescents 6 to 17 years of age

No data are currently available regarding concomitant use with other vaccines.

##### Adults 18 to 49 years of age

No data are available regarding concomitant use with other vaccines.

##### Adults aged 50 years and older

Prevenar 13 may be administered concomitantly with the seasonal trivalent inactivated influenza vaccine (TIV).

In two studies conducted in adults aged 50-59 and 65 years and older, it was demonstrated that Prevenar 13 may be given concomitantly with trivalent inactivated influenza vaccine (TIV). The responses to all three TIV antigens were comparable when TIV was given alone or concomitantly with Prevenar 13.

When Prevenar 13 was given concomitantly with TIV, the immune responses to Prevenar 13 were lower compared to when Prevenar 13 was given alone, however, there was no long-term impact on circulating antibody levels.

In a third study in adults aged 50-93 years, it was demonstrated that Prevenar 13 may be given concomitantly with the seasonal quadrivalent inactivated influenza vaccine (QIV). The immune responses to all four QIV strains were noninferior when Prevenar 13 was given concomitantly with QIV compared to when QIV was given alone.

The immune responses to Prevenar 13 were noninferior when Prevenar 13 was given concomitantly with QIV compared to when Prevenar 13 was given alone. As with concomitant administration with

trivalent vaccines, immune responses to some pneumococcal serotypes were lower when both vaccines were given concomitantly.

Concomitant use with other vaccines has not been investigated.

Different injectable vaccines should always be given at different vaccination sites.

Concomitant administration of Prevenar 13 and 23-valent pneumococcal polysaccharide vaccine has not been studied. In clinical studies when Prevenar 13 was given 1 year after 23-valent pneumococcal polysaccharide vaccine the immune responses were lower for all serotypes compared to when Prevenar 13 was given to subjects not previously immunised with 23-valent pneumococcal polysaccharide vaccine. The clinical significance of this is unknown.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

There are no data from the use of pneumococcal 13-valent conjugate vaccine in pregnant women. Therefore the use of Prevenar 13 should be avoided during pregnancy.

##### Breast-feeding

It is unknown whether pneumococcal 13-valent conjugate vaccine is excreted in human milk.

##### Fertility

Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3).

#### **4.7 Effects on ability to drive and use machines**

Prevenar 13 has no or negligible influence on the ability to drive and use machines. However, some of the effects mentioned under section 4.8 “Undesirable effects” may temporarily affect the ability to drive or use machines.

#### **4.8 Undesirable effects**

Analysis of postmarketing reporting rates suggests a potential increased risk of convulsions, with or without fever, and HHE when comparing groups which reported use of Prevenar 13 with Infanrix hexa to those which reported use of Prevenar 13 alone.

Adverse reactions reported in clinical studies or from the postmarketing experience for all age groups are listed in this section per system organ class, in decreasing order of frequency and seriousness. The frequency is defined as follows: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ), very rare ( $< 1/10,000$ ), not known (cannot be estimated from available data).

##### Infants and children aged 6 weeks to 5 years

The safety of the vaccine was assessed in controlled clinical studies where 14,267 doses were given to 4,429 healthy infants from 6 weeks of age at first vaccination and 11-16 months of age at booster dose. In all infant studies, Prevenar 13 was co-administered with routine paediatric vaccines (see section 4.5).

Safety in 354 previously unvaccinated children (7 months to 5 years of age) was also assessed.

The most commonly reported adverse reactions in children 6 weeks to 5 years of age were vaccination-site reactions, fever, irritability, decreased appetite, and increased and/or decreased sleep.

In a clinical study in infants vaccinated at 2, 3, and 4 months of age, fever  $\geq 38^{\circ}\text{C}$  was reported at higher rates among infants who received Prevenar (7-valent) concomitantly with Infanrix hexa (28.3% to 42.3%) than in infants receiving Infanrix hexa alone (15.6% to 23.1%). After a booster dose at 12 to 15 months of age, fever  $\geq 38^{\circ}\text{C}$  was reported in 50.0% of infants who received Prevenar (7-valent) and Infanrix hexa at the same time as compared to 33.6% of infants receiving Infanrix hexa alone. These reactions were mostly moderate (less than or equal to  $39^{\circ}\text{C}$ ) and transient.

An increase in vaccination-site reactions was reported in children older than 12 months compared to rates observed in infants during the primary series with Prevenar 13.

#### *Adverse reactions from clinical studies*

In clinical studies, the safety profile of Prevenar 13 was similar to Prevenar. The following frequencies are based on adverse reactions assessed in Prevenar 13 clinical studies:

#### Immune system disorders:

Rare: Hypersensitivity reaction including face oedema, dyspnoea, bronchospasm

#### Nervous system disorders:

Uncommon: Convulsions (including febrile convulsions)

Rare: Hypotonic-hyporesponsive episode

#### Gastrointestinal disorders:

Very common: Decreased appetite

Common: Vomiting; diarrhoea

#### Skin and subcutaneous tissue disorders:

Common: Rash

Uncommon: Urticaria or urticaria-like rash

#### General disorders and administration site conditions:

Very common: Pyrexia; irritability; any vaccination-site erythema, induration/swelling or pain/tenderness; somnolence; poor quality sleep

Vaccination-site erythema or induration/swelling 2.5 cm–7.0 cm (after the booster dose and in older children [age 2 to 5 years])

Common: Pyrexia  $> 39^{\circ}\text{C}$ ; vaccination-site movement impairment (due to pain); vaccination-site erythema or induration/swelling 2.5 cm–7.0 cm (after infant series)

Uncommon: Vaccination-site erythema, induration/swelling  $> 7.0$  cm; crying

#### *Adverse reactions from Prevenar 13 postmarketing experience*

Although the following adverse drug reactions were not observed in the Prevenar 13 clinical studies in infants and children, the following are considered adverse drug reactions for Prevenar 13 as they were reported in the postmarketing experience. Because these reactions were derived from spontaneous reports, the frequencies could not be determined and are thus considered as not known.

#### Blood and lymphatic system disorders:

Lymphadenopathy (localised to the region of the vaccination-site)

#### Immune system disorders:

Anaphylactic/anaphylactoid reaction including shock; angioedema

#### Skin and subcutaneous tissue disorders:

Erythema multiforme

General disorders and administration site conditions:

Vaccination-site urticaria; vaccination-site dermatitis; vaccination-site pruritus; flushing

Additional information in special populations:

Apnoea in very premature infants ( $\leq 28$  weeks of gestation) (see section 4.4).

Children and adolescents aged 6 to 17 years of age

Safety was evaluated in 592 children (294 children aged 5 to 10 years previously immunised with at least one dose of Prevenar and 298 children aged 10 to 17 years who had not received a pneumococcal vaccine).

The most common adverse events in children and adolescents 6 to 17 years of age were:

Nervous system disorders:

Common: Headaches

Gastrointestinal disorders:

Very common: Decreased appetite

Common: Vomiting; diarrhoea

Skin and subcutaneous tissue disorders:

Common: Rash; urticaria or urticaria-like rash

General disorders and administration site conditions:

Very common: Irritability; any vaccination-site erythema; induration/swelling or pain/tenderness; somnolence; poor quality sleep; vaccination-site tenderness (including impaired movement)

Common: Pyrexia

Other adverse events previously observed in infants and children 6 weeks to 5 years of age may also be applicable to this age group but were not seen in this study possibly due to the small sample size.

Additional information in special populations

Children and adolescents with sickle cell disease, HIV infection, or an haematopoietic stem cell transplant have similar frequencies of adverse reactions, except that headaches, vomiting, diarrhoea, pyrexia, fatigue, arthralgia, and myalgia were very common.

Adults  $\geq 18$  years and elderly

Safety was assessed in 7 clinical studies including 91,593 adults ranging in age from 18 to 101 years. Prevenar 13 was administered to 48,806 adults; 2,616 (5.4%) aged 50 to 64 years, and 45,291 (92.8%) aged 65 years and older. One of the 7 studies included a group of adults ( $n=899$ ) ranging from 18 to 49 years who received Prevenar 13 and who were not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine. Of the Prevenar 13 recipients 1,916 adults were previously vaccinated with the 23-valent pneumococcal polysaccharide vaccine at least 3 years prior to study vaccination, and 46,890 were 23-valent pneumococcal polysaccharide vaccine unvaccinated.

A trend to lower frequency of adverse reactions was associated with greater age; adults  $> 65$  years of age (regardless of prior pneumococcal vaccination status) reported fewer adverse reactions than younger adults, with adverse reactions generally most common in the youngest adults, 18 to 29 years of age.

Overall, the frequency categories were similar for all age groups, with the exception of vomiting which was very common ( $\geq 1/10$ ) in adults aged 18 to 49 years and common ( $\geq 1/100$  to  $< 1/10$ ) in all other age groups, and pyrexia was very common in adults aged 18 to 29 years and common in all other age groups. Severe vaccination-site pain/tenderness and severe limitation of arm movement was very common in adults 18 to 39 years and common in all other age groups.

Adverse reactions from clinical studies

Local reactions and systemic events were solicited daily after each vaccination for 14 days in 6 studies and 7 days in the remaining study. The following frequencies are based on adverse reactions assessed in Prevenar 13 clinical studies in adults:

Metabolism and nutrition disorders:

Very common: Decreased appetite

Nervous system disorders:

Very common: Headaches

Gastrointestinal disorders:

Very common: Diarrhoea; vomiting (in adults aged 18 to 49 years)

Common: Vomiting (in adults aged 50 years and over)

Uncommon: Nausea

Immune system disorders:

Uncommon: Hypersensitivity reaction including face oedema, dyspnoea, bronchospasm

Skin and subcutaneous tissue disorders:

Very common: Rash

General disorders and administration site conditions:

Very common: Chills; fatigue; vaccination-site erythema; vaccination-site induration/swelling; vaccination-site pain/tenderness (severe vaccination-site pain/tenderness very common in adults aged 18 to 39 years); limitation of arm movement (severe limitation of arm movements very common in adults aged 18 to 39 years)

Common: Pyrexia (very common in adults aged 18 to 29 years)

Uncommon: Lymphadenopathy localized to the region of the vaccination-site

Musculoskeletal and connective tissue disorders:

Very common: Arthralgia; myalgia

Overall, no significant differences in frequencies of adverse reactions were seen when Prevenar 13 was given to adults previously vaccinated with the pneumococcal polysaccharide vaccine.

Additional information in special populations

Adults with HIV infection have similar frequencies of adverse reactions, except that pyrexia and vomiting were very common and nausea common.

Adults with an haematopoietic stem cell transplant have similar frequencies of adverse reactions, except that pyrexia and vomiting were very common.

Higher frequency in some solicited systemic reactions was observed when Prevenar 13 was administered concomitantly with trivalent inactivated influenza vaccine (TIV) compared to TIV given alone (headache, chills, rash, decreased appetite, arthralgia, and myalgia) or Prevenar 13 given alone (headache, fatigue, chills, decreased appetite, and arthralgia).

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

#### **4.9 Overdose**

In infants and children there have been reports of overdose with Prevenar 13 defined as subsequent doses administered closer than recommended to the previous dose. In general, adverse events reported with overdose are consistent with those that have been reported with doses given in the recommended paediatric schedules of Prevenar 13.

### **5. PHARMACOLOGICAL PROPERTIES**

#### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: vaccines, pneumococcal vaccines; ATC code: J07AL02

Prevenar 13 contains the 7 pneumococcal capsular polysaccharides that are in Prevenar (4, 6B, 9V, 14, 18C, 19F, 23F) plus 6 additional polysaccharides (1, 3, 5, 6A, 7F, 19A) all conjugated to CRM<sub>197</sub> carrier protein.

#### Burden of disease

##### *Infants and children aged 6 weeks to 5 years*

Based on serotype surveillance in Europe performed before the introduction of Prevenar, Prevenar 13 is estimated to cover 73-100% (depending on the country) of serotypes causing invasive pneumococcal disease (IPD) in children less than 5 years of age. In this age group, serotypes 1, 3, 5, 6A, 7F, and 19A account for 15.6% to 59.7% of invasive disease, depending on the country, the time period studied, and the use of Prevenar.

Acute otitis media (AOM) is a common childhood disease with different aetiologies. Bacteria can be responsible for 60-70% of clinical episodes of AOM. *S. pneumoniae* is one of the most common causes of bacterial AOM worldwide.

Prevenar 13 is estimated to cover over 90% of serotypes causing antimicrobial-resistant IPD.

##### *Children and adolescents aged 6 to 17 years*

In children and adolescents aged 6 to 17 years, the incidence of pneumococcal disease is low, however, there is an increased risk of morbidity and mortality in those with underlying comorbidities.

##### *Adults ≥ 18 years and the elderly*

Pneumonia is the most common clinical presentation of pneumococcal disease in adults.

The reported incidence of community-acquired pneumonia (CAP) and IPD in Europe varies by country, increases with age from 50 years and is highest in individuals aged ≥ 65 years. *S. pneumoniae* is the most frequent cause of CAP, and is estimated to be responsible for approximately 30% of all CAP cases requiring hospitalisation in adults in developed countries.

Bacteraemic pneumonia (approximately 80% of IPD in adults), bacteraemia without a focus, and meningitis are the most common manifestations of IPD in adults. Based on surveillance data following the introduction of Prevenar but before the introduction of Prevenar 13 in childhood vaccination programmes, the pneumococcal serotypes in Prevenar 13 may be responsible for at least 50 – 76% (depending on country) of IPD in adults.

The risk for CAP and IPD in adults also increases with chronic underlying medical conditions, specifically, anatomical or functional asplenia, diabetes mellitus, asthma, chronic cardiovascular, pulmonary, kidney or liver disease, and it is highest in those who are immune-suppressed such as those with malignant haematological diseases or HIV infection.

#### Prevenar 13 immunogenicity clinical studies in infants, children and adolescents

The protective efficacy of Prevenar 13 against IPD has not been studied. As recommended by the World Health Organization (WHO) the assessment of potential efficacy against IPD in infants and young children has been based on a comparison of immune responses to the seven common serotypes shared between Prevenar 13 and Prevenar, for which protective efficacy has been proven (for Prevenar (7-valent) efficacy in infants and children, see below). Immune responses to the additional 6 serotypes were also measured.

#### Study of Prevenar 13 containing the preservative 2-phenoxyethanol (2-PE):

Safety and immunogenicity of Prevenar 13 containing the preservative 2-PE (presented in a multidose container) given to healthy infants at 8, 12 and 16 weeks of age were compared to those of Prevenar 13 without added preservative (250 infants per group).

Pneumococcal immune responses were compared using non-inferiority criteria, including the percentage of subjects with serum anti-polysaccharide serotype-specific IgG concentration  $\geq 0.35$   $\mu\text{g/ml}$  and the comparison of IgG GMCs one month after the infant series. In addition, OPA GMTs between subjects receiving Prevenar 13 with or without 2-PE were compared.

Non-inferiority for the proportion of subjects achieving an IgG concentration  $\geq 0.35$   $\mu\text{g/ml}$  was demonstrated for all 13 serotypes (lower bound of the 97.5% confidence interval (CI) for the difference in percentage of responders at 0.35  $\mu\text{g/ml}$  between groups was  $> -10\%$ ). Additionally, all 13 serotypes met the pre-defined non-inferiority criterion for IgG GMCs (lower bound of the 97.5% CI of GMC ratio [GMR] was greater than 0.5).

Correspondingly, OPA GMTs were similar in both groups, except for serotype 3, which was lower, and serotype 18C, which was higher, in the group that received Prevenar 13 with 2-PE.

#### Studies of Prevenar 13 not containing the preservative 2-PE

##### *Immune responses following a three-dose primary infant series*

Clinical studies have been conducted in a number of European countries and the US using a range of vaccination schedules, including two randomised non-inferiority studies (Germany using a 2, 3, 4 month primary series [006] and US using a 2, 4, 6 month primary series [004]). In these two studies pneumococcal immune responses were compared using a set of non-inferiority criteria including the percentage of subjects with serum anti-polysaccharide serotype-specific IgG  $\geq 0.35$   $\mu\text{g/ml}$  one month after the primary series and the comparison of IgG geometric mean concentrations (ELISA GMCs); in addition, functional antibody titres (OPA) between subjects receiving Prevenar 13 and Prevenar were compared. For the six additional serotypes, these values were compared with the lowest response among all of the seven common serotypes in the Prevenar recipients.

The non-inferiority immune response comparisons for study 006, based on the proportion of infants achieving anti-polysaccharide IgG concentrations  $\geq 0.35$   $\mu\text{g/ml}$ , are shown in Table 1. Results for study 004 were similar. Prevenar 13 non-inferiority (lower bound of the 95% CI for the difference in

percentage of responders at 0.35 µg/ml between groups was >-10%) was demonstrated for all 7 common serotypes, except for serotype 6B in study 006 and serotypes 6B and 9V in study 004, which missed by a small margin. All seven common serotypes met pre-defined non-inferiority criteria for IgG ELISA GMCs. Prevenar 13 elicited comparable, although slightly lower, antibody levels than Prevenar for the 7 common serotypes. The clinical relevance of these differences is not known.

Non-inferiority was met for the 6 additional serotypes based on the proportion of infants achieving antibody concentrations  $\geq 0.35$  µg/ml and comparison of IgG ELISA GMCs in study 006 and was met for 5 out of the 6 serotypes, with the exception of serotype 3 for study 004. For serotype 3, the percentages of Prevenar 13 recipients with serum IgG  $\geq 0.35$  µg/ml were 98.2% (study 006) and 63.5% (study 004).

<b>Table 1: Comparison of the proportion of subjects achieving a pneumococcal anti-polysaccharide IgG antibody concentration <math>\geq 0.35</math> µg/ml after dose 3 of the infant series – study 006</b>			
<b>Serotypes</b>	<b>Prevenar 13 % (N=282-285)</b>	<b>7-valent Prevenar % (N=277-279)</b>	<b>Difference (95 % CI)</b>
<b>7-valent Prevenar serotypes</b>			
4	98.2	98.2	0.0 (-2.5, 2.6)
6B	77.5	87.1	-9.6 (-16.0, -3.3)
9V	98.6	96.4	2.2 (-0.4, 5.2)
14	98.9	97.5	1.5 (-0.9, 4.1)
18C	97.2	98.6	-1.4 (-4.2, 1.2)
19F	95.8	96.0	-0.3 (-3.8, 3.3)
23F	88.7	89.5	-0.8 (-6.0, 4.5)
<b>Additional serotypes in Prevenar 13</b>			
1	96.1	87.1*	9.1 (4.5, 13.9)
3	98.2	87.1	11.2 (7.0, 15.8)
5	93.0	87.1	5.9 (0.8, 11.1)
6A	91.9	87.1	4.8 (-0.3, 10.1)
7F	98.6	87.1	11.5 (7.4, 16.1)
19A	99.3	87.1	12.2 (8.3, 16.8)

\*The serotype in Prevenar with the lowest percent response rate was 6B in study 006 (87.1 %).

Prevenar 13 elicited functional antibodies to all 13 vaccine serotypes in studies 004 and 006. For the 7 common serotypes there were no differences between groups in the proportion of subjects with OPA titres  $\geq 1:8$ . For each of the seven common serotypes, > 96% and > 90% of the Prevenar 13 recipients reached an OPA titre  $\geq 1:8$  one month after the primary series in studies 006 and 004, respectively.

For each of the 6 additional serotypes, Prevenar 13 elicited OPA titres  $\geq 1:8$  in 91.4% to 100% of vaccinees one month after the primary series in studies 004/006. The functional antibody (OPA) geometric mean titres for serotypes 1, 3 and 5 were lower than the titres for each of the other additional serotypes; the clinical relevance of this observation for protective efficacy is unknown.

#### *Immune responses following a two-dose primary infant series*

The immunogenicity after two doses in infants has been documented in four studies. The proportion of infants achieving a pneumococcal anti-capsular polysaccharide IgG concentration  $\geq 0.35$  µg/ml one month after the second dose ranged from 79.6% to 98.5% across 11 of the 13 vaccine serotypes. Smaller proportions of infants achieved this antibody concentration threshold for serotype 6B (27.9% to 57.3%) and 23F (55.8% to 68.1%) for all studies using a 2, 4 month regimen, compared to 58.4% for serotype 6B and 68.6% for 23F for a study using a 3, 5 month regimen. After the booster dose, all vaccine serotypes including 6B and 23F had immune responses consistent with adequate priming with a two-dose primary series. In a UK study, the functional antibody (OPA) responses were comparable for all serotypes including 6B and 23F in the Prevenar and Prevenar 13 arms after the primary series at two and four months of age and after the booster dose at 12 months of age. For Prevenar 13 recipients, the proportion of responders with an OPA titre  $\geq 1:8$  was at least 87% following the infant series, and

at least 93% following the booster dose. The OPA geometric mean titres for serotypes 1, 3 and 5 were lower than the titres for each of the other additional serotypes; the clinical relevance of this observation is unknown.

#### Booster responses following two-dose and three-dose primary infant series

Following the booster dose, antibody concentrations increased from the pre-booster level for all 13 serotypes. Post-booster antibody concentrations were higher for 12 serotypes than those achieved after the infant primary series. These observations are consistent with adequate priming (the induction of immunologic memory). The immune response for serotype 3 following the booster dose was not increased above the levels seen after the infant vaccination series; the clinical relevance of this observation regarding the induction of serotype 3 immune memory is unknown.

Antibody responses to booster doses following two-dose or three-dose infant primary series were comparable for all 13 vaccine serotypes.

For children aged from 7 months to 5 years, age appropriate catch-up immunisation schedules (as described in section 4.2) result in levels of anti-capsular polysaccharide IgG antibody responses to each of the 13 serotypes that are at least comparable to those of a three-dose primary series in infants.

Antibody persistence and immunological memory were evaluated in a study in healthy children who received a single dose of Prevenar 13 at least 2 years after they had been previously immunised with either 4 doses of Prevenar, a 3-dose infant series of Prevenar followed by Prevenar 13 at 12 months of age, or 4 doses of Prevenar 13.

The single dose of Prevenar 13, in children approximately 3.4 years of age regardless of previous vaccination history with Prevenar or Prevenar 13, induced a robust antibody response for both the 7 common serotypes and the 6 additional serotypes in Prevenar 13.

Since the introduction of 7-valent Prevenar in 2000, pneumococcal disease surveillance data have not shown that the immunity elicited by Prevenar in infancy has waned over time.

#### Preterm Infants

Safety and immunogenicity of Prevenar 13 given at 2, 3, 4 and 12 months was assessed in approximately 100 prematurely born infants (mean Estimated Gestational Age [EGA], 31 weeks; range, 26 to 36 weeks) and compared with approximately 100 infants born at term (mean EGA, 39 weeks; range, 37 to 42 weeks).

Immune responses in preterm and term infants were compared using the proportion of subjects achieving a pneumococcal polysaccharide IgG binding antibody concentration  $\geq 0.35$   $\mu\text{g/ml}$  1 month after the infant series, the approach used for immunogenicity comparisons of Prevenar 13 to Prevenar based on WHO guidelines.

More than 85% achieved a pneumococcal polysaccharide IgG binding antibody concentration  $\geq 0.35$   $\mu\text{g/ml}$  1 month after the infant series, except for serotypes 5 (71.7%), 6A (82.7%), and 6B (72.7%) in the preterm group. For these 3 serotypes, the proportion of responders among preterm infants was significantly lower than among term infants. Approximately one month after the toddler dose, the proportion of subjects in each group achieving this same antibody concentration threshold was  $>97\%$ , except for serotype 3 (71% in preterm infants and 79% in term infants). It is unknown whether immunological memory to all serotypes is induced in pre-term infants. In general, serotype-specific IgG GMCs were lower for preterm infants than term infants.

After the infant series, OPA GMTs were similar in preterm infants compared to term infants except for serotype 5, which was lower in preterm infants. OPA GMTs after the toddler dose relative to those after the infant series were similar or lower for 4 serotypes (4, 14, 18C, and 19F) and were statistically

significantly higher for 6 of 13 serotypes (1, 3, 5, 7F, 9V, and 19A) in preterm infants compared to 10 of 13 serotypes (1, 3, 4, 5, 6A, 7F, 9V, 18C, 19A, and 23F) in term infants.

#### Children (12-59 months) completely immunised with Prevenar (7-valent)

Following administration of a single dose of Prevenar 13 to children (12-59 months) who are considered completely immunised with Prevenar (7-valent) (either 2 or 3 dose primary series plus booster), the proportion achieving serum IgG levels  $\geq 0.35$   $\mu\text{g/ml}$  and OPA titres  $\geq 1:8$  was at least 90%. However, 3 (serotypes 1, 5 and 6A) of the 6 additional serotypes showed lower IgG GMC and OPA GMT when compared with children who had received at least one previous vaccination with Prevenar 13. The clinical relevance of the lower GMCs and GMTs is currently unknown.

#### Unvaccinated Children (12-23 months)

Studies in unvaccinated children (12 -23 months) with Prevenar (7-valent) demonstrated that 2 doses were required to achieve serum IgG concentrations for 6B and 23F similar to those induced by a 3-dose infant series.

#### Children and Adolescents 5 to 17 years of age

In an open-label study in 592 healthy children and adolescents including those with asthma (17.4%) who may be predisposed to pneumococcal infection, Prevenar 13 elicited immune responses to all 13 serotypes. A single dose of Prevenar 13 was given to children 5 to 10 years of age previously vaccinated with at least 1 dose of Prevenar, and children and adolescents 10 to 17 years of age who had never received a pneumococcal vaccine.

In both the children 5 to 10 years of age and children and adolescents aged 10 to 17 years, the immune response to Prevenar 13 was non inferior to Prevenar for the 7 common serotypes and to Prevenar 13 for the 6 additional serotypes compared to the immune response after the fourth dose in infants vaccinated at 2, 4, 6 and 12-15 months of age as measured by serum IgG.

In children and adolescents aged 10 to 17 years of age OPA GMTs 1 month after vaccination were noninferior to OPA GMTs in the 5 to 10 year old age group for 12 of the 13 serotypes (except serotype 3).

#### Immune responses after subcutaneous administration

Subcutaneous administration of Prevenar 13 was evaluated in a non-comparative study in 185 healthy Japanese infants and children who received 4 doses at 2, 4, 6 and 12-15 months of age. The study demonstrated that safety and immunogenicity were generally comparable with observations made in studies of intramuscular administration.

#### Prevenar 13 Effectiveness

##### *Invasive Pneumococcal Disease*

Data published by Public Health England showed that, four years after the introduction of Prevenar as a two dose primary infant series with booster dose in the second year of life and with a 94% vaccine uptake, there was a 98% (95% CI 95; 99) reduction in disease caused by the 7 vaccine serotypes in England and Wales. Subsequently, four years following the switch to Prevenar 13, the additional reduction in incidence of IPD due to the 7 serotypes in Prevenar ranged from 76% in children less than 2 years of age to 91% in children 5-14 years of age. The serotype specific reductions for each of the 5 additional serotypes in Prevenar 13 (no cases of serotype 5 IPD were observed) by age group are shown in Table 2 and ranged from 68% (serotype 3) to 100% (serotype 6A) for children less than 5 years of age. Significant incidence reductions were also observed in older age groups who had not been vaccinated with Prevenar 13 (indirect effect).

<b>Table 2: Serotype specific number of cases and incidence reductions of IPD in 2013/14 compared to 2008/09-2009/10 (2008/10) by age in England and Wales</b>									
	<b>&lt;5 years of age</b>			<b>5 to 64 years of age</b>			<b>≥65 years of age</b>		
	2008-10 <sup>§</sup>	2013/14 <sup>§</sup>	% Incidence reduction (95% CI*)	2008-10 <sup>§</sup>	2013/14 <sup>§</sup>	% Incidence reduction (95% CI*)	2008-10 <sup>§</sup>	2013/14 <sup>§</sup>	% Incidence reduction (95% CI*)
<b>Additional serotypes covered by Prevenar 13</b>									
<b>1</b>	59 (54)	5 (5)	91% (98%; 68%)**	458 (382)	77 (71)	83% (88%; 74%)**	102 (89)	13 (13)	87% (94%; 72%)**
<b>3</b>	26 (24)	8 (8)	68% (89%; 6%)	178 (148)	73 (68)	59% (72%; 38%)**	256 (224)	143 (146)	44% (57%; 27%)**
<b>6A</b>	10 (9)	0 (0)	100% (100%; 62%)**	53 (44)	5 (5)	90% (97%; 56%)**	94 (82)	5 (5)	95% (99%; 81%)**
<b>7F</b>	90 (82)	8 (8)	91% (97%; 74%)**	430 (361)	160 (148)	63% (71%; 50%)**	173 (152)	75 (77)	56% (70%; 37%)**
<b>19A</b>	85 (77)	7 (7)	91% (97%; 75%)**	225 (191)	104 (97)	54% (65%; 32%)**	279 (246)	97 (99)	65% (75%; 53%)**
<sup>§</sup> Corrected for proportion of samples serotyped, missing age, denominator compared with 2009/10, and for the trend in total invasive pneumococcal disease up to 2009/10 (after which no trend correction was applied). * 95% CI inflated from a Poisson interval based on over-dispersion of 2.1 seen from modelling of 2000-06 pre-Prevenar all IPD data. ** p<0.005 to cover 6A where p=0.002									

### *Otitis media (OM)*

In a published study performed in Israel, using a 2-dose primary series plus booster dose in the second year of life, the impact of Prevenar 13 on OM was documented in a population-based active-surveillance system with tympanocentesis culturing of middle ear fluid in Israeli children less than 2 years of age with OM.

Following the introduction of Prevenar and subsequently Prevenar 13 there was a decline in incidence from 2.1 to 0.1 cases per 1000 children (95%) for the Prevenar serotypes plus serotype 6A and a decline in incidence from 0.9 to 0.1 cases per 1000 children (89%) for the additional serotypes 1, 3, 5, 7F, and 19A in Prevenar 13. The annual overall pneumococcal incidence of OM declined from 9.6 to 2.1 cases per 1000 children (78%) between July 2004 (prior to the introduction of Prevenar) and June 2013 (post Prevenar 13 introduction).

### *Pneumonia*

In a multicenter observational study in France comparing the periods before and after the switch from Prevenar to Prevenar 13, there was 16% (2060 to 1725 cases) reduction in all community acquired pneumonia (CAP) cases in emergency departments in children 1 month to 15 years of age. Reductions were 53% (167 to 79 cases) ( $p < 0.001$ ) for CAP cases with pleural effusion and 63% (64 to 24 cases) ( $p < 0.001$ ) for microbiologically confirmed pneumococcal CAP cases. In the second year after the introduction of Prevenar 13 the total number of CAP cases due to the 6 additional vaccine serotypes in Prevenar 13 was reduced from 27 to 7 isolates (74%).

The decrease in all cause pneumonia cases was most pronounced in the younger vaccinated age groups with a decrease of 31.8% (757 to 516 cases) and 16.6% (833 to 695 cases) in the age groups <2 years and 2 to 5 years, respectively. The incidence in older, predominantly non-vaccinated children (>5 years) did not change over the duration of the study.

In an ongoing surveillance system (2004 to 2013) to document the impact of Prevenar and subsequently Prevenar 13 on CAP in children less than 5 years in Southern Israel using a 2 dose primary series with a booster dose in the second year of life, there was a reduction of 68% (95% CI 73; 61) in outpatient visits and 32% (95% CI 39; 22) in hospitalizations for alveolar CAP following the introduction of Prevenar 13 when compared to the period before the introduction of Prevenar.

### Effect on nasopharyngeal carriage

In a surveillance study in France in children presenting with acute otitis media, changes in nasopharyngeal (NP) carriage of pneumococcal serotypes were evaluated following the introduction of Prevenar (7-valent) and subsequently Prevenar 13. Prevenar 13 significantly reduced NP carriage of the 6 additional serotypes (and serotype 6C) combined and individual serotypes 6C, 7F, 19A when compared with Prevenar. A reduction in carriage was also seen for serotype 3 (2.5% vs 1.1%;  $p = 0.1$ ). There was no carriage of serotypes 1 and 5 observed.

The effect of pneumococcal conjugate vaccination on nasopharyngeal carriage was studied in a randomised double-blind study in which infants received either Prevenar 13 or Prevenar (7-valent) at 2, 4, 6 and 12 months of age in Israel. Prevenar 13 significantly reduced newly identified NP acquisition of the 6 additional serotypes (and serotype 6C) combined and of individual serotypes 1, 6A, 6C, 7F, 19A when compared with Prevenar. There was no reduction seen in serotype 3 and for serotype 5 the colonization was too infrequent to assess impact. For 6 of the remaining 7 common serotypes, similar rates of NP acquisition were observed in both vaccine groups; for serotype 19F a significant reduction was observed.

In this study, reductions of *S. pneumoniae* serotypes 19A, 19F, and 6A not susceptible to a number of antibiotics were documented. The reductions ranged between 34% and 62% depending on serotype and antibiotic.

### Prevenar (7-valent vaccine) protective efficacy in infants and children

The efficacy of 7-valent Prevenar was evaluated in two major studies – the Northern California Kaiser Permanente (NCKP) study and the Finnish Otitis Media (FinOM) study. Both studies were randomised, double-blind, active-control studies in which infants were randomised to receive either Prevenar or control vaccine (NCKP, meningococcal serogroup C CRM-conjugate [MnCC] vaccine; FinOM, hepatitis B vaccine) in a four-dose series at 2, 4, 6, and 12-15 months of age. The efficacy results from these studies (for invasive pneumococcal disease, pneumonia, and acute otitis media) are presented below (Table 3).

Test	N	VE <sup>2</sup>	95% CI
NCKP: Vaccine-serotype IPD <sup>3</sup>	30,258	97%	85, 100
NCKP: Clinical pneumonia with abnormal chest X-ray	23,746	35%	4, 56
NCKP: Acute Otitis Media (AOM) <sup>4</sup>	23,746		
Total episodes		7%	4, 10
Recurrent AOM (3 episodes in 6 months, or 4 episodes in 1 year)		9%	3, 15
Recurrent AOM (5 episodes in 6 months, or 6 episodes in 1 year)		23%	7, 36
Tympanostomy tube placement		20%	2, 35
FinOM: AOM	1,662		
Total episodes		6%	-4, 16
All pneumococcal AOM		34%	21, 45
Vaccine-serotype AOM		57%	44, 67
<sup>1</sup> Per protocol			
<sup>2</sup> Vaccine efficacy			
<sup>3</sup> October 1995 to April 20, 1999			
<sup>4</sup> October 1995 to April 30, 1998			

### Prevenar (7-valent) effectiveness

The effectiveness (both direct and indirect effect) of 7-valent Prevenar against pneumococcal disease has been evaluated in both three-dose and two-dose primary infant series immunisation programmes, each with booster doses (Table 4). Following the widespread use of Prevenar, the incidence of IPD has been consistently and substantially reduced.

Using the screening method, serotype-specific effectiveness estimates for 2 doses under the age of 1 year in the UK were 66 % (-29, 91 %) and 100 % (25, 100 %) for serotype 6B and 23F, respectively.

Country (year of introduction)	Recommended schedule	Disease reduction, %	95% CI
UK (England & Wales) <sup>1</sup> (2006)	2, 4, + 13 months	<u>Vaccine serotypes:</u> Two doses under age 1: 85%	49, 95%
USA (2000)	2, 4, 6, + 12-15 months		
Children < 5 <sup>2</sup>		Vaccine serotypes: 98% All serotypes: 77%	97, 99% 73, 79%
Persons ≥ 65 <sup>3</sup>		Vaccine serotypes: 76% All serotypes: 38%	NA NA
Canada (Quebec) <sup>4</sup> (2004)	2, 4, + 12 months	All serotypes: 73% <u>Vaccine serotypes:</u> 2-dose infant series: 99% Completed schedule: 100%	NA 92, 100% 82, 100%
<sup>1</sup> Children < 2 years of age. Calculated vaccine effectiveness as of June 2008 (Broome method).			
<sup>2</sup> 2005 data.			
<sup>3</sup> 2004 data.			
<sup>4</sup> Children < 5 years of age. January 2005 to December 2007. Complete effectiveness for routine 2+1 schedule not yet available.			

### *Acute Otitis Media*

Effectiveness of Prevenar in a 3+1 schedule has also been observed against acute otitis media and pneumonia since its introduction in a national immunisation programme. In a retrospective evaluation of a large US insurance database, AOM visits were reduced by 42.7 % (95 % CI, 42.4-43.1 %), and prescriptions for AOM by 41.9 % in children younger than 2 years of age, compared with a pre-licensure baseline (2004 vs. 1997-99). In a similar analysis, hospitalisations and ambulatory visits for all-cause pneumonia were reduced by 52.4 % and 41.1 %, respectively. For those events specifically identified as pneumococcal pneumonia, the observed reductions in hospitalisations and

ambulatory visits were 57.6 % and 46.9 %, respectively, in children younger than 2 years of age, compared with a pre-licensure baseline (2004 vs. 1997-99). While direct cause-and-effect cannot be inferred from observational analyses of this type, these findings suggest that Prevenar plays an important role in reducing the burden of mucosal disease (AOM and pneumonia) in the target population.

#### Efficacy study in adults 65 years and older

Efficacy against vaccine-type (VT) pneumococcal CAP and IPD was assessed in a large-scale randomised double-blind, placebo-controlled study (Community-Acquired Pneumonia Immunization Trial in Adults–CAPiTA) in the Netherlands. 84,496 subjects, 65 years and older received a single vaccination of either Prevenar 13 or placebo in a 1:1 randomisation.

The CAPiTA study enrolled volunteers  $\geq$  65 years of age whose demographic and health characteristics may differ from those seeking vaccination.

A first episode of hospitalised, chest X-ray confirmed pneumonia was identified in about 2% of this population (n=1,814 subjects) of which 329 cases were confirmed pneumococcal CAP and 182 cases were VT pneumococcal CAP in the per protocol and modified intent to treat (mITT) populations. For the primary endpoint (per protocol population), there were 139 (49 Prevenar 13: 90 Placebo) first episodes of VT-CAP resulting in an efficacy of 45.56% (95.2% CI, 21.82-62.49; p=0.0006).

Efficacy was also demonstrated for the two secondary endpoints in the per protocol population. For the non-bacteraemic/non-invasive (NB/NI) pneumococcal CAP secondary endpoint, there were 93 (33 Prevenar 13: 60 Placebo) first episodes of NB/NI VT pneumococcal CAP resulting in an efficacy of 45.00% (95.2% CI, 14.21-65.31; p=0.0067). For the IPD secondary endpoint, there were 35 (7 Prevenar 13: 28 Placebo) first episodes of VT-IPD, resulting in an efficacy of 75.00% (95.2% CI, 41.06-90.87; p=0.0005).

The duration of protective efficacy against a first episode of VT pneumococcal CAP, NB/NI VT pneumococcal CAP, and VT-IPD extended throughout the 4-year study.

The study was not designed to demonstrate efficacy in subgroups, and the number of subjects  $\geq$  85 years of age was not sufficient to demonstrate efficacy in this age group.

#### Immunogenicity studies in adults $\geq$ 18 years and the elderly

In adults, an antibody threshold of serotype-specific pneumococcal polysaccharide IgG binding antibody concentration associated with protection has not been defined. For all pivotal clinical trials, a serotype-specific opsonophagocytosis assay (OPA) was used as a surrogate to assess potential efficacy against invasive pneumococcal disease and pneumonia. OPA geometric mean titers (GMTs) measured 1-month after each vaccination were calculated. OPA titres are expressed as the reciprocal of the highest serum dilution that reduces survival of the pneumococci by at least 50 %.

Pivotal trials for Prevenar 13 were designed to show that functional OPA antibody responses for the 13 serotypes are non-inferior, and for some serotypes superior, to the 12 serotypes in common with the licensed 23-valent pneumococcal polysaccharide vaccine [1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F] one month after vaccine administration. The response to serotype 6A, which is unique to Prevenar 13, was assessed by demonstration of a 4-fold increase in the specific OPA titer above pre-immunised levels.

Five clinical studies were conducted in Europe and the USA evaluating the immunogenicity of Prevenar 13 in different age groups ranging from 18-95 years of age. Clinical studies with Prevenar 13 currently provide immunogenicity data in adults aged 18 years and older, including adults aged 65 and older previously vaccinated with one or more doses of 23-valent pneumococcal polysaccharide vaccine, 5 years prior to enrollment. Each study included healthy adults and immuno-competent adults with stable underlying conditions known to predispose individuals to pneumococcal infection (i.e.,

chronic cardiovascular disease, chronic pulmonary disease including asthma, renal disorders and diabetes mellitus, chronic liver disease including alcoholic liver disease), and adults with risk factors such as smoking and alcohol abuse.

Immunogenicity and safety of Prevenar 13 has been demonstrated in adults aged 18 years and older including those previously vaccinated with a pneumococcal polysaccharide vaccine.

*Adults not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine*

In a head-to-head, comparative trial conducted in adults aged 60-64 years, subjects received a single dose of either Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine. In the same study another group of adults aged 50-59 years and another group of adults aged 18-49 years received a single dose of Prevenar 13.

Table 5 compares the OPA GMTs, 1-month post-dose, in 60-64 year olds given either a single dose of Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine, and in 50-59 year olds given a single dose of Prevenar 13.

<b>Table 5: OPA GMTs in adults aged 60-64 years given Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine (PPSV23) and in adults aged 50-59 years given Prevenar 13<sup>a,b,c</sup></b>							
	<b>Prevenar 13</b>	<b>Prevenar 13</b>	<b>PPSV23</b>	<b>Prevenar 13</b>		<b>Prevenar 13 Relative</b>	
	<b>50-59 Years</b>	<b>60-64 Years</b>	<b>60-64 Years</b>	<b>50-59 Relative to</b>		<b>to PPSV23,</b>	
	<b>N=350-384</b>	<b>N=359-404</b>	<b>N=367-402</b>	<b>60-64 Years</b>		<b>60-64 Years</b>	
<b>Serotype</b>	<b>GMT</b>	<b>GMT</b>	<b>GMT</b>	<b>GMR</b>	<b>(95% CI)</b>	<b>GMR</b>	<b>(95% CI)</b>
1	200	146	104	1.4	(1.08, 1.73)	1.4	(1.10, 1.78)
3	91	93	85	1.0	(0.81, 1.19)	1.1	(0.90, 1.32)
4	2833	2062	1295	1.4	(1.07, 1.77)	1.6	(1.19, 2.13)
5	269	199	162	1.4	(1.01, 1.80)	1.2	(0.93, 1.62)
6A <sup>†</sup>	4328	2593	213	1.7	(1.30, 2.15)	12.1	(8.63, 17.08)
6B	3212	1984	788	1.6	(1.24, 2.12)	2.5	(1.82, 3.48)
7F	1520	1120	405	1.4	(1.03, 1.79)	2.8	(1.98, 3.87)
9V	1726	1164	407	1.5	(1.11, 1.98)	2.9	(2.00, 4.08)
14	957	612	692	1.6	(1.16, 2.12)	0.9	(0.64, 1.21)
18C	1939	1726	925	1.1	(0.86, 1.47)	1.9	(1.39, 2.51)
19A	956	682	352	1.4	(1.16, 1.69)	1.9	(1.56, 2.41)
19F	599	517	539	1.2	(0.87, 1.54)	1.0	(0.72, 1.28)
23F	494	375	72	1.3	(0.94, 1.84)	5.2	(3.67, 7.33)

<sup>a</sup> Non-inferiority was defined as the lower limit of the 2-sided 95% CI for GMR was greater than 0.5.  
<sup>b</sup> Statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR was greater than 1.  
<sup>c</sup> For serotype 6A<sup>†</sup>, which is unique to Prevenar 13, a statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR being greater than 2.

In adults aged 60-64 years, OPA GMTs to Prevenar 13 were non-inferior to the OPA GMTs elicited to the 23-valent pneumococcal polysaccharide vaccine for the twelve serotypes common to both vaccines. For 9 serotypes, the OPA titers were shown to be statistically significantly greater in Prevenar 13 recipients.

In adults aged 50-59 years, OPA GMTs to all 13 serotypes in Prevenar 13 were non-inferior to the Prevenar 13 responses in adults aged 60-64 years. For 9 serotypes, immune responses were related to age, with adults in the 50-59 years group showing statistically significantly greater responses than adults aged 60-64 years.

In all adults ≥ 50 years who received a single dose of Prevenar 13, the OPA titers to serotype 6A were significantly greater than in adults ≥ 60 years who received a single dose of 23-valent pneumococcal polysaccharide vaccine.

One year after vaccination with Prevenar 13 OPA titers had declined compared to one month after vaccination, however, OPA titers for all serotypes remained higher than levels at baseline:

	OPA GMT levels at baseline	OPA GMT levels one year after Prevenar 13
Adults 50-59 years not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine	5 to 45	20 to 1234
Adults 60-64 years not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine	5 to 37	19 to 733

Table 6 shows OPA GMTs 1-month after a single dose of Prevenar 13 in 18-49 year olds compared to 60-64 year olds.

<b>Table 6: OPA GMTs in adults aged 18-49 years and 60-64 years given Prevenar 13<sup>a,b</sup></b>				
	<b>18-49 Years N=836-866</b>	<b>60-64 Years N=359-404</b>	<b>18-49 Years Relative to 60-64 Years</b>	
<b>Serotype</b>	GMT <sup>b</sup>	GMT <sup>b</sup>	GMR	(95% CI <sup>c</sup> )
1	353	146	2.4	(2.03, 2.87)
3	91	93	1.0	(0.84, 1.13)
4	4747	2062	2.3	(1.92, 2.76)
5	386	199	1.9	(1.55, 2.42)
6A	5746	2593	2.2	(1.84, 2.67)
6B	9813	1984	4.9	(4.13, 5.93)
7F	3249	1120	2.9	(2.41, 3.49)
9V	3339	1164	2.9	(2.34, 3.52)
14	2983	612	4.9	(4.01, 5.93)
18C	3989	1726	2.3	(1.91, 2.79)
19A	1580	682	2.3	(2.02, 2.66)
19F	1533	517	3.0	(2.44, 3.60)
23F	1570	375	4.2	(3.31, 5.31)

<sup>a</sup> Non-inferiority was defined as the lower limit of the 2-sided 95% CI for GMR was greater than 0.5.  
<sup>b</sup> Statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR was greater than 1.  
<sup>c</sup> Confidence intervals (CIs) for the ratio are back transformations of a confidence interval based on the Student t distribution for the mean difference of the logarithms of the measures.

In adults aged 18-49 years, OPA GMTs to all 13 serotypes in Prevenar 13 were non-inferior to the Prevenar 13 responses in adults aged 60-64 years.

One year after vaccination with Prevenar 13 OPA titers had declined compared to one month after vaccination, however OPA titers for all serotypes remained higher than levels at baseline.

	OPA GMT levels at baseline	OPA GMT levels one year after Prevenar 13
Adults 18-49 years not previously vaccinated with 23-valent pneumococcal polysaccharide vaccine	5 to 186	23 to 2948

*Adults previously vaccinated with 23-valent pneumococcal polysaccharide vaccine*

Immune responses to Prevenar 13 and 23-valent pneumococcal polysaccharide vaccine were compared in a head to head trial in adults aged  $\geq 70$  years, who had received a single dose of pneumococcal polysaccharide vaccine at least 5 years before study vaccination.

Table 7 compares the OPA GMTs, 1-month post-dose, in pneumococcal polysaccharide vaccinated adults aged  $\geq 70$  years given a single dose of either Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine.

<b>Table 7 : OPA GMTs in pneumococcal polysaccharide vaccinated adults aged <math>\geq 70</math> years given either Prevenar 13 or 23-valent pneumococcal polysaccharide vaccine (PPSV23)<sup>a,b,c</sup></b>				
	<b>Prevenar 13 N=400-426</b>	<b>PPSV23 N=395-445</b>	<b>Prevenar OPA GMT Relative to PPSV23</b>	
<b>Serotype</b>	<b>OPA GMT</b>	<b>OPA GMT</b>	<b>GMR</b>	<b>(95% CI)</b>
1	81	55	1.5	(1.17, 1.88)
3	55	49	1.1	(0.91, 1.35)
4	545	203	2.7	(1.93, 3.74)
5	72	36	2.0	(1.55, 2.63)
6A <sup>†</sup>	903	94	9.6	(7.00, 13.26)
6B	1261	417	3.0	(2.21, 4.13)
7F	245	160	1.5	(1.07, 2.18)
9V	181	90	2.0	(1.36, 2.97)
14	280	285	1.0	(0.73, 1.33)
18C	907	481	1.9	(1.42, 2.50)
19A	354	200	1.8	(1.43, 2.20)
19F	333	214	1.6	(1.17, 2.06)
23F	158	43	3.7	(2.69, 5.09)

<sup>a</sup> Non-inferiority was defined as the lower limit of the 2-sided 95% CI for GMR was greater than 0.5.  
<sup>b</sup> Statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR was greater than 1.  
<sup>c</sup> For serotype 6A<sup>†</sup>, which is unique to Prevenar 13, a statistically significantly greater response was defined as the lower bound of the 2-sided 95% CI for the GMR greater than 2.

In adults vaccinated with pneumococcal polysaccharide vaccine at least 5 years prior to the clinical study, OPA GMTs to Prevenar 13 were non-inferior to the 23-valent pneumococcal polysaccharide vaccine responses for the 12 serotypes in common. Furthermore, in this study statistically significantly greater OPA GMTs were demonstrated for 10 of the 12 serotypes in common. Immune responses to serotype 6A were statistically significantly greater following vaccination with Prevenar 13 than after 23-valent pneumococcal polysaccharide vaccine.

One year after vaccination with Prevenar 13 in adults aged 70 years and over who were vaccinated with 23-valent pneumococcal polysaccharide vaccine, at least 5 years prior to study entry, OPA titers had declined compared to one month after vaccination, however, OPA titers for all serotypes remained higher than levels at baseline:

	<b>OPA GMT levels at baseline</b>	<b>OPA GMT levels one year after Prevenar 13</b>
Adults $\geq 70$ years vaccinated with 23-valent pneumococcal polysaccharide vaccine at least 5 years prior	9 to 122	18 to 381

### Immune responses in Special Populations

Individuals with the conditions described below have an increased risk of pneumococcal disease. The clinical relevance of the antibody levels elicited by Prevenar 13 in these special populations is unknown.

#### *Sickle cell disease*

An open label single arm study in France, Italy, UK, US, Lebanon, Egypt and Saudi Arabia with 2 doses of Prevenar 13 given 6 months apart was conducted in 158 children and adolescents  $\geq 6$  to  $< 18$  years of age with sickle cell disease who were previously vaccinated with one or more doses of

23-valent pneumococcal polysaccharide vaccine at least 6 months prior to enrollment. After the first vaccination, Prevenar 13 elicited antibody levels measured by both IgG GMCs and OPA GMTs that were statistically significantly higher when compared to levels prior to vaccination. After the second dose immune responses were comparable to those after the first dose. One year after the second dose, antibody levels measured by both IgG GMCs and OPA GMTs were higher than levels prior to the first dose of Prevenar 13, except for the IgG GMCs for serotypes 3 and 5 that were numerically similar.

*Additional Prevenar (7-valent) immunogenicity data: children with sickle cell disease*

The immunogenicity of Prevenar has been investigated in an open-label, multicentre study in 49 infants with sickle cell disease. Children were vaccinated with Prevenar (3 doses one month apart from the age of 2 months), 46 of these children also received a 23-valent pneumococcal polysaccharide vaccine at the age of 15-18 months. After primary immunisation, 95.6 % of the subjects had antibody levels of at least 0.35 µg/ml for all seven serotypes found in Prevenar. A significant increase was seen in the concentrations of antibodies against the seven serotypes after the polysaccharide vaccination, suggesting that immunological memory was well established.

*HIV infection*

*Children and adults not previously vaccinated with a pneumococcal vaccine*

HIV-infected children and adults with CD4  $\geq$  200 cells/ $\mu$ L (mean 717.0 cells/ $\mu$ L), viral load  $<$  50,000 copies/ml (mean 2090.0 copies/ml), free of active AIDS-related illness and not previously vaccinated with a pneumococcal vaccine received 3 doses of Prevenar 13. As per general recommendations, a single dose of 23-valent pneumococcal polysaccharide vaccine was subsequently administered. Vaccines were administered at 1 month intervals. Immune responses were assessed in 259-270 evaluable subjects approximately 1 month after each dose of vaccine. After the first dose, Prevenar 13 elicited antibody levels, measured by both IgG GMCs and OPA GMTs that were statistically significantly higher when compared to levels prior to vaccination. After the second and third dose of Prevenar 13, immune responses were similar or higher than those after the first dose.

*Adults previously vaccinated with 23-valent pneumococcal polysaccharide vaccine*

HIV-infected adults  $\geq$  18 years of age with CD4  $\geq$  200 cells/ $\mu$ L (mean 609.1 cells/ $\mu$ L) and viral load  $<$  50,000 copies/ml (mean 330.6 copies/ml), who were free of active AIDS-related illness and were previously vaccinated with 23-valent pneumococcal polysaccharide vaccine administered at least 6 months prior to enrollment, received 3 doses of Prevenar 13, at enrollment, 6 months, and 12 months after the first dose of Prevenar 13. Immune responses were assessed in 231-255 evaluable subjects approximately 1 month after each dose of Prevenar 13. After the first dose, Prevenar 13 elicited antibody levels measured by both IgG GMCs and OPA GMTs that were statistically significantly higher when compared to levels prior to vaccination. After the second and third dose of Prevenar 13, immune responses were comparable or higher than those after the first dose. In the study 162 subjects had received one prior dose of 23-valent pneumococcal polysaccharide vaccine, 143 subjects 2 prior doses and 26 subjects more than 2 prior doses of 23-valent polysaccharide vaccine. Subjects who received two or more previous doses of 23-valent pneumococcal polysaccharide vaccine showed a similar immune response compared with subjects who received a single previous dose.

*Haematopoietic stem cell transplant*

Children and adults with an allogeneic haematopoietic stem cell transplant (HSCT) at  $\geq$  2 years of age with complete haematologic remission of underlying disease or with very good partial remission in the case of lymphoma and myeloma received three doses of Prevenar 13 with an interval of at least 1 month between doses. The first dose was administered at 3 to 6 months after HSCT. A fourth (booster) dose of Prevenar 13 was administered 6 months after the third dose. As per general recommendations, a single dose of 23-valent pneumococcal polysaccharide vaccine was administered 1 month after the fourth dose of Prevenar 13. Immune responses as measured by IgG GMCs were assessed in 168-211 evaluable subjects approximately 1 month after vaccination. Prevenar 13 elicited increased antibody levels after each dose of Prevenar 13. Immune responses after the fourth dose of Prevenar 13 were significantly increased for all serotypes compared with after the third dose. Functional antibody titers (OPA titers) were not measured in this study.

## **5.2 Pharmacokinetic properties**

Not applicable.

## **5.3 Preclinical safety data**

Non-clinical data revealed no special hazard for humans based on conventional studies of safety pharmacology, single and repeated dose toxicity, local tolerance, and reproduction and developmental toxicity.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Sodium chloride  
Succinic acid  
Polysorbate 80  
2-phenoxyethanol  
Water for injections

For adjuvant, see section 2.

### **6.2 Incompatibilities**

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

2 years

#### After first use:

Once opened, the product may be stored for a maximum of 28 days at 2-8 °C. Other in-use storage times and conditions are the responsibility of the user.

### **6.4 Special precautions for storage**

Store in a refrigerator (2°C–8°C). Do not freeze.

For storage conditions after first opening of the medicinal product, see section 6.3.

### **6.5 Nature and contents of container**

2 ml (4 x 0.5 ml doses) suspension for injection in a container (Type I glass) with a latex-free grey chlorobutyl rubber stopper and sealed with an aluminum flip-off seal and a polypropylene flip-off cap.

Pack sizes of 1, 5, 10, 25 and 50.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal and other handling**

During storage, a white deposit and clear supernatant can be observed. This does not constitute a sign of deterioration.

The vaccine should be shaken well to obtain a homogeneous white suspension and should be inspected visually for any particulate matter and/or variation of physical aspect prior to administration. Do not use if the content appears otherwise.

No special requirements for disposal.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Pfizer Limited  
Ramsgate Road  
Sandwich  
Kent CT13 9NJ  
United Kingdom

## **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/09/590/012  
EU/1/09/590/013  
EU/1/09/590/014  
EU/1/09/590/015  
EU/1/09/590/016

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 09 December 2009  
Date of latest renewal: 18 September 2014

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## **ANNEX II**

- A. MANUFACTURER(S) OF THE BIOLOGICAL ACTIVE SUBSTANCE(S) AND MANUFACTURER (S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

**A. MANUFACTURER(S) OF THE BIOLOGICAL ACTIVE SUBSTANCE(S) AND  
MANUFACTURER (S) RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturers of the biological active substance

Wyeth BioPharma Division of Wyeth Pharmaceuticals Inc., a subsidiary of Pfizer Inc.  
One Burtt Road  
Andover, MA 01810  
USA

Pfizer Ireland Pharmaceuticals  
Grange Castle Business Park  
Clondalkin  
Dublin 22  
Ireland

Wyeth Pharmaceutical Division of Wyeth Holdings LLC, a subsidiary of Pfizer Inc.  
4300 Oak Park  
Sanford, NC 27330  
USA

Name and address of the manufacturers responsible for batch release

Pre-filled syringes:

Wyeth Pharmaceuticals  
New Lane  
Havant  
Hampshire PO9 2NG  
United Kingdom

Pfizer Manufacturing Belgium N.V.  
Rijksweg 12  
B-2870 Puurs  
Belgium

Single-dose vials:

Wyeth Pharmaceuticals  
New Lane  
Havant  
Hampshire PO9 2NG  
United Kingdom

Pfizer Manufacturing Belgium N.V.  
Rijksweg 12  
B-2870 Puurs  
Belgium

Multidose containers:

Pfizer Manufacturing Belgium N.V.  
Rijksweg 12  
B-2870 Puurs  
Belgium

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to medical prescription.

- **Official batch release**

In accordance with Article 114 Directive 2001/83/EC, the official batch release will be undertaken by a state laboratory or a laboratory designated for that purpose.

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic Safety Update Reports**

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**Pack of 1 or 10 pre-filled syringe, with or without needle – WITH BLUE BOX**

**1. NAME OF THE MEDICINAL PRODUCT**

Prevenar 13 suspension for injection  
Pneumococcal polysaccharide conjugate vaccine (13-valent, adsorbed)

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

1 dose (0.5 ml) contains 2.2 µg of polysaccharide for serotypes 1, 3, 4, 5, 6A, 7F, 9V, 14, 18C, 19A, 19F, 23F and 4.4 µg for serotype 6B.

**3. LIST OF EXCIPIENTS**

Sodium chloride, succinic acid, polysorbate 80 and water for injections

**4. PHARMACEUTICAL FORM AND CONTENTS**

Suspension for injection  
1 single-dose (0.5 ml) pre-filled syringe with separate needle  
1 single-dose (0.5 ml) pre-filled syringe without needle  
10 single-dose (0.5 ml) pre-filled syringes with separate needles  
10 single-dose (0.5 ml) pre-filled syringes without needle

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Intramuscular use.  
Shake well before use.  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator.

Do not freeze.

Prevenar 13 is stable at temperatures up to 25°C for four days. At the end of this period Prevenar 13 should be used or discarded.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Pfizer Limited  
Ramsgate Road  
Sandwich  
Kent CT13 9NJ  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/09/590/002 – pack of 1 with separate needle  
EU/1/09/590/001 – pack of 1 without needle  
EU/1/09/590/004 – pack of 10 with separate needles  
EU/1/09/590/003 – pack of 10 without needle

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

<b>18.    UNIQUE IDENTIFIER - HUMAN READABLE DATA</b>
---

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**Pack of 10 - for multipack of 50 (5 x 10) pre-filled syringes, with or without needle – WITHOUT BLUE BOX**

**1. NAME OF THE MEDICINAL PRODUCT**

Prevenar 13 suspension for injection  
Pneumococcal polysaccharide conjugate vaccine (13-valent, adsorbed)

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

1 dose (0.5 ml) contains 2.2 µg of polysaccharide for serotypes 1, 3, 4, 5, 6A, 7F, 9V, 14, 18C, 19A, 19F, 23F and 4.4 µg for serotype 6B.

**3. LIST OF EXCIPIENTS**

Sodium chloride, succinic acid, polysorbate 80 and water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Suspension for injection  
10 pre-filled syringes with separate needles, each containing 1 single-dose (0.5 ml). Component of a multipack, can't be sold separately.  
10 pre-filled syringes without needles, each containing 1 single-dose (0.5 ml). Component of a multipack, can't be sold separately.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Intramuscular use.  
Shake well before use.  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator.

Do not freeze.

Prevenar 13 is stable at temperatures up to 25°C for four days. At the end of this period Prevenar 13 should be used or discarded.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Pfizer Limited  
Ramsgate Road  
Sandwich  
Kent CT13 9NJ  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/09/590/006 – pack of 50 (5 x 10) with separate needles

EU/1/09/590/005 – pack of 50 (5 x 10) without needle

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

Not applicable.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

Not applicable.

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**Multipack of 50 (5 x 10) pre-filled syringes, with or without needle, (outer wrapper label to be applied on transparent foil) - WITH BLUE BOX**

**1. NAME OF THE MEDICINAL PRODUCT**

Prevenar 13 suspension for injection  
Pneumococcal polysaccharide conjugate vaccine (13-valent, adsorbed)

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

1 dose (0.5 ml) contains 2.2 µg of polysaccharide for serotypes 1, 3, 4, 5, 6A, 7F, 9V, 14, 18C, 19A, 19F, 23F and 4.4 µg for serotype 6B.

**3. LIST OF EXCIPIENTS**

Sodium chloride, succinic acid, polysorbate 80 and water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Suspension for injection  
Multipack comprising 5 packs, each containing 10 single-dose (0.5 ml) pre-filled syringes with separate needles.  
Multipack comprising 5 packs, each containing 10 single-dose (0.5 ml) pre-filled syringes without needle.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Intramuscular use.  
Shake well before use.  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator.

Do not freeze.

Prevenar 13 is stable at temperatures up to 25°C for four days. At the end of this period Prevenar 13 should be used or discarded.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Pfizer Limited  
Ramsgate Road  
Sandwich  
Kent CT13 9NJ  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/09/590/006 – pack of 50 (5 x 10) with separate needles

EU/1/09/590/005 – pack of 50 (5 x 10) without needle

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

<b>18.    UNIQUE IDENTIFIER - HUMAN READABLE DATA</b>
---

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1, 5, 10, 25, 50 single dose vials**

**1. NAME OF THE MEDICINAL PRODUCT**

Prevenar 13 suspension for injection in single dose vial  
Pneumococcal polysaccharide conjugate vaccine (13-valent, adsorbed)

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

1 dose (0.5 ml) contains 2.2 µg of polysaccharide for serotypes 1, 3, 4, 5, 6A, 7F, 9V, 14, 18C, 19A, 19F, 23F and 4.4 µg for serotype 6B.

**3. LIST OF EXCIPIENTS**

Sodium chloride, succinic acid, polysorbate 80 and water for injections

**4. PHARMACEUTICAL FORM AND CONTENTS**

Suspension for injection  
1 single-dose (0.5 ml) vial  
5 single-dose (0.5 ml) vials  
10 single-dose (0.5 ml) vials  
25 single-dose (0.5 ml) vials  
50 single-dose (0.5 ml) vials

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Intramuscular use.  
Shake well before use.  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator.  
Do not freeze.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Pfizer Limited  
Ramsgate Road  
Sandwich  
Kent CT13 9NJ  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/09/590/007 – pack of 1 vial  
EU/1/09/590/008 – pack of 5 vials  
EU/1/09/590/009 – pack of 10 vials  
EU/1/09/590/010 – pack of 25 vials  
EU/1/09/590/011 – pack of 50 vials

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING  
OUTER CARTON**

**1, 5, 10, 25, 50 multidose containers**

**1. NAME OF THE MEDICINAL PRODUCT**

Prevenar 13 suspension for injection in multidose container  
Pneumococcal polysaccharide conjugate vaccine (13-valent, adsorbed)

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

1 dose (0.5 ml) contains 2.2 µg of polysaccharide for serotypes 1, 3, 4, 5, 6A, 7F, 9V, 14, 18C, 19A, 19F, 23F and 4.4 µg for serotype 6B.

**3. LIST OF EXCIPIENTS**

Sodium chloride, succinic acid, polysorbate 80, 2-phenoxyethanol and water for injections

**4. PHARMACEUTICAL FORM AND CONTENTS**

Suspension for injection in multidose container  
1 multidose container (4 x 0.5 ml doses)  
5 multidose containers (4 x 0.5 ml doses per containers)  
10 multidose containers (4 x 0.5 ml doses per containers)  
25 multidose containers (4 x 0.5 ml doses per containers)  
50 multidose containers (4 x 0.5 ml doses per containers)

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Intramuscular use.  
Shake well before use.  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT  
OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator.

Do not freeze.

After first use, the product may be stored in a refrigerator for a maximum of 28 days.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Pfizer Limited  
Ramsgate Road  
Sandwich  
Kent CT13 9NJ  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/09/590/012 – pack of 1 multidose container  
EU/1/09/590/013– pack of 5 multidose containers  
EU/1/09/590/014– pack of 10 multidose containers  
EU/1/09/590/015 – pack of 25 multidose containers  
EU/1/09/590/016– pack of 50 multidose containers

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

<b>18.    UNIQUE IDENTIFIER - HUMAN READABLE DATA</b>
---

PC:  
SN:  
NN:

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**Pre-filled syringe**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Prevenar 13 suspension for injection  
IM

**2. METHOD OF ADMINISTRATION**

Shake well before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

1 dose (0.5 ml)

**6. OTHER**

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**Single Dose Vial Label**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Prevenar 13 suspension for injection in single dose vial  
IM

**2. METHOD OF ADMINISTRATION**

Shake well before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

1 dose (0.5 ml)

**6. OTHER**

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**Multidose Container Label**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Prevenar 13 suspension for injection in multidose container  
IM

**2. METHOD OF ADMINISTRATION**

Shake well before use.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

4 x 0.5 ml doses

**6. OTHER**

**B. PACKAGE LEAFLET**

## Package Leaflet: Information for the user

### Prevenar 13 suspension for injection

Pneumococcal polysaccharide conjugate vaccine (13 valent, adsorbed)

**Read all of this leaflet carefully before you or your child receives this vaccine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This vaccine has been prescribed for you or your child only. Do not pass it on to others.
- If you or your child get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Prevenar 13 is and what it is used for
2. What you need to know before you or your child receives Prevenar 13
3. How Prevenar 13 is given
4. Possible side effects
5. How to store Prevenar 13
6. Contents of the pack and other information

#### 1. What Prevenar 13 is and what it is used for

Prevenar 13 is a pneumococcal vaccine given to:

- **children from 6 weeks to 17 years** to help protect against diseases such as: meningitis (inflammation around the brain), sepsis or bacteraemia (bacteria in the blood stream), pneumonia (lung infection) and ear infections
- **adults aged 18 years and older** to help prevent disease such as: pneumonia (lung infection), sepsis or bacteraemia (bacteria in the blood stream) and meningitis (inflammation around the brain),

caused by 13 types of the bacteria *Streptococcus pneumoniae*.

Prevenar 13, provides protection against 13 types of *Streptococcus pneumoniae* bacteria, and replaces Prevenar, which provided protection against 7 types.

The vaccine works by helping the body to make its own antibodies, which protect you or your child against these diseases.

#### 2. What you need to know before you or your child receives Prevenar 13

##### Prevenar 13 should not be given:

- if you or your child is allergic (hypersensitive) to the active substances or to any of the other ingredients in this medicine (listed in section 6) or to any other vaccine that contains diphtheria toxoid.
- if you or your child has a severe infection with a high temperature (over 38°C). If this applies to you or your child, then the vaccination will be postponed until you or your child is feeling better. A minor infection, such as a cold, should not be a problem. However, talk to your doctor, pharmacist, or nurse first.

## Warnings and precautions

Talk to your doctor, pharmacist, or nurse before the vaccination if you or your child:

- has any present or past medical problems after any dose of Prevenar or Prevenar 13 such as an allergic reaction or problems with breathing.
- has any bleeding problems or bruises easily.
- has a weakened immune system (such as due to HIV infection), you/she/he may not get the full benefit from Prevenar 13.
- has experienced seizures as medicines to lower fever may need to be taken before Prevenar 13 is given. If your child should become unresponsive or experience seizures (fits) after the vaccination, please contact your doctor immediately. See also section 4.

Talk to your doctor, pharmacist, or nurse before the vaccination if your child was born very prematurely (at or before 28 weeks of gestation), as longer gaps than normal between breaths may occur for 2-3 days after vaccination. See also section 4.

As with any vaccine, Prevenar 13 will not protect all persons who are vaccinated.

Prevenar 13 will only protect against ear infections in children caused by the types of *Streptococcus pneumoniae* for which the vaccine has been developed. It will not protect against other infectious agents that can cause ear infections.

### Other medicines/vaccines and Prevenar 13

Your doctor may ask you to give your child paracetamol or other medicines that lower fever before Prevenar 13 is given. This will help to lower some of the side effects of Prevenar 13.

Tell your doctor, pharmacist or nurse if you or your child is taking, has recently taken or might take any other medicines, or has recently received any other vaccine.

### Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before receiving this medicine.

### Driving and using machines

Prevenar 13 has no or negligible influence on the ability to drive and use machines. However, some of the effects mentioned under section 4 "Possible side effects" may temporarily affect the ability to drive or use machines.

### Prevenar 13 contains sodium

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially 'sodium-free'.

## 3. How Prevenar 13 is given

The doctor or nurse will inject the recommended dose (0.5 ml) of the vaccine into your arm or your child's arm or leg muscle.

### Infants aged 6 weeks to 6 months of age

Typically, your child should receive an initial course of three injections of the vaccine followed by a booster dose.

- The first injection may be given from the age of six weeks.
- Each injection will be given at least one month apart.
- A fourth injection (booster) will be given between 11 and 15 months of age.
- You will be told when your child should come back for the next injection.

According to official recommendations in your country, an alternative schedule may be used by your healthcare provider. Please speak to your doctor, pharmacist, or nurse for more information.

#### Premature infants

Your child will receive an initial course of three injections. The first injection may be given as early as six weeks of age with at least one month between doses. Between 11 and 15 months of age, your child will receive a fourth injection (booster).

#### Unvaccinated infants, children, and adolescents over 7 months of age

Infants aged **7 to 11 months** should receive two injections. Each injection will be given at least one month apart. A third injection will be given in the second year of life.

Children aged **12 to 23 months** should receive two injections. Each injection will be given at least two months apart.

Children aged **2 to 17 years** should receive one injection.

#### Infants, children, and adolescents previously vaccinated with Prevenar

Infants and children who have previously received Prevenar may receive Prevenar 13 to complete the course of injections.

For children **1 to 5 years** of age previously vaccinated with Prevenar, your doctor or nurse will recommend how many injections of Prevenar 13 are required.

Children and adolescents **6 to 17 years** of age should receive one injection.

It is important to follow the instructions from the doctor, pharmacist, or nurse so that your child completes the course of injections.

If you forget to go back at the scheduled time, ask the doctor, pharmacist, or nurse for advice.

#### Adults

Adults should receive one injection.

Tell your doctor, pharmacist or nurse if you have been given a pneumococcal vaccine before.

If you have any further questions on the use of Prevenar 13, ask your doctor, pharmacist, or nurse.

#### Special Populations

Individuals considered to be at a higher risk of pneumococcal infection (such as those with sickle cell disease or HIV infection), including those previously vaccinated with the 23-valent pneumococcal polysaccharide vaccine, may receive at least one dose of Prevenar 13.

Individuals with a blood-forming stem cell transplant may receive three injections, with the first given at 3 to 6 months after the transplant and with an interval of at least 1 month between doses. A fourth injection (booster) is recommended 6 months after the third injection.

#### **4. Possible side effects**

Like all vaccines, Prevenar 13 can cause side effects, although not everybody gets them.

**The following side effects include those reported for Prevenar 13 in infants and children (6 weeks to 5 years of age):**

**The most common side effects** (these may occur with more than 1 in 10 doses of the vaccine) are:

- Decreased appetite
- Fever; irritability; pain, tenderness, redness, swelling or hardness at the vaccination-site; drowsiness; restless sleep
- Redness, hardness, swelling at the vaccination-site of 2.5 cm -7.0 cm (after the booster dose and in older children [aged 2 to 5 years])

**Common side effects** (these may occur with up to 1 in 10 doses of the vaccine) are:

- Vomiting; diarrhoea
- Fever of more than 39°C; tenderness at the vaccination-site interfering with movement, redness, hardness, swelling at the vaccination-site of 2.5 cm -7.0 cm (after the initial course of injections)
- Rash

**Uncommon side effects** (these may occur with up to 1 in 100 doses of the vaccine) are:

- Seizures (or fits), including those caused by a high temperature
- Hives (urticaria or urticaria-like rash)
- Redness, swelling, or hardness at the vaccination-site of more than 7 cm; crying

**Rare side effects** (these may occur with up to 1 in 1,000 doses of the vaccine) are:

- Collapse or shock-like state (hypotonic-hyporesponsive episode)
- Allergic (hypersensitivity) reaction, including swelling of the face and/or lips, difficulty in breathing

**The following side effects include those reported for Prevenar 13 in children and adolescents (6 to 17 years of age):**

**The most common side effects** (these may occur with more than 1 in 10 doses of the vaccine) are:

- Decreased appetite
- Irritability; pain, tenderness, redness, swelling or hardness at the vaccination-site; drowsiness; restless sleep; tenderness at the vaccination-site interfering with movement

**Common side effects** (these may occur with up to 1 in 10 doses of the vaccine) are:

- Headaches
- Vomiting; diarrhoea
- Rash; hives (urticaria or urticaria-like rash)
- Fever

Children and adolescents with either HIV infection, sickle cell disease or a blood-forming stem cell transplant had similar side effects however the frequencies of headaches, vomiting, diarrhoea, fever, fatigue, joint and muscle pain were very common.

**The following additional side effects have been seen with Prevenar 13 in infants and children up to 5 years of age in postmarketing experience**

- Severe allergic reaction including shock (cardiovascular collapse); angioedema (swelling of lips, face or throat)
- Hives (urticaria), redness and irritation (dermatitis) and itching (pruritus) at the vaccination-site; flushing
- Enlarged lymph nodes or glands (lymphadenopathy) near the vaccination-site, such as under the arm or in the groin
- A rash causing itchy red blotches (erythema multiforme)

In babies born very prematurely (at or before 28 weeks of gestation), longer gaps than normal between breaths may occur for 2-3 days after vaccination.

**The following side effects include those reported for Prevenar 13 in adults:**

**The most common side effects** (these may occur with more than 1 in 10 doses of the vaccine) are:

- Decreased appetite; headaches; diarrhoea; vomiting (for those 18 to 49 years of age)
- Chills; tiredness; rash; pain, redness, swelling hardness or tenderness at the vaccination-site, interfering with arm movement (severe pain or tenderness at vaccination-site for those 18-39 years of age and severe limitation of arm movements for those 18 to 39 years of age)
- Worsening or new pain in your joints, worsening or new pain in your muscles
- Fever (for those 18 to 29 years of age)

**Common side effects** (these may occur with up to 1 in 10 doses of the vaccine) are:

- Vomiting (for those 50 years and older); fever (for those 30 years and older)

**Uncommon side effects** (these may occur with up to 1 in 100 doses of the vaccine) are:

- Nausea
- Allergic (hypersensitivity) reaction, including swelling of the face and/or lips, difficulty in breathing
- Enlarged lymph nodes or glands (lymphadenopathy) near the vaccination-site, such as under the arm

Adults with HIV infection had similar side effects however the frequencies were very common for fever, vomiting and common for nausea.

Adults with a blood-forming stem cell transplant had similar side effects however the frequencies were very common for fever and vomiting.

**Reporting of side effects**

If you or your child get any side effects, talk to your doctor, pharmacist, or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via **the national reporting system listed in Appendix V**. By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Prevenar 13

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and label after EXP. The expiry date refers to the last day of that month.

Store in a refrigerator (2°C – 8°C).

Do not freeze.

Prevenar 13 is stable at temperatures up to 25°C for four days. At the end of this period Prevenar 13 should be used or discarded. These data are intended to guide health care professionals in case of temporary temperature excursions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Prevenar 13 contains

The active substances are polysaccharide CRM<sub>197</sub> conjugates consisting of:

- 2.2 µg of polysaccharide for serotypes 1, 3, 4, 5, 6A, 7F, 9V, 14, 18C, 19A, 19F and 23F
- 4.4 µg of polysaccharide for serotype 6B

1 dose (0.5 ml) contains approximately 32 µg CRM<sub>197</sub> carrier protein, adsorbed on aluminium phosphate (0.125 mg aluminium).

The other ingredients are sodium chloride, succinic acid, polysorbate 80 and water for injections.

### What Prevenar 13 looks like and contents of the pack

The vaccine is a white suspension for injection, provided in a single-dose, pre-filled syringe (0.5 ml). Pack sizes of 1 and 10, with or without needle, and a multipack each containing 5 packs of 10 pre-filled syringes, with or without needle. Not all pack sizes may be marketed.

### Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:  
Pfizer Limited  
Ramsgate Road  
Sandwich  
Kent CT13 9NJ  
United Kingdom

Manufacturer responsible for batch release:  
Wyeth Pharmaceuticals  
New Lane  
Havant  
Hampshire, PO9 2NG  
United Kingdom

Pfizer Manufacturing Belgium N.V.  
Rijksweg 12  
B-2870 Puurs  
Belgium

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**  
**Luxembourg/Luxemburg**  
Pfizer S.A./N.V.  
Tél/Tel: + 32 (0)2 554 62 11

**България**  
Пфайзер Люксембург САРЛ, Клон  
България  
Тел: +359 2 970 4333

**Česká Republika**  
Pfizer PFE, spol. s r.o.  
Tel: +420 283 004 111

**Danmark**  
Pfizer ApS  
Tlf: + 45 44 201 100

**Deutschland**  
Pfizer Pharma GmbH  
Tel: + 49 (0)30 550055-51000

**Eesti**  
Pfizer Luxembourg SARL Eesti filiaal  
Tel: +372 666 7500

**Ελλάδα**  
Pfizer Ελλάς Α.Ε.  
Τηλ.: +30 210 6785 800

**España**  
Pfizer, S.L.  
Télf:+34914909900

**France**  
Pfizer  
Tél +33 1 58 07 34 40

**Hrvatska**  
Pfizer Croatia d.o.o.  
Tel: + 385 1 3908 777

**Ireland**  
Pfizer Healthcare Ireland  
Tel: 1800 633 363 (toll free)  
+44 (0)1304 616161

**Ísland**  
Icepharma hf  
Simi: + 354 540 8000

**Lietuva**  
Pfizer Luxembourg SARL filialas Lietuvoje  
Tel. + 370 52 51 4000

**Magyarország**  
Pfizer Kft  
Tel: +36 1 488 3700

**Malta**  
Vivian Corporation Ltd.  
Tel: + 35621 344610

**Nederland**  
Pfizer BV  
Tel: +31 (0)10 406 43 01

**Norge**  
Pfizer Norge AS  
Tlf: +47 67 52 61 00

**Österreich**  
Pfizer Corporation Austria Ges.m.b.H  
Tel: + 43 (0)1 521 15-0

**Polska**  
Pfizer Polska Sp. z o.o.  
Tel.: +48 22 335 61 00

**Portugal**  
Pfizer Biofarmacêutica, Sociedade Unipessoal Lda  
Tel: +351 21 423 5500

**România**  
Pfizer Romania S.R.L  
Tel: +40 (0) 21 207 28 00

**Slovenija**  
Pfizer Luxembourg SARL  
Pfizer, podružnica za svetovanje s področja  
farmaceutvske dejavnosti, Ljubljana  
Tel.: + 386 (0) 1 52 11 400

**Slovenská republika**  
Pfizer Luxembourg SARL,  
organizačná zložka  
Tel: + 421 2 3355 5500

**Suomi/Finland**  
Pfizer Oy  
Puh/Tel: +358 (0)9 430 040

**Italia**

Pfizer S.r.l.  
Tel: +39 06 33 18 21

**Sverige**

Pfizer AB  
Tel: +46 (0)8 550 520 00

**Κύπρος**

Pfizer Ελλάς A.E. (Cyprus Branch)  
Τηλ: +357 22 817690

**United Kingdom**

Pfizer Limited  
Tel: +44 (0) 1304 616161

**Latvija**

Pfizer Luxembourg SARL filiāle Latvijā  
Tel.: + 371 670 35 775

**This leaflet was last revised in**

**Other sources of information**

Detailed information on this product is available on the European Medicines Agency website:  
<http://www.ema.europa.eu>.

---

**The following information is intended for healthcare professionals only:**

During storage, a white deposit and clear supernatant may be observed. This does not constitute a sign of deterioration.

Inspect visually for any foreign particulate matter and/or abnormal physical appearance; do not use if either are found.

Prior to expelling air from the syringe, shake well to obtain a homogeneous white suspension.

Administer the entire dose.

Prevenar 13 is for intramuscular use only. Do not administer intravascularly.

Prevenar 13 must not be mixed with any other vaccines in the same syringe.

Prevenar 13 can be given at the same time as other childhood vaccines; in this case, different vaccination sites should be used.

Prevenar 13 may be given to adults aged 50 years and older at the same time as the trivalent or quadrivalent inactivated influenza vaccine.

Any unused product or waste material should be disposed of in accordance with local requirements.

## Package Leaflet: Information for the user

### **Prevenar 13 suspension for injection in single dose vial** Pneumococcal polysaccharide conjugate vaccine (13 valent, adsorbed)

**Read all of this leaflet carefully before you or your child receives this vaccine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This vaccine has been prescribed for you or your child only. Do not pass it on to others.
- If you or your child get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Prevenar 13 is and what it is used for
2. What you need to know before you or your child receives Prevenar 13
3. How Prevenar 13 is given
4. Possible side effects
5. How to store Prevenar 13
6. Contents of the pack and other information

#### **1. What Prevenar 13 is and what it is used for**

Prevenar 13 is a pneumococcal vaccine given to:

- **children from 6 weeks to 17 years** to help protect against diseases such as: meningitis (inflammation around the brain), sepsis or bacteraemia (bacteria in the blood stream), pneumonia (lung infection) and ear infections
- **adults aged 18 years and older** to help prevent disease such as: pneumonia (lung infection), sepsis or bacteraemia (bacteria in the blood stream) and meningitis (inflammation around the brain),

caused by 13 types of the bacteria *Streptococcus pneumoniae*.

Prevenar 13, provides protection against 13 types of *Streptococcus pneumoniae* bacteria, and replaces Prevenar, which provided protection against 7 types.

The vaccine works by helping the body to make its own antibodies, which protect you or your child against these diseases.

#### **2. What you need to know before you or your child receives Prevenar 13**

##### **Prevenar 13 should not be given:**

- if you or your child is allergic (hypersensitive) to the active substances or to any of the other ingredients in this medicine (listed in section 6) or to any other vaccine that contains diphtheria toxoid.
- if you or your child has a severe infection with a high temperature (over 38°C). If this applies to you or your child, then the vaccination will be postponed until you or your child is feeling better. A minor infection, such as a cold, should not be a problem. However, talk to your doctor, pharmacist, or nurse first.

## Warnings and precautions

Talk to your doctor, pharmacist, or nurse before the vaccination if you or your child:

- has any present or past medical problems after any dose of Prevenar or Prevenar 13 such as an allergic reaction or problems with breathing.
- has any bleeding problems or bruises easily.
- has a weakened immune system (such as due to HIV infection), you/she/he may not get the full benefit from Prevenar 13.
- has experienced seizures as medicines to lower fever may need to be taken before Prevenar 13 is given. If your child should become unresponsive or experience seizures (fits) after the vaccination, please contact your doctor immediately. See also section 4.

Talk to your doctor, pharmacist, or nurse before the vaccination if your child was born very prematurely (at or before 28 weeks of gestation), as longer gaps than normal between breaths may occur for 2-3 days after vaccination. See also section 4.

As with any vaccine, Prevenar 13 will not protect all persons who are vaccinated.

Prevenar 13 will only protect against ear infections in children caused by the types of *Streptococcus pneumoniae* for which the vaccine has been developed. It will not protect against other infectious agents that can cause ear infections.

### Other medicines/vaccines and Prevenar 13

Your doctor may ask you to give your child paracetamol or other medicines that lower fever before Prevenar 13 is given. This will help to lower some of the side effects of Prevenar 13.

Tell your doctor, pharmacist or nurse if you or your child is taking, has recently taken or might take any other medicines, or has recently received any other vaccine.

### Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before receiving this medicine.

### Driving and using machines

Prevenar 13 has no or negligible influence on the ability to drive and use machines. However, some of the effects mentioned under section 4 “Possible side effects” may temporarily affect the ability to drive or use machines.

### Prevenar 13 contains sodium

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially ‘sodium-free’.

## 3. How Prevenar 13 is given

The doctor or nurse will inject the recommended dose (0.5 ml) of the vaccine into your arm or your child's arm or leg muscle.

### Infants aged 6 weeks to 6 months of age

Typically, your child should receive an initial course of three injections of the vaccine followed by a booster dose.

- The first injection may be given from the age of six weeks.
- Each injection will be given at least one month apart.
- A fourth injection (booster) will be given between 11 and 15 months of age.
- You will be told when your child should come back for the next injection.

According to official recommendations in your country, an alternative schedule may be used by your healthcare provider. Please speak to your doctor, pharmacist, or nurse for more information.

#### Premature infants

Your child will receive an initial course of three injections. The first injection may be given as early as six weeks of age with at least one month between doses. Between 11 and 15 months of age, your child will receive a fourth injection (booster).

#### Unvaccinated infants, children, and adolescents over 7 months of age

Infants aged **7 to 11 months** should receive two injections. Each injection will be given at least one month apart. A third injection will be given in the second year of life.

Children aged **12 to 23 months** should receive two injections. Each injection will be given at least two months apart.

Children aged **2 to 17 years** should receive one injection.

#### Infants, children, and adolescents previously vaccinated with Prevenar

Infants and children who have previously received Prevenar may receive Prevenar 13 to complete the course of injections.

For children **1 to 5 years** of age previously vaccinated with Prevenar, your doctor or nurse will recommend how many injections of Prevenar 13 are required.

Children and adolescents **6 to 17 years** of age should receive one injection.

It is important to follow the instructions from the doctor, pharmacist, or nurse so that your child completes the course of injections.

If you forget to go back at the scheduled time, ask the doctor, pharmacist, or nurse for advice.

#### Adults

Adults should receive one injection.

Tell your doctor, pharmacist or nurse if you have been given a pneumococcal vaccine before.

If you have any further questions on the use of Prevenar 13, ask your doctor, pharmacist, or nurse.

#### Special Populations

Individuals considered to be at a higher risk of pneumococcal infection (such as those with sickle cell disease or HIV infection), including those previously vaccinated with the 23-valent pneumococcal polysaccharide vaccine, may receive at least one dose of Prevenar 13.

Individuals with a blood-forming stem cell transplant may receive three injections, with the first given at 3 to 6 months after the transplant and with an interval of at least 1 month between doses. A fourth injection (booster) is recommended 6 months after the third injection.

#### **4. Possible side effects**

Like all vaccines, Prevenar 13 can cause side effects, although not everybody gets them.

**The following side effects include those reported for Prevenar 13 in infants and children (6 weeks to 5 years of age):**

**The most common side effects** (these may occur with more than 1 in 10 doses of the vaccine) are:

- Decreased appetite
- Fever; irritability; pain, tenderness, redness, swelling or hardness at the vaccination-site; drowsiness; restless sleep
- Redness, hardness, swelling at the vaccination-site of 2.5 cm -7.0 cm (after the booster dose and in older children [aged 2 to 5 years])

**Common side effects** (these may occur with up to 1 in 10 doses of the vaccine) are:

- Vomiting; diarrhoea
- Fever of more than 39°C; tenderness at the vaccination-site interfering with movement, redness, hardness, swelling at the vaccination-site of 2.5 cm -7.0 cm (after the initial course of injections)
- Rash

**Uncommon side effects** (these may occur with up to 1 in 100 doses of the vaccine) are:

- Seizures (or fits), including those caused by a high temperature
- Hives (urticaria or urticaria-like rash)
- Redness, swelling, or hardness at the vaccination-site of more than 7 cm; crying

**Rare side effects** (these may occur with up to 1 in 1,000 doses of the vaccine) are:

- Collapse or shock-like state (hypotonic-hyporesponsive episode)
- Allergic (hypersensitivity) reaction, including swelling of the face and/or lips, difficulty in breathing

**The following side effects include those reported for Prevenar 13 in children and adolescents (6 to 17 years of age):**

**The most common side effects** (these may occur with more than 1 in 10 doses of the vaccine) are:

- Decreased appetite
- Irritability; pain, tenderness, redness, swelling or hardness at the vaccination-site; drowsiness; restless sleep; tenderness at the vaccination-site interfering with movement

**Common side effects** (these may occur with up to 1 in 10 doses of the vaccine) are:

- Headaches
- Vomiting; diarrhoea
- Rash; hives (urticaria or urticaria-like rash)
- Fever

Children and adolescents with either HIV infection, sickle cell disease or a blood-forming stem cell transplant had similar side effects however the frequencies of headaches, vomiting, diarrhoea, fever, fatigue, joint and muscle pain were very common.

**The following additional side effects have been seen with Prevenar 13 in infants and children up to 5 years of age in postmarketing experience**

- Severe allergic reaction including shock (cardiovascular collapse); angioedema (swelling of lips, face or throat)
- Hives (urticaria), redness and irritation (dermatitis) and itching (pruritus) at the vaccination-site; flushing
- Enlarged lymph nodes or glands (lymphadenopathy) near the vaccination-site, such as under the arm or in the groin
- A rash causing itchy red blotches (erythema multiforme)

In babies born very prematurely (at or before 28 weeks of gestation), longer gaps than normal between breaths may occur for 2-3 days after vaccination.

**The following side effects include those reported for Prevenar 13 in adults:**

**The most common side effects** (these may occur with more than 1 in 10 doses of the vaccine) are:

- Decreased appetite; headaches; diarrhoea; vomiting (for those 18 to 49 years of age)
- Chills; tiredness; rash; pain, redness, swelling hardness or tenderness at the vaccination-site, interfering with arm movement (severe pain or tenderness at vaccination-site for those 18-39 years of age and severe limitation of arm movements for those 18 to 39 years of age)
- Worsening or new pain in your joints, worsening or new pain in your muscles
- Fever (for those 18 to 29 years of age)

**Common side effects** (these may occur with up to 1 in 10 doses of the vaccine) are:

- Vomiting (for those 50 years and older); fever (for those 30 years and older)

**Uncommon side effects** (these may occur with up to 1 in 100 doses of the vaccine) are:

- Nausea
- Allergic (hypersensitivity) reaction, including swelling of the face and/or lips, difficulty in breathing
- Enlarged lymph nodes or glands (lymphadenopathy) near the vaccination-site, such as under the arm

Adults with HIV infection had similar side effects however the frequencies were very common for fever, vomiting and common for nausea.

Adults with a blood-forming stem cell transplant had similar side effects however the frequencies were very common for fever and vomiting.

**Reporting of side effects**

If you or your child get any side effects, talk to your doctor, pharmacist, or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via **the national reporting system listed in Appendix V**. By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Prevenar 13

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and label after EXP. The expiry date refers to the last day of that month.

Store in a refrigerator (2°C – 8°C).

Do not freeze.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Prevenar 13 contains

The active substances are polysaccharide CRM<sub>197</sub> conjugates consisting of:

- 2.2 µg of polysaccharide for serotypes 1, 3, 4, 5, 6A, 7F, 9V, 14, 18C, 19A, 19F and 23F
- 4.4 µg of polysaccharide for serotype 6B

1 dose (0.5 ml) contains approximately 32 µg CRM<sub>197</sub> carrier protein, adsorbed on aluminium phosphate (0.125 mg aluminium).

The other ingredients are sodium chloride, succinic acid, polysorbate 80 and water for injections.

### What Prevenar 13 looks like and contents of the pack

The vaccine is a white suspension for injection, provided in a single-dose vial (0.5 ml). Pack sizes of 1, 5, 10, 25 and 50 vials. Not all pack sizes may be marketed.

### Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:  
Pfizer Limited  
Ramsgate Road  
Sandwich  
Kent CT13 9NJ  
United Kingdom

Manufacturer responsible for batch release:  
Wyeth Pharmaceuticals  
New Lane  
Havant  
Hampshire, PO9 2NG  
United Kingdom

Pfizer Manufacturing Belgium N.V.  
Rijksweg 12  
B-2870 Puurs  
Belgium

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**  
**Luxembourg/Luxemburg**  
Pfizer S.A./N.V.  
Tél/Tel: + 32 (0)2 554 62 11

**Lietuva**  
Pfizer Luxembourg SARL filialas Lietuvoje  
Tel. + 370 52 51 4000

**България**  
Пфайзер Люксембург САРЛ, Клон  
България  
Тел: +359 2 970 4333

**Magyarország**  
Pfizer Kft  
Tel: +36 1 488 3700

**Česká Republika**  
Pfizer PFE, spol. s r.o.  
Tel: +420 283 004 111

**Malta**  
Vivian Corporation Ltd.  
Tel: + 35621 344610

**Danmark**  
Pfizer ApS  
Tlf: + 45 44 201 100

**Nederland**  
Pfizer BV  
Tel: +31 (0)10 406 43 01

**Deutschland**  
Pfizer Pharma GmbH  
Tel: + 49 (0)30 550055-51000

**Norge**  
Pfizer Norge AS  
Tlf: +47 67 52 61 00

**Eesti**  
Pfizer Luxembourg SARL Eesti filiaal  
Tel: +372 666 7500

**Österreich**  
Pfizer Corporation Austria Ges.m.b.H  
Tel: + 43 (0)1 521 15-0

**Ελλάδα**  
Pfizer Ελλάς Α.Ε.  
Τηλ.: +30 210 6785 800

**Polska**  
Pfizer Polska Sp. z o.o.  
Tel.: +48 22 335 61 00

**España**  
Pfizer, S.L.  
Télf:+34914909900

**Portugal**  
Pfizer Biofarmacêutica, Sociedade Unipessoal Lda  
Tel: +351 21 423 5500

**France**  
Pfizer  
Tél +33 1 58 07 34 40

**România**  
Pfizer Romania S.R.L  
Tel: +40 (0) 21 207 28 00

**Hrvatska**  
Pfizer Croatia d.o.o.  
Tel: + 385 1 3908 777

**Slovenija**  
Pfizer Luxembourg SARL  
Pfizer, podružnica za svetovanje s področja  
farmacevtske dejavnosti, Ljubljana  
Tel.: + 386 (0) 1 52 11 400

**Ireland**  
Pfizer Healthcare Ireland  
Tel: 1800 633 363 (toll free)  
+44 (0)1304 616161

**Slovenská republika**  
Pfizer Luxembourg SARL,  
organizačná zložka  
Tel: + 421 2 3355 5500

**Ísland**  
Icepharma hf  
Sími: + 354 540 8000

**Suomi/Finland**  
Pfizer Oy  
Puh/Tel: +358 (0)9 430 040

**Italia**

Pfizer S.r.l.  
Tel: +39 06 33 18 21

**Sverige**

Pfizer AB  
Tel: +46 (0)8 550 520 00

**Κύπρος**

Pfizer Ελλάς A.E. (Cyprus Branch)  
Τηλ: +357 22 817690

**United Kingdom**

Pfizer Limited  
Tel: +44 (0) 1304 616161

**Latvija**

Pfizer Luxembourg SARL filiāle Latvijā  
Tel.: + 371 670 35 775

**This leaflet was last revised in**

**Other sources of information**

Detailed information on this product is available on the European Medicines Agency website:  
<http://www.ema.europa.eu>.

---

**The following information is intended for healthcare professionals only:**

During storage, a white deposit and clear supernatant may be observed. This does not constitute a sign of deterioration.

Inspect visually for any foreign particulate matter and/or abnormal physical appearance; do not use if either are found.

Shake well prior to use to obtain a homogeneous white suspension.

Administer the entire dose.

Prevenar 13 is for intramuscular use only. Do not administer intravascularly.

Prevenar 13 must not be mixed with any other vaccines in the same syringe.

Prevenar 13 can be given at the same time as other childhood vaccines; in this case, different vaccination sites should be used.

Prevenar 13 may be given to adults aged 50 years and older at the same time as the trivalent or quadrivalent inactivated influenza vaccine.

Any unused product or waste material should be disposed of in accordance with local requirements.

## Package Leaflet: Information for the user

### **Prevenar 13 suspension for injection in multidose container** Pneumococcal polysaccharide conjugate vaccine (13 valent, adsorbed)

**Read all of this leaflet carefully before you or your child receives this vaccine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This vaccine has been prescribed for you or your child only. Do not pass it on to others.
- If you or your child get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Prevenar 13 is and what it is used for
2. What you need to know before you or your child receives Prevenar 13
3. How Prevenar 13 is given
4. Possible side effects
5. How to store Prevenar 13
6. Contents of the pack and other information

#### **1. What Prevenar 13 is and what it is used for**

Prevenar 13 is a pneumococcal vaccine given to:

- **children from 6 weeks to 17 years** to help protect against diseases such as: meningitis (inflammation around the brain), sepsis or bacteraemia (bacteria in the blood stream), pneumonia (lung infection) and ear infections
- **adults aged 18 years and older** to help prevent disease such as: pneumonia (lung infection), sepsis or bacteraemia (bacteria in the blood stream) and meningitis (inflammation around the brain),

caused by 13 types of the bacteria *Streptococcus pneumoniae*.

Prevenar 13, provides protection against 13 types of *Streptococcus pneumoniae* bacteria, and replaces Prevenar, which provided protection against 7 types.

The vaccine works by helping the body to make its own antibodies, which protect you or your child against these diseases.

#### **2. What you need to know before you or your child receives Prevenar 13**

##### **Prevenar 13 should not be given:**

- if you or your child is allergic (hypersensitive) to the active substances or to any of the other ingredients in this medicine (listed in section 6) or to any other vaccine that contains diphtheria toxoid.
- if you or your child has a severe infection with a high temperature (over 38°C). If this applies to you or your child, then the vaccination will be postponed until you or your child is feeling better. A minor infection, such as a cold, should not be a problem. However, talk to your doctor, pharmacist, or nurse first.

## Warnings and precautions

Talk to your doctor, pharmacist, or nurse before the vaccination if you or your child:

- has any present or past medical problems after any dose of Prevenar or Prevenar 13 such as an allergic reaction or problems with breathing.
- has any bleeding problems or bruises easily.
- has a weakened immune system (such as due to HIV infection), you/she/he may not get the full benefit from Prevenar 13.
- has experienced seizures as medicines to lower fever may need to be taken before Prevenar 13 is given. If your child should become unresponsive or experience seizures (fits) after the vaccination, please contact your doctor immediately. See also section 4.

Talk to your doctor, pharmacist, or nurse before the vaccination if your child was born very prematurely (at or before 28 weeks of gestation), as longer gaps than normal between breaths may occur for 2-3 days after vaccination. See also section 4.

As with any vaccine, Prevenar 13 will not protect all persons who are vaccinated.

Prevenar 13 will only protect against ear infections in children caused by the types of *Streptococcus pneumoniae* for which the vaccine has been developed. It will not protect against other infectious agents that can cause ear infections.

### Other medicines/vaccines and Prevenar 13

Your doctor may ask you to give your child paracetamol or other medicines that lower fever before Prevenar 13 is given. This will help to lower some of the side effects of Prevenar 13.

Tell your doctor, pharmacist or nurse if you or your child is taking, has recently taken or might take any other medicines, or has recently received any other vaccine.

### Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before receiving this medicine.

### Driving and using machines

Prevenar 13 has no or negligible influence on the ability to drive and use machines. However, some of the effects mentioned under section 4 "Possible side effects" may temporarily affect the ability to drive or use machines.

### Prevenar 13 contains sodium

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially 'sodium-free'.

## 3. How Prevenar 13 is given

The doctor or nurse will inject the recommended dose (0.5 ml) of the vaccine into your arm or your child's arm or leg muscle.

### Infants aged 6 weeks to 6 months of age

Typically, your child should receive an initial course of three injections of the vaccine followed by a booster dose.

- The first injection may be given from the age of six weeks.
- Each injection will be given at least one month apart.
- A fourth injection (booster) will be given between 11 and 15 months of age.
- You will be told when your child should come back for the next injection.

According to official recommendations in your country, an alternative schedule may be used by your healthcare provider. Please speak to your doctor, pharmacist, or nurse for more information.

#### Premature infants

Your child will receive an initial course of three injections. The first injection may be given as early as six weeks of age with at least one month between doses. Between 11 and 15 months of age, your child will receive a fourth injection (booster).

#### Unvaccinated infants, children, and adolescents over 7 months of age

Infants aged **7 to 11 months** should receive two injections. Each injection will be given at least one month apart. A third injection will be given in the second year of life.

Children aged **12 to 23 months** should receive two injections. Each injection will be given at least two months apart.

Children aged **2 to 17 years** should receive one injection.

#### Infants, children, and adolescents previously vaccinated with Prevenar

Infants and children who have previously received Prevenar may receive Prevenar 13 to complete the course of injections.

For children **1 to 5 years** of age previously vaccinated with Prevenar, your doctor or nurse will recommend how many injections of Prevenar 13 are required.

Children and adolescents **6 to 17 years** of age should receive one injection.

It is important to follow the instructions from the doctor, pharmacist, or nurse so that your child completes the course of injections.

If you forget to go back at the scheduled time, ask the doctor, pharmacist, or nurse for advice.

#### Adults

Adults should receive one injection.

Tell your doctor, pharmacist or nurse if you have been given a pneumococcal vaccine before.

If you have any further questions on the use of Prevenar 13, ask your doctor, pharmacist, or nurse.

#### Special Populations

Individuals considered to be at a higher risk of pneumococcal infection (such as those with sickle cell disease or HIV infection), including those previously vaccinated with the 23-valent pneumococcal polysaccharide vaccine, may receive at least one dose of Prevenar 13.

Individuals with a blood-forming stem cell transplant may receive three injections, with the first given at 3 to 6 months after the transplant and with an interval of at least 1 month between doses. A fourth injection (booster) is recommended 6 months after the third injection.

#### **4. Possible side effects**

Like all vaccines, Prevenar 13 can cause side effects, although not everybody gets them.

**The following side effects include those reported for Prevenar 13 in infants and children (6 weeks to 5 years of age):**

**The most common side effects** (these may occur with more than 1 in 10 doses of the vaccine) are:

- Decreased appetite
- Fever; irritability; pain, tenderness, redness, swelling or hardness at the vaccination-site; drowsiness; restless sleep
- Redness, hardness, swelling at the vaccination-site of 2.5 cm -7.0 cm (after the booster dose and in older children [aged 2 to 5 years])

**Common side effects** (these may occur with up to 1 in 10 doses of the vaccine) are:

- Vomiting; diarrhoea
- Fever of more than 39°C; tenderness at the vaccination-site interfering with movement, redness, hardness, swelling at the vaccination-site of 2.5 cm -7.0 cm (after the initial course of injections)
- Rash

**Uncommon side effects** (these may occur with up to 1 in 100 doses of the vaccine) are:

- Seizures (or fits), including those caused by a high temperature
- Hives (urticaria or urticaria-like rash)
- Redness, swelling, or hardness at the vaccination-site of more than 7 cm; crying

**Rare side effects** (these may occur with up to 1 in 1,000 doses of the vaccine) are:

- Collapse or shock-like state (hypotonic-hyporesponsive episode)
- Allergic (hypersensitivity) reaction, including swelling of the face and/or lips, difficulty in breathing

**The following side effects include those reported for Prevenar 13 in children and adolescents (6 to 17 years of age):**

**The most common side effects** (these may occur with more than 1 in 10 doses of the vaccine) are:

- Decreased appetite
- Irritability; pain, tenderness, redness, swelling or hardness at the vaccination-site; drowsiness; restless sleep; tenderness at the vaccination-site interfering with movement

**Common side effects** (these may occur with up to 1 in 10 doses of the vaccine) are:

- Headaches
- Vomiting; diarrhoea
- Rash; hives (urticaria or urticaria-like rash)
- Fever

Children and adolescents with either HIV infection, sickle cell disease or a blood-forming stem cell transplant had similar side effects however the frequencies of headaches, vomiting, diarrhoea, fever, fatigue, joint and muscle pain were very common.

**The following additional side effects have been seen with Prevenar 13 in infants and children up to 5 years of age in postmarketing experience**

- Severe allergic reaction including shock (cardiovascular collapse); angioedema (swelling of lips, face or throat)
- Hives (urticaria), redness and irritation (dermatitis) and itching (pruritus) at the vaccination-site; flushing
- Enlarged lymph nodes or glands (lymphadenopathy) near the vaccination-site, such as under the arm or in the groin
- A rash causing itchy red blotches (erythema multiforme)

In babies born very prematurely (at or before 28 weeks of gestation), longer gaps than normal between breaths may occur for 2-3 days after vaccination.

**The following side effects include those reported for Prevenar 13 in adults:**

**The most common side effects** (these may occur with more than 1 in 10 doses of the vaccine) are:

- Decreased appetite; headaches; diarrhoea; vomiting (for those 18 to 49 years of age)
- Chills; tiredness; rash; pain, redness, swelling hardness or tenderness at the vaccination-site, interfering with arm movement (severe pain or tenderness at vaccination-site for those 18-39 years of age and severe limitation of arm movements for those 18 to 39 years of age)
- Worsening or new pain in your joints, worsening or new pain in your muscles
- Fever (for those 18 to 29 years of age)

**Common side effects** (these may occur with up to 1 in 10 doses of the vaccine) are:

- Vomiting (for those 50 years and older); fever (for those 30 years and older)

**Uncommon side effects** (these may occur with up to 1 in 100 doses of the vaccine) are:

- Nausea
- Allergic (hypersensitivity) reaction, including swelling of the face and/or lips, difficulty in breathing
- Enlarged lymph nodes or glands (lymphadenopathy) near the vaccination-site, such as under the arm

Adults with HIV infection had similar side effects however the frequencies were very common for fever, vomiting and common for nausea.

Adults with a blood-forming stem cell transplant had similar side effects however the frequencies were very common for fever and vomiting.

**Reporting of side effects**

If you or your child get any side effects, talk to your doctor, pharmacist, or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via **the national reporting system listed in Appendix V**. By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Prevenar 13

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and label after EXP. The expiry date refers to the last day of that month.

Store in a refrigerator (2°C – 8°C).

Do not freeze.

After first use, the product may be stored in a refrigerator for a maximum of 28 days.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Prevenar 13 contains

The active substances are polysaccharide CRM<sub>197</sub> conjugates consisting of:

- 2.2 µg of polysaccharide for serotypes 1, 3, 4, 5, 6A, 7F, 9V, 14, 18C, 19A, 19F and 23F
- 4.4 µg of polysaccharide for serotype 6B

1 dose (0.5 ml) contains approximately 32 µg CRM<sub>197</sub> carrier protein, adsorbed on aluminium phosphate (0.125 mg aluminium).

The other ingredients are sodium chloride, succinic acid, polysorbate 80, 2-phenoxyethanol and water for injections.

### What Prevenar 13 looks like and contents of the pack

The vaccine is a white suspension for injection, provided in a multidose container (4 x 0.5 ml doses). Pack sizes of 1, 5, 10, 25 and 50 vials. Not all pack sizes may be marketed.

### Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:  
Pfizer Limited  
Ramsgate Road  
Sandwich  
Kent CT13 9NJ  
United Kingdom

Manufacturer responsible for batch release:  
Pfizer Manufacturing Belgium N.V.  
Rijksweg 12  
B-2870 Puurs  
Belgium

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**  
**Luxembourg/Luxemburg**  
Pfizer S.A./N.V.  
Tél/Tel: + 32 (0)2 554 62 11

**Lietuva**  
Pfizer Luxembourg SARL filialas Lietuvoje  
Tel. + 370 52 51 4000

**България**  
Пфайзер Люксембург САРЛ, Клон  
България  
Тел: +359 2 970 4333

**Magyarország**  
Pfizer Kft  
Tel: +36 1 488 3700

**Česká Republika**

Pfizer PFE, spol. s r.o.  
Tel: +420 283 004 111

**Danmark**

Pfizer ApS  
Tlf: + 45 44 201 100

**Deutschland**

Pfizer Pharma GmbH  
Tel: + 49 (0)30 550055-51000

**Eesti**

Pfizer Luxembourg SARL Eesti filiaal  
Tel: +372 666 7500

**Ελλάδα**

Pfizer Ελλάς A.E.  
Τηλ.: +30 210 6785 800

**España**

Pfizer, S.L.  
Télf:+34914909900

**France**

Pfizer  
Tél +33 1 58 07 34 40

**Hrvatska**

Pfizer Croatia d.o.o.  
Tel: + 385 1 3908 777

**Ireland**

Pfizer Healthcare Ireland  
Tel: 1800 633 363 (toll free)  
+44 (0)1304 616161

**Ísland**

Icepharma hf  
Simi: + 354 540 8000

**Italia**

Pfizer S.r.l.  
Tel: +39 06 33 18 21

**Κύπρος**

Pfizer Ελλάς A.E. (Cyprus Branch)  
Τηλ: +357 22 817690

**Latvija**

Pfizer Luxembourg SARL filiāle Latvijā  
Tel.: + 371 670 35 775

**Malta**

Vivian Corporation Ltd.  
Tel: + 35621 344610

**Nederland**

Pfizer BV  
Tel: +31 (0)10 406 43 01

**Norge**

Pfizer Norge AS  
Tlf: +47 67 52 61 00

**Österreich**

Pfizer Corporation Austria Ges.m.b.H  
Tel: + 43 (0)1 521 15-0

**Polska**

Pfizer Polska Sp. z o.o.  
Tel.: +48 22 335 61 00

**Portugal**

Pfizer Biofarmacêutica, Sociedade Unipessoal Lda  
Tel: +351 21 423 5500

**România**

Pfizer Romania S.R.L  
Tel: +40 (0) 21 207 28 00

**Slovenija**

Pfizer Luxembourg SARL  
Pfizer, podružnica za svetovanje s področja  
farmacevtske dejavnosti, Ljubljana  
Tel.: + 386 (0) 1 52 11 400

**Slovenská republika**

Pfizer Luxembourg SARL,  
organizačná zložka  
Tel: + 421 2 3355 5500

**Suomi/Finland**

Pfizer Oy  
Puh/Tel: +358 (0)9 430 040

**Sverige**

Pfizer AB  
Tel: +46 (0)8 550 520 00

**United Kingdom**

Pfizer Limited  
Tel: +44 (0) 1304 616161

**This leaflet was last revised in**

**Other sources of information**

Detailed information on this product is available on the European Medicines Agency website:  
<http://www.ema.europa.eu>.

---

**The following information is intended for healthcare professionals only:**

During storage, a white deposit and clear supernatant may be observed. This does not constitute a sign of deterioration.

Inspect visually for any foreign particulate matter and/or abnormal physical appearance; do not use if either are found.

Shake well prior to use to obtain a homogeneous white suspension.

Administer the entire dose.

Prevenar 13 is for intramuscular use only. Do not administer intravascularly.

Prevenar 13 must not be mixed with any other vaccines in the same syringe.

Prevenar 13 can be given at the same time as other childhood vaccines; in this case, different vaccination sites should be used.

Prevenar 13 may be given to adults aged 50 years and older at the same time as the trivalent or quadrivalent inactivated influenza vaccine.

After first use, the product may be stored in a refrigerator for a maximum of 28 days.

Any unused product or waste material should be disposed of in accordance with local requirements.