

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## **1. NAME OF THE MEDICINAL PRODUCT**

Seebri Breezhaler 44 micrograms inhalation powder, hard capsules

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each capsule contains 63 micrograms of glycopyrronium bromide equivalent to 50 micrograms of glycopyrronium.

Each delivered dose (the dose that leaves the mouthpiece of the inhaler) contains 55 micrograms of glycopyrronium bromide equivalent to 44 micrograms of glycopyrronium.

Excipient(s) with known effect:

Each capsule contains 23.6 mg lactose (as monohydrate).

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Inhalation powder, hard capsule

Transparent orange capsules containing a white powder, with the product code “GPL50” printed in black above and the company logo (ℓ) printed in black below a black bar.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Seebri Breezhaler is indicated as a maintenance bronchodilator treatment to relieve symptoms in adult patients with chronic obstructive pulmonary disease (COPD).

### **4.2 Posology and method of administration**

#### Posology

The recommended dose is the inhalation of the content of one capsule once daily using the Seebri Breezhaler inhaler.

Seebri Breezhaler is recommended to be administered, at the same time of the day each day. If a dose is missed, the next dose should be taken as soon as possible. Patients should be instructed not to take more than one dose in a day.

#### Special populations

##### *Elderly population*

Seebri Breezhaler can be used at the recommended dose in elderly patients (75 years of age and older) (see section 4.8).

#### *Renal impairment*

Seebri Breezhaler can be used at the recommended dose in patients with mild to moderate renal impairment. In patients with severe renal impairment or end-stage renal disease requiring dialysis Seebri Breezhaler should be used only if the expected benefit outweighs the potential risk (see sections 4.4 and 5.2).

#### *Hepatic impairment*

No studies have been conducted in patients with hepatic impairment. Glycopyrronium is predominantly cleared by renal excretion and therefore no major increase in exposure is expected in patients with hepatic impairment.

#### *Paediatric population*

There is no relevant use of Seebri Breezhaler in the paediatric population (under 18 years) in the indication COPD.

#### Method of administration

For inhalation use only.

The capsules must be administered only using the Seebri Breezhaler inhaler (see section 6.6).

The capsules must not be swallowed.

Patients should be instructed on how to administer the product correctly. Patients who do not experience improvement in breathing should be asked if they are swallowing the medicine rather than inhaling it.

For instructions on use of the medicinal product before administration, see section 6.6.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

### **4.4 Special warnings and precautions for use**

#### Not for acute use

Seebri Breezhaler is a once-daily, long-term maintenance treatment and is not indicated for the initial treatment of acute episodes of bronchospasm, i.e. as a rescue therapy.

#### Hypersensitivity

Immediate hypersensitivity reactions have been reported after administration of Seebri Breezhaler. If signs suggesting allergic reactions occur, in particular, angioedema (including difficulties in breathing or swallowing, swelling of the tongue, lips, and face), urticaria or skin rash, Seebri Breezhaler should be discontinued immediately and alternative therapy instituted.

#### Paradoxical bronchospasm

In clinical studies with Seebri Breezhaler, paradoxical bronchospasm was not observed. However, paradoxical bronchospasm has been observed with other inhalation therapy and can be life-threatening. If this occurs, Seebri Breezhaler should be discontinued immediately and alternative therapy instituted.

#### Anticholinergic effect

Seebri Breezhaler should be used with caution in patients with narrow-angle glaucoma or urinary retention.

Patients should be informed about the signs and symptoms of acute narrow-angle glaucoma and should be informed to stop using Seebri Breezhaler and to contact their doctor immediately should any of these signs or symptoms develop.

#### Patients with severe renal impairment

A moderate mean increase in total system exposure ( $AUC_{last}$ ) of up to 1.4-fold was seen in subjects with mild and moderate renal impairment and up to 2.2-fold in subjects with severe renal impairment and end-stage renal disease. In patients with severe renal impairment (estimated glomerular filtration rate below 30 ml/min/1.73 m<sup>2</sup>), including those with end-stage renal disease requiring dialysis, Seebri Breezhaler should be used only if the expected benefit outweighs the potential risk (see section 5.2). These patients should be monitored closely for potential adverse reactions.

#### Patients with a history of cardiovascular disease

Patients with unstable ischaemic heart disease, left ventricular failure, history of myocardial infarction, arrhythmia (excluding chronic stable atrial fibrillation), a history of long QT syndrome or whose QTc (Fridericia method) was prolonged (>450 ms for males or >470 ms for females) were excluded from the clinical trials, and therefore the experience in these patient groups is limited. Seebri Breezhaler should be used with caution in these patient groups.

#### Excipients

Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

### **4.5 Interaction with other medicinal products and other forms of interaction**

The co-administration of Seebri Breezhaler with other anticholinergic-containing medicinal products has not been studied and is therefore not recommended.

Although no formal drug interaction studies have been performed, Seebri Breezhaler has been used concomitantly with other medicinal products commonly used in the treatment of COPD without clinical evidence of drug interactions. These include sympathomimetic bronchodilators, methylxanthines, and oral and inhaled steroids.

In a clinical study in healthy volunteers, cimetidine, an inhibitor of organic cation transport which is thought to contribute to the renal excretion of glycopyrronium, increased total exposure (AUC) to glycopyrronium by 22% and decreased renal clearance by 23%. Based on the magnitude of these changes, no clinically relevant drug interaction is expected when glycopyrronium is co-administered with cimetidine or other inhibitors of organic cation transport.

Concomitant administration of Seebri Breezhaler and orally inhaled indacaterol, a beta<sub>2</sub>-adrenergic agonist, under steady-state conditions of both active substances did not affect the pharmacokinetics of either medicinal product.

### **4.6 Fertility, pregnancy and lactation**

#### Pregnancy

There are no data from the use of Seebri Breezhaler in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3). Glycopyrronium should only be used during pregnancy if the expected benefit to the patient justifies the potential risk to the foetus.

#### Breast-feeding

It is unknown whether glycopyrronium bromide is excreted in human milk. However, glycopyrronium bromide (including its metabolites) was excreted in the milk of lactating rats (see section 5.3). The use of glycopyrronium by breast-feeding women should only be considered if the expected benefit to the woman is greater than any possible risk to the infant (see section 5.3).

#### Fertility

Reproduction studies and other data in animals do not indicate a concern regarding fertility in either males or females (see section 5.3).

#### 4.7 Effects on ability to drive and use machines

Glycopyrronium has no or negligible influence on the ability to drive and use machines.

#### 4.8 Undesirable effects

##### Summary of the safety profile

The most common anticholinergic adverse reaction was dry mouth (2.4%). The majority of the reports of dry mouth were suspected to be related to the medicinal product and were mild, with none being severe.

The safety profile is further characterised by other symptoms related to the anticholinergic effects, including signs of urinary retention, which were uncommon. Gastrointestinal effects including gastroenteritis and dyspepsia were also observed. Adverse reactions related to local tolerability included throat irritation, nasopharyngitis, rhinitis and sinusitis.

##### Tabulated summary of adverse reactions

Adverse reactions reported during the first six months of two pooled pivotal Phase III trials of 6 and 12 months duration are listed by MedDRA system organ class (Table 1). Within each system organ class, the adverse reactions are ranked by frequency, with the most frequent reactions first. Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness. In addition, the corresponding frequency category for each adverse reaction is based on the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data).

**Table 1 Adverse reactions reported in the pooled 6-month database**

<b>Adverse reactions</b>	<b>Frequency category</b>
<b>Infections and infestations</b>	
Nasopharyngitis <sup>1)</sup>	Common
Rhinitis	Uncommon
Cystitis	Uncommon
<b>Immune system disorders</b>	
Hypersensitivity	Uncommon
Angioedema <sup>2)</sup>	Uncommon
<b>Metabolism and nutrition disorders</b>	
Hyperglycaemia	Uncommon
<b>Psychiatric disorders</b>	
Insomnia	Common
<b>Nervous system disorders</b>	
Headache <sup>3)</sup>	Common
Hypoaesthesia	Uncommon
<b>Cardiac disorders</b>	
Atrial fibrillation	Uncommon
Palpitations	Uncommon

<b>Respiratory, thoracic and mediastinal disorders</b>	
Sinus congestion	Uncommon
Productive cough	Uncommon
Throat irritation	Uncommon
Epistaxis	Uncommon
Paradoxical bronchospasm <sup>2)</sup>	Not known
Dysphonia <sup>2)</sup>	Uncommon
<b>Gastrointestinal disorders</b>	
Dry mouth	Common
Gastroenteritis	Common
Dyspepsia	Uncommon
Dental caries	Uncommon
<b>Skin and subcutaneous tissue disorders</b>	
Rash	Uncommon
Pruritus <sup>2)</sup>	Uncommon
<b>Musculoskeletal and connective tissue disorders</b>	
Pain in extremity	Uncommon
Musculoskeletal chest pain	Uncommon
<b>Renal and urinary disorders</b>	
Urinary tract infection <sup>3)</sup>	Common
Dysuria	Uncommon
Urinary retention	Uncommon
<b>General disorders and administration site conditions</b>	
Fatigue	Uncommon
Asthenia	Uncommon

1) More frequent for glycopyrronium than placebo in the 12 months database only.

2) Reports have been received from post-approval marketing experience in association with the use of Seebri Breezhaler. These were reported voluntarily from a population of uncertain size, and it is therefore not always possible to reliably estimate the frequency or establish a causal relationship to drug exposure. Therefore the frequency was calculated from clinical trial experience.

3) Seen more frequently for glycopyrronium than placebo in elderly >75 years only.

#### Description of selected adverse reactions

In the pooled 6-month database the frequency of dry mouth was 2.2% versus 1.1%, of insomnia 1.0% versus 0.8%, and of gastroenteritis 1.4% versus 0.9%, for Seebri Breezhaler and placebo respectively.

Dry mouth was reported mainly during the first 4 weeks of treatment with a median duration of four weeks in the majority of patients. However in 40% of cases symptoms continued for the entire 6-month period. No new cases of dry mouth were reported in months 7-12.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

## 4.9 Overdose

High doses of glycopyrronium may lead to anticholinergic signs and symptoms for which symptomatic treatment may be indicated.

Acute intoxication by inadvertent oral ingestion of Seebri Breezhaler capsules is unlikely due to the low oral bioavailability (about 5%).

Peak plasma levels and total systemic exposure following intravenous administration of 150 micrograms glycopyrronium bromide (equivalent to 120 micrograms glycopyrronium) in healthy volunteers were respectively about 50-fold and 6-fold higher than the peak and total exposure at steady-state achieved with the recommended dose (44 micrograms once daily) of Seebri Breezhaler and were well tolerated.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Drugs for obstructive airway diseases, anticholinergics, ATC code: R03BB06

#### Mechanism of action

Seebri Breezhaler is an inhaled long-acting muscarinic receptor antagonist (anticholinergic) for once-daily maintenance bronchodilator treatment of COPD. Parasympathetic nerves are the major bronchoconstrictive neural pathway in airways, and cholinergic tone is the key reversible component of airflow obstruction in COPD. Glycopyrronium works by blocking the bronchoconstrictor action of acetylcholine on airway smooth muscle cells, thereby dilating the airways.

Glycopyrronium bromide is a high affinity muscarinic receptor antagonist. A greater than 4-fold selectivity for the human M3 receptors over the human M2 receptor has been demonstrated using radioligand binding studies. It has a rapid onset of action as evidenced by observed receptor association/dissociation kinetic parameters and the onset of action after inhalation in clinical studies.

The long duration of action can be partly attributed to sustained concentrations of active substance in the lung as reflected by the prolonged terminal elimination half-life of glycopyrronium after inhalation via the Seebri Breezhaler inhaler in contrast to the half life after intravenous administration (see section 5.2).

#### Pharmacodynamic effects

The clinical Phase III development programme included two phase III studies: a 6-month placebo-controlled study and a 12-month placebo and active-controlled (open label tiotropium 18 micrograms once daily) study, both in patients with clinical diagnosis of moderate to severe COPD.

#### *Effects on lung function*

Seebri Breezhaler 44 micrograms once daily provided consistently statistically significant improvement in lung function (forced expiratory volume in one second, FEV<sub>1</sub>, forced vital capacity, FVC, and inspiratory capacity, IC) in a number of clinical studies. In phase III studies, bronchodilator effects were seen within 5 minutes after the first dose and were maintained over the 24-hour dosing interval from the first dose. There was no attenuation of the bronchodilator effect over time in the 6- and 12-month studies. The magnitude of the effect was dependent on the degree of reversibility of airflow limitation at baseline (tested by administration of a short-acting muscarinic antagonist bronchodilator): Patients with the lowest degree of reversibility at baseline (<5%) generally exhibited a lower bronchodilator response than patients with a higher degree of reversibility at baseline (≥5%). At 12 weeks (primary endpoint), Seebri Breezhaler increased trough FEV<sub>1</sub> by 72 ml in patients with the lowest degree of reversibility (<5%) and by 113 ml in those patients with a higher degree of reversibility at baseline (≥5%) compared to placebo (both p<0.05).

In the 6-month study, Seebri Breezhaler increased FEV<sub>1</sub> after the first dose with an improvement of 93 ml within 5 minutes and 144 ml within 15 minutes of dosing, compared to placebo (both p<0.001). In the 12-month study, the improvements were 87 ml at 5 minutes and 143 ml at 15 minutes (both p<0.001). In the 12-month study, Seebri Breezhaler produced statistically significant improvements in FEV<sub>1</sub> compared to tiotropium in the first 4 hours after dosing on day 1 and at week 26, and numerically greater values for FEV<sub>1</sub> in the first 4 hours after dosing than tiotropium at week 12 and week 52.

The values for FEV<sub>1</sub> at the end of the dosing interval (24 h post dose) were similar between the first dose and those seen after 1 year of dosing. At 12 weeks (primary endpoint), Seebri Breezhaler increased trough FEV<sub>1</sub> by 108 ml in the 6-month study and by 97 ml in the 12-month study compared to placebo (both p<0.001). In the 12-month study, the improvement versus placebo for tiotropium was 83 ml (p<0.001).

#### *Symptomatic outcomes*

Seebri Breezhaler administered at 44 micrograms once daily statistically significantly reduced breathlessness as evaluated by the Transitional Dyspnoea Index (TDI). In a pooled analysis of the 6- and 12-month pivotal studies a statistically significantly higher percentage of patients receiving Seebri Breezhaler responded with a 1 point or greater improvement in the TDI focal score at week 26 compared to placebo (58.4% and 46.4% respectively, p<0.001). These findings were similar to those seen in patients receiving tiotropium, 53.4% of whom responded with 1 point or greater improvement (p=0.009 compared to placebo).

Seebri Breezhaler once daily has also shown a statistically significant effect on health-related quality of life measured using the St. George's Respiratory Questionnaire (SGRQ). A pooled analysis of the 6- and 12-month pivotal studies found a statistically significantly higher percentage of patients receiving Seebri Breezhaler responded with a 4 point or greater improvement in SGRQ compared to placebo at week 26 (57.8% and 47.6% respectively, p<0.001). For patients receiving tiotropium, 61.0% responded with a 4 point or greater improvement in SGRQ (p=0.004 compared to placebo).

#### *COPD exacerbations reduction*

COPD exacerbation data was collected in the 6- and 12-month pivotal studies. In both studies, the percentage of patients experiencing a moderate or severe exacerbation (defined as requiring treatment with systemic corticosteroids and/or antibiotics or hospitalisation) was reduced. In the 6-month study, the percentage of patients experiencing a moderate or severe exacerbation was 17.5% for Seebri Breezhaler and 24.2% for placebo (Hazard ratio: 0.69, p=0.023), and in the 12-month study it was 32.8% for Seebri Breezhaler and 40.2% for placebo (Hazard ratio: 0.66, p=0.001). In a pooled analysis of the first 6 months of treatment in the 6- and 12-month studies, compared to placebo Seebri Breezhaler statistically significantly prolonged time to first moderate or severe exacerbation and reduced the rate of moderate or severe COPD exacerbations (0.53 exacerbations/year versus 0.77exacerbations /year, p<0.001). The pooled analysis also showed fewer patients treated with Seebri Breezhaler than with placebo experienced an exacerbation requiring hospitalisation (1.7% versus 4.2%, p=0.003).

#### *Other effects*

Seebri Breezhaler once daily statistically significantly reduced the use of rescue medication (salbutamol) by 0.46 puffs per day (p=0.005) over 26 weeks and by 0.37 puffs per day (p=0.039) over 52 weeks, compared to placebo for the 6- and 12-month studies, respectively.

In a 3-week study where exercise tolerance was tested via cycle ergometer at submaximal (80%) workload (submaximal exercise tolerance test), Seebri Breezhaler, dosed in the morning, reduced dynamic hyperinflation and improved the length of time exercise could be maintained from the first dose onwards. On the first day of treatment inspiratory capacity under exercise was improved by 230 ml and exercise endurance time was improved by 43 seconds (an increase of 10%) compared to placebo. After three weeks of treatment the improvement in inspiratory capacity with Seebri Breezhaler was similar to the first day (200 ml), exercise endurance time however had increased by



89 seconds (an increase of 21%) compared to placebo. Seebri Breezhaler was found to decrease dyspnoea and leg discomfort when exercising as measured using Borg scales. Seebri Breezhaler also reduced dyspnoea at rest measured using the Transitional Dyspnoea Index.

#### *Secondary pharmacodynamic effects*

No change in mean heart rate or QTc interval was observed with Seebri Breezhaler in doses up to 176 micrograms in COPD patients. In a thorough QT study in 73 healthy volunteers, a single inhaled dose of glycopyrronium 352 micrograms (8 times the therapeutic dose) did not prolong the QTc interval and slightly reduced heart rate (maximal effect -5.9 bpm; average effect over 24 hours -2.8 bpm) when compared to placebo. The effect on heart rate and QTc interval of 150 micrograms glycopyrronium bromide (equivalent to 120 micrograms glycopyrronium) administered intravenously was investigated in young healthy subjects. Peak exposures ( $C_{max}$ ) about 50-fold higher than after inhalation of glycopyrronium 44 micrograms at steady state were achieved and did not result in tachycardia or QTc prolongation. A slight reduction in heart rate (mean difference over 24 h -2 bpm when compared to placebo), which is a known effect of low exposures to anticholinergic compounds in young healthy subjects, was observed.

#### Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with Seebri Breezhaler in all subsets of the paediatric population in COPD (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

### Absorption

Following oral inhalation using the Seebri Breezhaler inhaler, glycopyrronium was rapidly absorbed and reached peak plasma levels at 5 minutes post dose.

The absolute bioavailability of glycopyrronium inhaled via Seebri Breezhaler was estimated to be about 45% of the delivered dose. About 90% of systemic exposure following inhalation is due to lung absorption and 10% is due to gastrointestinal absorption.

In patients with COPD, pharmacokinetic steady-state of glycopyrronium was reached within one week of the start of treatment. The steady-state mean peak and trough plasma concentrations of glycopyrronium for a 44 micrograms once-daily dosing regimen were 166 picograms/ml and 8 picograms/ml, respectively. Steady-state exposure to glycopyrronium (AUC over the 24-hour dosing interval) was about 1.4- to 1.7-fold higher than after the first dose.

### Distribution

After intravenous dosing, the steady-state volume of distribution of glycopyrronium was 83 litres and the volume of distribution in the terminal phase was 376 litres. The apparent volume of distribution in the terminal phase following inhalation was almost 20-fold larger, which reflects the much slower elimination after inhalation. The *in vitro* human plasma protein binding of glycopyrronium was 38% to 41% at concentrations of 1 to 10 nanograms/ml.

### Biotransformation

*In vitro* metabolism studies showed consistent metabolic pathways for glycopyrronium bromide between animals and humans. Hydroxylation resulting in a variety of mono- and bis-hydroxylated metabolites and direct hydrolysis resulting in the formation of a carboxylic acid derivative (M9) were seen. *In vivo*, M9 is formed from the swallowed dose fraction of inhaled glycopyrronium bromide. Glucuronide and/or sulfate conjugates of glycopyrronium were found in urine of humans after repeated inhalation, accounting for about 3% of the dose.

Multiple CYP isoenzymes contribute to the oxidative biotransformation of glycopyrronium. Inhibition or induction of the metabolism of glycopyrronium is unlikely to result in a relevant change of systemic exposure to the active substance.

*In vitro* inhibition studies demonstrated that glycopyrronium bromide has no relevant capacity to inhibit CYP1A2, CYP2A6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1 or CYP3A4/5, the efflux transporters MDR1, MRP2 or MXR, and the uptake transporters OCT1 or OCT2. *In vitro* enzyme induction studies did not indicate a clinically relevant induction by glycopyrronium bromide for cytochrome P450 isoenzymes, or for UGT1A1 and the transporters MDR1 and MRP2.

#### Elimination

After intravenous administration of [<sup>3</sup>H]-labelled glycopyrronium bromide to humans, the mean urinary excretion of radioactivity in 48 hours amounted to 85% of the dose. A further 5% of the dose was found in the bile.

Renal elimination of parent drug accounts for about 60 to 70% of total clearance of systemically available glycopyrronium whereas non-renal clearance processes account for about 30 to 40%. Biliary clearance contributes to the non-renal clearance, but the majority of non-renal clearance is thought to be due to metabolism.

Mean renal clearance of glycopyrronium following inhalation was in the range of 17.4 and 24.4 litres/h. Active tubular secretion contributes to the renal elimination of glycopyrronium. Up to 23% of the delivered dose was found in urine as parent drug.

Glycopyrronium plasma concentrations declined in a multi-phasic manner. The mean terminal elimination half-life was much longer after inhalation (33 to 57 hours) than after intravenous (6.2 hours) and oral (2.8 hours) administration. The elimination pattern suggests sustained lung absorption and/or transfer of glycopyrronium into the systemic circulation at and beyond 24 hours after inhalation.

#### Linearity/non-linearity

In COPD patients both systemic exposure and total urinary excretion of glycopyrronium at pharmacokinetic steady state increased about dose-proportionally over the dose range of 44 to 176 micrograms.

#### Special populations

A population pharmacokinetic analysis of data in COPD patients identified body weight and age as factors contributing to inter-patient variability in systemic exposure. Seebri Breezhaler 44 micrograms once daily can be safely used in all age and body weight groups.

Gender, smoking status and baseline FEV<sub>1</sub> had no apparent effect on systemic exposure.

There were no major differences in total systemic exposure (AUC) between Japanese and Caucasian subjects following inhalation of glycopyrronium bromide. Insufficient pharmacokinetic data is available for other ethnicities or races.

#### *Patients with hepatic impairment*

Clinical studies have not been conducted in patients with hepatic impairment. Glycopyrronium is cleared predominantly from the systemic circulation by renal excretion. Impairment of the hepatic metabolism of glycopyrronium is not thought to result in a clinically relevant increase of systemic exposure.

#### *Patients with renal impairment*

Renal impairment has an impact on the systemic exposure to glycopyrronium bromide. A moderate mean increase in total systemic exposure (AUC<sub>last</sub>) of up to 1.4-fold was seen in subjects with mild and moderate renal impairment and up to 2.2-fold in subjects with severe renal impairment and end-stage renal disease. In COPD patients with mild and moderate renal impairment (estimated glomerular filtration rate, eGFR ≥30 ml/min/1.73 m<sup>2</sup>) Seebri Breezhaler can be used at the recommended dose. In patients with severe renal impairment (eGFR <30 ml/min/1.73 m<sup>2</sup>), including those with end-stage renal disease requiring dialysis, Seebri Breezhaler should only be used in the expected benefit outweighs the potential risk (see section 4.4).

### **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction and development.

Effects attributable to the muscarinic receptor antagonist properties of glycopyrronium bromide included mild to moderate increases in heart rate in dogs, lens opacities in rats and, reversible changes associated with reduced glandular secretions in rats and dogs. Mild irritancy or adaptive changes in the respiratory tract were seen in rats. All these findings occurred at exposures sufficiently in excess of those anticipated in humans.

Glycopyrronium was not teratogenic in rats or rabbits following inhalation administration. Fertility and pre- and post-natal development were not affected in rats. Glycopyrronium bromide and its metabolites did not significantly cross the placental barrier of pregnant mice, rabbits and dogs. Glycopyrronium bromide (including its metabolites) was excreted into the milk of lactating rats and reached up to 10-fold higher concentrations in the milk than in the blood of the dam.

Genotoxicity studies did not reveal any mutagenic or clastogenic potential for glycopyrronium bromide. Carcinogenicity studies in transgenic mice using oral administration and in rats using inhalation administration revealed no evidence of carcinogenicity at systemic exposures (AUC) of approximately 53-fold higher in mice and 75-fold higher in rats than the maximum recommended dose of 44 micrograms once daily for humans.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Capsule content

Lactose monohydrate

Magnesium stearate

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

2 years

Each inhaler should be disposed of after 30 days of use.

### **6.4 Special precautions for storage**

Do not store above 25°C.

The capsules must always be stored in the blister to protect from moisture and only removed immediately before use.

## 6.5 Nature and contents of container

Seebri Breezhaler is a single-dose inhaler. Inhaler body and cap are made from acrylonitrile butadiene styrene, push buttons are made from methyl methacrylate acrylonitrile butadiene styrene. Needles and springs are made from stainless steel. Each blister contains either 6 or 10 hard capsules.

PA/Alu/PVC – Alu perforated unit-dose blister

Single pack containing 6x1, 10x1, 12x1 or 30x1 hard capsules, together with one inhaler.

Multipacks containing 90 (3 packs of 30x1) hard capsules and 3 inhalers.

Multipacks containing 96 (4 packs of 24x1) hard capsules and 4 inhalers.

Multipacks containing 150 (15 packs of 10x1) hard capsules and 15 inhalers.

Multipacks containing 150 (25 packs of 6x1) hard capsules and 25 inhalers.


Not all pack sizes may be marketed.

## 6.6 Special precautions for disposal and other handling


The inhaler provided with each new prescription should be used. Each inhaler should be disposed of after 30 days of use.

### Instructions for handling and use

#### How to use your inhaler








**Pull off the cap.**



**Open inhaler:**

Hold the base of the inhaler firmly and tilt the mouthpiece. This opens the inhaler.



**Prepare capsule:**

Separate one of the blisters from the blister card by tearing along the perforation.  
Take one blister and peel away the protective backing to expose the capsule.

Do not push capsule through foil.

**Remove a capsule:**

Capsules should always be stored in the blister and only removed immediately before use.  
With dry hands, remove capsule from the blister.  
Do not swallow the capsule.

**Insert capsule:**

Place the capsule into the capsule chamber.





**Never place a capsule directly into the mouthpiece.**

**Close the inhaler:**

Close the inhaler until you hear a “click”.

**Pierce the capsule:**

- Hold the inhaler upright with the mouthpiece pointing up.
- Pierce the capsule by firmly pressing together both side buttons at the same time.  
**Do this only once.**
- You should hear a “click” as the capsule is being pierced.



**Release the side buttons fully.**

**Breathe out:**

Before placing the mouthpiece in your mouth, breathe out fully.

**Do not blow into the mouthpiece.**

**Inhale the medicine:**

To breathe the medicine deeply into your airways:

- Hold the inhaler as shown in the picture. The side buttons should be facing left and right. Do not press the side buttons.
- Place the mouthpiece in your mouth and close your lips firmly around it.
- Breathe in rapidly but steadily, as deeply as you can. **Do not press the side buttons.**

**Note:**


As you breathe in through the inhaler, the capsule spins around in the chamber and you should hear a whirring noise. You will experience a sweet flavour as the medicine goes into your lungs.

**If you do not hear a whirring noise:**

The capsule may be stuck in the capsule chamber.

If this happens:

- Open the inhaler and carefully loosen the capsule by tapping the base of the inhaler. **Do not press the side buttons.**
- Inhale the medicine again by repeating steps 9 and 10.



**Hold breath:**

**After you have inhaled the medicine:**

- **Hold your breath** for at least 5-10 seconds or as long as you comfortably can while taking the inhaler out of your mouth.
- Then breathe out.
- Open the inhaler to see if any powder is left in the capsule.


**If there is powder left in the capsule:**

- Close the inhaler.
- Repeat steps 9 to 12.

Most people are able to empty the capsule with one or two inhalations.

**Additional information**

Some people occasionally cough briefly soon after inhaling a medicine. If you do, don't worry. As long as the capsule is empty, you have received enough of your medicine.



**After you have finished taking your daily dose of Seebri Breezhaler:**

- Open the mouthpiece again, and remove the empty capsule by tipping it out of the capsule chamber. Put the empty capsule in your household waste.
- Close the inhaler and replace the cap.

**Do not store the capsules in the Seebri Breezhaler inhaler.**

**7. MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Frimley Business Park  
Camberley GU16 7SR  
United Kingdom

**8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/12/788/001-008

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

28.09.2012

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>



## **ANNEX II**

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

**A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer responsible for batch release

Novartis Pharma GmbH  
Roonstraße 25  
D-90429 Nuremberg  
Germany

**B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to medical prescription.

**C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

• **Periodic Safety Update Reports**

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

**D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

• **Risk Management Plan (RMP)**

The MAH shall perform the pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency.
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

• **Obligation to conduct post-authorisation measures**

The MAH shall complete, within the stated timeframe, the below measures:

<b>Description</b>	<b>Due date</b>
Post-authorisation safety study on cardio- and cerebrovascular outcomes (Multinational database cohort study to assess adverse cardiovascular outcomes in association with inhaled glycopyrronium in Europe).	Final report 5 years after launch.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON OF UNIT PACK**

**1. NAME OF THE MEDICINAL PRODUCT**

Seebri Breezhaler 44 micrograms inhalation powder, hard capsules  
glycopyrronium (glycopyrronium bromide)

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each capsule contains 50 micrograms glycopyrronium. The amount of glycopyrronium inhaled is 44 micrograms.

**3. LIST OF EXCIPIENTS**

Also contains: lactose and magnesium stearate.  
See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

6x1 capsules + 1 inhaler  
10x1 capsules + 1 inhaler  
12x1 capsules + 1 inhaler  
30x1 capsules + 1 inhaler

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For use only with the inhaler provided in the pack.  
Do not swallow capsules.  
Read the package leaflet before use.  
Inhalation use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

Each inhaler should be disposed of after 30 days of use.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C.

Store the capsules in the original package in order to protect from moisture and do not remove until immediately before use.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Frimley Business Park  
Camberley GU16 7SR  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/12/788/001	6 capsules + 1 inhaler
EU/1/12/788/007	10 capsules + 1 inhaler
EU/1/12/788/002	12 capsules + 1 inhaler
EU/1/12/788/003	30 capsules + 1 inhaler

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Seebri Breezhaler

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON OF MULTIPACK (INCLUDING BLUE BOX)**

**1. NAME OF THE MEDICINAL PRODUCT**

Seebri Breezhaler 44 micrograms inhalation powder, hard capsules  
glycopyrronium (glycopyrronium bromide)

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each capsule contains 50 micrograms glycopyrronium. The amount of glycopyrronium inhaled is 44 micrograms.

**3. LIST OF EXCIPIENTS**

Also contains: lactose and magnesium stearate.  
See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

Multipack: 90 (3 packs of 30x1) hard capsules + 3 inhalers.

Multipack: 96 (4 packs of 24x1) hard capsules + 4 inhalers.

Multipack: 150 (15 packs of 10x1) hard capsules + 15 inhalers.

Multipack: 150 (25 packs of 6x1) hard capsules + 25 inhalers.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For use only with the inhaler provided in the pack.

Do not swallow capsules.

Read the package leaflet before use.

Inhalation use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**



**8. EXPIRY DATE**

EXP

Each inhaler should be disposed of after 30 days of use.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C.

Store the capsules in the original package in order to protect from moisture and do not remove until immediately before use.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Frimley Business Park  
Camberley GU16 7SR  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/12/788/004	Multipack comprising 3 packs (30 capsules + 1 inhaler)
EU/1/12/788/005	Multipack comprising 4 packs (24 capsules + 1 inhaler)
EU/1/12/788/008	Multipack comprising 15 packs (10 capsules + 1 inhaler)
EU/1/12/788/006	Multipack comprising 25 packs (6 capsules + 1 inhaler)

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Seebri Breezhaler

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**INTERMEDIATE CARTON OF MULTIPACK (WITHOUT BLUE BOX)**

**1. NAME OF THE MEDICINAL PRODUCT**

Seebri Breezhaler 44 micrograms inhalation powder, hard capsules  
glycopyrronium (glycopyrronium bromide)

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each capsule contains 50 micrograms glycopyrronium. The amount of glycopyrronium inhaled is 44 micrograms micrograms.

**3. LIST OF EXCIPIENTS**

Also contains: lactose and magnesium stearate.  
See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Inhalation powder, hard capsule

30x1 capsules + 1 inhaler. Component of a multipack. Not to be sold separately.  
24x1 capsules + 1 inhaler. Component of a multipack. Not to be sold separately.  
10x1 capsules + 1 inhaler. Component of a multipack. Not to be sold separately.  
6x1 capsules + 1 inhaler. Component of a multipack. Not to be sold separately.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For use only with the inhaler provided in the pack.  
Do not swallow capsules.  
Read the package leaflet before use.  
Inhalation use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

Each inhaler should be disposed of after 30 days of use.

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 25°C.

Store the capsules in the original package in order to protect from moisture and do not remove until immediately before use.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited  
Frimley Business Park  
Camberley GU16 7SR  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/12/788/004	Multipack comprising 3 packs (30 capsules + 1 inhaler)
EU/1/12/788/005	Multipack comprising 4 packs (24 capsules + 1 inhaler)
EU/1/12/788/008	Multipack comprising 15 packs (10 capsules + 1 inhaler)
EU/1/12/788/006	Multipack comprising 25 packs (6 capsules + 1 inhaler)

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Seebri Breezhaler

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**INNER LID OF OUTER CARTON OF UNIT PACK AND OF INTERMEDIATE CARTON OF MULTIPACK**

**1. OTHER**

Do not push capsule through foil.

(1) Tear along perforations, (2) then peel back foil and (3) remove capsule.

Do not swallow capsules.

Read the leaflet before use.

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Seebri Breezhaler 44 mcg inhalation powder  
glycopyrronium

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Novartis Europharm Limited

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

Inhalation use only

**B. PACKAGE LEAFLET**

## Package leaflet: Information for the user

### Seebri Breezhaler 44 micrograms inhalation powder, hard capsules glycopyrronium (glycopyrronium bromide)

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

#### **Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Seebri Breezhaler is and what it is used for
2. What you need to know before you use Seebri Breezhaler
3. How to use Seebri Breezhaler
4. Possible side effects
5. How to store Seebri Breezhaler
6. Contents of the pack and other information

#### **1. What Seebri Breezhaler is and what it is used for**

##### **What Seebri Breezhaler is**

This medicine contains an active substance called glycopyrronium bromide. This belongs to a group of medicines called bronchodilators.

##### **What Seebri Breezhaler is used for**

This medicine is used to make breathing easier for adult patients who have breathing difficulties due to a lung disease called chronic obstructive pulmonary disease (COPD). When you inhale it, it helps you breathe more easily.

If you use this medicine once a day, it will help to reduce the effects of COPD on your everyday life.

In COPD the muscles around the airways tighten. This makes breathing difficult. This medicine blocks the tightening of these muscles in the lungs, making it easier for air to get in and out of the lungs.

#### **2. What you need to know before you use Seebri Breezhaler**

##### **Do not use Seebri Breezhaler**

- if you are allergic to glycopyrronium bromide or any of the other ingredients of this medicine (listed in section 6).

##### **Warnings and precautions**

Talk to your doctor before using Seebri Breezhaler, if any of the following applies to you:

- you have kidney problems.
- you have an eye problem called narrow-angle glaucoma.
- you have difficulty passing urine.



**During treatment with Seebri Breezhaler**, stop taking this medicine and tell your doctor immediately:

- if you experience tightness of the chest, coughing, wheezing or breathlessness immediately after using Seebri Breezhaler (signs of bronchospasm).
- if you experience difficulties in breathing or swallowing, swelling of the tongue, lips or face, skin rash, itching and hives (signs of allergic reaction).
- if you experience eye pain or discomfort, temporary blurring of vision, visual halos or coloured images in association with red eyes. These may be signs of an acute attack of narrow-angle glaucoma.

Seebri Breezhaler is used as a maintenance treatment for your COPD. Do not use this medicine to treat a sudden attack of breathlessness or wheezing.

### **Children and adolescents**

Do not give this medicine to children or adolescents below the age of 18 years.

### **Other medicines and Seebri Breezhaler**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes medicines similar to Seebri Breezhaler used for your lung disease, such as ipratropium, oxitropium or tiotropium (so called anticholinergics).

No specific side effects have been reported when Seebri Breezhaler has been taken together with other products used to treat COPD such as reliever inhalers (e.g. salbutamol), methylxanthines (e.g. theophylline) and/or oral and inhaled steroids (e.g. prednisolone).

### **Pregnancy, breast-feeding and fertility**

There are no data from the use of this medicine in pregnant women and it is not known whether the active substance of this medicine passes into human milk.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

### **Driving and using machines**

It is unlikely that this medicine will affect your ability to drive and use machines.

### **Seebri Breezhaler contains lactose**

This medicine contains lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

Ask your doctor or pharmacist for advice before taking any medicine.

## **3. How to use Seebri Breezhaler**

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

### **How much Seebri Breezhaler to use**

The usual dose is to inhale the content of one capsule each day.

You only need to inhale once a day because the effect of this medicine lasts for 24 hours.

Do not use more than your doctor tells you to use.

### **Older people**

You can use this medicine if you are aged 75 years and over at the same dose as for other adults.

### **When to inhale Seebri Breezhaler**

Use this medicine at the same time each day. This will also help you to remember to use it.

### **Seebri Breezhaler with food and drink**

You can inhale this medicine anytime before or after food or drink.

### **How to inhale Seebri Breezhaler**

- In this pack, you will find an inhaler and capsules (in blisters) that contain the medicine as inhalation powder. Only use the capsules with the inhaler provided in this pack (Seebri Breezhaler inhaler). The capsules should remain in the blister until you need to use them.
- Do not push the capsule through the foil.
- When you start a new pack, use the new Seebri Breezhaler inhaler that is supplied in the pack.
- Dispose of each inhaler after 30 days of use.
- Do not swallow the capsules.
- Please read the instructions at the end of this leaflet for more information on how to use the inhaler.

### **If you use more Seebri Breezhaler than you should**

If you have inhaled too much of this medicine or if someone else accidentally uses your capsules, you must immediately either tell your doctor or go to the nearest emergency unit. Show the pack of Seebri Breezhaler. Medical attention may be needed.

### **If you forget to use Seebri Breezhaler**

If you forget to inhale a dose, take one as soon as possible. However, do not take two doses on the same day. Then take the next dose as usual.

### **How long to continue your treatment with Seebri Breezhaler**

- Keep using this medicine for as long as your doctor tells you.
- COPD is a long-term disease and you should use this medicine every day and not only when you have breathing problems or other symptoms of COPD.

If you have questions about how long to continue your treatment with this medicine, talk to your doctor or pharmacist.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

### **Some side effects may be serious but are uncommon**

- Irregular heart beat
- High level of blood sugar (hyperglycaemia: typical symptoms include excessive thirst or hunger and frequent urination)
- Rash, itching, hives, difficulty breathing or swallowing, dizziness (possible signs of allergic reaction)
- Swelling mainly of the tongue, lips, face or throat (possible signs of angioedema)

If you get any of these side effects, tell your doctor immediately.

### **Some side effects may be serious, but the frequency of these side effects is unknown**

- Difficulty breathing with wheezing or coughing (signs of paradoxical bronchospasm)

**Some side effects are common**

*(may affect up to 1 in 10 people)*

- Dry mouth
- Difficulty sleeping
- Runny or stuffy nose, sneezing, sore throat
- Diarrhoea or stomach ache

**Some side effects are uncommon**

*(may affect up to 1 in 100 people)*

- Difficulty and pain when passing urine
- Painful and frequent urination
- Palpitations
- Rash
- Numbness
- Cough with sputum
- Dental caries
- Feeling of pressure or pain in the cheeks and forehead
- Nose bleeds
- Pain in arms or legs
- Pain in muscles, bones or joints of the chest
- Stomach discomfort after meals
- Throat irritation
- Tiredness
- Weakness
- Itching
- Voice alteration (hoarseness)

Some elderly patients above 75 years of age experienced headache (frequency common) and urinary tract infection (frequency common).

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Seebri Breezhaler**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after “EXP”. The expiry date refers to the last day of that month.

Do not store above 25°C.

Store the capsules in the original package in order to protect from moisture and do not remove until immediately before use.

Each inhaler should be disposed of after 30 days of use.

Do not use this medicine if you notice that the pack is damaged or shows signs of tampering.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Seebri Breezhaler contains

- The active substance is glycopyrronium bromide. Each capsule contains 63 micrograms of glycopyrronium bromide (equivalent to 50 micrograms glycopyrronium). The delivered dose (the dose that leaves the mouthpiece of the inhaler) is equivalent to 44 micrograms of glycopyrronium.
- The other ingredients of the inhalation powder are lactose monohydrate and magnesium stearate.

### What Seebri Breezhaler looks like and contents of the pack

In this pack, you will find a device called an inhaler, together with capsules in blisters. Each blister contains either 6 or 10 hard capsules. The capsules are transparent and orange and contain a white powder. They have the product code “GPL50” printed in black above and a company logo (Ⓛ) printed in black below a black bar.

The following pack sizes are available:

Single pack containing 6x1, 10x1, 12x1 or 30x1 hard capsules, together with one inhaler.

Multipacks containing 90 (3 packs of 30x1) hard capsules and 3 inhalers.

Multipacks containing 96 (4 packs of 24x1) hard capsules and 4 inhalers.

Multipacks containing 150 (15 packs of 10x1) hard capsules and 15 inhalers.

Multipacks containing 150 (25 packs of 6x1) hard capsules and 25 inhalers.

Not all pack sizes may be available in your country.

### Marketing Authorisation Holder

Novartis Europharm Limited  
Frimley Business Park  
Camberley GU16 7SR  
United Kingdom

### Manufacturer

Novartis Pharma GmbH  
Roonstraße 25  
D-90429 Nuremberg  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### België/Belgique/Belgien

Novartis Pharma N.V.  
Tél/Tel: +32 2 246 16 11

#### Lietuva

Novartis Pharma Services Inc.  
Tel: +370 5 269 16 50

#### България

Novartis Pharma Services Inc.  
Тел.: +359 2 489 98 28

#### Luxembourg/Luxemburg

Novartis Pharma N.V.  
Tél/Tel: +32 2 246 16 11

#### Česká republika

Novartis s.r.o.  
Tel: +420 225 775 111

#### Magyarország

Novartis Hungária Kft. Pharma  
Tel.: +36 1 457 65 00

**Danmark**

Novartis Healthcare A/S  
Tlf: +45 39 16 84 00

**Deutschland**

Novartis Pharma GmbH  
Tel: +49 911 273 0

**Eesti**

Novartis Pharma Services Inc.  
Tel: +372 66 30 810

**Ελλάδα**

Novartis (Hellas) A.E.B.E.  
Τηλ: +30 210 281 17 12

**España**

Novartis Farmacéutica, S.A.  
Tel: +34 93 306 42 00

**France**

Novartis Pharma S.A.S.  
Tél: +33 1 55 47 66 00

**Hrvatska**

Novartis Hrvatska d.o.o.  
Tel. +385 1 6274 220

**Ireland**

Novartis Ireland Limited  
Tel: +353 1 260 12 55

**Ísland**

Vistor hf.  
Sími: +354 535 7000

**Italia**

Novartis Farma S.p.A.  
Tel: +39 02 96 54 1

**Κύπρος**

Novartis Pharma Services Inc.  
Τηλ: +357 22 690 690

**Latvija**

Novartis Pharma Services Inc.  
Tel: +371 67 887 070

**Malta**

Novartis Pharma Services Inc.  
Tel: +356 2122 2872

**Nederland**

Novartis Pharma B.V.  
Tel: +31 26 37 82 111

**Norge**

Novartis Norge AS  
Tlf: +47 23 05 20 00

**Österreich**

Novartis Pharma GmbH  
Tel: +43 1 86 6570

**Polska**

Novartis Poland Sp. z o.o.  
Tel.: +48 22 375 4888

**Portugal**

Novartis Farma - Produtos Farmacêuticos, S.A.  
Tel: +351 21 000 8600

**România**

Novartis Pharma Services Romania SRL  
Tel: +40 21 31299 01

**Slovenija**

Novartis Pharma Services Inc.  
Tel: +386 1 300 75 50

**Slovenská republika**

Novartis Slovakia s.r.o.  
Tel: +421 2 5542 5439

**Suomi/Finland**

Novartis Finland Oy  
Puh/Tel: +358 (0)10 6133 200

**Sverige**

Novartis Sverige AB  
Tel: +46 8 732 32 00

**United Kingdom**

Novartis Pharmaceuticals UK Ltd.  
Tel: +44 1276 698370

**This leaflet was last revised in**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency website:  
<http://www.ema.europa.eu>

## Instructions for use of Seebri Breezhaler inhaler


Please read the following instructions carefully to learn how to use this medicine.

- Only use the Seebri Breezhaler inhaler contained in this pack. Do not use Seebri Breezhaler capsules with any other inhaler, and do not use Seebri Breezhaler inhaler to take any other capsule medicine.
- Do not push the capsule through the foil to remove it from the blister.
- When you start a new pack, only use the new Seebri Breezhaler inhaler that is supplied in the pack.
- Each inhaler should be disposed of after 30 days of use. Ask your pharmacist how to dispose of medicines and inhalers no longer required.
- Do not swallow the capsules. The powder in the capsules is for you to inhale.


### Your Seebri Breezhaler pack

One Seebri Breezhaler pack contains:


- one Seebri Breezhaler inhaler
- one or more blisters, each containing either 6 or 10 Seebri Breezhaler capsules to be used in the inhaler.



**Inhaler**




**Blister card**




**Inhaler base**

### How to use your inhaler




**Pull off the cap.**



**Open inhaler:**

Hold the base of the inhaler firmly and tilt the mouthpiece. This opens the inhaler.




**Prepare capsule:**

Separate one of the blisters from the blister card by tearing along the perforation.

Take one blister and peel away the protective backing to expose the capsule.


Do not push capsule through foil.



**Remove a capsule:**

Capsules should always be stored in the blister and only removed immediately before use.

With dry hands, remove capsule from the blister. Do not swallow the capsule.



**Insert capsule:**





Place the capsule into the capsule chamber.

**Never place a capsule directly into the mouthpiece.**



**Close the inhaler:**

Close the inhaler until you hear a “click”.



**Pierce the capsule:**

- Hold the inhaler upright with the mouthpiece pointing up.
- Pierce the capsule by firmly pressing together both side buttons at the same time.  
**Do this only once.**
- You should hear a “**click**” as the capsule is being pierced.

**Release the side buttons fully.**

**Breathe out:**

Before placing the mouthpiece in your mouth, breathe out fully.




**Do not blow into the mouthpiece.**

**Inhale the medicine:**

To breathe the medicine deeply into your airways:

- Hold the inhaler as shown in the picture. The side buttons should be facing left and right. Do not press the side buttons.
- Place the mouthpiece in your mouth and close your lips firmly around it.
- Breathe in rapidly but steadily, as deeply as you can. **Do not press the side buttons.**





**Note:**

As you breathe in through the inhaler, the capsule spins around in the chamber and you should hear a whirring noise. You will experience a sweet flavour as the medicine goes into your lungs.

**If you do not hear a whirring noise:**

The capsule may be stuck in the capsule chamber. If this happens:

- Open the inhaler and carefully loosen the capsule by tapping the base of the inhaler. **Do not press the side buttons.**
- Inhale the medicine again by repeating steps 9 and 10.

**Hold breath:**

**After you have inhaled the medicine:**

- **Hold your breath** for at least 5-10 seconds or as long as you comfortably can while taking the inhaler out of your mouth.
- Then breathe out.
- Open the inhaler to see if any powder is left in the capsule.

**If there is powder left in the capsule:**

- Close the inhaler.
- Repeat steps 9 to 12.

Most people are able to empty the capsule with one or two inhalations.

**Additional information**

Some people occasionally cough briefly soon after inhaling a medicine. If you do, don't worry. As long as the capsule is empty, you have received enough of your medicine.

**After you have finished taking your daily dose of Seebri Breezhaler:**

- Open the mouthpiece again, and remove the empty capsule by tipping it out of the capsule chamber. Put the empty capsule in your household waste.
- Close the inhaler and replace the cap.

**Do not store the capsules in the Seebri Breezhaler inhaler.**

**Additional information**

Occasionally, very small pieces of the capsule can get past the screen and enter your mouth. If this happens, you may be able to feel these pieces on your tongue. It is not harmful if these pieces are swallowed or inhaled. The chances of the capsule shattering will be increased if the capsule is pierced more than once (step 7).

**How to clean your inhaler**

Never wash your inhaler with water. If you want to clean your inhaler, wipe the mouthpiece inside and outside with a clean, dry, lint-free cloth to remove any powder residue. Keep the inhaler dry.