

## Summary of the risk management plan (RMP) for Ivabradine Anpharm (ivabradine)

This is a summary of the risk management plan (RMP) for Ivabradine Anpharm, which details the measures to be taken in order to ensure that Ivabradine Anpharm is used as safely as possible. For more information on RMP summaries, see [here](#).

This RMP summary should be read in conjunction with the EPAR summary and the product information for Ivabradine Anpharm, which can be found on [Ivabradine Anpharm's EPAR page](#).

### Overview of disease epidemiology

Ivabradine Anpharm is a medicine used for two long-term (chronic) heart conditions: to treat symptoms of chronic stable angina pectoris and to treat chronic heart failure.

Stable angina pectoris (angina) is a heart disease which happens when blood supply to the heart muscle is inadequate, so the heart does not receive enough oxygen. It usually appears between 40 and 50 years of age. The most common symptom of angina is pain in the chest, jaw and back, particularly when the heart beats faster in situations such as exercise, emotion, exposure to the cold or after eating. It is estimated that between 2 and 4% of the population in the EU suffer from stable angina pectoris.

Chronic heart failure is a heart disease in which the heart cannot pump enough blood to the rest of the body. The most common symptoms of heart failure are breathlessness, tiredness and ankle swelling due to fluid build-up. It is estimated to affect between 1 and 2% of the population in the EU.

### Summary of treatment benefits

Ivabradine Anpharm has been studied in five main studies involving over 4,000 adults with long-term stable angina. The medicine was compared with placebo (a dummy treatment) in 360 patients, atenolol (a beta-blocker) in 939 patients and amlodipine (another medicine used to treat angina) in 1,195 patients. It was also compared with placebo as an add-on to atenolol in 889 patients and as an add-on to amlodipine in 728 patients. Each study lasted three to four months. The main measure of effectiveness was how long the patients could exercise on a bicycle or a treadmill, measured at the start and the end of each study. The studies showed that the medicine was more effective than placebo at improving exercise capacity and was as effective as atenolol and amlodipine. Ivabradine Anpharm was also more effective than placebo when added to atenolol. However, adding it to amlodipine did not provide an additional benefit.

Ivabradine Anpharm has also been compared with placebo in one main study involving 6,558 patients with long-term moderate to severe heart failure. The main measure of effectiveness was the time until death due to disease of the heart or blood vessels, or hospitalisation due to worsening heart failure. It was more effective than placebo at preventing death due to disease of the heart or blood vessels or hospitalisation due to worsening heart failure: 24.5% (793 out of 3,241) of patients treated with

Ivabradine Anpharm died or were hospitalised for the first time due to worsening heart failure, compared with 28.7% (937 out of 3,264) of patients treated with placebo.

Another study compared Ivabradine Anpharm with placebo in 19,102 patients with coronary heart disease and without clinical heart failure. The main measure of effectiveness was a reduction in the risk of death due to heart problems and non-fatal heart attack. In this study a specific subgroup of patients who had symptomatic angina had a small but significant increase in the combined risk of cardiovascular death or non-fatal heart attack with Ivabradine Anpharm compared with placebo (3.4% vs 2.9% yearly incidence rates). However it should be noted that patients in this study were given doses higher than the recommended dose (up to 10 mg twice a day).

## Unknowns relating to treatment benefits

There is limited experience of using Ivabradine Anpharm in patients with reduced kidney and liver function and in patients suffering from chronic heart failure who have certain problems with electrical conduction in the heart (such as so-called bundle branch blocks).

## Summary of safety concerns

### *Important identified risks*

<b>Risk</b>	<b>What is known</b>	<b>Preventability</b>
Decrease in heart rate (bradycardia)	Decrease in heart rate is a common side effect with Ivabradine Anpharm (seen in between 1 and 10 patients in 100), especially in the first 2 to 3 months after starting treatment.  This may lead to dizziness, low blood pressure, fainting, feeling of weakness, feeling of faintness or feeling unwell.	Doctors should follow the dosing recommendations in the product information and not exceed the maximum recommended dose. Treatment must only be started in patients with heart rate above 70 beats per minute, and must be avoided in patients also taking the medicines verapamil or diltiazem.
Luminous visual phenomena (phosphenes/ blurred vision)	Luminous visual phenomena or phosphenes (brief moments of increased brightness, coloured flashes or halos) are very common side effects with Ivabradine Anpharm (seen in more than 1 patient in 10). They are most often caused by sudden changes in light intensity. They generally occur within the first two months of treatment.	The product information contains warnings for doctors and patients about the possibility of luminous visual phenomena, which are linked to the way the medicine works. Patients should be informed that phosphenes resolve during or after treatment with no after effects. If affected, patients should be warned to be careful when driving or using machines at times when there could be sudden changes in light intensity, especially when driving at night.
Irregular heartbeats (2 <sup>nd</sup> and 3 <sup>rd</sup> degree atrioventricular blocks (AVB II and III))	Irregular heartbeats due to interference with the electrical conduction of the heart are very rare side effects with Ivabradine Anpharm (seen in less than 1 patient in 10,000).	Patients receiving Ivabradine Anpharm will normally have their heart function regularly monitored, particularly when starting treatment or changing dose. The product information contains a warning for doctors that Ivabradine Anpharm is

<b>Risk</b>	<b>What is known</b>	<b>Preventability</b>
		not recommended in patients with existing 2 <sup>nd</sup> degree atrioventricular block and must be avoided in those with existing 3 <sup>rd</sup> degree block.
Uncontrolled blood pressure (increase in blood pressure in hypertensive patients)	Uncontrolled blood pressure is a common side effect with Ivabradine Anpharm (seen in between 1 and 10 patients in 100). This may happen in patients suffering from hypertension (high blood pressure), especially after a change in their blood pressure treatment.	The product information for Ivabradine Anpharm warns doctors that when blood pressure treatment is modified in chronic heart failure patients treated with Ivabradine Anpharm, their blood pressure should be monitored at an appropriate interval.
Irregular rapid contractions of the upper chambers of the heart (atrial fibrillation)	Atrial fibrillation (a rapid, irregular rhythm of the upper chambers of the heart) is a common side effect with Ivabradine Anpharm (seen in between 1 and 10 patients in 100).	The product information states that patients should be informed of signs and symptoms of atrial fibrillation (such as high pulse rate when resting, or irregular pulse rate) and be advised to contact their doctor if these occur.  If atrial fibrillation develops during treatment, the balance of benefits and risks of continued ivabradine treatment should be carefully reconsidered.
Abnormal ECG heart tracing (prolonged QT interval on ECG)	A type of abnormal ECG heart tracing known as QT prolongation (reflecting a change in the electrical activity of the heart) is an uncommon side effect with Ivabradine Anpharm (seen between 1 and 10 patients in 1,000). This event is linked to the effect of ivabradine in lowering heart rate.	The product information warns doctors that use of Ivabradine Anpharm in patients with inborn alteration in QT interval (congenital QT syndrome) or with other medicines that may also prolong QT interval should be avoided if possible. If Ivabradine Anpharm is used in such circumstances close monitoring of heart function is needed.

### ***Important potential risks***

<b>Risk</b>	<b>What is known</b>
Irregular contractions of the upper chambers of the heart (supra-ventricular tachyarrhythmia other than atrial fibrillation)	Based on clinical trial data, supra-ventricular tachyarrhythmia (fast, irregular contractions of the upper chambers of the heart) of types other than atrial fibrillation could occur. However, studies performed in current medical practice reported a similar risk in patients treated with Ivabradine Anpharm and patients given placebo (dummy treatment).
Immune disorders	Based on data in animals, Ivabradine Anpharm may cause shrinkage of the thymus gland, which plays a role in the immune system (the body's natural

<b>Risk</b>	<b>What is known</b>
	defences). Studies in humans have not so far shown any role of the medicine in the occurrence of immune disorders.
Severe irregular contractions of the lower chambers of the heart (severe ventricular arrhythmia)	Based on post-marketing data, severe ventricular arrhythmias may happen in patients with different risk factors or taking other drugs. However, clinical studies in humans have not so far shown an increased risk in patients treated with Ivabradine Anpharm.
Heart attack (myocardial infarction)	Based on the preliminary results of a large clinical study called SIGNIFY, in which Ivabradine Anpharm was given differently than currently authorised, severe slowing of heart rate (bradycardia) caused by the medicine could lead to myocardial infarction.

### ***Missing information***

<b>Risk</b>	<b>What is known</b>
Limited information on use in children under 18 years old	The effectiveness and safety of Ivabradine Anpharm have not been studied in this population.
Limited information on use in pregnancy and breastfeeding women	The effectiveness and safety of Ivabradine Anpharm have not been studied in this population. Ivabradine Anpharm must not be used during pregnancy or breastfeeding as studies in animals suggest it can harm the unborn child. Women who are able to have children should use appropriate contraceptive measures while taking the medicine.
Limited information on use in patients with severely reduced liver function (hepatic insufficiency)	The effectiveness and safety of Ivabradine Anpharm have not been studied in this population and it is recommended that they should not be given the medicine,
Limited information on use in patients with severely reduced kidney function (renal impairment)	The effectiveness and safety of Ivabradine Anpharm have not been studied in this population and it should be used with caution.
Limited information on use in chronic heart failure patients with intra-ventricular conduction defects	The effectiveness and safety of Ivabradine Anpharm have not been fully studied in this population.

## Summary of risk minimisation measures by safety concern

All medicines have a summary of product characteristics (SmPC) which provides physicians, pharmacists and other healthcare professionals with details on how to use the medicine, and also describes the risks and recommendations for minimising them. Information for patients is available in lay language in the package leaflet. The measures listed in these documents are known as 'routine risk minimisation measures'.

The SmPC and the package leaflet are part of the medicine's product information. The product information for Ivabradine Anpharm can be found on [Ivabradine Anpharm's EPAR page](#).

This medicine has no additional risk minimisation measures.

## Planned post-authorisation development plan

### *List of studies in post-authorisation development plan*

<b>Study/activity (including study number)</b>	<b>Objectives</b>	<b>Safety concerns /efficacy issue addressed</b>	<b>Status</b>	<b>Planned date for submission of (interim and) final results</b>
CL3-067	Documentation of the long-term (3 years) ophthalmic safety of ivabradine in patients with chronic stable angina pectoris with specific examinations.	Long-term ocular safety	Ongoing	Final report : Q3.2016
Drug Utilisation study	Describing the characteristics of ivabradine users, as well as the patterns of use of ivabradine, and adherence to the risk minimisation measures.	Effectiveness of measures to reduce the risks of treatment	Ongoing	Final report: 06.2018

### *Studies which are a condition of the marketing authorisation*

The drug utilisation study is a condition of the marketing authorisation

## Summary of changes to the risk management plan over time

### *Major changes to the Risk Management Plan over time*

Not applicable.

This summary was last updated in 09-2015.