



17 January 2013  
EMA/CHMP/30243/2013  
Committee for Medicinal Products for Human Use (CHMP)

## Summary of opinion<sup>1</sup> (post authorisation)

---

### Humira

#### adalimumab

On 17 January 2013, the Committee for Medicinal Products for Human Use (CHMP) adopted a positive opinion recommending a variation to the terms of the marketing authorisation for the medicinal product Humira. The marketing authorisation holder for this medicinal product is AbbVie Ltd. They may request a re-examination of the CHMP opinion, provided that they notify the European Medicines Agency in writing of their intention within 15 days of receipt of the opinion.

The CHMP adopted a change to an indication as follows<sup>2</sup>:

#### Polyarticular juvenile idiopathic arthritis

Humira in combination with methotrexate is indicated for the treatment of active polyarticular juvenile idiopathic arthritis, in children and adolescents aged **2** to 17 years who have had an inadequate response to one or more disease-modifying anti-rheumatic drugs (DMARDs). Humira can be given as monotherapy in case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate (for the efficacy in monotherapy see section 5.1). Humira has not been studied in children aged less than **2** years.

Detailed conditions for the use of this product will be described in the updated summary of product characteristics (SmPC), which will be published in the revised European public assessment report (EPAR), and will be available in all official European Union languages after the variation to the marketing authorisation has been granted by the European Commission.

For information, the full indications for Humira will be as follows:

#### Rheumatoid arthritis

Humira in combination with methotrexate, is indicated for:

- the treatment of moderate to severe, active rheumatoid arthritis in adult patients when the response to disease-modifying anti-rheumatic drugs including methotrexate has been inadequate.

---

<sup>1</sup> Summaries of positive opinion are published without prejudice to the Commission decision, which will normally be issued within 44 days (Type II variations) and 67 days (Annex II applications) from adoption of the opinion.

<sup>2</sup> The text in bold represents the new or the amended indication.



- the treatment of severe, active and progressive rheumatoid arthritis in adults not previously treated with methotrexate.

Humira can be given as monotherapy in case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate.

Humira has been shown to reduce the rate of progression of joint damage as measured by X-ray and to improve physical function, when given in combination with methotrexate.

#### Polyarticular juvenile idiopathic arthritis

Humira in combination with methotrexate is indicated for the treatment of active polyarticular juvenile idiopathic arthritis, in children and adolescents aged 2 to 17 years who have had an inadequate response to one or more disease-modifying anti-rheumatic drugs (DMARDs). Humira can be given as monotherapy in case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate. (for the efficacy in monotherapy see section 5.1). Humira has not been studied in children aged less than 2 years.

#### Axial spondyloarthritis

##### Ankylosing spondylitis(AS)

Humira is indicated for the treatment of adults with severe active ankylosing spondylitis who have had an inadequate response to conventional therapy.

##### Axial spondyloarthritis without radiographic evidence of AS

Humira is indicated for the treatment of adults with severe axial spondyloarthritis without radiographic evidence of AS but with objective signs of inflammation by elevated CRP and / or MRI, who have had an inadequate response to, or are intolerant to nonsteroidal anti-inflammatory drugs.

#### Psoriatic arthritis

Humira is indicated for the treatment of active and progressive psoriatic arthritis in adults when the response to previous disease-modifying anti-rheumatic drug therapy has been inadequate. Humira has been shown to reduce the rate of progression of peripheral joint damage as measured by X-ray in patients with polyarticular symmetrical subtypes of the disease (see Section 5.1) and to improve physical function.

#### Psoriasis

Humira is indicated for the treatment of moderate to severe chronic plaque psoriasis in adult patients who failed to respond to or who have a contraindication to, or are intolerant to other systemic therapy including cyclosporine, methotrexate or PUVA.

#### Crohn's disease

Humira is indicated for treatment of moderately to severely active Crohn's disease, in adult patients who have not responded despite a full and adequate course of therapy with a corticosteroid and/or an immunosuppressant; or who are intolerant to or have medical contraindications for such therapies.

#### Paediatric Crohn's Disease

Humira is indicated for the treatment of severe active Crohn's disease in paediatric patients (6 to 17 years of age) who have had an inadequate response to conventional therapy including primary nutrition therapy, a corticosteroid, and an immunomodulator, or who are intolerant to or have contraindications for such therapies.

#### Ulcerative colitis

Humira is indicated for treatment of moderately to severely active ulcerative colitis in adult patients who have had an inadequate response to conventional therapy including corticosteroids and 6-mercaptopurine (6-MP) or azathioprine (AZA), or who are intolerant to or have medical contraindications for such therapies.