



EUROPEAN MEDICINES AGENCY  
SCIENCE MEDICINES HEALTH

05 June 2018  
EMA/HMPC/444244/2015  
Committee on Herbal Medicinal Products (HMPC)

## European Union herbal monograph on *Pelargonium sidoides* DC and/or *Pelargonium reniforme* Curt., radix

**This document was valid from 05 June 2018 until May 2024. It is now superseded by a new version adopted by the HMPC on 29 May 2024 and published on the EMA website.**

<b>Initial assessment</b>	
Discussion in Working Party on Community monographs and Community list (MLWP)	September 2010 November 2010 January 2011 March 2011
Adopted by Committee on Herbal Medicinal Products (HMPC) for release for consultation	31 March 2011
End of consultation (deadline for comments).	15 August 2011
Re-discussion in MLWP	September 2011 November 2011 January 2012 May 2012
Adoption by HMPC Monograph (EMA/HMPC/560961/2010) AR (EMA/HMPC/560962/2010) List of references (EMA/HMPC/560963/2010) Overview of comments received during the public consultation (EMA/HMPC/748350/2011)> HMPC Opinion (EMA/HMPC/742263/2012)	20 November 2012
<b>First review</b>	
Discussion in MLWP	September 2014 July 2015
Adopted by HMPC for release for consultation	29 September 2015
End of consultation (deadline for comments <sup>1</sup> )	31 January 2016
Re-discussion in MLWP	April 2016 May/June 2016 July 2016

<sup>1</sup> No comments were received during the period of public consultation. Therefore the final monograph is published together with the final assessment report and list of references, without an 'Overview of comments received during the public consultation'.



	November 2016 March 2017 May 2017 July 2017 September 2017 November 2017 January 2017 March 2018
Adoption by HMPC	05 June 2018

Keywords	Herbal medicinal products; HMPC; Community herbal monographs; traditional use; <i>Pelargonium sidoides</i> DC and/or <i>Pelargonium reniforme</i> Curt., radix; <i>Pelargonii</i> radix; pelargonium root
----------	---

BG (bългарски): Пеларгониум, корен CS (čeština): pelargoniový kořen DA (dansk): Pelargonierod DE (Deutsch): Pelargoniumwurzel EL (elliniká): πελαργονίου ρίζα EN (English): pelargonium root ES (español): pelargonio, raíz de ET (eesti keel): pelargoonijuur FI (suomi): pelargoni, juuri FR (français): pélargonium (racine de) HU (magyar): muskátligyökér IT (italiano): Pelargonio radice	LT (lietuvių kalba): Pelargonijų šaknys LV (latviešu valoda): Pelargonijas saknes MT (malti): għerq tal-ġeranju NL (nederlands): Geranium PL (polski): Korzeń pelargonii PT (português): pelargónio, raiz RO (română): rădăcină de muscata SK (slovenčina): koreň muškátu SL (slovenščina): korenina pelargonije SV (svenska): pelargon, rot IS (íslenska): NO (norsk): pelargoniumrot
--	---

# European Union herbal monograph on *Pelargonium sidoides* DC and/or *Pelargonium reniforme* Curt., radix

## 1. Name of the medicinal product

To be specified for the individual finished product.

## 2. Qualitative and quantitative composition<sup>2, 3</sup>

Well-established use	Traditional use
	With regard to the registration application of Article 16d(1) of Directive 2001/83/EC  <i>Pelargonium sidoides</i> DC and/or <i>Pelargonium reniforme</i> Curt., radix (pelargonium root)  i) Herbal substance  Not applicable  ii) Herbal preparations  a) Liquid extract (DER 1:8-10), extraction solvent ethanol 11% (m/m)  b) Dry extract, (DER 4-25:1), extraction solvent ethanol 11% (m/m)

## 3. Pharmaceutical form

Well-established use	Traditional use
	Herbal preparations in liquid or solid dosage forms for oral use.  The pharmaceutical form should be described by the European Pharmacopoeia full standard term.

## 4. Clinical particulars

### 4.1. Therapeutic indications

Well-established use	Traditional use
	Traditional herbal medicinal product for the symptomatic treatment of common cold.  The product is a traditional herbal medicinal

<sup>2</sup> The material complies with the Ph. Eur. monograph (ref.: 2264).

<sup>3</sup> The declaration of the active substance(s) for an individual finished product should be in accordance with relevant herbal quality guidance.

Well-established use	Traditional use
	product for use in the specified indication exclusively based upon long-standing use.

#### 4.2. Posology and method of administration

Well-established use	Traditional use
	<p><b>Posology</b></p> <p>Single dose</p> <p><i>Adolescents over the age of 12 years, adults and elderly</i></p> <p>a) Liquid extract: 1.19-1.25 ml, 3 times daily.</p> <p>b) Dry extract: 20 mg, 3 times daily.</p> <p><i>Children between 6-12 years</i></p> <p>a) Liquid extract: 0.79-0.83 ml, 3 times daily.</p> <p>b) Dry extract: 20 mg, 2 times daily.</p> <p>The use in children under 6 years of age is not recommended (see section 4.4 'Special warnings and precautions for use').</p> <p><b>Duration of use</b></p> <p>If the symptoms persist longer than 1 week during the use of the medicinal product, a doctor or a qualified health care practitioner should be consulted.</p> <p><b>Method of administration</b></p> <p>Oral use</p>

#### 4.3. Contraindications

Well-established use	Traditional use
	Hypersensitivity to the active substance(s).

#### 4.4. Special warnings and precautions for use

Well-established use	Traditional use
	<p>The use in children under 6 years of age has not been established due to lack of adequate data.</p> <p>Hepatotoxicity and hepatitis cases were reported in association with the administration of the</p>

Well-established use	Traditional use
	<p>medicinal product. In case signs of hepatotoxicity occur, the administration of the medicinal product should be stopped immediately and a medical doctor should be consulted.</p> <p>If the symptoms worsen during the use of the medicinal product, a doctor or a qualified health care practitioner should be consulted.</p> <p>For liquid extracts containing ethanol, the appropriate labelling for ethanol, taken from the 'Guideline on excipients in the label and package leaflet of medicinal products for human use', must be included.</p>

#### **4.5. Interactions with other medicinal products and other forms of interaction**

Well-established use	Traditional use
	None reported

#### **4.6. Pregnancy and lactation**

Well-established use	Traditional use
	<p>No fertility data available.</p> <p>Safety during pregnancy and lactation has not been established. In the absence of sufficient data, the use during pregnancy and lactation is not recommended.</p>

#### **4.7. Effects on ability to drive and use machines**

Well-established use	Traditional use
	No studies on the effect on the ability to drive and use machines have been performed.

#### **4.8. Undesirable effects**

Well-established use	Traditional use
	Mild gastrointestinal complaints (diarrhea, epigastric discomfort, nausea or vomiting, dysphagia), mild nasal and gingival bleeding and allergic reactions have been reported. The

Well-established use	Traditional use
	<p>frequency was very rare.</p> <p>Hepatotoxicity has been reported. The frequency is not known.</p> <p>If other adverse reactions not mentioned above occur, a doctor or a qualified health care practitioner should be consulted.</p>

#### 4.9. Overdose

Well-established use	Traditional use
	No case of overdose has been reported.

## 5. Pharmacological properties

### 5.1. Pharmacodynamic properties

Well-established use	Traditional use
	Not required as per Article 16c(1)(a)(iii) of Directive 2001/83/EC.

### 5.2. Pharmacokinetic properties

Well-established use	Traditional use
	Not required as per Article 16c(1)(a)(iii) of Directive 2001/83/EC.

### 5.3. Preclinical safety data

Well-established use	Traditional use
	<p>Not required as per Article 16c(1)(a)(iii) of Directive 2001/83/EC, unless necessary for the safe use of the product.</p> <p>Adequate tests on reproductive toxicity, genotoxicity and carcinogenicity are not available.</p>

## 6. Pharmaceutical particulars

Well-established use	Traditional use
	Not applicable

## 7. Date of compilation/last revision

05 June 2018

Superseded