



EUROPEAN MEDICINES AGENCY  
SCIENCE MEDICINES HEALTH

15 April 2011  
EMA/HMPC/143182/2010  
Committee on Herbal Medicinal Products (HMPC)

## List of references supporting the assessment of *Lavandula angustifolia* Mill., aetheroleum and *Lavandula angustifolia* Mill., flos

Draft

**The Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.**

Akhondzadeh S, Kashani I, Fotouhi A, Jarvandi S, Mohaseri M, Moin M, et al. Comparison of *Lavandula angustifolia* Mill. tincture and imipramine in the treatment of mild to moderate depression: a double-blind, randomized trial. *Prog Neuropsychopharmacol Biol Psychiat* 2003; 27: 123-127.

Akutsu H, Kikusui T, Takeuchi Y, Sano K, Hatanaka A, Mori Y. Alleviating effects of plant-derived fragrances on stress-induced hyperthermia in rats. *Physiol Behav* 2002; 75: 355-360.

Atassanova-Shopova S, Roussinov KS. On certain central neurotropic effects of lavender essential oil. *Bulletin of the Institute of Physiology (Bulgarian Academy of Sciences)* 1970; 13: 69-77.

Atsumi T, Tonosaki K. Smelling lavender and rosemary increases free radical scavenging activity and decreases cortisol level in saliva. *Psychiatry Research* 2007; 150(1): 89-96.

Austin LM, Alford C. The sedating effects of lavender oil and the stimulating effects of jasmine oil on sleep. *J Psychopharmacol Oxford* 1997; 11(3): A55, abstract 2171997.

Bickers D, Calow P, Greim h, Hanifin JM, Rogers AE, Saurat JH, Sipes IG, Smith RL, Tagami H. A toxicological and dermatological assessment of linalool and related esters when used as fragrance ingredients. *Food and Chemical Toxicology* 2003; 41: 919-942.

Blumenthal M, Goldberg A, Brinckmann J, Foster S. Herbal Medicine. Expanded Commission E Monographs. American Botanical Council, Austin Texas 2000; 226-229.

Bradley BF, Brown SL, Chu S, Lea RW. Effects of orally administered lavender essential on responses to anxiety-provoked film clips. *Hum Psychopharmacol* 2009; 24(4): 319-30.

Bradley BF, Starkey NJ, Brown SL, Lea RW. Anxiolytic effects of *Lavandula angustifolia* odour on the Mongolian gerbil elevated plus maze. *J Ethnopharmacology* 2007; 111(3): 517-25.

British Herbal Pharmacopoeia, 2<sup>nd</sup> ed. Part 2. Cowling, British Herbal Medicine Association 1979.

7 Westferry Circus • Canary Wharf • London E14 4HB • United Kingdom  
**Telephone** +44 (0)20 7418 8400 **Facsimile** +44 (0)20 7523 7051  
**E-mail** info@ema.europa.eu **Website** www.ema.europa.eu

An agency of the European Union



- British Herbal Pharmacopoeia. British Herbal Medicine Association, Bournemouth 1983; 128-129.
- Bruneton J. Pharmacognosie: Phytochimie Plantes Médicinales. Ed. Tec & Doc, Paris 1999; 529-530.
- Buchbauer G, Jirovetz L, Jäger W, Dietrich H, Plank C, Karamat E. Aromatherapy: evidence for sedative effects of the essential oil of lavender after inhalation. *Z Naturforsch* 1991; 46c: 1067-1072.
- Buchbauer G, Jirovetz L, Jäger W, Plank C, Dietrich H. Fragrance compounds and essential oils with sedative effects upon inhalation. *J Pharm Sci* 1993a; 82: 660-664.
- Buchbauer G, Jäger W, Jirovetz L, Limberger J, Dietrich H. Therapeutic properties of essential oils and fragrances. In: Teranishi R, Buttery RG, Sugisama H, editors. Bioactive Volatile Compounds from Plants. AC Symposium Series 525. Washington DC: American Chemical Society, 1993b.
- Büyükkuroglu ME, Gepdiremen A, Hacimuftüoğlu A, Oktay M. The effects of aqueous extract of *Lavandula angustifolia* flowers in glutamate-induced neurotoxicity of cerebellar granular cell culture of rat pups. *J Ethnopharmacol* 2003; 84: 91-94.
- Coulson IH, Khan ASA. Facial 'pillows' dermatitis due to lavender oil allergy. *Contact Dermatitis* 1999; 41: 111.
- Dale A, Cornwell S. The role of lavender oil in relieving perineal discomfort following childbirth: a blind randomized clinical trial. *J Advanced Nurs* 1994; 19: 89-96.
- Delaveau P, Guillemain J, Narcisse G, Rousseau A. The Neurodepressive Properties of Essential Oil of Lavender. *C R Seances Soc Biol Ses Fil* 1989; 183(4): 342-48.
- Diego MA, Jones NA, Filed T, Hernandez-Reif M, Schanberg S, Kuhn C, et al. Aromatherapy positively affects mood, EEG patterns of alertness and math computations. *Int J Neurosci* 1998; 96: 217-224.
- Dodonaeus R. Herbarius oft Cruydt-Boeck. Plantijnsche Druckerije, Antwerpen 1608; 470-473.
- Duan X, Tashiro M, Wu D, Yambe T, et al. Autonomic nervous function and localization of cerebral activity during lavender aromatic immersion. *Technol Health Care* 2007; 15(2): 69-78.
- Dunn C, Sleep J, Collett D. Sensing an improvement: an experimental study to evaluate the use of aromatherapy, massage and periods of rest in an intensive care unit. *J Advanced Nursing* 1995; 21: 34-40.
- Eder E, Neudecker T, Lutz D, Henschler D. Mutagenic potential of allyl and allylic compounds. Structure activity relationship as determined by alkylating and direct in vitro mutagenic properties. *Biochem Pharmacol* 1980; 29: 993-998.
- Eder E, Henschler D, Neudecker T. Mutagenic properties of allylic and alpha,beta-unsaturated compounds: consideration of alkylating mechanisms. *Xenobiotica* 1982a; 12: 831-848.
- Eder E, Neudecker T, Lutz D, Henschler D. Correlation of alkylating and mutagenic activities of allyl and allylic compounds: standard alkylation test vs. kinetic investigation. *Chem-Biol Interact* 1982b; 38: 303-315.
- Elhajili M, Baddouri K, Elkabbaj S, Meiouat F, Settaf A. Effet diurétique de l'infusion de fleurs de *Lavandula officinalis*. *Reprod Nutr Dev* 2001; 41: 393-399.
- ESCOP Monographs 2<sup>nd</sup> ed. *Lavandulae Flos*, *Lavandulae Aetheroleum*. European Scientific Cooperative on Phytotherapy, editor. Supplement 2009. Georg Thieme Verlag, Stuttgart 2009; 147-156.
- European Pharmacopoeia 6.0 (2008a) Lavender floweroil. Council of Europe. 1/2008: 1534.
- European Pharmacopoeia 6.0 (2008b) Lavender oil. Council of Europe. 1/2008: 1338.

- Fernandez M, Hernandez Reif M, Field T, Diego M, Sanders C, Roca A. EEG during lavender and rosemary exposure in infants of depressed and non-depressed mothers. *Infant Behavior & Development* 2004; 27(1): 91-100.
- Fujii M, Hatakeyama R, Fukukoa Y, et al. Lavender aromatherapy for behaviour and psychological symptoms in dementia patients. *Geriatr Gerontol Int* 2008; 8: 136-138.
- Gedney JJ, Glover TL, Fillingim RB. Sensory and affective pain discrimination after inhalation of essential oils. *Psychosomatic Med* 2004; 66: 599-606.
- Ghelardini C, Galeotti N, Salvatore G, Mazzanti G. Local anaesthetic activity of essential oil of *Lavandula angustifolia*. *Planta Medica* 1999; 65: 700-703.
- Guillemain J, Rousseau A, Delaveau P. Effets neurodépresseurs de l'huile essentielle de *Lavandula angustifolia* Mill *Ann Pharmaceutiques Françaises* 1989; 47(6): 337-343.
- Hänsel R, Keller K, Rimpler H, Schneider G. Hager's Handbuch der Pharmazeutischen Praxis. Springer Verlag, Berlin 1993; 630-644.
- Hardy M, Kirk Smith MD, Stretch DD. Replacement of drug treatment for insomnia by ambient odor. *Lancet* 1995; 346: 701.
- Hausen BM, Vieluf IK. *Lavandula angustifolia* Miller subsp. *angustifolia*. Echte Lavendel. In: Allergiepflanzen – Handbuch und Atlas. Kontaktallergene, Allergische Frühreaktionen. 2<sup>nd</sup> ed. Nikol. Verlagsgesellschaft, Hamburg 1997; 175-177.
- Hawrelak JA, Cattley T, Myers SP. Essential oils in the treatment of intestinal dysbiosis: a preliminary in vitro study. *Altern Med Rev* 2009; 14(4): 380-4.
- Henley DV, Lipson N, Korach KS, Bloch CA. Prepubertal gynecomastia linked to lavender and tea tree oils. *N Engl J Med* 2007; 356(5): 479-485.
- Holmes C, Hopkins V, Hensford C, MacLaughlin V, Wilkinson D, Rosenvinge H. Lavender oil as a treatment for agitated behaviour in severe dementia: a placebo controlled study. *Int J Geriatric Psychiatry* 2002; 17: 305-308.
- Hudson R. The value of lavender for rest and activity in the elderly patient. *Complement Ther Med* 1996; 4(1): 52-7.
- Hwang JH. The effects of the inhalation method using essential oils on blood pressure and stress responses of clients with essential hypertension. *Taehan Kanho Hakhoe Chi* 2006; 36(7): 1123-34.
- Ishidate M, Sofuni T, Yoshikawa K, Hayashi M, Nohmi T, Sawada M, Matsuoka A. Primary mutagenicity screening of food additives currently used in Japan. *Food Chem Toxicol* 1984; 22: 623-636.
- Jäger W, Buchbauer G, Jirovetz L, Fritzer M. Percutaneous absorption of lavender oil from a massage oil. *J Soc Cosmet Chem* 1992; 43: 49-54.
- Jirovetz L, Buchbauer G, Jäger W, Raverdinov V, Nikiforov A. Determination of lavender oil fragrance compounds in blood samples. *Fresenius J Anal Chem* 1990; 338: 922-923.
- Kane FMA, Brodie EE, Coull A, Coyne I, Howd A, Milne A, et al. The analgesic effect of odor and music upon dressing change. *Br J Nurs* 2004; 13(19): 4-12.
- Kawakami K, Takai-Kawakami K, Okazaki Y, Kurihara H, Shimizu Y, Yanaihara T. The effect of odors on human newborn infants under stress. *Infant Behavior & Development* 1997; 20(4): 531-535.

- Kim HM, Cho SH. Lavender oil inhibits immediate-type allergic reaction in mice and rats. *J Pharm Pharmacol* 1999; 55: 221-226.
- Kim JT, Ren CJ, Fielding GA, Pitti A, Kasumi T, Wajda M, et al. Treatment with lavender aromatherapy in the post-anesthesia care unit reduces opioid requirements of morbidly obese patients undergoing laparoscopic adjustable gastric banding. *Obes Surg* 2007; 17: 920-925.
- Kim JT, Wajda M, Cuff G, Serota D, Schlame M, Axelrod DM, Guth AA, Bekker AY. Evaluation of aromatherapy in treating postoperative pain: pilot study. *Pain Practice: the Official Journal of World Institute of Pain* 2006; 6(4): 273-7.
- Koto R, Imamura M, Watanabe C, Obayashi S, Shiraishi Mi, Sasaki Y, Azuma H. Linalyl acetate as a major ingredient of lavender essential oil relaxes the rabbit vascular smooth muscle through dephosphorylation of myosin light chain. *J Cardiovasc Pharmacol* 2006; 48(1): 850-6.
- Kritsidima M, Newton T, Asimakopoulou K. The effects of lavender scent on dental patient anxiety levels: a cluster randomized-controlled trial. *Community Dent Oral Epidemiol* 2010; 38(1): 83-87.
- Kubo I, Muroi H, Kubo A. Antibacterial activity of long-chain alcohols against *Streptococcus mutans*. *J Agric Food Chem* 1993; 41: 2447-2450.
- Leclerc H. Précis de phytothérapie. Masson Eds, Paris 1966 ; 204-205.
- Lee IS, Lee GJ. Effects of lavender aromatherapy on insomnia and depression in women college students. *Taehan Kanho Hakhoe Chi* 2006; 36(1): 136-43.
- Lis-Balchin M, Hart S. A preliminary study of the effect of essential oils on skeletal and smooth muscle in vitro. *J Ethnopharmacol* 1997; 58: 183-187.
- Lis-Balchin M, Hart S. Studies on the mode of action of the essential oil of lavender (*Lavandula angustifolia* P. Miller). *Phytotherapy Research* 1999; 13: 540-542.
- Louis M, Kowalski SD. Use of aromatherapy with hospice patients to decrease pain, anxiety, and depression and to promote an increasing sense of well-being. *Am J Hospice Palliative Care* 2002; 19: 381-386.
- Madaus Lehrbuch der biologischen Heilmittel.  
<http://www.henriettesherbal.com/eclectic/madaus/lavandula.html> (last access on 4 January 2011).
- Maddocks-Jennings W. Critical incident: idiosyncratic allergic reactions to essential oils. *Complement Ther Nurs Midwifery* 2004; 10: 58-60.
- Masago R, Matsuda T, Kikuchi Y, Miyazaki Y, Iwanaga K, Harada H, Katsuura T. Effects of inhalation of essential oils on EEG activity and sensory evaluation. *J Physiol Antropol* 2000; 19: 35-42.
- Morris N. The effects of lavender (*Lavendula angustifolium*) baths on psychological well-being: two exploratory randomised control trials. *Complementary Therapies in Medicine* 2002; 10(4): 223-228.
- Moss M, Cook J, Wesnes K, Duckett P. Aromas of rosemary and lavender essential oils differentially affect cognition and mood in healthy adults. *Int J Neurosci* 2003; 113: 15-38.
- Motomura N, Sakurai A, Yotsuya Y. Reduction of mental stress with lavender odorant. *Percept Motor Skills* 2001; 93: 713-718.
- Muzzarelli L, Force M, Sebold M. Aromatherapy and reducing preprocedural anxiety: A controlled prospective study. *Gastroenterol Nurs* 2006; 29(6): 466-71.

Nord D, Belew J. Effectiveness of the essential oils lavender and ginger in promotion children's comfort in a perianesthesia setting. *J Perianesth Nurs* 2009; 24(5): 307-12.

Opdyke DLJ. Monographs on fragrance raw materials: Lavender oil. *Food Cosmet Toxicol* 1976; 14: 451.

Parke DV, Rahman KMQ, Walker R. Effect of linalool on hepatic drug-metabolising enzymes in the rat. *Biochem Soc Trans* 1974; 2: 615-618.

Peana AT, D'Aquila PS, Panin F, Serra G, Pippia P, Moretti MDL. Anti-inflammatory activity of linalool and linalyl acetate constituents of essential oils. *Phytomedicine* 2002; 9: 721-726.

Peana AT, D'Aquila PS, Chessa ML, Moretti MDL, Serra G, Pippia P. (-)-Linalool produces antinociception in two experimental models of pain. *Eur J Pharmacology* 2003; 460: 37-41.

Peana AT, De Montis MG, Sechi S, Sircana G, D'Aquila PS, Pippia P. Effects of (-)-linalool in the acute hyperalgesia induced by carrageenan, L-glutamate and prostaglandin E. *Eur J Pharmacol* 2004a; 497: 279-284.

Peana AT, De Montis MG, Nieddu E, Spano MT, D'Aquila PS, Pippia P. Profile of spinal and supraspinal antinociception of (-)-linalool. *Eur J Pharmacol* 2004b; 485: 165-174.

Pulla Reddy AC, Lokesh BR. Studies on anti-inflammatory activity of spice principles and dietary n-3-polyunsaturated fatty acids on carrageenan-induced inflammation in rats. *Ann Nutr Metabol* 1994; 38: 349-358.

Rademaker M. Allergic contact dermatitis from lavender fragrance in Difflam® gel. *Contact Derm* 1994; 31: 58.

Rho KH, Han SH, Kim KS, Lee MS. Effects of aromatherapy massage on anxiety and self-esteem in Korean elderly women: a pilot study. *The International Journal of Neuroscience* 2006; 116(12): 1447-55.

Roffey SJ, Walker R, Gibson GG. Hepatic peroxisomal and microsomal enzyme induction by citral and linalool in rats. *Food Chem Toxic* 1990; 28: 403-408.

Romine IJ, Bush AM, Geist CR. Lavender aromatherapy in recovery from exercise. *Percept Motor Skills* 1999; 88: 756-758.

Saeki Y. The effects of footbath with or without the essential oil of lavender on the autonomic nervous system: a randomized trial. *Complement Ther Med* 2000; 8: 2-7.

Sanders C, Diego M, Fernandez M, Field T, Hernandez-Reil M, Roca A. EEG asymmetry responses to lavender and rosemary aromas in adults and infants. *Int J Neurosci* 2002; 112: 305-320.

Schulz H, Jobert M, Huebner WD. The quantitative EEG as a screening instrument to identify sedative effects of single doses of plant extracts in comparison with diazepam. *Phytomedicine* 1998; 5(6): 449-58.

Shaw D, Annett JM, Doherty B, Leslie JC. Anxiolytic affects of lavender oil inhalation on open-field behavior in rats. *Phytomedicine* 2007; 14(9): 613-20.

Shimizu K, Gyokusen M, Kitamura S, Kawabe T, Kozaki T, Ishibashi K, et al. Essential oil of lavender inhibited decreased attention during a long-term task in humans. *Biosci Biotechnol Biochem* 2008; 72: 1944-1947.

Snow AL. A controlled trial of aromatherapy for agitation in nursing home patients with dementia. *Journal of Alternative and Complementary Medicine* 2004; 10(3): 431-437.

- Sweetman S. Martindale the Complete Drug Reference. 36<sup>th</sup> ed. Pharmaceutical Press, London 2009; 2331.
- Tašev T, Taléva P, Balabanova V. Effet neuro-psychique des huiles essentielles bulgares de rose, de lavande et de géranium. *Folia Medica* 1969; XI(V): 307-316.
- Umezū T, Nagano K, Ito H, Kosakai K, Sakaniwa M, Morita M. Anticonflict effects of lavender oil and identification of its active constituents. *Pharmacology Biochemistry and Behavior* 2006; 85(4): 713-21.
- Valnet J. L'Aromathérapie. Maloine S.A. Editeur (since 1964), Vigot, Paris 2001; 272-278.
- von Skramlik E. Über die Giftigkeit und Verdrächlichkeit von ätherischen Ölen. *Pharmazie* 1959; 14: 435-445.
- Weiss RF, Fintelmann V. Lehrbuch der Phytotherapie. Hippokrates Verlag, Stuttgart 1999; 298.
- Williams TI. Evaluating effects of aromatherapy massage on sleep in children with autism: a pilot study. *Evidence-based Complementary and Alternative Medicine* 2006; 3(3): 373-7.
- Woelk H, Schläfke S. A multi-center, double-blind, randomized study of the lavender oil preparation Silexan in comparison to lorazepam for generalized anxiety. *Phytomedicine* 2010; 17(2): 94-9.
- Yamada K, Nimaki Y, Sashida Y. Anticonvulsive effects of inhaling lavender oil vapour. *Biol Pharm Bulletin* 1994; 17: 359-360.
- Yasukawa K, Takido M, Takeuchi M, Nakagawa S. Effect of chemical constituents from plants on 12-O-tetradecanoylphorbol-13-acetate induced inflammation in mice. *Chem Pharm Bull* 1989; 37: 1071-1073.
- Yip YB, Tse SHM. The effectiveness of relaxation acupoint stimulation and acupressure with aromatic lavender essential oil for non-specific low back pain in Hong Kong: a randomized controlled trial. *Complement Ther Med* 2004; 12: 28-37.
- Yip YB, Tse SHM. An experimental study on the effectiveness of acupressure with aromatic lavender essential oil for sub-acute, non-specific neck pain in Hong Kong. *Complementary Therapies in Clinical Practice* 2006; 12(1): 18-26.