

| EU (MA) Number | Invented Name | Strength | Pharmaceutical Form | Target species | Route of administration | Packaging | Content | Package size | Withdrawal period |
|-----------------------|----------------------|-----------------|----------------------------|-------------------------|--|----------------------------|----------------|----------------------|--|
| EU/2/11/134/001 | Inflacam | 1.5 mg/ml | Oral suspension | Dogs | Oral use | Bottle (HDPE) syringe (PP) | 15 ml | 1 bottle + 1 syringe | Not applicable |
| EU/2/11/134/002 | Inflacam | 1.5 mg/ml | Oral suspension | Dogs | Oral use | Bottle (PET) syringe (PP) | 42 ml | 1 bottle + 1 syringe | Not applicable |
| EU/2/11/134/003 | Inflacam | 1.5 mg/ml | Oral suspension | Dogs | Oral use | Bottle (PET) syringe (PP) | 100 ml | 1 bottle + 1 syringe | Not applicable |
| EU/2/11/134/004 | Inflacam | 1.5 mg/ml | Oral suspension | Dogs | Oral use | Bottle (PET) syringe (PP) | 200 ml | 1 bottle + 1 syringe | Not applicable |
| EU/2/11/134/005 | Inflacam | 20 mg/ml | Solution for injection | Cattle, pigs and horses | Subcutaneous use, intramuscular use or intravenous use | Vial (glass) | 20 ml | 1 vial | Meat and offal: 15 days (Cattle), Milk: 5 days (Cattle), Meat and offal: 5 days. (Horses) Meat and offal: 5 days (Pigs) |
| EU/2/11/134/006 | Inflacam | 20 mg/ml | Solution for injection | Cattle, pigs and horses | Subcutaneous use, intramuscular use or intravenous use | Vial (glass) | 50 ml | 1 vial | Meat and offal: 15 days (Cattle), Milk: 5 days (Cattle), Meat and offal: 5 days. (Horses) Meat and offal: 5 days (Pigs) |
| EU/2/11/134/007 | Inflacam | 20 mg/ml | Solution for injection | Cattle, pigs and horses | Subcutaneous use, intramuscular use or intravenous use | Vial (glass) | 100 ml | 1 vial | Meat and offal: 15 days (Cattle), Milk: 5 days (Cattle), Meat and offal: 5 days. (Horses) Meat and offal: 5 days (Pigs) |

| | | | | | | | | | |
|-----------------|----------|----------|------------------------|-------------------------|---|----------------------------|--------|-----------------------------|---|
| EU/2/11/134/008 | Inflacam | 20 mg/ml | Solution for injection | Cattle, pigs and horses | Subcutaneous use, intramuscular use or intravenous use | Vial (glass) | 250 ml | 1 vial | Meat and offal: 15 days (Cattle), Milk: 5 days (Cattle), Meat and offal: 5 days. (Horses) Meat and offal: 5 days (Pigs) |
| EU/2/11/134/009 | Inflacam | 15 mg/ml | Oral suspension | Horses | Oral use | Bottle (HDPE) syringe (PP) | 100 ml | 1 bottle + 1 syringe | Meat and offal: 3 days |
| EU/2/11/134/010 | Inflacam | 15 mg/ml | Oral suspension | Horses | Oral use | Bottle (HDPE) syringe (PP) | 250 ml | 1 bottle + 1 syringe | Meat and offal: 3 days |
| EU/2/11/134/011 | Inflacam | 1 mg | Chewable tablet | Dogs | Oral use | Blister (PVC/PVDC) | | 20 x 1 tablets (Unit dose) | Not applicable |
| EU/2/11/134/012 | Inflacam | 1 mg | Chewable tablet | Dogs | Oral use | Blister (PVC/PVDC) | | 100 x 1 tablets (Unit dose) | Not applicable |
| EU/2/11/134/013 | Inflacam | 2.5 mg | Chewable tablet | Dogs | Oral | Blister (PVC/PVDC) | | 20 x 1 tablets (Unit dose) | Not applicable |
| EU/2/11/134/014 | Inflacam | 2.5 mg | Chewable tablet | Dogs | Oral | Blister (PVC/PVDC) | | 100 x 1 tablets (Unit dose) | Not applicable |
| EU/2/11/134/015 | Inflacam | 5 mg/ml | Solution for injection | Dogs and cats | Dogs: intravenous or subcutaneous Cats: subcutaneous | Vial (glass) | 10 ml | 1 vial | Not applicable |
| EU/2/11/134/016 | Inflacam | 5 mg/ml | Solution for injection | Dog and cats | Dogs: intravenous or subcutaneous Cats: subcutaneous | Vial (glass) | 20 ml | 1 vial | Not applicable |
| EU/2/11/134/017 | Inflacam | 5 mg/ml | Solution for injection | Dogs and cats | Dogs: intravenous or subcutaneous Cats: subcutaneous | Vial (glass) | 100 ml | 1 vial | Not applicable |

| | | | | | | | | | |
|-----------------|----------|-----------|------------------------|-----------------|--|--------------------------------|--------|----------------------|--|
| EU/2/11/134/018 | Inflacam | 5 mg/ml | Solution for injection | Cattle and pigs | Cattle: intravenous or subcutaneous Pigs: intramuscular | Vial (glass) | 20 ml | 1 vial | Meat and offal: 15 days (Cattle), Meat and offal: 5 days (Pigs) |
| EU/2/11/134/019 | Inflacam | 5 mg/ml | Solution for injection | Cattle and pigs | Cattle: intravenous or subcutaneous Pigs: intramuscular | Vial (glass) | 50 ml | 1 vial | Meat and offal: 15 days (Cattle), Meat and offal: 5 days (Pigs) |
| EU/2/11/134/020 | Inflacam | 5 mg/ml | Solution for injection | Cattle and pigs | Cattle: intravenous or subcutaneous Pigs: intramuscular | Vial (glass) | 100 ml | 1 vial | Meat and offal: 15 days (Cattle), Meat and offal: 5 days (Pigs) |
| EU/2/11/134/021 | Inflacam | 330 mg | Granules in sachet | Horses | In-feed use | Sachet (paper/PE/Alu/PE) | | 100 sachets | Meat and offal: 3 days |
| EU/2/11/134/022 | Inflacam | 330 mg | Granules in sachet | Horses | In-feed use | Sachet (paper/PE/Alu/PE) | | 20 sachets | Meat and offal: 3 days |
| EU/2/11/134/023 | Inflacam | 0.5 mg/ml | Oral suspension | Cats | Oral use | Bottle (HDPE), Syringe (PP) | 10 ml | 1 bottle + 1 syringe | Not applicable |
| EU/2/11/134/024 | Inflacam | 0.5 mg/ml | Oral suspension | Cats | Oral use | Bottle (HDPE), Syringe (PP) | 15 ml | 1 bottle + 1 syringe | Not applicable |
| EU/2/11/134/025 | Inflacam | 0.5 mg/ml | Oral suspension | Cats | Oral use | Bottle (PP), Syringe (PP) | 3 ml | 1 bottle + 1 syringe | Not applicable |
| EU/2/11/134/026 | Inflacam | 0.5 mg/ml | Oral suspension | Cats | Oral use | Bottle (PP), Syringe (PP) | 5 ml | 1 bottle + 1 syringe | Not applicable |