

European Medicines Agency Evaluation of Medicines for Human Use

> London, 5 July 2007 Doc. Ref. EMEA/HMPC/179283/2007

COMMITTEE ON HERBAL MEDICINAL PRODUCTS (HMPC)

DRAFT

COMMUNITY LIST ENTRY ON CALENDULA OFFICINALIS L., FLOS

DISCUSSION IN WORKING PARTY ON COMMUNITY MONOGRAPHS AND COMMUNITY LIST (MLWP)	July 2007
ADOPTION BY HMPC FOR RELEASE FOR CONSULTATION	5 July 2007
END OF CONSULTATION (DEADLINE FOR COMMENTS)	15 October 2007
REDISCUSSION IN WORKING PARTY ON COMMUNITY MONOGRAPHS AND COMMUNITY LIST (MLWP)	
ADOPTION BY HMPC	

Comments should be provided using this <u>template</u> to <u>hmpc.secretariat@emea.europa.eu</u> Fax: +44 20 75 23 70 51

KEYWORDS	Herbal medicinal products; HMPC; Community list of herbal substances,	
	preparations and combinations thereof for use in traditional herbal	
	medicinal products; traditional use; Calendula officinalis L.; Calendulae	
	flos; calendula flower	

Page 1/4

ENTRY TO LIST OF HERBAL SUBSTANCES, PREPARATIONS AND COMBINATIONS THEREOF FOR USE IN TRADITIONAL HERBAL MEDICINAL PRODUCTS:

CALENDULA OFFICINALIS L., FLOS

Scientific name of the plant

Calendula officinalis L., flos

Botanical family

Asteraceae

Herbal substance

Calendula flower

Common name in all EU official languages of herbal substance

BG (bălgarski): CS (čeština): měsíčkový květ DA (dansk): DE (Deutsch): Ringelblumenblüten EL (elliniká): EN (English): calendula flower ES (espanol): Caléndula, flor de ET (eesti keel): saialilleõisik FI (suomi): kehäkukka FR (français): Souci HU (magyar): IT (italiano): LT (lietuvių kalba): LV (latviešu valoda): MT (malti): NL (nederlands): Goudsbloem PL (polski): kwiat nagietka PT (português): RO (română): SK (slovenčina): Nechtíkový kvet SL (slovenščina): cvet vrtnega ognjiča SV (svenska):

IS (*íslenska*): morgunfrú,blóm NO (norsk):

Herbal preparation(s)

A) Liquid extract (DER 1:1, ethanol 40-50% v/v)

B) Tincture (DER 1:5, ethanol 70-90% v/v)

C) Liquid extract (DER 1:10, fatty vegetable oil, e.g. olive oil)

D) Ointment (DER 1:5 - 1:25, hardened vegetable fat, petroleum jelly¹)

E) Comminuted herbal substance for infusion

European Pharmacopoeia monograph reference

Calendula flower - Calendulae flos 01/2005:1297

¹ Calendula ointment is prepared by gentle digestion of the herbal substance in the melted ointment base for up to 16 hours and subsequent filtration and congealment during fall in temperature

Indication(s)

- a) Traditional herbal medicinal product for the symptomatic treatment of minor inflammations of the skin (such as sunburn) and as an aid in healing of minor wounds.
- b) Traditional herbal medicinal product for the symptomatic treatment of minor inflammations in the mouth or the throat.

The product is a traditional herbal medicinal product for use in specified indications exclusively based upon long-standing use.

Type of tradition

European

Specified strength

Herbal substance:

Infusion for topical application: 1-2 g / 150 ml of water. The still warm infusion is used as such to rinse or gargle for the treatment of inflammations in the mouth or the throat or to prepare compresses.

Herbal preparations:

- A) Liquid extract (DER 1:1, ethanol 40-50% v/v)
- In semi-solid dosage forms: amount equivalent to 2-10% herbal substance
- B) Tincture

In compresses diluted at least 1:3 with freshly boiled water; in semi-solid dosage forms: amount equivalent to 2-10% herbal substance

- C) Liquid extract (DER 1:10, fatty vegetable oil e.g. olive oil) In semi-solid dosage forms: amount equivalent to 2-8% herbal substance
- D) Ointment
 - Equivalent to 4-20% herbal substance

Specified posology

Please see 'Specified strength'

Indication a)

The use is not recommended in children under 6 years of age (see below Special warnings and precautions for use).

Indication b)

The use is not recommended in children under 12 years of age (see below Special warnings and precautions for use).

Route of administration

Topical application, 2 to 4 times daily.

Duration of use or any restrictions on the duration of use

Compresses: remove after 30-60 minutes

If the symptoms persist after 1 week during the use of the medicinal product a doctor or a qualified health care practitioner should be consulted.

Any other information necessary for the safe use

Contraindications

Hypersensitivity to members of the Asteraceae family (Compositae family).

Special warnings and precautions for use

Indication a)

The use in children under 6 years of age is not recommended because there is no experience available.

Indication b)

The use in children under 12 years of age is not recommended because there is no experience available.

If signs of skin infection are observed, medical advice should be sought.

Interactions with other medicinal products and other forms of interaction None reported.

Pregnancy and lactation

Safety during pregnancy and lactation has not been established. In the absence of sufficient data, the use during pregnancy and lactation is not recommended.

Effects on ability to drive and use machines

Not relevant.

Undesirable effects

Skin sensitization. The frequency is not known.

If other adverse reactions not mentioned above occur, a doctor or a qualified health care practitioner should be consulted.

Overdose None reported.

Pharmaceutical particulars [If necessary] Not applicable.

Pharmacological effects or efficacy plausible on the basis of long-standing use and experience [**If necessary for the safe use of the product**] Not applicable.