

Medicinal product no longer authorised

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

BTVPUR AlSap 1 suspension for injection for sheep and cattle.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each dose of 1 ml contains:

Active substances:

Inactivated Bluetongue Virus Serotype 1..... $\geq 1.9 \log_{10}$ pixels*

(*) Antigen content (VP2 protein) by immuno-assay

Adjuvants:

. Al³⁺ (as hydroxide) 2.7 mg

. Saponin 30 HU**

(**) Haemolytic units

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Homogeneous milky white suspension for injection.

4. CLINICAL PARTICULARS

4.1 Target species

Sheep and cattle

4.2 Indications for use, specifying the target species

Active immunisation of sheep and cattle to prevent viraemia* and to reduce clinical signs caused by bluetongue virus serotype 1.

*(below the level of detection by the validated RT-PCR method at $3.68 \log_{10}$ RNA copies/ml, indicating no infectious virus transmission)

Onset of immunity has been demonstrated 3 weeks after the primary vaccination course.

The duration of immunity for cattle and sheep is 1 year after the primary vaccination course.

4.3 Contraindications

None.

4.4 Special warnings for each target species

If used in other domestic and wild ruminant species that are considered at risk of infection, its use in these species should be undertaken with care and it is advisable to test the vaccine on a small number of animals prior to mass vaccination. The level of efficacy for other species may differ from that observed in sheep and cattle.

4.5 Special precautions for use

Special precautions for use in animals

Vaccinate healthy animals only.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

None

4.6 Adverse reactions (frequency and seriousness)

In very rare cases it has been observed a small local swelling at the injection site (at most 32 cm² in cattle and 24 cm² in sheep) which becomes residual 35 days later (≤ 1 cm²).

In very rare cases it has been observed a transient increase in body temperature, normally not exceeding an average of 1.1 °C, may occur within 24 hours after vaccination.

The frequency of adverse reactions is defined using the following convention:

- Very common (more than 1 in 10 animals displaying adverse reaction(s) during the course of one treatment).
- Common (more than 1 but less than 10 animals in 100 animals).
- Uncommon (more than 1 but less than 10 animals in 1,000 animals).
- Rare (more than 1 but less than 10 animals in 10,000 animals).
- Very rare (less than 1 animal in 10,000 animals, including isolated reports).

4.7 Use during pregnancy, lactation or lay

Can be used during pregnancy in ewes. Can be used during pregnancy and lactation in cows. The safety and the efficacy of the vaccine have not been established in breeding males. In this category of animals the vaccine should be used only according to the benefit/risk assessment by the responsible veterinarian and/or national Competent Authorities on the current vaccination policies against Bluetongue Virus (BTV).

4.8 Interaction with other medicinal products and other forms of interaction

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

4.9 Amounts to be administered and administration route

Apply usual aseptic procedures.

Shake gently immediately before use. Avoid bubble formation, as this can be irritating at the site of injection. The entire content of the bottle should be used immediately after broaching and during the same procedure. Avoid multiple vial broaching.

Administer one dose of 1 ml subcutaneously according to the following vaccination scheme:

• Primary vaccination

In sheep

- First injection: from 1 month of age in naive animals (or from 2.5 months of age in young animals born to immune sheep).
- Second injection: after 3-4 weeks

In cattle

- First injection: from 1 month of age in naive animals (or from 2.5 months of age in young animals born to immune cattle).

- Second injection: after 3-4 weeks

- **Revaccination**

Annual.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Very rare and transient apathy can be observed after the administration of a double-dose of the vaccine. No other adverse reactions except those mentioned in section 4.6 were observed.

4.11 Withdrawal period

Zero days.

5. IMMUNOLOGICAL PROPERTIES

Pharmacotherapeutic group: bluetongue virus vaccine,
ATCvet code QI04AA02 (sheep) and QI02AA08 (cattle).

The vaccine contains inactivated Bluetongue Virus Serotype 1 with aluminium hydroxide and saponin adjuvants. It induces an active and specific immunity against bluetongue virus serotype 1 in the vaccinated animal.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Aluminium hydroxide
Saponin
Silicon antifoam
Phosphate buffer
Glycine buffer

6.2 Incompatibilities

Do not mix with any other veterinary medicinal product.

6.3 Shelf life

Shelf life of the veterinary medicinal product as package for sale (100 ml bottle): 2 years
Shelf life of the veterinary medicinal product as package for sale (50 ml bottle): 2 years
Shelf life of the veterinary medicinal product as package for sale (10 ml bottle): 18 months
Shelf life after first opening the immediate packaging: use immediately

6.4 Special precautions for storage

Store and transport refrigerated (2 °C-8 °C)
Do not freeze
Protect from light

6.5 Nature and composition of immediate packaging

Polypropylene bottle of 50 or 100 ml with butyl elastomere closure
Box of 1 bottle of 100 doses (1 x 100 ml)
Box of 10 bottles of 100 doses (10 x 100 ml)

Box of 1 bottle of 50 doses (1 x 50 ml)
Box of 10 bottles of 50 doses (10 x 50 ml)

Type I glass bottle of 10 ml with butyl elastomere closure
Box of 1 bottle of 10 doses (1 x 10 ml)

6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

MERIAL
29, avenue Tony Garnier
69007 Lyon
FRANCE

8. MARKETING AUTHORISATION NUMBER(S)

EU/2/10/112/001
EU/2/10/112/002
EU/2/10/112/003
EU/2/10/112/004
EU/2/10/112/005

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of the first authorisation: 17/12/2010
Date of the last renewal:

10. DATE OF REVISION OF THE TEXT

Detailed information of this product is available on the website of the European Medicines Agency
<http://www.ema.europa.eu/>

PROHIBITION OF SALE, SUPPLY AND/OR USE

The manufacture, import, possession sale, supply and/or use of BTVPUR AlSap 1 may be prohibited in a Member State on the whole or part of their territory pursuant to national legislation. Any person intending to manufacture, import, possess, sell, supply and/or use BTVPUR AlSap 1 must consult the relevant Member State's competent authority on the current vaccination policies prior to the manufacture, import, possession, sale, supply and/or use.

ANNEX II

- A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND
MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY OR USE**
- C. STATEMENT OF THE MRLs**

**A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND
MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer of the biological active substance

MERIAL Animal Health Limited
Biological Laboratory, Ash Road,
Pirbright, Woking, Surrey GU24 0NQ
United Kingdom

MERIAL Laboratoire de Lyon Gerland
254, rue Marcel Mérieux
69342 LYON CEDEX 07
France

Name and address of the manufacturer responsible for batch release

MERIAL
Laboratory of Lyon Porte des Alpes
Rue de l'Aviation,
69800 Saint-Priest
France

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY OR USE

Veterinary medicinal product subject to prescription.

According to Article 71 of Directive 2001/82/EC of the European Parliament and of the Council as amended, a Member State may, in accordance with its national legislation, prohibit the manufacture, import, possession, sale, supply and/or use of immunological veterinary medicinal products on the whole or part of its territory if it is established that:

- a) the administration of the product to animals will interfere with the implementation of a national programme for the diagnosis, control or eradication of animal diseases, or will cause difficulties in certifying the absence of contamination in live animals or in foodstuffs or other products obtained from treated animals.
- b) the disease to which the product is intended to confer immunity is largely absent from the territory in question.

The use of this veterinary medicinal product is only allowed under the particular conditions established by European Community legislation on the control of Bluetongue.

C. STATEMENT OF THE MRLs

The active substance being a principle of biological origin intended to produce active immunity is not within the scope of Regulation (EC) No 470/2009.

The excipients (including adjuvants) listed in section 6.1 of the SPC are either allowed substances for which table 1 of the annex to Commission Regulation (EU) No 37/2010 indicates that no MRLs are required.

Medicinal product no longer authorised

ANNEX III
LABELLING AND PACKAGE LEAFLET

Medicinal product no longer authorised

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGE

**Box of 1 bottle of 10 ml,
Box of 1 bottle of 50 ml,
Box of 10 bottles of 50 ml,
Box of 1 bottle of 100 ml,
Box of 10 bottles of 100 ml**

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

BTVPUR AlSap 1 suspension for injection for sheep and cattle

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Each dose of 1 ml contains:

Inactivated BTV1 $\geq 1.9 \log_{10}$ pixels*

Aluminium hydroxide, Saponin, qs 1 dose (*)

(*) see package leaflet

3. PHARMACEUTICAL FORM

Suspension for injection

4. PACKAGE SIZE

1 bottle of 10 doses (1 x 10 ml)

1 bottle of 50 doses (1 x 50 ml)

10 bottles of 50 doses (10 x 50 ml)

1 bottle of 100 doses (1 x 100 ml)

10 bottles of 100 doses (10 x 100 ml)

5. TARGET SPECIES

Sheep and cattle

6. INDICATION(S)**7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Subcutaneous use

Read the package leaflet before use.

8. WITHDRAWAL PERIOD

Withdrawal period: zero days

9. SPECIAL WARNING(S), IF NECESSARY**10. EXPIRY DATE**

EXP {month/year}

Once broached, use immediately.

11. SPECIAL STORAGE CONDITIONS

Store and transport refrigerated.

Do not freeze.

Protect from light.

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only. To be supplied only on veterinary prescription

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

MERIAL
29, avenue Tony Garnier
69007 Lyon
FRANCE

16. MARKETING AUTHORISATION NUMBER(S)

EU/2/10/112/005
EU/2/10/112/003
EU/2/10/112/004
EU/2/10/112/001
EU/2/10/112/002

17. MANUFACTURER'S BATCH NUMBER
--

Lot {number}

Medicinal product no longer authorised

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**Bottle of 10 and 50 ml****1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

BTVPUR AlSap 1 suspension for injection for sheep and cattle

2. QUANTITY OF THE ACTIVE SUBSTANCE(S)Inactivated BTV 1 $\geq 1,9 \log_{10}$ pixels**3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

10 doses (10 ml)

50 doses (50 ml)

4. ROUTE(S) OF ADMINISTRATION

SC

5. WITHDRAWAL PERIOD

Withdrawal period: zero days

6. BATCH NUMBER

Lot {number}

7. EXPIRY DATE

EXP {month/year}

Once broached, use immediately

8. THE WORDS “FOR ANIMAL TREATMENT ONLY”

For animal treatment only.

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE

Bottle of 100 ml

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

BTVPUR AlSap 1 suspension for injection for sheep and cattle

2. STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Each dose of 1 ml contains Bluetongue Virus

Inactivated BTV 1 $\geq 1.9 \log_{10}$ pixels

3. PHARMACEUTICAL FORM

Suspension for injection

4. PACKAGE SIZE

100 doses (100 ml)

5. TARGET SPECIES

Sheep and cattle

6. INDICATION(S)

7. METHOD AND ROUTE(S) OF ADMINISTRATION

SC

Read the package leaflet before use.

8. WITHDRAWAL PERIOD

Withdrawal period: zero days

9. SPECIAL WARNING(S), IF NECESSARY

10. EXPIRY DATE

EXP {month/year}

Once broached, use immediately

11. SPECIAL STORAGE CONDITIONS

Store and transport refrigerated.
Do not freeze
Protect from light.

12. SPECIFIC PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY

13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, if applicable

For animal treatment only. To be supplied only on veterinary prescription

14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”

Keep out of the sight and reach of children.

15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

MERIAL
29, avenue Tony Garnier
69007 Lyon
FRANCE

16. MARKETING AUTHORISATION NUMBER(S)

EU/2/10/112/001
EU/2/10/112/002

17. MANUFACTURER'S BATCH NUMBER

Lot {number}

Medicinal product no longer authorised

B. PACKAGE LEAFLET

PACKAGE LEAFLET FOR:
BTVPUR AlSap 1 suspension for injection for sheep and cattle

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder:

MERIAL

29 avenue Tony Garnier

69007 Lyon,

France

Manufacturer responsible for batch release:

MERIAL

Laboratory of Lyon Porte des Alpes

Rue de l'Aviation,

69800 Saint-Priest

France

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

BTVPUR AlSap 1 suspension for injection for sheep and cattle

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each dose of 1 ml of vaccine (homogeneous milky white suspension) contains:

Active substance:

.Inactivated Bluetongue Virus Serotype 1 $\geq 1.9 \log_{10}$ pixels *

(*) Antigen content (VP2 protein) by immuno-assay

Adjuvants:

.Al3+ (as hydroxide) 2.7 mg

.Saponin 30 HU**

(**) Haemolytic units

4. INDICATION(S)

Active immunisation of sheep and cattle to prevent viraemia* and to reduce clinical signs caused by Bluetongue Virus Serotype 1.

*(below the level of detection by the validated RT-PCR method at $3.68 \log_{10}$ RNA copies/ml, indicating no infectious virus transmission).

Onset of immunity has been demonstrated 3 weeks after the primary vaccination course.

The duration of immunity for cattle and sheep is 1 year after the primary vaccination course.

5. CONTRAINDICATIONS

None.

6. ADVERSE REACTIONS

In very rare cases it has been observed a small local swelling at the injection site (at most 32 cm² in cattle and 24 cm² in sheep) which becomes residual 35 days later (≤ 1 cm²).

In very rare cases it has been observed a transient increase in body temperature, normally not exceeding an average of 1.1 °C, may occur within 24 hours after vaccination.

The frequency of adverse reactions is defined using the following convention:

- Very common (more than 1 in 10 animals displaying adverse reaction(s) during the course of one treatment).
- Common (more than 1 but less than 10 animals in 100 animals).
- Uncommon (more than 1 but less than 10 animals in 1,000 animals).
- Rare (more than 1 but less than 10 animals in 10,000 animals).
- Very rare (less than 1 animal in 10,000 animals, including isolated reports).

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

7. TARGET SPECIES

Sheep and cattle.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Administer one dose of 1 ml subcutaneously according to the following vaccination scheme:

• Primary vaccination

In sheep

- 1st injection: from 1 month of age in naive animals (or from 2.5 months of age in young animals born to immune sheep).
- 2nd injection: after 3-4 weeks

In cattle

- 1st injection: from 1 month of age in naive animals (or from 2.5 months of age in young animals born to immune cattle).
- 2nd injection: after 3-4 weeks.

• Revaccination

Annual.

9. ADVICE ON CORRECT ADMINISTRATION

Apply usual aseptic procedures.

Shake gently immediately before use. Avoid bubble formation, as this can be irritating at the site of injection. The entire content of the bottle should be used immediately after broaching and during the same procedure. Avoid multiple vial broaching.

10. WITHDRAWAL PERIOD

Zero days.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Store and transport refrigerated (2 °C-8 °C).

Do not freeze.

Protect from light.

Shelf life after first opening the immediate packaging: immediately

Do not use this veterinary medicinal product after the expiry date which is stated on the carton after EXP:

12. SPECIAL WARNINGS

Special warnings for each target species:

If used in other domestic and wild ruminant species that are considered at risk of infection, its use in these species should be undertaken with care and it is advisable to test the vaccine on a small number of animals prior to mass vaccination. The level of efficacy for other species may differ from that observed in sheep and cattle.

Special precautions for use in animals:

Vaccinate healthy animals only.

Pregnancy and lactation:

Can be used during pregnancy in ewes. Can be used during pregnancy and lactation in cows

Fertility:

The safety and the efficacy of the vaccine have not been established in breeding males. In this category of animals the vaccine should be used only according to the benefit/risk assessment by the responsible veterinarian and/ or national Competent Authorities on the current vaccination policies against Bluetongue Virus (BTV).

Interaction with other medicinal products and other forms of interaction:

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be decided on a case by case basis.

Overdose (symptoms, emergency procedures, antidotes):

Very rare and transient apathy can be observed after the administration of a double-dose of the vaccine. No other adverse reactions except those mentioned in section 'Adverse Reactions' were observed.

Incompatibilities:

Do not mix with any other veterinary medicinal product.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Medicines should not be disposed of via wastewater or household waste.

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

Detailed information on this product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>.

15. OTHER INFORMATION

The vaccine contains inactivated Bluetongue Virus Serotype 1 with aluminium hydroxide and saponin adjuvants. It induces an active and specific immunity against Bluetongue Virus Serotype 1 in the vaccinated animal.

Not all pack sizes may be marketed

Box of 1 bottle of 10 doses (1 x 10 ml)

Box of 1 bottle of 50 doses (1 x 50 ml)

Box of 10 bottles of 50 doses (10 x 50 ml)

Box of 1 bottle of 100 doses (1 x 100 ml)

Box of 10 bottles of 100 doses (10 x 100 ml)

The manufacture, import, possession, sale, supply and/or use of BTVPUR AlSap 1 may be prohibited in a certain Member State on the whole or part of their territory pursuant to national legislation. Any person intending to manufacture, import, possess, sell, supply and/or use BTVPUR AlSap 1 must consult the relevant Member State's competent authority on the current vaccination policies prior to the manufacture, import, possession, sale, supply and/or use.