

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Deferasirox Mylan 90 mg film-coated tablets  
Deferasirox Mylan 180 mg film-coated tablets  
Deferasirox Mylan 360 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

### Deferasirox Mylan 90 mg film-coated tablets

Each film-coated tablet contains 90 mg deferasirox.

### Deferasirox Mylan 180 mg film-coated tablets

Each film-coated tablet contains 180 mg deferasirox.

### Deferasirox Mylan 360 mg film-coated tablets

Each film-coated tablet contains 360 mg deferasirox.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablets (Tablets)

### Deferasirox Mylan 90 mg film-coated tablets

A blue, film-coated, modified capsule shaped, biconvex tablet debossed with “ $\cap$ ” on one side of the tablet and ‘DF’ on the other side.

Approximate tablet dimensions 10.00 mm × 4.5 mm.

### Deferasirox Mylan 180 mg film-coated tablets

A blue, film-coated, modified capsule shaped, biconvex tablet debossed with “ $\cap$ ” on one side of the tablet and ‘DF 1’ on the other side.

Approximate tablet dimensions 12.8 mm × 6.00 mm.

### Deferasirox Mylan 360 mg film-coated tablets

A blue, film-coated, modified capsule shaped, biconvex tablet debossed with “ $\cap$ ” on one side of the tablet and ‘DF 2’ on the other side.

Approximate tablet dimensions 17 mm × 6.7 mm.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Deferasirox Mylan is indicated for the treatment of chronic iron overload due to frequent blood transfusions ( $\geq 7$  ml/kg/month of packed red blood cells) in patients with beta thalassaemia major aged 6 years and older.

Deferasirox Mylan is also indicated for the treatment of chronic iron overload due to blood transfusions when deferoxamine therapy is contraindicated or inadequate in the following patient groups:

- in paediatric patients with beta thalassaemia major with iron overload due to frequent blood transfusions ( $\geq 7$  ml/kg/month of packed red blood cells) aged 2 to 5 years,
- in adult and paediatric patients with beta thalassaemia major with iron overload due to infrequent blood transfusions ( $< 7$  ml/kg/month of packed red blood cells) aged 2 years and older,
- in adult and paediatric patients with other anaemias aged 2 years and older.

Deferasirox Mylan is also indicated for the treatment of chronic iron overload requiring chelation therapy when deferoxamine therapy is contraindicated or inadequate in patients with non-transfusion-dependent thalassaemia syndromes aged 10 years and older.

## **4.2 Posology and method of administration**

All references to the dispersible tablet formulation throughout the SmPC refer to medicinal products by different Marketing Authorisation Holders with dispersible tablet formulations of the active ingredient deferasirox.

Treatment with Deferasirox Mylan should be initiated and maintained by physicians experienced in the treatment of chronic iron overload.

### Posology

#### Transfusional iron overload

It is recommended that treatment be started after the transfusion of approximately 20 units (about 100 ml/kg) of packed red blood cells (PRBC) or when there is evidence from clinical monitoring that chronic iron overload is present (e.g. serum ferritin  $> 1,000$   $\mu\text{g/l}$ ). Doses (in mg/kg) must be calculated and rounded to the nearest whole tablet size.

The goals of iron chelation therapy are to remove the amount of iron administered in transfusions and, as required, to reduce the existing iron burden.

Caution should be taken during chelation therapy to minimise the risk of over-chelation in all patients (see section 4.4).

In the EU, medicines containing deferasirox are available as film-coated tablets and dispersible tablets marketed under different tradenames as generic alternatives of deferasirox. Due to different pharmacokinetic profiles, a 30% lower dose of deferasirox film-coated tablets is needed in comparison to the recommended dose for deferasirox dispersible tablets (see section 5.1).

**Table 1** Recommended doses for transfusional iron overload

	<b>Film-coated tablets</b>	<b>Transfusions</b>	<b>Serum ferritin</b>
<b>Starting dose</b>	<b>14 mg/kg/day</b>	After 20 units (about 100 ml/kg) of PRBC	or >1,000µg/l
<b>Alternative starting doses</b>	21 mg/kg/day	>14 ml/kg/month of PRBC (approx. >4 units/month for an adult)	
	7 mg/kg/day	<7 ml/kg/month of PRBC (approx. <2 units/month for an adult)	
For patients well managed on deferoxamine	One third of deferoxamine dose		
<b>Monitoring</b>			<b>Monthly</b>
<b>Target range</b>			<b>500 – 1,000 µg/l</b>
<b>Adjustment Steps</b>	<b>Increase</b>		
(every 3 – 6 months)	3.5 – 7 mg/kg/day		>2,500 µg/l
	Up to 28 mg/kg/day		
	<b>Decrease</b>		
	3.5 – 7 mg/kg/day		<2,500 µg/l
	In patients treated with doses >21 mg/kg/day		
	– When target is reached		500 – 1,000 µg/l
<b>Maximum dose</b>	<b>28 mg/kg/day</b>		
<b>Consider interruption</b>			<b>&lt;500 µg/l</b>

*Starting dose*

The recommended initial daily dose of Deferasirox Mylan film-coated tablets is 14 mg/kg body weight.

An initial daily dose of 21 mg/kg may be considered for patients who require reduction of elevated body iron levels and who are also receiving more than 14 ml/kg/month of packed red blood cells (approximately >4 units/month for an adult).

An initial daily dose of 7 mg/kg may be considered for patients who do not require reduction of body iron levels and who are also receiving less than 7 ml/kg/month of packed red blood cells (approximately <2 units/month for an adult). The patient’s response must be monitored and a dose increase should be considered if sufficient efficacy is not obtained (see section 5.1).

For patients, already well managed on treatment with deferoxamine, a starting dose of Deferasirox Mylan film-coated tablets that is numerically one third that of the deferoxamine dose could be considered (e.g. a patient receiving 40 mg/kg/day of deferoxamine for 5 days per week (or equivalent) could be transferred to a starting daily dose of 14 mg/kg/day of Deferasirox Mylan film-coated tablets). When this results in a daily dose less than 14 mg/kg body weight, the patient’s response must be monitored and a dose increase should be considered if sufficient efficacy is not obtained (see section 5.1).

### *Dose adjustment*

It is recommended that serum ferritin be monitored every month and that the dose of Deferasirox be adjusted, if necessary, every 3 to 6 months based on the trends in serum ferritin. Dose adjustments may be made in steps of 3.5 to 7 mg/kg and are to be tailored to the individual patient's response and therapeutic goals (maintenance or reduction of iron burden). In patients not adequately controlled with doses of 21 mg/kg (e.g. serum ferritin levels persistently above 2,500 µg/l and not showing a decreasing trend over time), doses of up to 28 mg/kg may be considered. The availability of long-term efficacy and safety data from clinical studies conducted with deferasirox dispersible tablets used at doses above 30 mg/kg is currently limited (264 patients followed for an average of 1 year after dose escalation). If only very poor haemosiderosis control is achieved at doses up to 21 mg/kg (film-coated tablet dose equivalent of 30 mg/kg of dispersible tablets), a further increase (to a maximum of 28 mg/kg) may not achieve satisfactory control, and alternative treatment options may be considered. If no satisfactory control is achieved at doses above 21 mg/kg, treatment at such doses should not be maintained and alternative treatment options should be considered whenever possible. Doses above 28 mg/kg are not recommended because there is only limited experience with doses above this level (see section 5.1).

In patients treated with doses greater than 21 mg/kg, dose reductions in steps of 3.5 to 7 mg/kg should be considered when control has been achieved (e.g. serum ferritin levels persistently  $\leq$  2,500 µg/l and showing a decreasing trend over time). In patients whose serum ferritin level has reached the target (usually between 500 and 1,000 µg/l), dose reductions in steps of 3.5 to 7 mg/kg should be considered to maintain serum ferritin levels within the target range and to minimise the risk of over-chelation. If serum ferritin falls consistently below 500 µg/l, an interruption of treatment should be considered (see section 4.4).

### *Non-transfusion-dependent thalassaemia syndromes*

Chelation therapy should only be initiated when there is evidence of iron overload (liver iron concentration [LIC]  $\geq$  5 mg Fe/g dry weight [dw] or serum ferritin consistently  $>$  800 µg/l). LIC is the preferred method of iron overload determination and should be used wherever available. Caution should be taken during chelation therapy to minimise the risk of over-chelation in all patients (see section 4.4).

In the EU, medicines containing deferasirox are available as film-coated tablets and dispersible tablets marketed under different tradenames as generic alternatives of deferasirox. Due to different pharmacokinetic profiles, a 30% lower dose of deferasirox film-coated tablets is needed in comparison to the recommended dose for deferasirox dispersible tablets (see section 5.1).

Table 2 Recommended doses for non-transfusion-dependent thalassaemia syndromes

	<b>Film-coated tablets</b>	<b>Liver iron concentration (LIC)*</b>	<b>Serum ferritin</b>
<b>Starting dose</b>	<b>7 mg/kg/day</b>	$\geq$ 5 mg Fe/g dw	or $>$ 800 µg/l
<b>Monitoring</b>			<b>Monthly</b>
<b>Adjustment steps</b> (every 3 – 6 months)	<b>Increase</b> 3.5 – 7 mg/kg/day	$\geq$ 7 mg Fe/g dw	or $>$ 2,000 µg/l
	<b>Decrease</b> 3.5 – 7 mg/kg/day	$<$ 7 mg Fe/g dw	or $\leq$ 2,000 µg/l
<b>Maximum dose</b>	<b>14 mg/kg/day</b> For adult patients		
	<b>7 mg/kg/day</b> For paediatric patients		
	<b>7 mg/kg/day</b>		

	For both adult and paediatric patients	not assessed	and $\leq 2,000 \mu\text{g/l}$
<b>Interruption</b>		<b>&lt;3 mg Fe/g dw</b>	or <b>&lt;300 <math>\mu\text{g/l}</math></b>
<b>Retreatment</b>		<b>Not recommended</b>	

\* LIC is the preferred method of iron overload determination.

### *Starting dose*

The recommended initial daily dose of Deferasirox Mylan film-coated tablets in patients with non-transfusion-dependent thalassaemia syndromes is 7 mg/kg body weight.

### *Dose adjustment*

It is recommended that serum ferritin be monitored every month to assess the patient's response to therapy and to minimise the risk of over-chelation (see section 4.4). After every 3 to 6 months of treatment, a dose increase in increments of 3.5 to 7 mg/kg should be considered if the patient's LIC is  $\geq 7$  mg Fe/g dw, or if serum ferritin is consistently  $>2,000$   $\mu\text{g/l}$  and not showing a downward trend, and the patient is tolerating the medicinal product well. Doses above 14 mg/kg are not recommended because there is no experience with doses above this level in patients with non-transfusion-dependent thalassaemia syndromes.

In both paediatric and adult patients in whom LIC was not assessed and serum ferritin is  $\leq 2,000$   $\mu\text{g/l}$ , dosing should not exceed 7 mg/kg.

For patients in whom the dose was increased to  $>7$  mg/kg, dose reduction to 7 mg/kg or less is recommended when LIC is  $<7$  mg Fe/g dw or serum ferritin is  $\leq 2,000$   $\mu\text{g/l}$ .

### *Treatment cessation*

Once a satisfactory body iron level has been achieved (LIC  $<3$  mg Fe/g dw or serum ferritin  $<300$   $\mu\text{g/l}$ ), treatment should be stopped. There are no data available on the retreatment of patients who reaccumulate iron after having achieved a satisfactory body iron level and therefore retreatment cannot be recommended.

### *Special populations*

#### *Elderly patients ( $\geq 65$ years of age)*

The dosing recommendations for elderly patients are the same as described above. In clinical studies, elderly patients experienced a higher frequency of adverse reactions than younger patients (in particular, diarrhoea) and should be monitored closely for adverse reactions that may require a dose adjustment.

#### *Paediatric population*

##### Transfusional iron overload:

The dosing recommendations for paediatric patients aged 2 to 17 years with transfusional iron overload are the same as for adult patients (see section 4.2). It is recommended that serum ferritin be monitored every month to assess the patient's response to therapy and to minimise the risk of over-chelation (see section 4.4). Changes in weight of paediatric patients over time must be taken into account when calculating the dose.

In children with transfusional iron overload aged between 2 and 5 years, exposure is lower than in adults (see section 5.2). This age group may therefore require higher doses than are necessary in adults. However, the initial dose should be the same as in adults, followed by individual titration.

##### Non-transfusion-dependent thalassaemia syndromes:

In paediatric patients with non-transfusion-dependent thalassaemia syndromes, dosing should not exceed 7 mg/kg. In these patients, closer monitoring of LIC and serum ferritin is essential to avoid over-chelation (see section 4.4). In addition to monthly serum ferritin assessments, LIC should be monitored every three months when serum ferritin is  $\leq 800$   $\mu\text{g/l}$ .

##### Children from birth to 23 months:

The safety and efficacy of Deferasirox Mylan in children from birth to 23 months of age have not been established. No data are available.

#### *Patients with renal impairment*

Deferasirox Mylan has not been studied in patients with renal impairment and is contraindicated in patients with estimated creatinine clearance <60 ml/min (see sections 4.3 and 4.4).

#### *Patients with hepatic impairment*

Deferasirox Mylan is not recommended in patients with severe hepatic impairment (Child-Pugh Class C). In patients with moderate hepatic impairment (Child-Pugh Class B), the dose should be considerably reduced followed by progressive increase up to a limit of 50% of recommended treatment dose for patients with normal hepatic function (see sections 4.4 and 5.2), and Deferasirox Mylan must be used with caution in such patients. Hepatic function in all patients should be monitored before treatment, every 2 weeks during the first month and then every month (see section 4.4).

#### Method of administration

For oral use.

The film-coated tablets should be swallowed whole with some water. For patients who are unable to swallow whole tablets, the film-coated tablets may be crushed and administered by sprinkling the full dose onto soft food, e.g. yogurt or apple sauce (pureed apple). The dose should be immediately and completely consumed, and not stored for future use.

The film-coated tablets should be taken once a day, preferably at the same time each day, and may be taken on an empty stomach or with a light meal (see sections 4.5 and 5.2).

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1. Combination with other iron chelator therapies as the safety of such combinations has not been established (see section 4.5).

Patients with estimated creatinine clearance <60 ml/min.

### **4.4 Special warnings and precautions for use**

#### Renal function

Deferasirox has been studied only in patients with baseline serum creatinine within the age-appropriate normal range.

During clinical studies, increases in serum creatinine of >33% on  $\geq 2$  consecutive occasions, sometimes above the upper limit of the normal range, occurred in about 36% of patients. These were dose-dependent. About two-thirds of the patients showing serum creatinine increase returned below the 33% level without dose adjustment. In the remaining third the serum creatinine increase did not always respond to a dose reduction or a dose interruption. In some cases, only a stabilisation of the serum creatinine values has been observed after dose reduction. Cases of acute renal failure have been reported following post-marketing use of deferasirox (see section 4.8). In some post-marketing cases, renal function deterioration has led to renal failure requiring temporary or permanent dialysis.

The causes of the rises in serum creatinine have not been elucidated. Particular attention should therefore be paid to monitoring of serum creatinine in patients who are concomitantly receiving medicinal products that depress renal function, and in patients who are receiving high doses of deferasirox and/or low rates of transfusion (<7 ml/kg/month of packed red blood cells or <2 units/month for an adult). While no increase in renal adverse events was observed after dose escalation of deferasirox dispersible tablets to doses above 30 mg/kg in clinical studies, an increased risk of renal adverse events with film-coated tablet doses above 21 mg/kg cannot be excluded.



It is recommended that serum creatinine be assessed in duplicate before initiating therapy. **Serum creatinine, creatinine clearance** (estimated with the Cockcroft-Gault or MDRD formula in adults and with the Schwartz formula in children) and/or plasma cystatin C levels **should be monitored prior to therapy, weekly in the first month after initiation or modification of therapy with deferasirox (including switch of formulation), and monthly thereafter.** Patients with pre-existing renal conditions and patients who are receiving medicinal products that depress renal function may be more at risk of complications. Care should be taken to maintain adequate hydration in patients who develop diarrhoea or vomiting.

There have been post-marketing reports of metabolic acidosis occurring during treatment with deferasirox. The majority of these patients had renal impairment, renal tubulopathy (Fanconi syndrome) or diarrhoea, or conditions where acid-base imbalance is a known complication. Acid-base balance should be monitored as clinically indicated in these populations. Interruption of deferasirox therapy should be considered in patients who develop metabolic acidosis.

Post-marketing cases of severe forms of renal tubulopathy (such as Fanconi syndrome) and renal failure associated with changes in consciousness in the context of hyperammonaemic encephalopathy have been reported in patients treated with deferasirox, mainly in children. It is recommended that hyperammonaemic encephalopathy be considered and ammonia levels measured in patients who develop unexplained changes in mental status while on Deferasirox Mylan therapy.

**Table 3** Dose adjustment and interruption of treatment for renal monitoring

	<b>Serum creatinine</b>		<b>Creatinine clearance</b>
<b>Before initiation of therapy</b>	Twice (2×)	and	Once (1×)
<b>Contraindicated</b>			<b>&lt;60 ml/min</b>
<b>Monitoring</b>			
– First month after start of therapy or dose modification (including switch of formulation)	Weekly	and	Weekly
– Thereafter	Monthly	and	Monthly
<b>Reduction of daily dose by 7 mg/kg/day</b> (film-coated tablet formulation), <i>if following renal parameters are observed at two consecutive visits and cannot be attributed to other causes</i>			
Adult patients	>33% above pre-treatment average	and	Decreases <LLN* (<90 ml/min)
Paediatric patients	> age appropriate ULN**	and/or	Decreases <LLN* (<90 ml/min)
<b>After dose reduction, interrupt treatment, if</b>			
Adult and paediatric	Remains >33% above pre-treatment average	and/or	Decreases <LLN* (<90 ml/min)
* LLN: lower limit of the normal range			
** ULN: upper limit of the normal range			

Treatment may be reinitiated depending on the individual clinical circumstances.

Dose reduction or interruption may be also considered if abnormalities occur in levels of markers of renal tubular function and/or as clinically indicated:

- Proteinuria (test should be performed prior to therapy and monthly thereafter)
- Glycosuria in non-diabetics and low levels of serum potassium, phosphate, magnesium or urate, phosphaturia, aminoaciduria (monitor as needed).

Renal tubulopathy has been mainly reported in children and adolescents with beta-thalassaemia treated with Deferasirox Mylan.

Patients should be referred to a renal specialist, and further specialised investigations (such as renal biopsy) may be considered if the following occur despite dose reduction and interruption:

- Serum creatinine remains significantly elevated and
- Persistent abnormality in another marker of renal function (e.g. proteinuria, Fanconi Syndrome).

### Hepatic function

Liver function test elevations have been observed in patients treated with deferasirox. Post-marketing cases of hepatic failure, sometimes fatal, have been reported in patients treated with deferasirox. Severe forms associated with changes in consciousness in the context of hyperammonaemic encephalopathy, may occur in patients treated with deferasirox, particularly in children. It is recommended that hyperammonaemic encephalopathy be considered and ammonia levels measured in patients who develop unexplained changes in mental status while on Deferasirox Mylan. Care should be taken to maintain adequate hydration in patients who experience volume-depleting events (such as diarrhoea or vomiting), particularly in children with acute illness. Most reports of hepatic failure involved patients with significant comorbidities including pre-existing chronic liver conditions (including cirrhosis and hepatitis C) and multi-organ failure. The role of deferasirox as a contributing or aggravating factor cannot be excluded (see section 4.8).

It is recommended that serum transaminases, bilirubin and alkaline phosphatase be checked before the initiation of treatment, every 2 weeks during the first month and monthly thereafter. If there is a persistent and progressive increase in serum transaminase levels that cannot be attributed to other causes, deferasirox should be interrupted. Once the cause of the liver function test abnormalities has been clarified or after return to normal levels, cautious re-initiation of treatment at a lower dose followed by gradual dose escalation may be considered.

Deferasirox Mylan is not recommended in patients with severe hepatic impairment (Child-Pugh Class C) (see section 5.2).

<b>Test</b>	<b>Frequency</b>
Serum creatinine	In duplicate prior to therapy. Weekly during first month of therapy or after dose modification (including switch of formulation). Monthly thereafter.
Creatinine clearance and/or plasma cystatin C	Prior to therapy. Weekly during first month of therapy or after dose modification (including switch of formulation). Monthly thereafter.
Proteinuria	Prior to therapy. Monthly thereafter.
Other markers of renal tubular function (such as glycosuria in non-diabetics and low levels of serum potassium, phosphate, magnesium or urate, phosphaturia, aminoaciduria)	As needed.
Serum transaminases, bilirubin, alkaline phosphatase	Prior to therapy. Every 2 weeks during first month of therapy. Monthly thereafter.
Auditory and ophthalmic testing	Prior to therapy. Annually thereafter.
Body weight, height and sexual development	Prior to therapy. Annually in paediatric patients.

In patients with a short life expectancy (e.g. high-risk myelodysplastic syndromes), especially when co-morbidities could increase the risk of adverse events, the benefit of Deferasirox Mylan might be limited and may be inferior to risks. As a consequence, treatment with Deferasirox Mylan is not recommended in these patients.

Caution should be used in elderly patients due to a higher frequency of adverse reactions (in particular, diarrhoea).

Data in children with non-transfusion-dependent thalassaemia are very limited (see section 5.1). As a consequence, Deferasirox Mylan therapy should be closely monitored to detect adverse reactions and to follow iron burden in the paediatric population. In addition, before treating heavily iron-overloaded children with non-transfusion-dependent thalassaemia with Deferasirox Mylan, the physician should be aware that the consequences of long-term exposure in such patients are currently not known.

#### Gastrointestinal disorders

Upper gastrointestinal ulceration and haemorrhage have been reported in patients, including children and adolescents, receiving deferferasirox. Multiple ulcers have been observed in some patients (see section 4.8). There have been reports of ulcers complicated with digestive perforation. Also, there have been reports of fatal gastrointestinal haemorrhages, especially in elderly patients who had haematological malignancies and/or low platelet counts. Physicians and patients should remain alert for signs and symptoms of gastrointestinal ulceration and haemorrhage during Deferasirox Mylan therapy. In case of gastrointestinal ulceration or haemorrhage, Deferasirox Mylan should be discontinued and additional evaluation and treatment must be promptly initiated. Caution should be exercised in patients who are taking Deferasirox Mylan in combination with substances that have known ulcerogenic potential, such as NSAIDs, corticosteroids, or oral bisphosphonates, in patients

receiving anticoagulants and in patients with platelet counts below 50,000/mm<sup>3</sup> ( $50 \times 10^9/l$ ) (see section 4.5).

### Skin disorders

Skin rashes may appear during Deferasirox Mylan treatment. The rashes resolve spontaneously in most cases. When interruption of treatment may be necessary, treatment may be reintroduced after resolution of the rash, at a lower dose followed by gradual dose escalation. In severe cases, this reintroduction could be conducted in combination with a short period of oral steroid administration. Severe cutaneous adverse reactions (SCARs) including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS), which could be life-threatening or fatal, have been reported. If any SCAR is suspected, Deferasirox Mylan should be discontinued immediately and should not be reintroduced. At the time of prescription, patients should be advised of the signs and symptoms of severe skin reactions, and be closely monitored.

### Hypersensitivity reactions

Cases of serious hypersensitivity reactions (such as anaphylaxis and angioedema) have been reported in patients receiving deferasirox, with the onset of the reaction occurring in the majority of cases within the first month of treatment (see section 4.8). If such reactions occur, Deferasirox Mylan should be discontinued and appropriate medical intervention instituted. Deferasirox should not be reintroduced in patients who have experienced a hypersensitivity reaction due to the risk of anaphylactic shock (see section 4.3).

### Vision and hearing

Auditory (decreased hearing) and ocular (lens opacities) disturbances have been reported (see section 4.8). Auditory and ophthalmic testing (including fundoscopy) is recommended before the start of treatment and at regular intervals thereafter (every 12 months). If disturbances are noted during the treatment, dose reduction or interruption may be considered.

### Blood disorders

There have been post-marketing reports of leukopenia, thrombocytopenia or pancytopenia (or aggravation of these cytopenias) and of aggravated anaemia in patients treated with deferasirox. Most of these patients had pre-existing haematological disorders that are frequently associated with bone marrow failure. However, a contributory or aggravating role cannot be excluded. Interruption of treatment should be considered in patients who develop unexplained cytopenia.

### Other considerations

Monthly monitoring of serum ferritin is recommended in order to assess the patient's response to therapy and to avoid over-chelation (see section 4.2). Dose reduction or closer monitoring of renal and hepatic function, and serum ferritin levels are recommended during periods of treatment with high doses and when serum ferritin levels are close to the target range. If serum ferritin falls consistently below 500 µg/l (in transfusional iron overload) or below 300 µg/l (in non-transfusion-dependent thalassaemia syndromes), an interruption of treatment should be considered.

The results of the tests for serum creatinine, serum ferritin and serum transaminases should be recorded and regularly assessed for trends.

In two clinical studies, growth and sexual development of paediatric patients treated with deferasirox for up to 5 years were not affected (see section 4.8). However, as a general precautionary measure in the management of paediatric patients with transfusional iron overload, body weight, height and sexual development should be monitored prior to therapy and at regular intervals (every 12 months).

Cardiac dysfunction is a known complication of severe iron overload. Cardiac function should be monitored in patients with severe iron overload during long-term treatment with Deferasirox Mylan.

#### Sodium content

Deferasirox Mylan contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

The safety of deferasirox in combination with other iron chelators has not been established. Therefore, it must not be combined with other iron chelator therapies (see section 4.3).

#### Interaction with food

The  $C_{max}$  of deferasirox film-coated tablets was increased (by 29%) when taken with a high-fat meal. Deferasirox Mylan film-coated tablets may be taken either on an empty stomach or with a light meal, preferably at the same time each day (see sections 4.2 and 5.2).

#### Agents that may decrease Deferasirox Mylan systemic exposure

Deferasirox metabolism depends on UGT enzymes. In a healthy volunteer study, the concomitant administration of deferasirox (single dose of 30 mg/kg, dispersible tablet formulation) and the potent UGT inducer, rifampicin, (repeated dose of 600 mg/day) resulted in a decrease of deferasirox exposure by 44% (90% CI: 37% – 51%). Therefore, the concomitant use of Deferasirox Mylan with potent UGT inducers (e.g. rifampicin, carbamazepine, phenytoin, phenobarbital, ritonavir) may result in a decrease in Deferasirox Mylan efficacy. The patient's serum ferritin should be monitored during and after the combination, and the dose of Deferasirox Mylan adjusted if necessary.

Cholestyramine significantly reduced the deferasirox exposure in a mechanistic study to determine the degree of enterohepatic recycling (see section 5.2).

#### Interaction with midazolam and other agents metabolised by CYP3A4

In a healthy volunteer study, the concomitant administration of deferasirox dispersible tablets and midazolam (a CYP3A4 probe substrate) resulted in a decrease of midazolam exposure by 17% (90% CI: 8% – 26%). In the clinical setting, this effect may be more pronounced. Therefore, due to a possible decrease in efficacy, caution should be exercised when deferasirox is combined with substances metabolised through CYP3A4 (e.g. ciclosporin, simvastatin, hormonal contraceptive agents, bepridil, ergotamine).

#### Interaction with repaglinide and other agents metabolised by CYP2C8

In a healthy volunteer study, the concomitant administration of deferasirox as a moderate CYP2C8 inhibitor (30 mg/kg daily, dispersible tablet formulation), with repaglinide, a CYP2C8 substrate, given as a single dose of 0.5 mg, increased repaglinide AUC and  $C_{max}$  about 2.3-fold (90% CI [2.03–2.63]) and 1.6-fold (90% CI [1.42–1.84]), respectively. Since the interaction has not been established with dosages higher than 0.5 mg for repaglinide, the concomitant use of deferasirox with repaglinide should be avoided. If the combination appears necessary, careful clinical and blood glucose monitoring should be performed (see section 4.4). An interaction between deferasirox and other CYP2C8 substrates like paclitaxel cannot be excluded.

#### Interaction with theophylline and other agents metabolised by CYP1A2

In a healthy volunteer study, the concomitant administration of deferasirox as a CYP1A2 inhibitor (repeated dose of 30 mg/kg/day, dispersible tablet formulation) and the CYP1A2 substrate theophylline (single dose of 120 mg) resulted in an increase of theophylline AUC by 84% (90% CI:

73% to 95%). The single dose  $C_{max}$  was not affected, but an increase of theophylline  $C_{max}$  is expected to occur with chronic dosing. Therefore, the concomitant use of deferasirox with theophylline is not recommended. If deferasirox and theophylline are used concomitantly, monitoring of theophylline concentration and theophylline dose reduction should be considered. An interaction between deferasirox and other CYP1A2 substrates cannot be excluded. For substances that are predominantly metabolised by CYP1A2 and that have a narrow therapeutic index (e.g. clozapine, tizanidine), the same recommendations apply as for theophylline.

#### Other information

The concomitant administration of deferasirox and aluminium-containing antacid preparations has not been formally studied. Although deferasirox has a lower affinity for aluminium than for iron, it is not recommended to take deferasirox tablets with aluminium-containing antacid preparations.

The concomitant administration of deferasirox with substances that have known ulcerogenic potential, such as NSAIDs (including acetylsalicylic acid at high dosage), corticosteroids or oral bisphosphonates may increase the risk of gastrointestinal toxicity (see section 4.4). The concomitant administration of deferasirox with anticoagulants may also increase the risk of gastrointestinal haemorrhage. Close clinical monitoring is required when deferasirox is combined with these substances.

Concomitant administration of deferasirox and busulfan resulted in an increase of busulfan exposure (AUC), but the mechanism of the interaction remains unclear. If possible, evaluation of the pharmacokinetics (AUC, clearance) of a busulfan test dose should be performed to allow dose adjustment.

### **4.6 Fertility, pregnancy and lactation**

#### Pregnancy

No clinical data on exposed pregnancies are available for deferasirox. Studies in animals have shown some reproductive toxicity at maternally toxic doses (see section 5.3). The potential risk for humans is unknown.

As a precaution, it is recommended that Deferasirox Mylan is not used during pregnancy unless clearly necessary.

Deferasirox Mylan may decrease the efficacy of hormonal contraceptives (see section 4.5). Women of childbearing potential are recommended to use additional or alternative non-hormonal methods of contraception when using Deferasirox Mylan.

#### Breast-feeding

In animal studies, deferasirox was found to be rapidly and extensively secreted into maternal milk. No effect on the offspring was noted. It is not known if deferasirox is secreted into human milk. Breast-feeding while taking Deferasirox Mylan is not recommended.

#### Fertility

No fertility data is available for humans. In animals, no adverse effects on male or female fertility were found (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

Deferasirox Mylan has minor influence on the ability to drive and use machines. Patients experiencing the uncommon adverse reaction of dizziness should exercise caution when driving or operating machines (see section 4.8).

## 4.8 Undesirable effects

### Summary of the safety profile

The most frequent reactions reported during chronic treatment in clinical studies conducted with deferasirox dispersible tablets in adult and paediatric patients include gastrointestinal disturbances (mainly nausea, vomiting, diarrhoea or abdominal pain) and skin rash. Diarrhoea is reported more commonly in paediatric patients aged 2 to 5 years and in the elderly. These reactions are dose-dependent, mostly mild to moderate, generally transient and mostly resolve even if treatment is continued.

During clinical studies dose-dependent increases in serum creatinine occurred in about 36% of patients, though most remained within the normal range. Decreases in mean creatinine clearance have been observed in both paediatric and adult patients with beta-thalassemia and iron overload during the first year of treatment, but there is evidence that this does not decrease further in subsequent years of treatment. Elevations of liver transaminases have been reported. Safety monitoring schedules for renal and liver parameters are recommended. Auditory (decreased hearing) and ocular (lens opacities) disturbances are uncommon, and yearly examinations are also recommended (see section 4.4).

Severe cutaneous adverse reactions (SCARs), including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported with the use of Deferasirox Mylan (see section 4.4).

### Tabulated list of adverse reactions

Adverse reactions are ranked below using the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Table 5

<b>Blood and lymphatic system disorders</b>	
Not known:	Pancytopenia <sup>1</sup> , thrombocytopenia <sup>1</sup> , anaemia aggravated <sup>1</sup> , neutropenia <sup>1</sup>
<b>Immune system disorders</b>	
Not known:	Hypersensitivity reactions (including anaphylactic reactions and angioedema) <sup>1</sup>
<b>Metabolism and nutrition disorders</b>	
Not known:	Metabolic acidosis <sup>1</sup>
<b>Psychiatric disorders</b>	
Uncommon:	Anxiety, sleep disorder
<b>Nervous system disorders</b>	
Common:	Headache
Uncommon:	Dizziness
<b>Eye disorders</b>	
Uncommon:	Cataract, maculopathy
Rare:	Optic neuritis
<b>Ear and labyrinth disorders</b>	
Uncommon:	Deafness
<b>Respiratory, thoracic and mediastinal disorders</b>	
Uncommon:	Laryngeal pain

<b>Gastrointestinal disorders</b>	
Common:	Diarrhoea, constipation, vomiting, nausea, abdominal pain, abdominal distension, dyspepsia
Uncommon:	Gastrointestinal haemorrhage, gastric ulcer (including multiple ulcers), duodenal ulcer, gastritis
Rare:	Oesophagitis
Not known:	Gastrointestinal perforation <sup>1</sup> , acute pancreatitis <sup>1</sup>
<b>Hepatobiliary disorders</b>	
Common:	Transaminases increased
Uncommon:	Hepatitis, cholelithiasis
Not known:	Hepatic failure <sup>1,2</sup>
<b>Skin and subcutaneous tissue disorders</b>	
Common:	Rash, pruritus
Uncommon:	Pigmentation disorder
Rare:	Drug reaction with eosinophilia and systemic symptoms (DRESS)
Not known:	Stevens-Johnson syndrome <sup>1</sup> , hypersensitivity vasculitis <sup>1</sup> , urticaria <sup>1</sup> , erythema multiforme <sup>1</sup> , alopecia <sup>1</sup> , toxic epidermal necrolysis (TEN) <sup>1</sup>
<b>Renal and urinary disorders</b>	
Very common:	Blood creatinine increased
Common:	Proteinuria
Uncommon:	Renal tubular disorder <sup>2</sup> (acquired Fanconi syndrome), glycosuria
Not known:	Acute renal failure <sup>1,2</sup> , tubulointerstitial nephritis <sup>1</sup> , nephrolithiasis <sup>1</sup> , renal tubular necrosis <sup>1</sup>
<b>General disorders and administration site conditions</b>	
Uncommon:	Pyrexia, oedema, fatigue

<sup>1</sup> Adverse reactions reported during post-marketing experience. These are derived from spontaneous reports for which it is not always possible to reliably establish frequency or a causal relationship to exposure to the medicinal product.

<sup>2</sup> Severe forms associated with changes in consciousness in the context of hyperammonaemic encephalopathy have been reported.

### Description of selected adverse reactions

Gallstones and related biliary disorders were reported in about 2% of patients. Elevations of liver transaminases were reported as an adverse reaction in 2% of patients. Elevations of transaminases greater than 10 times the upper limit of the normal range, suggestive of hepatitis, were uncommon (0.3%). During post-marketing experience, hepatic failure, sometimes fatal, has been reported with deferasirox (see section 4.4). There have been post-marketing reports of metabolic acidosis. The majority of these patients had renal impairment, renal tubulopathy (Fanconi syndrome) or diarrhoea, or conditions where acid-base imbalance is a known complication (see section 4.4). Cases of serious acute pancreatitis were observed without documented underlying biliary conditions. As with other iron chelator treatment, high-frequency hearing loss and lenticular opacities (early cataracts) have been uncommonly observed in patients treated with deferasirox (see section 4.4).

### Creatinine clearance in transfusional iron overload

In a retrospective meta-analysis of 2,102 adult and paediatric beta-thalassaemia patients with transfusional iron overload treated with deferasirox dispersible tablets in two randomised and four open label studies of up to five years' duration, a mean creatinine clearance decrease of 13.2% in adult patients (95% CI: -14.4% to -12.1%; n=935) and 9.9% (95% CI: -11.1% to -8.6%; n=1,142) in paediatric patients was observed during the first year of treatment. In 250 patients who were followed for up to five years, no further decrease in mean creatinine clearance levels was observed.



### Clinical study in patients with non-transfusion-dependent thalassaemia syndromes

In a 1-year study in patients with non-transfusion-dependent thalassaemia syndromes and iron overload (dispersible tablets at a dose of 10 mg/kg/day), diarrhoea (9.1%), rash (9.1%), and nausea (7.3%) were the most frequent study drug-related adverse events. Abnormal serum creatinine and creatinine clearance values were reported in 5.5% and 1.8% of patients, respectively. Elevations of liver transaminases greater than 2 times the baseline and 5 times the upper limit of normal were reported in 1.8% of patients.

### Paediatric population

In two clinical studies, growth and sexual development of paediatric patients treated with deferasirox for up to 5 years were not affected (see section 4.4).

Diarrhoea is reported more commonly in paediatric patients aged 2 to 5 years than in older patients.

Renal tubulopathy has been mainly reported in children and adolescents with beta-thalassaemia treated with deferasirox. In post-marketing reports, a high proportion of cases of metabolic acidosis occurred in children in the context of Fanconi syndrome.

Acute pancreatitis has been reported, particularly in children and adolescents.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via **the national reporting system** listed in [Appendix V](#).

## **4.9 Overdose**

Early signs of acute overdose are digestive effects such as abdominal pain, diarrhoea, nausea and vomiting. Hepatic and renal disorders have been reported, including cases of liver enzyme and creatinine increased with recovery after treatment discontinuation. An erroneously administered single dose of 90 mg/kg led to Fanconi syndrome which resolved after treatment.

There is no specific antidote for deferasirox. Standard procedures for management of overdose may be indicated as well as symptomatic treatment, as medically appropriate.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Iron chelating agents, ATC code: V03AC03

### Mechanism of action

Deferasirox is an orally active chelator that is highly selective for iron (III). It is a tridentate ligand that binds iron with high affinity in a 2:1 ratio. Deferasirox promotes excretion of iron, primarily in the faeces. Deferasirox has low affinity for zinc and copper, and does not cause constant low serum levels of these metals.

### Pharmacodynamic effects

In an iron-balance metabolic study in iron-overloaded adult thalassaemic patients, deferasirox at daily doses of 10, 20 and 40 mg/kg (dispersible tablet formulation) induced the mean net excretion of 0.119, 0.329 and 0.445 mg Fe/kg body weight/day, respectively.

## Clinical efficacy and safety

Clinical efficacy studies were conducted with deferasirox dispersible tablets. Compared to the deferasirox dispersible tablet formulation, the dose of the deferasirox film coated tablets is 30% lower than the dose of the deferasirox dispersible tablets, rounded to the nearest whole tablet (see section 5.2).

Deferasirox has been investigated in 411 adult (age  $\geq 16$  years) and 292 paediatric patients (aged 2 to  $< 16$  years) with chronic iron overload due to blood transfusions. Of the paediatric patients 52 were aged 2 to 5 years. The underlying conditions requiring transfusion included beta-thalassaemia, sickle cell disease and other congenital and acquired anaemias (myelodysplastic syndromes, Diamond-Blackfan syndrome, aplastic anaemia and other very rare anaemias).

Daily treatment with the deferasirox dispersible tablet formulation at doses of 20 and 30 mg/kg for one year in frequently transfused adult and paediatric patients with beta-thalassaemia led to reductions in indicators of total body iron; liver iron concentration was reduced by about  $-0.4$  and  $-8.9$  mg Fe/g liver (biopsy dry weight (dw)) on average, respectively, and serum ferritin was reduced by about  $-36$  and  $-926$   $\mu\text{g/l}$  on average, respectively. At these same doses, the ratios of iron excretion: iron intake were 1.02 (indicating net iron balance) and 1.67 (indicating net iron removal), respectively. Deferasirox induced similar responses in iron-overloaded patients with other anaemias. Daily doses of 10 mg/kg (dispersible tablet formulation) for one year could maintain liver iron and serum ferritin levels and induce net iron balance in patients receiving infrequent transfusions or exchange transfusions. Serum ferritin assessed by monthly monitoring reflected changes in liver iron concentration indicating that trends in serum ferritin can be used to monitor response to therapy. Limited clinical data (29 patients with normal cardiac function at baseline) using MRI indicate that treatment with deferasirox 10 – 30 mg/kg/day (dispersible tablet formulation) for 1 year may also reduce levels of iron in the heart (on average, MRI T2\* increased from 18.3 to 23.0 milliseconds).

The principal analysis of the pivotal comparative study in 586 patients suffering from beta-thalassaemia and transfusional iron overload did not demonstrate non-inferiority of deferasirox dispersible tablets to deferoxamine in the analysis of the total patient population. It appeared from a post-hoc analysis of this study that, in the subgroup of patients with liver iron concentration  $\geq 7$  mg Fe/g dw treated with deferasirox dispersible tablets (20 and 30 mg/kg) or deferoxamine (35 to  $\geq 50$  mg/kg), the non-inferiority criteria were achieved. However, in patients with liver iron concentration  $< 7$  mg Fe/g dw treated with deferasirox dispersible tablets (5 and 10 mg/kg) or deferoxamine (20 to 35 mg/kg), non-inferiority was not established due to imbalance in the dosing of the two chelators. This imbalance occurred because patients on deferoxamine were allowed to remain on their pre-study dose even if it was higher than the protocol specified dose. Fifty-six patients under the age of 6 years participated in this pivotal study, 28 of them receiving deferasirox dispersible tablets.

It appeared from preclinical and clinical studies that deferasirox dispersible tablets could be as active as deferoxamine when used in a dose ratio of 2:1 (i.e. a dose of deferasirox dispersible tablets that is numerically half of the deferoxamine dose). For deferasirox film-coated tablets, a dose ratio of 3:1 can be considered (i.e. a dose of deferasirox film-coated tablets that is numerically one third of the deferoxamine dose). However, this dosing recommendation was not prospectively assessed in the clinical studies.

In addition, in patients with liver iron concentration  $\geq 7$  mg Fe/g dw with various rare anaemias or sickle cell disease, deferasirox dispersible tablets up to 20 and 30 mg/kg produced a decrease in liver iron concentration and serum ferritin comparable to that obtained in patients with beta-thalassaemia.

A placebo-controlled randomised study was performed in 225 patients with MDS (Low/Int-1 risk) and transfusional iron overload. The results of this study suggest that there is a positive impact of deferasirox on event-free survival (EFS, a composite endpoint including non-fatal cardiac or liver events) and serum ferritin levels. The safety profile was consistent with previous studies in adult MDS patients.

In a 5-year observational study in which 267 children aged 2 to  $< 6$  years (at enrollment) with

transfusional haemosiderosis received deferasirox, there were no clinically meaningful differences in the safety and tolerability profile of deferasirox in paediatric patients aged 2 to <6 years compared to the overall adult and older paediatric population, including increases in serum creatinine of >33% and above the upper limit of normal on  $\geq 2$  consecutive occasions (3.1%), and elevation of alanine aminotransferase (ALT) greater than 5 times the upper limit of normal (4.3%). Single events of increase in ALT and aspartate aminotransferase were reported in 20.0% and 8.3%, respectively, of the 145 patients who completed the study.

In a study to assess the safety of deferasirox film-coated and dispersible tablets, 173 adult and paediatric patients with transfusion dependent thalassaemia or myelodysplastic syndrome were treated for 24 weeks. A comparable safety profile for film-coated and dispersible tablets was observed.

An open label 1:1 randomised study was performed in 224 paediatric patients aged 2 to <18 years old with transfusion-dependant anaemia and iron overload to evaluate treatment compliance, efficacy and safety of the deferasirox granule formulation compared to the dispersible tablet formulation. The majority of patients (142, 63.4%) in the study had beta-thalassemia major, 108 (48.2%) patients were naïve to iron chelation therapy (ICT) (median age 2 years, 92.6% aged 2 to <10 years) and 116 (51.8%) were ICT pre-treated (median age 7.5 years, 71.6% aged 2 to <10 years) of whom 68.1% had previously received deferasirox. In the primary analysis performed in ICT naïve patients after 24 weeks of treatment, the compliance rate was 84.26% and 86.84% in the deferasirox dispersible tablets arm and in the deferasirox granules arm, respectively, with no statistically significant difference. Similarly, there was no statistically significant difference in mean changes from baseline in serum ferritin (SF) values between the two treatment arms ( 171.52  $\mu\text{g/l}$  [95% CI: 517.40, 174.36] for dispersible tablets [DT] and 4.84  $\mu\text{g/l}$  [95% CI: -333.58, 343.27] for the granule formulation, difference between means [granules – DT] 176.36  $\mu\text{g/l}$  [95% CI: 129.00, 481.72], two sided p-value = 0.25). The study concluded that treatment compliance and efficacy were no different between deferasirox granules and deferasirox dispersible tablet arms at different time points (24 and 48 weeks). The safety profile was overall comparable between the granules and the dispersible tablet formulations.

In patients with non-transfusion-dependent thalassaemia syndromes and iron overload, treatment with deferasirox dispersible tablets was assessed in a 1-year, randomised, double-blind, placebo-controlled study. The study compared the efficacy of two different deferasirox dispersible tablet regimens (starting doses of 5 and 10 mg/kg/day, 55 patients in each arm) and of matching placebo (56 patients). The study enrolled 145 adult and 21 paediatric patients. The primary efficacy parameter was the change in liver iron concentration (LIC) from baseline after 12 months of treatment. One of the secondary efficacy parameters was the change in serum ferritin between baseline and fourth quarter. At a starting dose of 10 mg/kg/day, deferasirox dispersible tablets led to reductions in indicators of total body iron. On average, liver iron concentration decreased by 3.80 mg Fe/g dw in patients treated with deferasirox dispersible tablets (starting dose 10 mg/kg/day) and increased by 0.38 mg Fe/g dw in patients treated with placebo ( $p < 0.001$ ). On average, serum ferritin decreased by 222.0  $\mu\text{g/l}$  in patients treated with deferasirox dispersible tablets (starting dose 10 mg/kg/day) and increased by 115  $\mu\text{g/l}$  in patients treated with placebo ( $p < 0.001$ ).

## 5.2 Pharmacokinetic properties

Deferasirox film-coated tablets demonstrate higher bioavailability compared to deferasirox dispersible tablet formulations. After adjustment of the strength, the film-coated tablet formulation (360 mg strength) was equivalent to deferasirox dispersible tablets (500 mg strength) with respect to the mean area under the plasma concentration time curve (AUC) under fasting conditions. The  $C_{\text{max}}$  was increased by 30% (90% CI: 20.3% – 40.0%); however a clinical exposure/response analysis revealed no evidence of clinically relevant effects of such an increase.

### Absorption

Deferasirox (dispersible tablet formulation) is absorbed following oral administration with a median time to maximum plasma concentration ( $t_{\text{max}}$ ) of about 1.5 to 4 hours. The absolute bioavailability (AUC) of deferasirox (dispersible tablet formulation) is about 70% compared to an intravenous dose.

The absolute bioavailability of the film-coated tablet formulation has not been determined. Bioavailability of deferasirox film-coated tablets was 36% greater than that with dispersible tablets.

A food-effect study involving administration of the film-coated tablets to healthy volunteers under fasting conditions and with a low-fat (fat content <10% of calories) or high-fat (fat content >50% of calories) meal indicated that the AUC and  $C_{max}$  were slightly decreased after a low-fat meal (by 11% and 16%, respectively). After a high-fat meal, AUC and  $C_{max}$  were increased (by 18% and 29%, respectively). The increases in  $C_{max}$  due to the change in formulation and due to the effect of a high-fat meal may be additive and therefore, it is recommended that the film-coated tablets should be taken either on an empty stomach or with a light meal.

### Distribution

Deferasirox is highly (99%) protein bound to plasma proteins, almost exclusively serum albumin, and has a small volume of distribution of approximately 14 litres in adults.

### Biotransformation

Glucuronidation is the main metabolic pathway for deferasirox, with subsequent biliary excretion. Deconjugation of glucuronidates in the intestine and subsequent reabsorption (enterohepatic recycling) is likely to occur: in a healthy volunteer study, the administration of cholestyramine after a single dose of deferasirox resulted in a 45% decrease in deferasirox exposure (AUC).

Deferasirox is mainly glucuronidated by UGT1A1 and to a lesser extent UGT1A3. CYP450-catalysed (oxidative) metabolism of deferasirox appears to be minor in humans (about 8%). No inhibition of deferasirox metabolism by hydroxyurea was observed *in vitro*.

### Elimination

Deferasirox and its metabolites are primarily excreted in the faeces (84% of the dose). Renal excretion of deferasirox and its metabolites is minimal (8% of the dose). The mean elimination half-life ( $t_{1/2}$ ) ranged from 8 to 16 hours. The transporters MRP2 and MXR (BCRP) are involved in the biliary excretion of deferasirox.

### Linearity / non-linearity

The  $C_{max}$  and  $AUC_{0-24h}$  of deferasirox increase approximately linearly with dose under steady-state conditions. Upon multiple dosing exposure increased by an accumulation factor of 1.3 to 2.3.

### Characteristics in patients

#### *Paediatric patients*

The overall exposure of adolescents (12 to  $\leq 17$  years) and children (2 to <12 years) to deferasirox after single and multiple doses was lower than that in adult patients. In children younger than 6 years old exposure was about 50% lower than in adults. Since dosing is individually adjusted according to response this is not expected to have clinical consequences.

#### *Gender*

Females have a moderately lower apparent clearance (by 17.5%) for deferasirox compared to males. Since dosing is individually adjusted according to response this is not expected to have clinical consequences.

#### *Elderly patients*

The pharmacokinetics of deferasirox have not been studied in elderly patients (aged 65 or older).

#### *Renal or hepatic impairment*

The pharmacokinetics of deferasirox have not been studied in patients with renal impairment. The pharmacokinetics of deferasirox were not influenced by liver transaminase levels up to 5 times the upper limit of the normal range.

In a clinical study using single doses of 20 mg/kg deferasirox dispersible tablets, the average exposure was increased by 16% in subjects with mild hepatic impairment (Child-Pugh Class A) and by 76% in subjects with moderate hepatic impairment (Child-Pugh Class B) compared to subjects with normal hepatic function. The average  $C_{max}$  of deferasirox in subjects with mild or moderate hepatic impairment was increased by 22%. Exposure was increased 2.8-fold in one subject with severe hepatic impairment (Child-Pugh Class C) (see sections 4.2 and 4.4).

### 5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity or carcinogenic potential. The main findings were kidney toxicity and lens opacity (cataracts). Similar findings were observed in neonatal and juvenile animals. The kidney toxicity is considered mainly due to iron deprivation in animals that were not previously overloaded with iron.

Tests of genotoxicity *in vitro* were negative (Ames test, chromosomal aberration test) while deferasirox caused formation of micronuclei *in vivo* in the bone marrow, but not liver, of non-iron-loaded rats at lethal doses. No such effects were observed in iron-preloaded rats. Deferasirox was not carcinogenic when administered to rats in a 2-year study and transgenic p53+/- heterozygous mice in a 6-month study.

The potential for toxicity to reproduction was assessed in rats and rabbits. Deferasirox was not teratogenic, but caused increased frequency of skeletal variations and stillborn pups in rats at high doses that were severely toxic to the non-iron-overloaded mother. Deferasirox did not cause other effects on fertility or reproduction.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

#### Tablet core:

Cellulose, microcrystalline  
Crospovidone (Type A)  
Povidone (K30)  
Magnesium stearate  
Silica, colloidal anhydrous  
Poloxamer (P188)

#### Coating material:

Hypromellose  
Indigo Carmine Aluminium Lake (E132)  
Titanium dioxide (E171)  
Macrogol/PEG (6000)  
Talc

### 6.2 Incompatibilities

Not applicable.

### 6.3 Shelf life

3 years

### 6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

#### **6.5 Nature and contents of container**

Clear transparent PVC/PVDC/Aluminium blisters containing 30 or 90 film-coated tablets and unit dose blisters of 30 × 1 tablets.

Deferasirox Mylan 360 mg film-coated tablets are also available in blister pack of 300 tablets.

White HDPE bottle with white opaque polypropylene (PP) screw cap with aluminum seal containing 90 or 300 film-coated tablets.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

## **8. MARKETING AUTHORISATION NUMBER(S)**

### Deferasirox Mylan 90 mg film-coated tablets

EU/1/19/1386/001  
EU/1/19/1386/002  
EU/1/19/1386/003  
EU/1/19/1386/004  
EU/1/19/1386/005

### Deferasirox Mylan 180 mg film-coated tablets

EU/1/19/1386/006  
EU/1/19/1386/007  
EU/1/19/1386/008  
EU/1/19/1386/009  
EU/1/19/1386/010

### Deferasirox Mylan 360 mg film-coated tablets

EU/1/19/1386/011  
EU/1/19/1386/012  
EU/1/19/1386/013  
EU/1/19/1386/014  
EU/1/19/1386/015  
EU/1/19/1386/016

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 26 September 2019

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## **ANNEX II**

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**



## **A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer(s) responsible for batch release

Mylan Hungary Kft.  
Mylan utca 1  
2900 Komarom  
HUNGARY

Mylan Germany GmbH  
Zweigniederlassung Bad Homburg v. d. Hoehe  
Benzstrasse 1, Bad Homburg v. d. Hoehe, Hessen, 61352  
GERMANY

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

- **Additional risk minimisation measures**

Prior to launch of Deferasirox Mylan in each Member State the Marketing Authorisation Holder (MAH) must agree about the content and format of the educational programme, including communication media, distribution modalities, and any other aspects of the programme, with the National Competent Authority.

The educational programme is aimed to inform healthcare professionals and patients to minimise the risks of:

- Non-compliance of the posology and biological monitoring
- Medication errors due to switching between formulations available on the market by different MAHs (film-coated tablets/granules and generic versions of dispersible tablets).

The risk of medication error is due to switching between Deferasirox film coated tablets/granules and generic deferasirox dispersible tablet formulations available on the market by different MAHs and as appropriate depending on the coexistence of these formulations at a national level. The MAH shall ensure that, at launch, in each Member State where Deferasirox Mylan is marketed, all healthcare professionals and patients who are expected to prescribe, dispense and use Deferasirox Mylan are provided with the following educational package for all available formulations (e.g. dispersible tablets, Deferasirox film-coated tablets and Deferasirox granules) for all indications:

- Physician educational material
- Patient information pack

Additional periodic distributions should be performed, notably after substantial safety modifications of the product information justifying educational material updates.

The physician educational material should contain:

- The Summary of Product Characteristics
- Guide for healthcare professionals (which also includes a prescriber checklist).

**The Guide for healthcare professionals** shall contain the following key elements as appropriate depending on the coexistence of Deferasirox formulations at a national level:

- Description of available deferasirox formulations on the market (e.g. dispersible tablets, film-coated tablets and granules) in the EU.
  - Different posology regimen
  - Different conditions of administration
  - Dose conversion table of Deferasirox filmcoated tablets/granules and Deferasirox dispersible tablets as a reference when switching from one formulation to another
- The recommended doses and the rules for starting treatment
- The need to monitor serum ferritin monthly
- That deferasirox causes rises in serum creatinine in some patients
  - The need to monitor serum creatinine
    - On two occasions prior to initiation of treatment
    - Every week during the first month of initiation of treatment or after therapy modification
    - Monthly thereafter
  - The need to reduce by 7 mg/kg the dose if serum creatinine rises:
    - Adults: >33% above baseline and creatinine clearance <LLN (90 ml/min)
    - Paediatrics: either >ULN or creatinine clearance falls to <LLN at two consecutive visits.
  - The need to interrupt treatment after a dose reduction if serum creatinine rises:
    - Adults and Paediatrics: remain >33% above baseline or creatinine clearance <LLN (90 ml/min)
  - The need to consider renal biopsy:

- When serum creatinine is elevated and if another abnormality has been detected (e.g. proteinuria, signs of Fanconi syndrome).
- The importance of measuring creatinine clearance
- Brief overview of methods of measuring creatinine clearance

- That rises in serum transaminases may occur in patients treated with Deferasirox Mylan
  - The need for liver function tests prior to prescription, then at monthly intervals or more often if clinically indicated
  - Not to prescribe to patients with pre-existing severe hepatic disease
  - The need to interrupt treatment if persistent and progressive increase in liver enzyme were noted.
- The need for annual auditory and ophthalmic testing
- The need for a guidance table highlighting pre-treatment measurements of serum creatinine, creatinine clearance, proteinuria, hepatic enzymes, ferritin, such as:

Before initiating treatment	
Serum creatinine at Day – X	Value 1
Serum creatinine at Day – Y	Value 2

X and Y are the days (to be determined) when pre-treatment measurements should be performed.

- A warning on the risk of overchelation and on the necessity of close monitoring of serum ferritin levels and renal and hepatic function.
- The rules for treatment dose adjustments and interruption when target serum ferritin +/- liver iron concentration are reached.
- Recommendations for treatment of non-transfusion-dependent thalassaemia (NTDT) syndromes:
  - Information that only one course of treatment is proposed for NTDT patients
  - A warning on the necessity of closer monitoring of liver iron concentration and serum ferritin in the paediatric population
  - A warning on the currently unknown safety consequences of long-term treatment in the paediatric population

**The patient information pack** should contain:

- Patient information leaflet
- Patient guide

Patient guide should contain the following key elements:

- Information on the need for regular monitoring, and when it should be carried out, of serum creatinine, creatinine clearance, proteinuria, hepatic enzymes, ferritin
- Information that renal biopsy may be considered if significant renal abnormalities occur
- Availability of several oral formulations (e.g. dispersible tablets, film-coated tablets and granules) and the main differences associated with these formulations (i.e., different posology regimen, different conditions of administration notably with food)

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON (BLISTER AND BOTTLE)**

**1. NAME OF THE MEDICINAL PRODUCT**

Deferasirox Mylan 90 mg film-coated tablets  
deferasirox

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 90 mg deferasirox.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet (tablet)

*[Blisters]*

30 film-coated tablets

90 film-coated tablets

*[Unit dose blisters]*

30 × 1 film-coated tablets

*[Bottles]:*

90 film-coated tablets

300 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Ltd  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/19/1386/001  
EU/1/19/1386/002  
EU/1/19/1386/003  
EU/1/19/1386/004  
EU/1/19/1386/005

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Deferasirox Mylan 90 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included



**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON (BLISTER AND BOTTLE)**

**1. NAME OF THE MEDICINAL PRODUCT**

Deferasirox Mylan 180 mg film-coated tablets  
deferasirox

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 180 mg deferasirox

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet (tablet)

*[Blisters]*

30 film-coated tablets

90 film-coated tablets

*[Unit dose blisters]*

30 × 1 film-coated tablets

*[Bottles]*

90 film-coated tablets

300 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Ltd  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/19/1386/006  
EU/1/19/1386/007  
EU/1/19/1386/008  
EU/1/19/1386/009  
EU/1/19/1386/010

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Deferasirox Mylan 180 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON (BLISTER AND BOTTLE)**

**1. NAME OF THE MEDICINAL PRODUCT**

Deferasirox Mylan 360 mg film-coated tablets  
deferasirox

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 360 mg deferasirox

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet (tablet)

*[Blisters]*

30 film-coated tablets

90 film-coated tablets

300 film-coated tablets

*[Unit dose blisters]*

30 × 1 film-coated tablets

*[Bottles]*

90 film-coated tablets

300 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Ltd  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/19/1386/011  
EU/1/19/1386/012  
EU/1/19/1386/013  
EU/1/19/1386/014  
EU/1/19/1386/015  
EU/1/19/1386/016

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Deferasirox Mylan 360 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**BOTTLE LABEL**

**1. NAME OF THE MEDICINAL PRODUCT**

Deferasirox Mylan 90 mg film-coated tablets  
deferasirox

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 90 mg deferasirox.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet (tablet)

90 film-coated tablets  
300 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**



**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Ltd  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/19/1386/004  
EU/1/19/1386/005

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**BOTTLE LABEL**

**1. NAME OF THE MEDICINAL PRODUCT**

Deferasirox Mylan 180 mg film-coated tablets  
deferasirox

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 180 mg deferasirox

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet (tablet)

90 film-coated tablets  
300 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Ltd  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/19/1386/009  
EU/1/19/1386/010

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**BOTTLE LABEL**

**1. NAME OF THE MEDICINAL PRODUCT**

Deferasirox Mylan 360 mg film-coated tablets  
deferasirox

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 360 mg deferasirox

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet (tablet)

90 film-coated tablets

300 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Ltd  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/19/1386/015  
EU/1/19/1386/016

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Deferasirox Mylan 90 mg film-coated tablets  
deferasirox

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Ltd

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Deferasirox Mylan 180 mg film-coated tablets  
deferasirox

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Ltd

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Deferasirox Mylan 360 mg film-coated tablets  
deferasirox

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Ltd

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**



**B. PACKAGE LEAFLET**

## Package leaflet: Information for the user

**Deferasirox Mylan 90 mg film-coated tablets**  
**Deferasirox Mylan 180 mg film-coated tablets**  
**Deferasirox Mylan 360 mg film-coated tablets**  
deferasirox

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed only for you or your child. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### **What is in this leaflet**

1. What Deferasirox Mylan is and what it is used for
2. What you need to know before you take Deferasirox Mylan
3. How to take Deferasirox Mylan
4. Possible side effects
5. How to store Deferasirox Mylan
6. Contents of the pack and other information

### **1. What Deferasirox Mylan is and what it is used for**

#### **What Deferasirox Mylan is**

Deferasirox Mylan contains an active substance called deferasirox. It is an iron chelator which is a medicine used to remove the excess iron from the body (also called iron overload). It traps and removes excess iron which is then excreted mainly in the stools.

#### **What Deferasirox Mylan is used for**

Repeated blood transfusions may be necessary in patients with various types of anaemia (for example thalassaemia, sickle cell disease or myelodysplastic syndromes (MDS)). However, repeated blood transfusions can cause a build-up of excess iron. This is because blood contains iron and your body does not have a natural way to remove the excess iron you get with your blood transfusions. In patients with non-transfusion-dependent thalassaemia syndromes, iron overload may also develop over time, mainly due to increased absorption of dietary iron in response to low blood cell counts. Over time, the excess iron can damage important organs such as the liver and heart. Medicines called *iron chelators* are used to remove the excess iron and reduce the risk of it causing organ damage.

Deferasirox Mylan is used to treat chronic iron overload caused by frequent blood transfusions in patients with beta thalassaemia major aged 6 years and older.

Deferasirox Mylan is also used to treat chronic iron overload when deferoxamine therapy is contraindicated or inadequate in patients with beta thalassaemia major with iron overload caused by infrequent blood transfusions, in patients with other types of anaemias, and in children aged 2 to 5 years.

Deferasirox Mylan is also used when deferoxamine therapy is contraindicated or inadequate to treat patients aged 10 years or older who have iron overload associated with their thalassaemia syndromes, but who are not transfusion dependent.

## 2. What you need to know before you take Deferasirox Mylan

### Do not take Deferasirox Mylan

- if you are allergic to deferasirox or any of the other ingredients of this medicine (listed in section 6). If this applies to you, **tell your doctor before taking Deferasirox Mylan**. If you think you may be allergic, ask your doctor for advice.
- if you have moderate or severe kidney disease.
- if you are currently taking any other iron chelator medicines.

### Deferasirox Mylan is not recommended

- if you are at an advanced stage of myelodysplastic syndrome (MDS; decreased production of blood cells by the bone marrow) or have advanced cancer.

### Warnings and precautions

Talk to your doctor or pharmacist before taking Deferasirox Mylan:

- if you have a kidney or liver problem.
- if you have a cardiac problem due to iron overload.
- if you notice a marked decrease in your urine output (sign of kidney problem).
- if you develop a severe rash, or difficulty breathing and dizziness or swelling mainly of the face and throat (signs of severe allergic reaction, see also section 4 “Possible side effects”).
- if you experience a combination of any of the following symptoms: rash, red skin, blistering of the lips, eyes or mouth, skin peeling, high fever, flu-like symptoms, enlarged lymph nodes (signs of severe skin reaction, see also section 4 “Possible side effects”).
- if you experience a combination of drowsiness, upper right abdominal pain, yellowing or increased yellowing of your skin or eyes and dark urine (signs of liver problems).
- if you experience difficulty thinking, remembering information, or solving problems, being less alert or aware or feeling very sleepy with low energy (signs of a high level of ammonia in your blood, which may be associated with liver or renal problems, see also section 4 “Possible side effects”).
- if you vomit blood and/or have black stools.
- if you experience frequent abdominal pain, particularly after eating or taking Deferasirox Mylan.
- if you experience frequent heartburn.
- if you have a low level of platelets or white blood cells in your blood test.
- if you have blurred vision
- if you have diarrhoea or vomiting.

If any of these apply to you, tell your doctor straight away.

### Monitoring your Deferasirox Mylan treatment

You will have regular blood and urine tests during treatment. These will monitor the amount of iron in your body (blood level of *ferritin*) to see how well Deferasirox Mylan is working. The tests will also monitor your kidney function (blood level of creatinine, presence of protein in the urine) and liver function (blood level of transaminases). Your doctor may require you to undergo a kidney biopsy, if he/she suspects significant kidney damage. You may also have MRI (magnetic resonance imaging) tests to determine the amount of iron in your liver. Your doctor will take these tests into consideration when deciding on the dose of Deferasirox Mylan most suitable for you and will also use these tests to decide when you should stop taking Deferasirox Mylan.

Your eyesight and hearing will be tested each year during treatment as a precautionary measure.

### Other medicines and Deferasirox Mylan

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes in particular:

- other iron chelators, which must not be taken with Deferasirox Mylan,
- antacids (medicines used to treat heartburn) containing aluminium, which should not be taken at the same time of day as Deferasirox Mylan,

- ciclosporin (used to prevent the body rejecting a transplanted organ or for other conditions, such as rheumatoid arthritis or atopic dermatitis),
- simvastatin (used to lower cholesterol),
- certain painkillers or anti-inflammatory medicines (e.g. aspirin, ibuprofen, corticosteroids),
- oral bisphosphonates (used to treat osteoporosis),
- anticoagulant medicines (used to prevent or treat blood clotting),
- hormonal contraceptive agents (birth control medicines),
- bepridil, ergotamine (used for heart problems and migraines),
- repaglinide (used to treat diabetes),
- rifampicin (used to treat tuberculosis),
- phenytoin, phenobarbital, carbamazepine (used to treat epilepsy),
- ritonavir (used in the treatment of HIV infection),
- paclitaxel (used in cancer treatment),
- theophylline (used to treat respiratory diseases such as asthma),
- clozapine (used to treat psychiatric disorders such as schizophrenia),
- tizanidine (used as a muscle relaxant),
- cholestyramine (used to lower cholesterol levels in the blood),
- busulfan (used as a treatment prior to transplantation in order to destroy the original bone marrow before the transplant),
- midazolam (used to relieve anxiety and/or trouble sleeping).

Additional tests may be required to monitor the blood levels of some of these medicines.

#### **Older people (age 65 years and over)**

Deferasirox Mylan can be used by people aged 65 years and over at the same dose as for other adults. Elderly patients may experience more side effects (in particular diarrhoea) than younger patients. They should be monitored closely by their doctor for side effects that may require a dose adjustment.

#### **Children and adolescents**

Deferasirox Mylan can be used in children and adolescents receiving regular blood transfusions aged 2 years and over and in children and adolescents not receiving regular blood transfusions aged 10 years and over. As the patient grows the doctor will adjust the dose.

Deferasirox Mylan is not recommended for children aged under 2 years.

#### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

Deferasirox Mylan is not recommended during pregnancy unless clearly necessary.

If you are currently using a hormonal contraceptive to prevent pregnancy, you should use an additional or different type of contraception (e.g. condom), as Deferasirox Mylan may reduce the effectiveness of hormonal contraceptives.

Breast-feeding is not recommended during treatment with Deferasirox Mylan.

#### **Driving and using machines**

If you feel dizzy after taking Deferasirox Mylan, do not drive or operate any tools or machines until you are feeling normal again.

**Deferasirox Mylan contains less than 1 mmol sodium** (23 mg) per tablet, that is to say essentially 'sodium-free'.

### 3. How to take Deferasirox Mylan

Treatment with Deferasirox Mylan will be overseen by a doctor who is experienced in the treatment of iron overload caused by blood transfusions.

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

#### How much Deferasirox Mylan to take

The dose of Deferasirox Mylan is related to body weight for all patients. Your doctor will calculate the dose you need and tell you how many tablets to take each day.

- The usual daily dose for Deferasirox Mylan film-coated tablets at the start of the treatment for patients receiving regular blood transfusions is 14 mg per kilogram body weight. A higher or lower starting dose may be recommended by your doctor based on your individual treatment needs.
- The usual daily dose for Deferasirox Mylan film-coated tablets at the start of the treatment for patients not receiving regular blood transfusions is 7 mg per kilogram body weight.
- Depending on how you respond to treatment, your doctor may later adjust your treatment to a higher or lower dose.

The maximum recommended daily dose for Deferasirox Mylan film-coated tablets is:

- 28 mg per kilogram body weight for patients receiving regular blood transfusions,
- 14 mg per kilogram body weight for adult patients not receiving regular blood transfusions,
- 7 mg per kilogram body weight for children and adolescents not receiving regular blood transfusions.

Deferasirox also comes as “dispersible” tablets. If you are switching from the dispersible tablets to these film-coated tablets, you will need an adjustment of the dose.

#### When to take Deferasirox Mylan

- Take Deferasirox Mylan once a day, every day, at about the same time each day with some water.
- Take Deferasirox Mylan film-coated tablets either on an empty stomach or with a light meal.

Taking Deferasirox Mylan at the same time each day will also help you remember when to take your tablets.

For patients who are unable to swallow whole tablets, Deferasirox Mylan film-coated tablets may be crushed and taken by sprinkling the full dose onto soft food such as yogurt or apple sauce (pureed apple). The food should be immediately and completely consumed. Do not store it for future use.

#### How long to take Deferasirox Mylan

**Continue taking Deferasirox Mylan every day for as long as your doctor tells you.** This is a long-term treatment, possibly lasting for months or years. Your doctor will regularly monitor your condition to check that the treatment is having the desired effect (see also section 2: “Monitoring your Deferasirox Mylan treatment”).

If you have questions about how long to take Deferasirox Mylan, talk to your doctor.

#### If you take more Deferasirox Mylan than you should

If you have taken too much Deferasirox Mylan, or if someone else accidentally takes your tablets, contact your doctor or hospital for advice straight away. Show the doctor the pack of tablets. Urgent medical treatment may be necessary. You may experience effects such as abdominal pain, diarrhoea, nausea and vomiting and kidney or liver problems that can be serious.

#### If you forget to take Deferasirox Mylan

If you miss a dose, take it as soon as you remember on that day. Take your next dose as scheduled. Do not take a double dose on the next day to make up for the forgotten tablet(s).

### **If you stop taking Deferasirox Mylan**

Do not stop taking Deferasirox Mylan unless your doctor tells you to. If you stop taking it, the excess iron will no longer be removed from your body (see also above section “How long to take Deferasirox Mylan”).

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them. Most of the side effects are mild to moderate and will generally disappear after a few days to a few weeks of treatment.

### **Some side effects could be serious and need immediate medical attention.**

*These side effects are **uncommon** (may affect up to 1 in 100 people) or **rare** (may affect up to 1 in 1,000 people).*

- If you get a severe rash, or difficulty breathing and dizziness or swelling mainly of the face and throat (signs of severe allergic reaction),
- If you experience a combination of any of the following symptoms: rash, red skin, blistering of the lips, eyes or mouth, skin peeling, high fever, flu-like symptoms, enlarged lymph nodes, (signs of severe skin reactions),
- If you notice a marked decrease in your urine output (sign of kidney problem),
- If you experience a combination of drowsiness, upper right abdominal pain, yellowing or increased yellowing of your skin or eyes and dark urine (signs of liver problems),
- If you experience difficulty thinking, remembering information, or solving problems, being less alert or aware or feeling very sleepy with low energy (signs of a high level of ammonia in your blood, which may be associated with liver or renal problems and lead to a change in your brain function),
- If you vomit blood and/or have black stools,
- If you experience frequent abdominal pain, particularly after eating or taking Deferasirox Mylan,
- If you experience frequent heartburn,
- If you experience partial loss of vision,
- If you experience severe upper stomach pain (pancreatitis),

**stop taking this medicine and tell your doctor straight away.**

### **Some side effects could become serious.**

*These side effects are **uncommon**.*

- If you get blurred or cloudy eyesight,
- If you get reduced hearing,

**tell your doctor as soon as possible.**

### **Other side effects**

***Very common** (may affect more than 1 in 10 people)*

- Disturbance in kidney function tests.

***Common** (may affect up to 1 in 10 people)*

- Gastrointestinal disorders, such as nausea, vomiting, diarrhoea, pain in the abdomen, bloating, constipation, indigestion
- Rash
- Headache
- Disturbance in liver function tests
- Itching
- Disturbance in urine test (protein in the urine)

If any of these affects you severely, tell your doctor.

**Uncommon** (may affect up to 1 in 100 people)

- Dizziness
- Fever
- Sore throat
- Swelling of arms or legs
- Change in the colour of the skin
- Anxiety
- Sleep disorder
- Tiredness

If any of these affects you severely, tell your doctor.

**Frequency not known** (cannot be estimated from the available data)

- A decrease in the number of cells involved in blood clotting (thrombocytopenia), in the number of red blood cells (anaemia aggravated), in the number of white blood cells (neutropenia) or in the number of all kinds of blood cells (pancytopenia)
- Hair loss
- Kidney stones
- Low urine output
- Tear in stomach or intestine wall that can be painful and cause nausea
- Severe upper stomach pain (pancreatitis)
- Abnormal level of acid in blood

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Deferasirox Mylan**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister/label and the carton after 'EXP'. The expiry date refers to the last day of that month.

Do not use any pack that is damaged or shows signs of tampering.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Deferasirox Mylan contains**

The active substance is deferasirox.

- Each film-coated tablet of Deferasirox Mylan 90 mg contains 90 mg deferasirox.
- Each film-coated tablet of Deferasirox Mylan 180 mg contains 180 mg deferasirox.
- Each film-coated tablet of Deferasirox Mylan 360 mg contains 360 mg deferasirox.

The other ingredients are microcrystalline cellulose, crospovidone, povidone, magnesium stearate, colloidal anhydrous silica and poloxamer. The tablet coating material contains: hypromellose, titanium dioxide (E171), macrogol/PEG (6000), talc, indigo carmine aluminium Lake (E132).

### **What Deferasirox Mylan looks like and contents of the pack**

Deferasirox Mylan is supplied as film-coated tablets.

- Deferasirox Mylan 90 mg film-coated tablets are blue, film-coated, modified capsule shaped, biconvex tablets debossed with “ $\cap$ ” on one side of the tablet and ‘DF’ on the other side.

- Deferasirox Mylan 180 mg film-coated tablets are blue, film-coated, modified capsule shaped, biconvex tablets debossed with “M” on one side of the tablet and ‘DF 1’ on the other side.
- Deferasirox Mylan 360 mg film-coated tablets are blue, film-coated, modified capsule shaped, biconvex tablets debossed with “M” on one side of the tablet and ‘DF 2’ on the other side.

Deferasirox Mylan is available in clear, transparent PVC/PVdC /aluminium blister packs containing 30 or 90 film-coated tablets, in unit dose blister pack of 30 tablets and in white, plastic bottles with white opaque screw cap with aluminium seal containing 90 and 300 tablets.

Deferasirox Mylan 360 mg film-coated tablets are also available in blister packs of 300 tablets.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park, Mulhuddart, Dublin 15, DUBLIN, Ireland

### **Manufacturer**

Mylan Hungary Kft., Mylan utca 1, Komárom 2900, Hungary

Mylan Germany GmbH, Zweigniederlassung Bad Homburg v. d. Hoehe  
Benzstrasse 1, Bad Homburg v. d. Hoehe, Hessen, 61352, Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

#### **België/Belgique/Belgien**

Viatrix  
Tél/Tel: + 32 (0)2 658 61 00

#### **Lietuva**

Viatrix UAB  
Tel: +370 5 205 1288

#### **България**

Майлан ЕООД  
Тел: +359 2 44 55 400

#### **Luxembourg/Luxemburg**

Viatrix  
Tél/Tel: + 32 (0)2 658 61 00  
(Belgique/Belgien)

#### **Česká republika**

Viatrix CZ, s.r.o.  
Tel: + 420 222 004 400

#### **Magyarország**

Viatrix Healthcare Kft  
Tel: + 36 1 465 2100

#### **Danmark**

Viatrix ApS  
Tlf: +45 28 11 69 32

#### **Malta**

V.J. Salomone Pharma Ltd  
Tel: + 356 21 22 01 74

#### **Deutschland**

Viatrix Healthcare GmbH  
Tel: +49 800 0700 800

#### **Nederland**

Mylan BV  
Tel: +31 (0)20 426 3300



**Eesti**

Viatriis OU  
Tel: + 372 6363 052

**Ελλάδα**

Viatriis Hellas Ltd  
Τηλ: +30 2100 100 02

**España**

Viatriis Pharmaceuticals, S.L.  
Tel: + 34 900 102 712

**France**

Viatriis Santé  
Tél : +33 4 37 25 75 00

**Hrvatska**

Viatriis Hrvatska d.o.o.  
Tel: +385 1 23 50 599

**Ireland**

Viatriis Limited  
Tel: +353 1 8711600

**Ísland**

Icepharma hf  
Sími: +354 540 8000

**Italia**

Viatriis Italia S.r.l.  
Tel: + 39 02 612 46921

**Κύπρος**

CPO Pharmaceuticals Ltd  
Τηλ: + 357 22863100

**Latvija**

Viatriis SIA  
Tel: +371 676 055 80

**Norge**

Viatriis AS  
Tlf: +47 66 75 33 00

**Österreich**

Viatriis Austria GmbH  
Tel: +43 1 863904

**Polska**

Viatriis Healthcare Sp. z.o.o  
.Tel: + 48 22 546 64 00

**Portugal**

Mylan, Lda.  
Tel: + 351 214 127 200

**România**

BGP Products SRL  
Tel: + 40 372 579 000

**Slovenija**

Viatriis d.o.o.  
Tel: + 386 1 23 63 180

**Slovenská republika**

Viatriis Slovakia s.r.o.  
Tel: +421 2 32 199 100

**Suomi/Finland**

Viatriis Oy  
Puh/Tel: +358 20 720 9555

**Sverige**

Viatriis AB  
Tel: + 46 (0) 8 630 19 00

**This leaflet was last revised in {MM/YYYY}.**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency website:  
<http://www.ema.europa.eu>.