

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatriis 2.5 mg film-coated tablets
Olanzapine Viatriis 5 mg film-coated tablets
Olanzapine Viatriis 7.5 mg film-coated tablets
Olanzapine Viatriis 10 mg film-coated tablets
Olanzapine Viatriis 15 mg film-coated tablets
Olanzapine Viatriis 20 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Olanzapine Viatriis 2.5 mg film-coated tablets

Each film-coated tablet contains 2.5 mg olanzapine.

Excipients with known effect

Each film-coated tablet contains 76 mg lactose (as monohydrate).

The film-coating of each 2.5 mg tablet contains 0.06 mg soya lecithin.

Olanzapine Viatriis 5 mg film-coated tablets

Each film-coated tablet contains 5 mg olanzapine.

Excipients with known effect

Each film-coated tablet contains 152 mg lactose (as monohydrate).

The film-coating of each 5 mg tablet contains 0.12 mg soya lecithin.

Olanzapine Viatriis 7.5 mg film-coated tablets

Each film-coated tablet contains 7.5 mg olanzapine.

Excipients with known effect

Each film-coated tablet contains 228 mg lactose (as monohydrate).

The film-coating of each 7.5 mg tablet contains 0.18 mg soya lecithin.

Olanzapine Viatriis 10 mg film-coated tablets

Each film-coated tablet contains 10 mg olanzapine.

Excipients with known effect

Each film-coated tablet contains 304 mg lactose (as monohydrate).

The film-coating of each 10 mg tablet contains 0.24 mg soya lecithin.

Olanzapine Viatriis 15 mg film-coated tablets

Each film-coated tablet contains 15 mg olanzapine.

Excipients with known effect

Each film-coated tablet contains 183 mg lactose (as monohydrate).

The film-coating of each 15 mg tablet contains 0.15 mg soya lecithin.

Olanzapine Viatriis 20 mg film-coated tablets

Each film-coated tablet contains 20 mg olanzapine.

Excipients with known effect

Each film-coated tablet contains 244 mg lactose (as monohydrate).

The film-coating of each 20 mg tablet contains 0.20 mg soya lecithin.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet (tablet).

Olanzapine Viatriis 2.5 mg film-coated tablets

Approximately 7.0 mm, round, normal convex, white film-coated tablets debossed “OZ” over “2.5” on one side and “G” on the other side.

Olanzapine Viatriis 5 mg film-coated tablets

Approximately 8.0 mm, round, normal convex, white film-coated tablets debossed “OZ” over “5” on one side and “G” on the other side.

Olanzapine Viatriis 7.5 mg film-coated tablets

Approximately 9.0 mm, round, normal convex, white film-coated tablets debossed “OZ” over “7.5” on one side and “G” on the other side.

Olanzapine Viatriis 10 mg film-coated tablets

Approximately 10.2 mm, round, normal convex, white film-coated tablets debossed “OZ” over “10” on one side and “G” on the other side.

Olanzapine Viatriis 15 mg film-coated tablets

Approximately 12.2 mm x 6.7 mm, ellipse-shaped, normal convex, white film-coated tablets debossed “OZ 15” on one side and “G” on the other side.

Olanzapine Viatriis 20 mg film-coated tablets

Approximately 13.4 mm x 7.3 mm, ellipse-shaped, normal convex, white film-coated tablets debossed “OZ 20” on one side and “G” on the other side.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Adults

Olanzapine is indicated for the treatment of schizophrenia.

Olanzapine is effective in maintaining the clinical improvement during continuation therapy in patients who have shown an initial treatment response.

Olanzapine is indicated for the treatment of moderate to severe manic episode.

In patients whose manic episode has responded to olanzapine treatment, olanzapine is indicated for the prevention of recurrence in patients with bipolar disorder (see section 5.1).

4.2 Posology and method of administration

Adults

Schizophrenia: The recommended starting dose for olanzapine is 10 mg/day.

Manic episode: The starting dose is 15 mg as a single daily dose in monotherapy or 10 mg daily in combination therapy (see section 5.1).

Preventing recurrence in bipolar disorder: The recommended starting dose is 10 mg/day. For patients who have been receiving olanzapine for treatment of manic episode, continue therapy for preventing recurrence at the same dose. If a new manic, mixed, or depressive episode occurs, olanzapine treatment should be continued (with dose optimisation as needed), with supplementary therapy to treat mood symptoms, as clinically indicated.

During treatment for schizophrenia, manic episode and recurrence prevention in bipolar disorder, daily dosage may subsequently be adjusted on the basis of individual clinical status within the range 5-20 mg/day. An increase to a dose greater than the recommended starting dose is advised only after appropriate clinical reassessment and should generally occur at intervals of not less than 24 hours.

Olanzapine can be given without regards for meals as absorption is not affected by food. Gradual tapering of the dose should be considered when discontinuing olanzapine.

Special populations

Elderly

A lower starting dose (5 mg/day) is not routinely indicated but should be considered for those 65 and over when clinical factors warrant (see section 4.4).

Renal and/or hepatic impairment

A lower starting dose (5 mg) should be considered for such patients. In cases of moderate hepatic insufficiency (cirrhosis, Child-Pugh Class A or B), the starting dose should be 5 mg and only increased with caution.

Smokers

The starting dose and dose range need not be routinely altered for non-smokers relative to smokers. The metabolism of olanzapine may be induced by smoking. Clinical monitoring is recommended and an increase of olanzapine dose may be considered if necessary (see section 4.5).

When more than one factor is present which might result in slower metabolism (female gender, geriatric age, non-smoking status), consideration should be given to decreasing the starting dose. Dose escalation, when indicated, should be conservative in such patients (see sections 4.5 and 5.2).

Paediatric population

Olanzapine is not recommended for use in children and adolescents below 18 years of age due to a lack of data on safety and efficacy. A greater magnitude of weight gain, lipid and prolactin alterations has been reported in short term studies of adolescent patients than in studies of adult patients (see sections 4.4, 4.8, 5.1 and 5.2).

4.3 Contraindications

Hypersensitivity to the active substance, peanut or soya, or to any of the excipients listed in section 6.1.

Patients with known risk of narrow-angle glaucoma.

4.4 Special warnings and precautions for use

During antipsychotic treatment, improvement in the patient's clinical condition may take several days to some weeks. Patients should be closely monitored during this period.

Dementia-related psychosis and/or behavioural disturbances

Olanzapine is not recommended for use in patients with dementia-related psychosis and/or behavioural disturbances because of an increase in mortality and the risk of cerebrovascular accident. In placebo-controlled clinical trials (6-12 weeks duration) of elderly patients (mean age 78 years) with dementia-related psychosis and/or disturbed behaviours, there was a 2-fold increase in the incidence of death in olanzapine-treated patients compared to patients treated with placebo (3.5 % vs. 1.5 %, respectively). The higher incidence of death was not associated with olanzapine dose (mean daily dose 4.4 mg) or duration of treatment. Risk factors that may predispose this patient population to increased mortality include age >65 years, dysphagia, sedation, malnutrition and dehydration, pulmonary conditions (e.g., pneumonia, with or without aspiration), or concomitant use of benzodiazepines. However, the incidence of death was higher in olanzapine-treated than in placebo-treated patients independent of these risk factors.

In the same clinical trials, cerebrovascular adverse events (CVAE e.g., stroke, transient ischaemic attack), including fatalities, were reported. There was a 3-fold increase in CVAE in patients treated with olanzapine compared to patients treated with placebo (1.3 % vs. 0.4 %, respectively). All olanzapine- and placebo-treated patients who experienced a cerebrovascular event had pre-existing

risk factors. Age >75 years and vascular/mixed type dementia were identified as risk factors for CVAE in association with olanzapine treatment. The efficacy of olanzapine was not established in these trials.

Parkinson's disease

The use of olanzapine in the treatment of dopamine agonist associated psychosis in patients with Parkinson's disease is not recommended. In clinical trials, worsening of Parkinsonian symptomatology and hallucinations were reported very commonly and more frequently than with placebo (see section 4.8), and olanzapine was not more effective than placebo in the treatment of psychotic symptoms. In these trials, patients were initially required to be stable on the lowest effective dose of anti-Parkinsonian medicinal products (dopamine agonist) and to remain on the same anti-Parkinsonian medicinal products and dosages throughout the study. Olanzapine was started at 2.5 mg/day and titrated to a maximum of 15 mg/day based on investigator judgement.

Neuroleptic Malignant Syndrome (NMS)

NMS is a potentially life-threatening condition associated with antipsychotic medicinal products. Rare cases reported as NMS have also been received in association with olanzapine. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status, and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional signs may include elevated creatine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure. If a patient develops signs and symptoms indicative of NMS, or presents with unexplained high fever without additional clinical manifestations of NMS, all antipsychotic medicines, including olanzapine must be discontinued.

Hyperglycaemia and diabetes

Hyperglycaemia and/or development or exacerbation of diabetes occasionally associated with ketoacidosis or coma has been reported uncommonly, including some fatal cases (see section 4.8). In some cases, a prior increase in body weight has been reported which may be a predisposing factor. Appropriate clinical monitoring is advisable in accordance with utilised antipsychotic guidelines, e.g. measuring of blood glucose at baseline, 12 weeks after starting olanzapine treatment and annually thereafter. Patients treated with any antipsychotic medicines, including olanzapine, should be observed for signs and symptoms of hyperglycaemia (such as polydipsia, polyuria, polyphagia, and weakness) and patients with diabetes mellitus or with risk factors for diabetes mellitus should be monitored regularly for worsening of glucose control. Weight should be monitored regularly, e.g. at baseline, 4, 8 and 12 weeks after starting olanzapine treatment and quarterly thereafter.

Lipid alterations

Undesirable alterations in lipids have been observed in olanzapine-treated patients in placebo-controlled clinical trials (see section 4.8). Lipid alterations should be managed as clinically appropriate, particularly in dyslipidemic patients and in patients with risk factors for the development of lipids disorders. Patients treated with any antipsychotic medicines, including olanzapine, should be monitored regularly for lipids in accordance with utilised antipsychotic guidelines, e.g. at baseline, 12 weeks after starting olanzapine treatment and every 5 years thereafter.

Anticholinergic activity

While olanzapine demonstrated anticholinergic activity *in vitro*, experience during the clinical trials revealed a low incidence of related events. However, as clinical experience with olanzapine in patients with concomitant illness is limited, caution is advised when prescribing for patients with prostatic hypertrophy, or paralytic ileus and related conditions.

Hepatic function

Transient, asymptomatic elevations of hepatic aminotransferases, ALT, AST have been seen commonly, especially in early treatment. Caution should be exercised and follow-up organised in patients with elevated ALT and/or AST, in patients with signs and symptoms of hepatic impairment, in patients with pre-existing conditions associated with limited hepatic functional reserve, and in patients who are being treated with potentially hepatotoxic medicines. In cases where hepatitis (including hepatocellular, cholestatic or mixed liver injury) has been diagnosed, olanzapine treatment should be discontinued.

Neutropenia

Caution should be exercised in patients with low leukocyte and/or neutrophil counts for any reason, in patients receiving medicines known to cause neutropenia, in patients with a history of drug-induced bone marrow depression/toxicity, in patients with bone marrow depression caused by concomitant illness, radiation therapy or chemotherapy and in patients with hypereosinophilic conditions or with myeloproliferative disease. Neutropenia has been reported commonly when olanzapine and valproate are used concomitantly (see section 4.8).

Discontinuation of treatment

Acute symptoms such as sweating, insomnia, tremor, anxiety, nausea, or vomiting have been reported rarely ($\geq 0.01\%$ and $< 0.1\%$) when olanzapine is stopped abruptly.

QT interval

In clinical trials, clinically meaningful QTc prolongations (Fridericia QT correction [QTcF] ≥ 500 milliseconds [msec] at any time post baseline in patients with baseline QTcF < 500 msec) were uncommon (0.1% to 1%) in patients treated with olanzapine, with no significant differences in associated cardiac events compared to placebo. However, caution should be exercised when olanzapine is prescribed with medicines known to increase QTc interval, especially in the elderly, in patients with congenital long QT syndrome, congestive heart failure, heart hypertrophy, hypokalaemia or hypomagnesaemia.

Thromboembolism

Temporal association of olanzapine treatment and venous thromboembolism has been reported uncommonly ($\geq 0.1\%$ and $< 1\%$). A causal relationship between the occurrence of venous thromboembolism and treatment with olanzapine has not been established. However, since patients with schizophrenia often present with acquired risk factors for venous thromboembolism all possible risk factors of VTE e.g. immobilisation of patients, should be identified and preventive measures undertaken.

General CNS activity

Given the primary CNS effects of olanzapine, caution should be used when it is taken in combination with other centrally acting medicines and alcohol. As it exhibits *in vitro* dopamine antagonism, olanzapine may antagonise the effects of direct and indirect dopamine agonists.

Seizures

Olanzapine should be used cautiously in patients who have a history of seizures or are subject to factors which may lower the seizure threshold. Seizures have been reported to occur uncommonly in patients when treated with olanzapine. In most of these cases, a history of seizures or risk factors for seizures were reported.

Tardive Dyskinesia

In comparator studies of one year or less duration, olanzapine was associated with a statistically significant lower incidence of treatment emergent dyskinesia. However the risk of tardive dyskinesia increases with long term exposure, and therefore if signs or symptoms of tardive dyskinesia appear in a patient on olanzapine, a dose reduction or discontinuation should be considered.

These symptoms can temporally deteriorate or even arise after discontinuation of treatment.

Postural hypotension

Postural hypotension was infrequently observed in the elderly in olanzapine clinical trials. It is recommended that blood pressure is measured periodically in patients over 65 years.

Sudden cardiac death

In postmarketing reports with olanzapine, the event of sudden cardiac death has been reported in patients with olanzapine. In a retrospective observational cohort study, the risk of presumed sudden cardiac death in patients treated with olanzapine was approximately twice the risk in patients not using

antipsychotics. In the study, the risk of olanzapine was comparable to the risk of atypical antipsychotics included in a pooled analysis.

Paediatric population

Olanzapine is not indicated for use in the treatment of children and adolescents. Studies in patients aged 13-17 years showed various adverse reactions, including weight gain, changes in metabolic parameters and increases in prolactin levels (see sections 4.8 and 5.1).

Lactose

Olanzapine tablets contain lactose monohydrate. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

Interaction studies have only been performed in adults.

Potential interactions affecting olanzapine

Since olanzapine is metabolised by CYP1A2, substances that can specifically induce or inhibit this isoenzyme may affect the pharmacokinetics of olanzapine.

Induction of CYP1A2

The metabolism of olanzapine may be induced by smoking and carbamazepine, which may lead to reduced olanzapine concentrations. Only slight to moderate increase in olanzapine clearance has been observed. The clinical consequences are likely to be limited, but clinical monitoring is recommended and an increase of olanzapine dose may be considered if necessary (see section 4.2).

Inhibition of CYP1A2

Fluvoxamine, a specific CYP1A2 inhibitor, has been shown to significantly inhibit the metabolism of olanzapine. The mean increase in olanzapine C_{max} following fluvoxamine was 54 % in female non-smokers and 77 % in male smokers. The mean increase in olanzapine AUC was 52 % and 108 % respectively. A lower starting dose of olanzapine should be considered in patients who are using fluvoxamine or any other CYP1A2 inhibitors, such as ciprofloxacin. A decrease in the dose of olanzapine should be considered if treatment with an inhibitor of CYP1A2 is initiated.

Decreased bioavailability

Activated charcoal reduces the bioavailability of oral olanzapine by 50 % to 60 % and should be taken at least 2 hours before or after olanzapine.

Fluoxetine (a CYP2D6 inhibitor), single doses of antacid (aluminium, magnesium) or cimetidine have not been found to significantly affect the pharmacokinetics of olanzapine.

Potential for olanzapine to affect other medicinal products

Olanzapine may antagonise the effects of direct and indirect dopamine agonists.

Olanzapine does not inhibit the main CYP450 isoenzymes *in vitro* (e.g. 1A2, 2D6, 2C9, 2C19, 3A4). Thus no particular interaction is expected as verified through *in vivo* studies where no inhibition of metabolism of the following active substances was found: tricyclic antidepressant (representing mostly CYP2D6 pathway), warfarin (CYP2C9), theophylline (CYP1A2) or diazepam (CYP3A4 and 2C19).

Olanzapine showed no interaction when co-administered with lithium or biperiden.

Therapeutic monitoring of valproate plasma levels did not indicate that valproate dosage adjustment is required after the introduction of concomitant olanzapine.

General CNS activity

Caution should be exercised in patients who consume alcohol or receive medicinal products that can cause central nervous system depression.

The concomitant use of olanzapine with anti-Parkinsonian medicinal products in patients with Parkinson's disease and dementia is not recommended (see section 4.4).

QTc interval

Caution should be used if olanzapine is being administered concomitantly with medicinal products known to increase QTc interval (see section 4.4).

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no adequate and well-controlled studies in pregnant women. Patients should be advised to notify their physician if they become pregnant or intend to become pregnant during treatment with olanzapine. Nevertheless, because human experience is limited, olanzapine should be used in pregnancy only if the potential benefit justifies the potential risk to the foetus.

New born infants exposed to antipsychotics (including olanzapine) during the third trimester of pregnancy are at risk of adverse reactions including extrapyramidal and/or withdrawal symptoms that may vary in severity and duration following delivery. There have been reports of agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress, or feeding disorder. Consequently, newborns should be monitored carefully.

Breast-feeding

In a study in breast-feeding, healthy women, olanzapine was excreted in breast milk. Mean infant exposure (mg/kg) at steady state was estimated to be 1.8 % of the maternal olanzapine dose (mg/kg). Patients should be advised not to breast feed an infant if they are taking olanzapine.

Fertility

Effects on fertility are unknown (see section 5.3 for preclinical information).

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. Because olanzapine may cause somnolence and dizziness, patients should be cautioned about operating machinery, including motor vehicles.

4.8 Undesirable effects

Summary of the safety profile

Adults

The most frequently (seen in ≥ 1 % of patients) reported adverse reactions associated with the use of olanzapine in clinical trials were somnolence, weight gain, eosinophilia, elevated prolactin, cholesterol, glucose and triglyceride levels (see section 4.4), glucosuria, increased appetite, dizziness, akathisia, parkinsonism, leukopenia, neutropenia (see section 4.4), dyskinesia, orthostatic hypotension, anticholinergic effects, transient asymptomatic elevations of hepatic aminotransferases (see section 4.4), rash, asthenia, fatigue, pyrexia, arthralgia, increased alkaline phosphatase, high gamma glutamyltransferase, high uric acid, high creatine phosphokinase and oedema.

Tabulated list of adverse reactions

The following table lists the adverse reactions and laboratory investigations observed from spontaneous reporting and in clinical trials. Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness. The frequency terms listed are defined as follows: Very

common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), very rare ($< 1/10,000$), not known (cannot be estimated from the data available).

Very common	Common	Uncommon	Rare	Not known
Blood and lymphatic system disorders				
	Eosinophilia Leukopenia ¹⁰ Neutropenia ¹⁰		Thrombocytopenia ¹¹	
Immune system disorders				
		Hypersensitivity ¹¹		
Metabolism and nutrition disorders				
Weight gain ¹	Elevated cholesterol levels ^{2,3} Elevated glucose levels ⁴ Elevated triglyceride levels ^{2,5} Glucosuria Increased appetite	Development or exacerbation of diabetes occasionally associated with ketoacidosis or coma, including some fatal cases (see section 4.4) ¹¹	Hypothermia ¹²	
Nervous system disorders				
Somnolence	Dizziness Akathisia ⁶ Parkinsonism ⁶ Dyskinesia ⁶	Seizures where in most cases a history of seizures or risk factors for seizures were reported ¹¹ Dystonia (including oculogyration) ¹¹ Tardive dyskinesia ¹¹ Amnesia ⁹ Dysarthria Stuttering ¹¹ Restless legs syndrome ¹¹	Neuroleptic malignant syndrome (see section 4.4) ¹² Discontinuation symptoms ^{7, 12}	
Cardiac disorders				
		Bradycardia QT _c prolongation (see section 4.4)	Ventricular tachycardia/fibrillation, sudden death (see section 4.4) ¹¹	
Vascular disorders				
Orthostatic hypotension ¹⁰		Thromboembolism (including pulmonary embolism and deep vein thrombosis) (see section 4.4)		
Respiratory, thoracic and mediastinal disorders				
		Epistaxis ⁹		
Gastrointestinal disorders				
	Mild, transient anticholinergic effects	Abdominal distension ⁹ Salivary hypersecretion ¹¹	Pancreatitis ¹¹	

	including constipation and dry mouth			
Hepatobiliary disorders				
	Transient, asymptomatic elevations of hepatic aminotransferases (ALT, AST), especially in early treatment (see section 4.4)		Hepatitis (including hepatocellular, cholestatic or mixed liver injury) ¹¹	
Skin and subcutaneous tissue disorders				
	Rash	Photosensitivity reaction Alopecia		Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)
Musculoskeletal and connective tissue disorders				
	Arthralgia ⁹		Rhabdomyolysis ¹¹	
Renal and urinary disorders				
		Urinary incontinence, urinary retention Urinary hesitation ¹¹		
Pregnancy, puerperium and perinatal conditions				
				Drug withdrawal syndrome neonatal (see section 4.6)
Reproductive system and breast disorders				
	Erectile dysfunction in males Decreased libido in males and females	Amenorrhea Breast enlargement Galactorrhea in females Gynaecomastia/breast enlargement in males	Priapism ¹²	
General disorders and administration site conditions				
	Asthenia Fatigue Oedema Pyrexia ¹⁰			
Investigations				
Elevated plasma prolactin levels ⁸	Increased alkaline phosphatase ¹⁰ High creatine phosphokinase ¹¹ High Gamma Glutamyltransferase ¹⁰ High Uric Acid ¹⁰	Increased total bilirubin		

¹ Clinically significant weight gain was observed across all baseline Body Mass Index (BMI) categories. Following short term treatment (median duration 47 days), weight gain ≥ 7 % of baseline body weight was very common (22.2 %), ≥ 15 % was common (4.2 %) and ≥ 25 % was uncommon (0.8 %). Patients gaining ≥ 7 %, ≥ 15 % and ≥ 25 % of their baseline body weight with long-term exposure (at least 48 weeks) were very common (64.4 %, 31.7 % and 12.3 % respectively).

² Mean increases in fasting lipid values (total cholesterol, LDL cholesterol, and triglycerides) were greater in patients without evidence of lipid dysregulation at baseline.

³ Observed for fasting normal levels at baseline (< 5.17 mmol/l) which increased to high (≥ 6.2 mmol/l). Changes in total fasting cholesterol levels from borderline at baseline (≥ 5.17 -< 6.2 mmol) to high (≥ 6.2 mmol) were very common.

⁴ Observed for fasting normal levels at baseline (< 5.56 mmol/l) which increased to high (≥ 7 mmol/l). Changes in fasting glucose from borderline at baseline (≥ 5.56 -< 7 mmol/l) to high (≥ 7 mmol/l) were very common.

⁵ Observed for fasting normal levels at baseline (< 1.69 mmol/l) which increased to high (≥ 2.26 mmol/l). Changes in fasting triglycerides from borderline at baseline (≥ 1.69 mmol/l-< 2.26 mmol/l) to high (≥ 2.26 mmol/l) were very common.

⁶ In clinical trials, the incidence of Parkinsonism and dystonia in olanzapine-treated patients was numerically higher, but not statistically significantly different from placebo. Olanzapine-treated patients had a lower incidence of Parkinsonism, akathisia and dystonia compared with titrated doses of haloperidol. In the absence of detailed information on the pre-existing history of individual acute and tardive extrapyramidal movement disorders, it cannot be concluded at present that olanzapine produces less tardive dyskinesia and/or other tardive extrapyramidal syndromes.

⁷ Acute symptoms such as sweating, insomnia, tremor, anxiety, nausea and vomiting have been reported when olanzapine is stopped abruptly.

⁸ In clinical trials of up to 12 weeks, plasma prolactin concentrations exceeded the upper limit of normal range in approximately 30 % of olanzapine treated patients with normal baseline prolactin value. In the majority of these patients the elevations were generally mild, and remained below two times the upper limit of normal range.

⁹ Adverse event identified from clinical trials in the Olanzapine Integrated Database.

¹⁰ As assessed by measured values from clinical trials in the Olanzapine Integrated Database.

¹¹ Adverse event identified from spontaneous post-marketing reporting with frequency determined utilising the Olanzapine Integrated Database.

¹² Adverse event identified from spontaneous post-marketing reporting with frequency estimated at the upper limit of the 95 % confidence interval utilising the Olanzapine Integrated Database.

Long-term exposure (at least 48 weeks)

The proportion of patients who had adverse, clinically significant changes in weight gain, glucose, total/LDL/HDL cholesterol or triglycerides increased over time. In adult patients who completed 9-12 months of therapy, the rate of increase in mean blood glucose slowed after approximately 6 months.

Additional information on special populations

In clinical trials in elderly patients with dementia, olanzapine treatment was associated with a higher incidence of death and cerebrovascular adverse reactions compared to placebo (see section 4.4). Very common adverse reactions associated with the use of olanzapine in this patient group were abnormal gait and falls. Pneumonia, increased body temperature, lethargy, erythema, visual hallucinations and urinary incontinence were observed commonly.

In clinical trials in patients with drug-induced (dopamine agonist) psychosis associated with Parkinson's disease, worsening of Parkinsonian symptomatology and hallucinations were reported very commonly and more frequently than with placebo.

In one clinical trial in patients with bipolar mania, valproate combination therapy with olanzapine resulted in an incidence of neutropenia of 4.1 %; a potential contributing factor could be high plasma valproate levels. Olanzapine administered with lithium or valproate resulted in increased levels (≥ 10 %) of tremor, dry mouth, increased appetite, and weight gain. Speech disorder was also reported commonly. During treatment with olanzapine in combination with lithium or divalproex, an increase of ≥ 7 % from baseline body weight occurred in 17.4 % of patients during acute treatment (up to 6 weeks). Long-term olanzapine treatment (up to 12 months) for recurrence prevention in patients with bipolar disorder was associated with an increase of ≥ 7 % from baseline body weight in 39.9 % of patients.

Paediatric population

Olanzapine is not indicated for the treatment of children and adolescent patients below 18 years. Although no clinical studies designed to compare adolescents to adults have been conducted, data from the adolescent trials were compared to those of the adult trials.

The following table summarises the adverse reactions reported with a greater frequency in adolescent patients (aged 13-17 years) than in adult patients or adverse reactions only identified during short-term clinical trials in adolescent patients. Clinically significant weight gain (≥ 7 %) appears to occur more frequently in the adolescent population compared to adults with comparable exposures. The magnitude of weight gain and the proportion of adolescent patients who had clinically significant weight gain were greater with long-term exposure (at least 24 weeks) than with short-term exposure.

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness. The frequency terms listed are defined as follows: Very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$).

Metabolism and nutrition disorders <i>Very common:</i> Weight gain ¹³ , elevated triglyceride levels ¹⁴ , increased appetite. <i>Common:</i> Elevated cholesterol levels ¹⁵
Nervous system disorders <i>Very common:</i> Sedation (including: hypersomnia, lethargy, somnolence).
Gastrointestinal disorders <i>Common:</i> Dry mouth
Hepatobiliary disorders <i>Very common:</i> Elevations of hepatic aminotransferases (ALT/AST; see section 4.4).
Investigations <i>Very common:</i> Decreased total bilirubin, increased GGT, elevated plasma prolactin levels ¹⁶ .

¹³ Following short term treatment (median duration 22 days), weight gain ≥ 7 % of baseline body weight (kg) was very common (40.6 %), ≥ 15 % of baseline body weight was common (7.1 %) and ≥ 25 % was common (2.5 %). With long-term exposure (at least 24 weeks), 89.4 % gained ≥ 7 %, 55.3 % gained ≥ 15 % and 29.1 % gained ≥ 25 %, of their baseline body weight.

¹⁴ Observed for fasting normal levels at baseline (< 1.016 mmol/l) which increased to high (≥ 1.467 mmol/l) and changes in fasting triglycerides from borderline at baseline (≥ 1.016 mmol/l - < 1.467 mmol/l) to high (≥ 1.467 mmol/l).

¹⁵ Changes in total fasting cholesterol levels from normal at baseline (< 4.39 mmol/l) to high (≥ 5.17 mmol/l) were observed commonly. Changes in total fasting cholesterol levels from borderline at baseline (≥ 4.39 - < 5.17 mmol/l) to high (≥ 5.17 mmol/l) were very common.

¹⁶ Elevated plasma prolactin levels were reported in 47.4 % of adolescent patients.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare

professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

4.9 Overdose

Signs and symptoms

Very common symptoms in overdose (> 10 % incidence) include tachycardia, agitation/aggressiveness, dysarthria, various extrapyramidal symptoms, and reduced level of consciousness ranging from sedation to coma.

Other medically significant sequelae of overdose include delirium, convulsion, coma, possible neuroleptic malignant syndrome, respiratory depression, aspiration, hypertension or hypotension, cardiac arrhythmias (< 2 % of overdose cases) and cardiopulmonary arrest. Fatal outcomes have been reported for acute overdoses as low as 450 mg but survival has also been reported following acute overdose of approximately 2 g of oral olanzapine.

Management

There is no specific antidote for olanzapine. Induction of emesis is not recommended. Standard procedures for management of overdose may be indicated (i.e. gastric lavage, administration of activated charcoal). The concomitant administration of activated charcoal was shown to reduce the oral bioavailability of olanzapine by 50 to 60 %.

Symptomatic treatment and monitoring of vital organ function should be instituted according to clinical presentation, including treatment of hypotension and circulatory collapse and support of respiratory function. Do not use epinephrine, dopamine, or other sympathomimetic agents with beta agonist activity since beta stimulation may worsen hypotension. Cardiovascular monitoring is necessary to detect possible arrhythmias. Close medical supervision and monitoring should continue until the patient recovers.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: psycholeptics, diazepines, oxazepines, thiazepines and oxepines, ATC code: N05AH03.

Pharmacodynamic effects

Olanzapine is an antipsychotic, antimanic and mood stabilising agent that demonstrates a broad pharmacologic profile across a number of receptor systems.

In preclinical studies, olanzapine exhibited a range of receptor affinities ($K_i < 100$ nM) for serotonin 5 HT_{2A/2C}, 5 HT₃, 5 HT₆; dopamine D₁, D₂, D₃, D₄, D₅; cholinergic muscarinic receptors M₁-M₅; α_1 adrenergic; and histamine H₁ receptors. Animal behavioural studies with olanzapine indicated 5HT, dopamine, and cholinergic antagonism, consistent with the receptor-binding profile. Olanzapine demonstrated a greater *in vitro* affinity for serotonin 5HT₂ than dopamine D₂ receptors and greater 5 HT₂ than D₂ activity *in vivo* models. Electrophysiological studies demonstrated that olanzapine selectively reduced the firing of mesolimbic (A10) dopaminergic neurons, while having little effect on the striatal (A9) pathways involved in motor function. Olanzapine reduced a conditioned avoidance response, a test indicative of antipsychotic activity, at doses below those producing catalepsy, an effect indicative of motor side-effects. Unlike some other antipsychotic agents, olanzapine increases responding in an “anxiolytic” test.

In a single oral dose (10 mg) Positron Emission Tomography (PET) study in healthy volunteers, olanzapine produced a higher 5 HT_{2A} than dopamine D₂ receptor occupancy. In addition, a Single Photon Emission Computed Tomography (SPECT) imaging study in schizophrenic patients revealed

that olanzapine-responsive patients had lower striatal D₂ occupancy than some other antipsychotic- and risperidone-responsive patients, while being comparable to clozapine-responsive patients.

Clinical efficacy

In two of two placebo and two of three comparator controlled trials with over 2,900 schizophrenic patients presenting with both positive and negative symptoms, olanzapine was associated with statistically significantly greater improvements in negative as well as positive symptoms.

In a multinational, double-blind, comparative study of schizophrenia, schizoaffective, and related disorders which included 1,481 patients with varying degrees of associated depressive symptoms (baseline mean of 16.6 on the Montgomery-Asberg Depression Rating Scale), a prospective secondary analysis of baseline to endpoint mood score change demonstrated a statistically significant improvement ($p=0.001$) favouring olanzapine (-6.0) versus haloperidol (-3.1).

In patients with a manic or mixed episode of bipolar disorder, olanzapine demonstrated superior efficacy to placebo and valproate semisodium (divalproex) in reduction of manic symptoms over 3 weeks. Olanzapine also demonstrated comparable efficacy results to haloperidol in terms of the proportion of patients in symptomatic remission from mania and depression at 6 and 12 weeks. In a co-therapy study of patients treated with lithium or valproate for a minimum of 2 weeks, the addition of olanzapine 10 mg (co-therapy with lithium or valproate) resulted in a greater reduction in symptoms of mania than lithium or valproate monotherapy after 6 weeks.

In a 12-month recurrence prevention study in manic episode patients who achieved remission on olanzapine and were then randomised to olanzapine or placebo, olanzapine demonstrated statistically significant superiority over placebo on the primary endpoint of bipolar recurrence. Olanzapine also showed a statistically significant advantage over placebo in terms of preventing either recurrence into mania or recurrence into depression.

In a second 12 month recurrence prevention study in manic episode patients who achieved remission with a combination of olanzapine and lithium and were then randomised to olanzapine or lithium alone, olanzapine was statistically non-inferior to lithium on the primary endpoint of bipolar recurrence (olanzapine 30.0 %, lithium 38.3 %; $p = 0.055$).

In an 18-month co-therapy study in manic or mixed episode patients stabilised with olanzapine plus a mood stabiliser (lithium or valproate), long-term olanzapine co-therapy with lithium or valproate was not statistically significantly superior to lithium or valproate alone in delaying bipolar recurrence, defined according to syndromic (diagnostic) criteria.

Paediatric population

Controlled efficacy data in adolescents (ages 13 to 17 years) are limited to short term studies in schizophrenia (6 weeks) and mania associated with bipolar I disorder (3 weeks), involving less than 200 adolescents. Olanzapine was used as a flexible dose starting with 2.5 and ranging up to 20 mg/day. During treatment with olanzapine, adolescents gained significantly more weight compared with adults. The magnitude of changes in fasting total cholesterol, LDL cholesterol, triglycerides, and prolactin (see sections 4.4 and 4.8) were greater in adolescents than in adults. There are no controlled data on maintenance of effect or long term safety (see sections 4.4 and 4.8).

Information on long term safety is primarily limited to open-label, uncontrolled data.

5.2 Pharmacokinetic properties

Absorption

Olanzapine is well absorbed after oral administration, reaching peak plasma concentrations within 5 to 8 hours. The absorption is not affected by food. Absolute oral bioavailability relative to intravenous administration has not been determined.

Distribution

The plasma protein binding of olanzapine was about 93 % over the concentration range of about 7 to about 1000 ng/ml. Olanzapine is bound predominantly to albumin and α_1 -acid-glycoprotein.

Biotransformation

Olanzapine is metabolised in the liver by conjugative and oxidative pathways. The major circulating metabolite is the 10-N-glucuronide, which does not pass the blood brain barrier. Cytochromes P450-CYP1A2 and P450-CYP2D6 contribute to the formation of the N-desmethyl and 2-hydroxymethyl metabolites, both exhibited significantly less *in vivo* pharmacological activity than olanzapine in animal studies. The predominant pharmacologic activity is from the parent olanzapine.

Elimination

After oral administration, the mean terminal elimination half-life of olanzapine in healthy subjects varied on the basis of age and gender.

In healthy elderly (65 and over) versus non-elderly subjects, the mean elimination half-life was prolonged (51.8 versus 33.8 hr) and the clearance was reduced (17.5 versus 18.2 l/hr). The pharmacokinetic variability observed in the elderly is within the range for the non-elderly. In 44 patients with schizophrenia > 65 years of age, dosing from 5 to 20 mg/day was not associated with any distinguishing profile of adverse events.

In female versus male subjects the mean elimination half life was somewhat prolonged (36.7 versus 32.3 hr) and the clearance was reduced (18.9 versus 27.3 l/hr). However, olanzapine (5-20 mg) demonstrated a comparable safety profile in female (n=467) as in male patients (n=869).

Renal impairment

In renally impaired patients (creatinine clearance < 10 ml/min) versus healthy subjects, there was no significant difference in mean elimination half-life (37.7 versus 32.4 hr) or clearance (21.2 versus 25.0 l/hr). A mass balance study showed that approximately 57 % of radiolabelled olanzapine appeared in urine, principally as metabolites.

Hepatic impairment

A small study of the effect of impaired liver function in 6 subjects with clinically significant (Childs Pugh Classification A (n = 5) and B (n = 1)) cirrhosis revealed little effect on the pharmacokinetics of orally administered olanzapine (2.5 – 7.5 mg single dose): Subjects with mild to moderate hepatic dysfunction had slightly increased systemic clearance and faster elimination half-time compared to subjects with no hepatic dysfunction (n = 3). There were more smokers among subjects with cirrhosis (4/6; 67 %) than among subjects with no hepatic dysfunction (0/3; 0 %).

Smoking

In non-smoking versus smoking subjects (males and females) the mean elimination half-life was prolonged (38.6 versus 30.4 hr) and the clearance was reduced (18.6 versus 27.7 l/hr).

The plasma clearance of olanzapine is lower in elderly versus young subjects, in females versus males, and in non-smokers versus smokers. However, the magnitude of the impact of age, gender, or smoking on olanzapine clearance and half-life is small in comparison to the overall variability between individuals.

In a study of Caucasians, Japanese, and Chinese subjects, there were no differences in the pharmacokinetic parameters among the three populations.

Paediatric population

Adolescents (ages 13 to 17 years): The pharmacokinetics of olanzapine are similar between adolescents and adults. In clinical studies, the average olanzapine exposure was approximately 27 % higher in adolescents. Demographic differences between the adolescents and adults include a lower average body weight and fewer adolescents were smokers. Such factors possibly contribute to the higher average exposure observed in adolescents.

5.3 Preclinical safety data

Acute (single-dose) toxicity

Signs of oral toxicity in rodents were characteristic of potent neuroleptic compounds: hypoactivity, coma, tremors, clonic convulsions, salivation, and depressed weight gain. The median lethal doses were approximately 210 mg/kg (mice) and 175 mg/kg (rats). Dogs tolerated single oral doses up to 100 mg/kg without mortality. Clinical signs included sedation, ataxia, tremors, increased heart rate, laboured respiration, miosis, and anorexia. In monkeys, single oral doses up to 100 mg/kg resulted in prostration and, at higher doses, semi-consciousness.

Repeated-dose toxicity

In studies up to 3 months duration in mice and up to 1 year in rats and dogs, the predominant effects were CNS depression, anticholinergic effects, and peripheral haematological disorders. Tolerance developed to the CNS depression. Growth parameters were decreased at high doses. Reversible effects consistent with elevated prolactin in rats included decreased weights of ovaries and uterus and morphologic changes in vaginal epithelium and in mammary gland.

Haematologic toxicity

Effects on haematology parameters were found in each species, including dose-related reductions in circulating leukocytes in mice and non-specific reductions of circulating leukocytes in rats; however, no evidence of bone marrow cytotoxicity was found. Reversible neutropenia, thrombocytopenia, or anaemia developed in a few dogs treated with 8 or 10 mg/kg/day (total olanzapine exposure [AUC] is 12- to 15-fold greater than that of a man given a 12 mg dose). In cytopenic dogs, there were no adverse effects on progenitor and proliferating cells in the bone marrow.

Reproductive toxicity

Olanzapine had no teratogenic effects. Sedation affected mating performance of male rats. Oestrous cycles were affected at doses of 1.1 mg/kg (3 times the maximum human dose) and reproduction parameters were influenced in rats given 3 mg/kg (9 times the maximum human dose). In the offspring of rats given olanzapine, delays in foetal development and transient decreases in offspring activity levels were seen.

Mutagenicity

Olanzapine was not mutagenic or clastogenic in a full range of standard tests, which included bacterial mutation tests and *in vitro* and *in vivo* mammalian tests.

Carcinogenicity

Based on the results of studies in mice and rats, it was concluded that olanzapine is not carcinogenic.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core

Lactose monohydrate
Maize starch
Maize starch, pregelatinised
Crospovidone (Type A)
Magnesium stearate

Tablet coat

Polyvinyl alcohol
Titanium dioxide (E171)
Talc (E553b)
Soya lecithin (E322)

Xanthan gum (E415)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

Blisters 3 years.

Bottles 3 years. After first opening use within 90 days.

6.4 Special precautions for storage

Do not store above 25 °C.

6.5 Nature and contents of container

Cold-formed aluminium/aluminium blister in cartons of 7 (10 mg only), 10, 28, 30, 35, 56, 70 and multipacks containing 70 (2 packs of 35) film-coated tablets.

Cold-formed aluminium/aluminium perforated unit dose blisters in cartons of 28 x 1, 56 x 1 (7.5 mg only), 98 x 1 (5 mg, 7.5 mg and 10 mg only) and 100 x 1 (7.5 mg only) film-coated tablets.

High Density Polyethylene (HDPE) bottle with polypropylene screw cap containing 100 (7.5 mg, 10 mg, 15 mg and 20 mg only), 250 (2.5 mg and 5 mg only) and 500 (2.5 mg, 5 mg and 10 mg only) film-coated tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

No special requirements.

7. MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/08/475/001

EU/1/08/475/002

EU/1/08/475/003

EU/1/08/475/004

EU/1/08/475/005

EU/1/08/475/006

EU/1/08/475/007

EU/1/08/475/008

EU/1/08/475/009

EU/1/08/475/010

EU/1/08/475/011

EU/1/08/475/012
EU/1/08/475/013
EU/1/08/475/014
EU/1/08/475/015
EU/1/08/475/016
EU/1/08/475/017
EU/1/08/475/018
EU/1/08/475/019
EU/1/08/475/020
EU/1/08/475/021
EU/1/08/475/022
EU/1/08/475/023
EU/1/08/475/024
EU/1/08/475/025
EU/1/08/475/026
EU/1/08/475/027
EU/1/08/475/028
EU/1/08/475/029
EU/1/08/475/030
EU/1/08/475/031
EU/1/08/475/032
EU/1/08/475/033
EU/1/08/475/034
EU/1/08/475/035
EU/1/08/475/036
EU/1/08/475/037
EU/1/08/475/038
EU/1/08/475/039
EU/1/08/475/040
EU/1/08/475/041
EU/1/08/475/042
EU/1/08/475/043
EU/1/08/475/044
EU/1/08/475/045
EU/1/08/475/046
EU/1/08/475/047
EU/1/08/475/048
EU/1/08/475/049
EU/1/08/475/050
EU/1/08/475/051
EU/1/08/475/052
EU/1/08/475/053
EU/1/08/475/054
EU/1/08/475/055
EU/1/08/475/056
EU/1/08/475/057
EU/1/08/475/058
EU/1/08/475/059
EU/1/08/475/060
EU/1/08/475/061
EU/1/08/475/062
EU/1/08/475/063

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 7 October 2008

Date of latest renewal: 22 May 2013

10. DATE OF REVISION OF THE TEXT

{MM/YYYY}

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

ANNEX II

- A. MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturers responsible for batch release

McDermott Laboratories Ltd. t/a Gerard Laboratories
35/36 Baldoyle Industrial Estate
Grange road
Dublin 13
Ireland

Mylan Hungary Kft.
Mylan utca 1.
Komárom, 2900
Hungary

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to medical prescription.

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

Not applicable.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**CARTON OF SINGLE PACK****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viatris 2.5 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 2.5 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

10 film-coated tablets
28 film-coated tablets
30 film-coated tablets
35 film-coated tablets
56 film-coated tablets
70 film-coated tablets
28 x 1 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBERS

EU/1/08/475/001
EU/1/08/475/002
EU/1/08/475/003
EU/1/08/475/004
EU/1/08/475/035
EU/1/08/475/036
EU/1/08/475/056

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE**16. INFORMATION IN BRAILLE**

Olanzapine 2.5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**OUTER CARTON OF MULTIPACK (WITH BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viartis 2.5 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 2.5 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

Multipack 70 (2 packs of 35) film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE
--

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/047

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY
--

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 2.5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**INNER CARTON OF MULTIPACK (WITHOUT BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viatriis 2.5 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 2.5 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

35 film-coated tablets
Component of a multipack, not to be sold separately.

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/047

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 2.5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viartis 2.5 mg film-coated tablets
olanzapine

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Viartis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. OTHER

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

UNIT-DOSE BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatriis 2.5 mg tablets
olanzapine

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Viatriis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**CARTON OF SINGLE PACK****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viartis 5 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 5 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

10 film-coated tablets
28 film-coated tablets
30 film-coated tablets
35 film-coated tablets
56 film-coated tablets
70 film-coated tablets
28 x 1 film-coated tablets
98 x 1 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBERS

EU/1/08/475/007
EU/1/08/475/008
EU/1/08/475/009
EU/1/08/475/010
EU/1/08/475/037
EU/1/08/475/038
EU/1/08/475/053
EU/1/08/475/061

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE**16. INFORMATION IN BRAILLE**

Olanzapine 5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**OUTER CARTON OF MULTIPACK (WITH BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viatris 5 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 5 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

Multipack 70 (2 packs of 35) film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE
--

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/048

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY
--

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**INNER CARTON OF MULTIPACK (WITHOUT BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viartis 5 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 5 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

35 film-coated tablets
Component of a multipack, not to be sold separately

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE
--

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/048

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY
--

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatriis 5 mg film-coated tablets
olanzapine

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Viatriis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. OTHER

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

UNIT-DOSE BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatriis 5 mg tablets
olanzapine

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Viatriis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**CARTON OF SINGLE PACK****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viartis 7.5 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 7.5 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS**Film-coated tablet**

10 film-coated tablets
28 film-coated tablets
30 film-coated tablets
35 film-coated tablets
56 film-coated tablets
70 film-coated tablets
28 x 1 film-coated tablets
56 x 1 film-coated tablets
98 x 1 film-coated tablets
100 x 1 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBERS

EU/1/08/475/013
EU/1/08/475/014
EU/1/08/475/015
EU/1/08/475/016
EU/1/08/475/039
EU/1/08/475/040
EU/1/08/475/054
EU/1/08/475/055
EU/1/08/475/057
EU/1/08/475/062

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE**16. INFORMATION IN BRAILLE**

Olanzapine 7.5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**OUTER CARTON OF MULTIPACK (WITH BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viartis 7.5 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 7.5 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

Multipack 70 (2 packs of 35) film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/049

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 7.5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**INNER CARTON OF MULTIPACK (WITHOUT BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viartis 7.5 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 7.5 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

35 film-coated tablets
Component of a multipack, not to be sold separately

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/049

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 7.5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatriis 7.5 mg film-coated tablets
olanzapine

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Viatriis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. OTHER

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

UNIT-DOSE BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatriis 7.5 mg tablets
olanzapine

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Viatriis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**CARTON OF SINGLE PACK****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viartis 10 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 10 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

7 film-coated tablets
10 film-coated tablets
28 film-coated tablets
30 film-coated tablets
35 film-coated tablets
56 film-coated tablets
70 film-coated tablets
28 x 1 film-coated tablets
98 x 1 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBERS

EU/1/08/475/018
EU/1/08/475/019
EU/1/08/475/020
EU/1/08/475/021
EU/1/08/475/022
EU/1/08/475/041
EU/1/08/475/042
EU/1/08/475/058
EU/1/08/475/063

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE**16. INFORMATION IN BRAILLE**

Olanzapine 10 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**OUTER CARTON OF MULTIPACK (WITH BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viartis 10 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 10 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

Multipack 70 (2 packs of 35) film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE
--

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/050

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY
--

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 10 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**INNER CARTON OF MULTIPACK (WITHOUT BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viartis 10 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 10 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

35 film-coated tablets
Component of a multipack, not to be sold separately

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE
--

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/050

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY
--

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 10 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatriis 10 mg film-coated tablets
olanzapine

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Viatriis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. OTHER

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

UNIT-DOSE BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viartis 10 mg tablets
olanzapine

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Viartis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**CARTON OF SINGLE PACK****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viatriis 15 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 15 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

10 film-coated tablets
28 film-coated tablets
30 film-coated tablets
35 film-coated tablets
56 film-coated tablets
70 film-coated tablets
28 x 1 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBERS

EU/1/08/475/025
EU/1/08/475/026
EU/1/08/475/027
EU/1/08/475/028
EU/1/08/475/043
EU/1/08/475/044
EU/1/08/475/059

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE**16. INFORMATION IN BRAILLE**

Olanzapine 15 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**OUTER CARTON OF MULTIPACK (WITH BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viartis 15 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 15 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

Multipack 70 (2 packs of 35) film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/051

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 15 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**INNER CARTON OF MULTIPACK (WITHOUT BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viartis 15 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 15 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

35 film-coated tablets
Component of a multipack, not to be sold separately

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/051

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 15 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viartis 15 mg film-coated tablets
olanzapine

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Viartis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. OTHER

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

UNIT-DOSE BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatriis 15 mg tablets
olanzapine

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Viatriis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**CARTON OF SINGLE PACK****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viatriis 20 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 20 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

10 film-coated tablets
28 film-coated tablets
30 film-coated tablets
35 film-coated tablets
56 film-coated tablets
70 film-coated tablets
28 x 1 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBERS

EU/1/08/475/030
EU/1/08/475/031
EU/1/08/475/032
EU/1/08/475/033
EU/1/08/475/045
EU/1/08/475/046
EU/1/08/475/060

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE**16. INFORMATION IN BRAILLE**

Olanzapine 20 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**OUTER CARTON OF MULTIPACK (WITH BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viartis 20 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 20 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

Multipack 70 (2 packs of 35) film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE
--

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/052

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY
--

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 20 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**INNER CARTON OF MULTIPACK (WITHOUT BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Olanzapine Viatriis 20 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 20 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

35 film-coated tablets
Component of a multipack, not to be sold separately

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE
--

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/052

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY
--

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 20 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA
--

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatriis 20 mg film-coated tablets
olanzapine

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Viatriis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. OTHER

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS
--

UNIT-DOSE BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viartis 20 mg tablets
olanzapine

2. NAME OF THE MARKETING AUTHORISATION HOLDER
--

Viartis Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Batch

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING

BOTTLE LABEL AND CARTON LABEL

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatris 2.5 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 2.5 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

250 film-coated tablets
500 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

After first opening use within 90 days.

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/08/475/005
EU/1/08/475/006

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE**16. INFORMATION IN BRAILLE**

Olanzapine 2.5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

[Outer packaging only]

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

[Outer packaging only]

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING

BOTTLE LABEL AND CARTON LABEL

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatris 5 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each film-coated tablet contains 5 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

250 film-coated tablets
500 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

After first opening use within 90 days.

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBERS

EU/1/08/475/011
EU/1/08/475/012

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE**16. INFORMATION IN BRAILLE**

Olanzapine 5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

[Outer packaging only]

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

[Outer packaging only]

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING

BOTTLE LABEL AND CARTON LABEL

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatrix 7.5 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 7.5 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

100 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

After first opening use within 90 days.

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBER

EU/1/08/475/017

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 7.5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

[Outer packaging only]

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

[Outer packaging only]

PC
SN

NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING

BOTTLE LABEL AND CARTON LABEL

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viartis 10 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 10 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

100 film-coated tablets
500 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

After first opening use within 90 days.

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBERS

EU/1/08/475/023
EU/1/08/475/024

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 10 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

[Outer packaging only]

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

[Outer packaging only]

PC

SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING

BOTTLE LABEL AND CARTON LABEL

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatriis 15 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 15 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

100 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

After first opening use within 90 days.

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBERS

EU/1/08/475/029

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 15 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

[Outer packaging only]

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

[Outer packaging only]

PC
SN

NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING

BOTTLE LABEL AND CARTON LABEL

1. NAME OF THE MEDICINAL PRODUCT

Olanzapine Viatriis 20 mg film-coated tablets
olanzapine

2. STATEMENT OF ACTIVE SUBSTANCE

Each film-coated tablet contains 20 mg olanzapine.

3. LIST OF EXCIPIENTS

Also contains lactose monohydrate and soya lecithin.
See package leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

100 film-coated tablets

5. METHOD AND ROUTE OF ADMINISTRATION

Oral use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

After first opening use within 90 days.

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Viartis Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

12. MARKETING AUTHORISATION NUMBERS

EU/1/08/475/034

13. BATCH NUMBER

Batch

14. GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription.

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Olanzapine 20 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

[Outer packaging only]

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER – HUMAN READABLE DATA

[Outer packaging only]

PC
SN

NN

B. PACKAGE LEAFLET

Package leaflet: Information for the user

Olanzapine Viatris 2.5 mg film-coated tablets
Olanzapine Viatris 5 mg film-coated tablets
Olanzapine Viatris 7.5 mg film-coated tablets
Olanzapine Viatris 10 mg film-coated tablets
Olanzapine Viatris 15 mg film-coated tablets
Olanzapine Viatris 20 mg film-coated tablets

olanzapine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Olanzapine Viatris is and what it is used for
2. What you need to know before you take Olanzapine Viatris
3. How to take Olanzapine Viatris
4. Possible side effects
5. How to store Olanzapine Viatris
6. Contents of the pack and other information

1. What Olanzapine Viatris is and what it is used for

Olanzapine Viatris contains the active substance olanzapine. Olanzapine belongs to a group of medicines called antipsychotics and is used to treat the following conditions:

- Schizophrenia, a disease with symptoms such as hearing, seeing or sensing things which are not there, mistaken beliefs, unusual suspiciousness, and becoming withdrawn. People with this disease may also feel depressed, anxious or tense.
- Moderate to severe manic episodes, a condition with symptoms of excitement or euphoria.

Olanzapine Viatris has been shown to prevent recurrence of these symptoms in patients with bipolar disorder whose manic episode has responded to olanzapine treatment.

2. What you need to know before you take Olanzapine Viatris

Do not take Olanzapine Viatris

- if you are allergic (hypersensitive) to olanzapine, peanut or soya or any of the other ingredients of this medicine (listed in section 6). An allergic reaction may be recognised as a rash, itching, a swollen face, swollen lips or shortness of breath. If this has happened to you, tell your doctor.
- if you have been previously diagnosed with eye problems such as certain kinds of glaucoma (increased pressure in the eye).

Warnings and precautions

Talk to your doctor or pharmacist before you take Olanzapine Viatris.

- The use of Olanzapine Viatris in elderly patients with dementia is not recommended as it may have serious side effects.

- Medicines of this type may cause unusual movements mainly of the face or tongue. If this happens after you have been given Olanzapine Viatris tell your doctor.
- Very rarely, medicines of this type cause a combination of fever, faster breathing, sweating, muscle stiffness and drowsiness or sleepiness. If this happens, contact your doctor at once.
- Weight gain has been seen in patients taking Olanzapine Viatris. You and your doctor should check your weight regularly. Consider referral to a dietician or help with a diet plan if necessary.
- High blood sugar and high levels of fat (triglycerides and cholesterol) have been seen in patients taking Olanzapine Viatris. Your doctor should do blood tests to check blood sugar and certain fat levels before you start taking Olanzapine Viatris and regularly during treatment.
- Tell the doctor if you or someone else in your family has a history of blood clots, as medicines like these have been associated with the formation of blood clots.

If you suffer from any of the following illnesses tell your doctor as soon as possible:

- Stroke or “mini” stroke (temporary symptoms of stroke)
- Parkinson’s disease
- Prostate problems
- A blocked intestine (Paralytic ileus)
- Liver or kidney disease
- Blood disorders
- Heart disease
- Diabetes
- Seizures
- If you know that you may have salt depletion as a result of prolonged severe diarrhoea and vomiting (being sick) or usage of diuretics (water tablets)

If you suffer from dementia, you or your carer/relative should tell your doctor if you have ever had a stroke or “mini” stroke.

As a routine precaution, if you are over 65 years your blood pressure may be monitored by your doctor.

Children and adolescents

Olanzapine Viatris is not for patients who are under 18 years.

Other medicines and Olanzapine Viatris

Only take other medicines while you are on Olanzapine Viatris if your doctor tells you that you can. You might feel drowsy if Olanzapine Viatris is taken in combination with antidepressants or medicines taken for anxiety or to help you sleep (tranquillisers).

Tell your doctor if you are taking, have recently taken or might take any other medicines.

In particular, tell your doctor if you are taking:

- medicines for Parkinson’s disease.
- carbamazepine (an anti-epileptic and mood stabiliser), fluvoxamine (an antidepressant) or ciprofloxacin (an antibiotic) - it may be necessary to change your Olanzapine Viatris dose.

Olanzapine Viatris with alcohol

Do not drink any alcohol if you have been given Olanzapine Viatris as together with alcohol it may make you feel drowsy.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. You should not be given this medicine when breast-feeding, as small amounts of olanzapine can pass into breast milk.

The following symptoms may occur in newborn babies of mothers that have used Olanzapine Viatris in the last trimester (last three months of their pregnancy): shaking, muscle stiffness and/or weakness, sleepiness, agitation, breathing problems, and difficulty in feeding. If your baby develops any of these symptoms you may need to contact your doctor.

Driving and using machines

There is a risk of feeling drowsy when you are given Olanzapine Viatris. If this happens do not drive or operate any tools or machines. Tell your doctor.

Olanzapine Viatris contains lactose and soya lecithin

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product. The tablet film-coating contains soya lecithin. If you are allergic to peanut or soya do not take these tablets.

3. How to take Olanzapine Viatris

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Your doctor will tell you how many Olanzapine Viatris tablets to take and how long you should continue to take them. The daily dose of olanzapine is between 5 mg and 20 mg. Consult your doctor if your symptoms return but do not stop taking Olanzapine Viatris unless your doctor tells you to.

You should take your Olanzapine Viatris tablets once a day following the advice of your doctor. Try to take your tablets at the same time each day. It does not matter whether you take them with or without food. Olanzapine Viatris tablets are for oral use. You should swallow the Olanzapine Viatris tablets whole with water.

If you take more Olanzapine Viatris than you should

Patients who have taken more olanzapine than they should have experienced the following symptoms: rapid beating of the heart, agitation/aggressiveness, problems with speech, unusual movements (especially of the face or tongue) and reduced level of consciousness. Other symptoms may be: acute confusion, seizures (epilepsy), coma, a combination of fever, faster breathing, sweating, muscle stiffness and drowsiness or sleepiness, slowing of the breathing rate, aspiration, high blood pressure or low blood pressure, abnormal rhythms of the heart. Contact your doctor or hospital straight away if you experience any of the above symptoms. Show the doctor your pack of tablets.

If you forget to take Olanzapine Viatris

Take your tablets as soon as you remember. Do not take a double dose to make up for the forgotten tablet.

If you stop taking Olanzapine Viatris

Do not stop taking your tablets just because you feel better. It is important that you carry on taking Olanzapine Viatris for as long as your doctor tells you.

If you suddenly stop taking Olanzapine Viatris, symptoms such as sweating, unable to sleep, tremor, anxiety or nausea and vomiting might occur. Your doctor may suggest you to reduce the dose gradually before stopping treatment.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Tell your doctor immediately if you have:

- unusual movement (a common side effect that may affect up to 1 in 10 people) mainly of the face or tongue;
- blood clots in the veins (an uncommon side effect that may affect up to 1 in 100 people) especially in the legs (symptoms include swelling, pain, and redness in the leg), which may travel through blood vessels to the lungs causing chest pain and difficulty in breathing. If you notice any
- of these symptoms seek medical advice immediately;
- a combination of fever, faster breathing, sweating, muscle stiffness and drowsiness or sleepiness
- (the frequency of this side effect cannot be estimated from the available data)

Very common side effects (may affect more than 1 in 10 people) include weight gain; sleepiness; and increases in levels of prolactin in the blood. In the early stages of treatment, some people may feel dizzy or faint (with a slow heart rate), especially when getting up from a lying or sitting position. This will usually pass on its own but if it does not, tell your doctor.

Common side effects (may affect up to 1 in 10 people) include changes in the levels of some blood cells, circulating fats and early in treatment, temporary increases in liver enzymes; increases in the level of sugars in the blood and urine; increases in levels of uric acid and creatine phosphokinase in the blood; feeling more hungry; dizziness; restlessness; tremor; unusual movements (dyskinesias); constipation; dry mouth; rash; loss of strength; extreme tiredness; water retention leading to swelling of the hands, ankles or feet; fever; joint pain; and sexual dysfunctions such as decreased libido in males and females or erectile dysfunction in males.

Uncommon side effects (may affect up to 1 in 100 people) include hypersensitivity (e.g. swelling in the mouth and throat, itching, rash); diabetes or the worsening of diabetes, occasionally associated with ketoacidosis (ketones in the blood and urine) or coma; seizures, usually associated with a history of seizures (epilepsy); muscle stiffness or spasms (including eye movements); restless legs syndrome; problems with speech; stuttering; slow heart rate; sensitivity to sunlight; bleeding from the nose; abdominal distension; drooling; memory loss or forgetfulness; urinary incontinence; lack of ability to urinate; hair loss; absence or decrease in menstrual periods; and changes in breasts in males and females such as an abnormal production of breast milk or abnormal growth.

Rare side effects (may affect up to 1 in 1 000 people) include lowering of normal body temperature; abnormal rhythms of the heart; sudden unexplained death; inflammation of the pancreas causing severe stomach pain, fever and sickness; liver disease appearing as yellowing of the skin and white parts of the eyes; muscle disease presenting as unexplained aches and pains; and prolonged and/or painful erection.

Very rare side effects include serious allergic reactions such as Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS). DRESS appears initially as flu-like symptoms with a rash on the face and then with an extended rash, high temperature, enlarged lymph nodes, increased levels of liver enzymes seen in blood tests and an increase in a type of white blood cells (eosinophilia).

While taking olanzapine, elderly patients with dementia may suffer from stroke, pneumonia, urinary incontinence, falls, extreme tiredness, visual hallucinations, a rise in body temperature, redness of the skin and have trouble walking. Some fatal cases have been reported in this particular group of patients.

In patients with Parkinson's disease Olanzapine Viatris may worsen the symptoms.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Olanzapine Viatris

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date, which is stated on the carton or label after EXP. The expiry date refers to the last day of that month.

Do not store above 25°C.

Bottles: After first opening use within 90 days.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Olanzapine Viatris contains

- The active substance is olanzapine. Each Olanzapine Viatris tablet contains either 2.5 mg, 5 mg, 7.5 mg, 10 mg, 15 mg or 20 mg of the active substance. The exact amount is shown on your Olanzapine Viatris tablet pack.
- The other ingredients are:
(tablet core) lactose monohydrate (see section 2 ‘Olanzapine contains lactose’), maize starch, pregelatinised maize starch, croscopovidone type A, magnesium stearate and
(tablet coating) polyvinyl alcohol, titanium dioxide (E171), talc (E553b), soya lecithin (E322) (see section 2 ‘Olanzapine Viatris contains soya lecithin’), xanthan gum (E415).

What Olanzapine Viatris looks like and contents of the pack

Olanzapine Viatris 2.5 mg are round, white film-coated tablets (tablets) with sides that curve outwards, marked with “OZ over 2.5” on one side and “G” on the other side.

Olanzapine Viatris 5 mg are round, white film-coated tablets (tablets) with sides that curve outwards, marked with “OZ over 5” on one side and “G” on the other side.

Olanzapine Viatris 7.5 mg are round, white film-coated tablets (tablets) with sides that curve outwards, marked with “OZ over 7.5” on one side and “G” on the other side.

Olanzapine Viatris 10 mg are round, white film-coated tablets (tablets) with sides that curve outwards, marked with “OZ over 10” on one side and “G” on the other side.

Olanzapine Viatris 15 mg are oval-shaped, white film-coated tablets (tablets) with sides that curve outwards, marked with “OZ 15” on one side and “G” on the other side.

Olanzapine Viatris 20 mg are oval-shaped, white film-coated tablets (tablets) with sides that curve outwards, marked with “OZ 20” on one side and “G” on the other side.

Blisters:

Olanzapine Viatris 2.5 mg, 5 mg, 7.5 mg, 15 mg and 20 mg is available in packs containing 10, 28, 30, 35, 56, 70 (2 x 35 multipack) and 70 film-coated tablets.

Olanzapine Viatris 10 mg is available in packs containing 7, 10, 28, 30, 35, 56, 70 (2 x 35) (multipack) and 70 film-coated tablets.

Perforated unit dose blisters:

Olanzapine Viatris 2.5 mg, 15 mg and 20 mg is available in packs containing 28 x 1 film-coated tablets.

Olanzapine Viatris 5 mg and 10 mg is available in packs containing 28 x 1 and 98 x 1 film-coated tablets.

Olanzapine Viatris 7.5 mg is available in packs containing 28 x 1, 56 x 1, 98 x 1 and 100 x 1 film-coated tablets.

Bottles:

Olanzapine Viatris 2.5 mg and 5.0 mg is available in packs containing 250 and 500 film-coated tablets.

Olanzapine Viatris 7.5 mg, 15 mg and 20 mg is available in packs containing 100 film-coated tablets.

Olanzapine Viatris 10 mg is available in packs containing 100 and 500 film-coated tablets.

Marketing Authorisation Holder

Viatriis Limited, Damastown Industrial Park, Mulhuddart, Dublin 15, DUBLIN, Ireland.

Manufacturer

Gerard Laboratories, 35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13, Ireland.

Mylan Hungary Kft., Mylan utca 1., Komárom, 2900, Hungary.

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

Viatriis

Tél/Tel: + 32 (0)2 658 61 00

Lietuva

Viatriis UAB

Tel: +370 5 205 1288

България

Майлан ЕООД

Тел: +359 2 44 55 400

Luxembourg/Luxemburg

Viatriis

Tél/Tel: +32 (0)2 658 61 00
(Belgique/Belgien)

Česká republika

Viatriis CZ s.r.o.

Tel: + 420 222 004 400

Magyarország

Viatriis Healthcare Kft.

Tel.: + 36 1 465 2100

Danmark

Viatriis ApS

Tlf: +45 28 11 69 32

Malta

V.J. Salomone Pharma Ltd

Tel: + 356 21 22 01 74

Deutschland

Viatriis Healthcare GmbH

Tel: +49 800 0700 800

Nederland

Mylan BV

Tel: +31 (0)20 426 3300

Eesti

Viatriis OÜ

Tel: + 372 6363 052

Norge

Viatriis AS

Tlf: + 47 66 75 33 00

Ελλάδα

Viatriis Hellas Ltd

Τηλ: +30 2100 100 002

Österreich

Viatriis Austria GmbH

Tel: +43 1 86390

España

Viatriis Pharmaceuticals, S.L.

Tel: + 34 900 102 712

Polska

Viatriis Healthcare Sp. z o.o.

Tel.: + 48 22 546 64 00

France

Viatriis Santé

Tél: +33 4 37 25 75 00

Portugal

Mylan, Lda.

Tel: + 351 214 127 200

Hrvatska

Viatriis Hrvatska d.o.o.

România

BGP Products SRL

Tel: +385 1 23 50 599

Ireland

Viatriis Limited

Tel: +353 1 8711600

Ísland

Icepharma hf.

Sími: +354 540 8000

Italia

Viatriis Italia S.r.l.

Tel: + 39 (0) 2 612 46921

Κύπρος

CPO Pharmaceuticals Limited

Τηλ: +357 22863100

Latvija

Viatriis SIA

Tel: +371 676 055 80

Tel: +40 372 579 000

Slovenija

Viatriis d.o.o.

Tel: + 386 1 23 63 180

Slovenská republika

Viatriis Slovakia s r.o.

Tel: +421 2 32 199 100

Suomi/Finland

Viatriis Oy

Puh/Tel: +358 20 720 9555

Sverige

Viatriis AB

Tel: +46 (0)8 630 19 00

This leaflet was last revised in {month YYYY}.

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu>.