

Medicinal Product no longer authorised

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Pregabalin Mylan Pharma 25 mg hard capsules  
Pregabalin Mylan Pharma 50 mg hard capsules  
Pregabalin Mylan Pharma 75 mg hard capsules  
Pregabalin Mylan Pharma 100 mg hard capsules  
Pregabalin Mylan Pharma 150 mg hard capsules  
Pregabalin Mylan Pharma 200 mg hard capsules  
Pregabalin Mylan Pharma 225 mg hard capsules  
Pregabalin Mylan Pharma 300 mg hard capsules

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Pregabalin Mylan Pharma 25 mg hard capsules

Each hard capsule contains 25 mg of pregabalin.

Pregabalin Mylan Pharma 50 mg hard capsules

Each hard capsule contains 50 mg of pregabalin.

Pregabalin Mylan Pharma 75 mg hard capsules

Each hard capsule contains 75 mg of pregabalin.

Pregabalin Mylan Pharma 100 mg hard capsules

Each hard capsule contains 100 mg of pregabalin.

Pregabalin Mylan Pharma 150 mg hard capsules

Each hard capsule contains 150 mg of pregabalin.

Pregabalin Mylan Pharma 200 mg hard capsules

Each hard capsule contains 200 mg of pregabalin.

Pregabalin Mylan Pharma 225 mg hard capsules

Each hard capsule contains 225 mg of pregabalin.

Pregabalin Mylan Pharma 300 mg hard capsules

Each hard capsule contains 300 mg of pregabalin.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Hard capsule.

Pregabalin Mylan Pharma 25 mg hard capsules

No. 4, light peach opaque cap and white opaque body, hard-shell gelatin capsule filled with white to off-white powder. The capsule is axially printed with MYLAN over PB25 in black ink on cap and body.

#### Pregabalin Mylan Pharma 50 mg hard capsules

The 50 mg capsules are No. 3, dark peach opaque cap and white opaque body, hard-shell gelatin capsule filled with white to off-white powder. The capsule is axially printed with MYLAN over PB50 in black ink on cap and body.

#### Pregabalin Mylan Pharma 75 mg hard capsules

The 75 mg capsules are No. 4, light peach opaque cap and light peach opaque body, hard-shell gelatin capsule filled with white to off-white powder. The capsule is axially printed with MYLAN over PB75 in black ink on cap and body.

#### Pregabalin Mylan Pharma 100 mg hard capsules

The 100 mg capsules are No. 3, dark peach opaque cap and dark peach opaque body, hard-shell gelatin capsules filled with white to off-white powder. The capsule is axially printed with MYLAN over PB100 in black ink on cap and body.

#### Pregabalin Mylan Pharma 150 mg hard capsules

The 150 mg capsules are No. 2, light peach opaque cap and white opaque body, hard-shell gelatin capsules filled with white to off-white powder. The capsule is axially printed with MYLAN over PB150 in black ink on cap and body.

#### Pregabalin Mylan Pharma 200 mg hard capsules

The 200 mg capsules are No. 1, light peach opaque cap and light peach opaque body, hard-shell gelatin capsules filled with white to off-white powder. The capsule is axially printed with MYLAN over PB200 in black ink on cap and body.

#### Pregabalin Mylan Pharma 225 mg hard capsules

The 225 mg capsules are No. 1, dark peach opaque cap and dark peach opaque body, hard-shell gelatin capsules filled with white to off-white powder. The capsule is axially printed with MYLAN over PB225 in black ink on cap and body.

#### Pregabalin Mylan Pharma 300 mg hard capsules

The 300 mg capsules are No. 0, light peach opaque cap and white opaque body, hard-shell gelatin capsules filled with white to off-white powder. The capsule is axially printed with MYLAN over PB300 in black ink on cap and body.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

#### Epilepsy

Pregabalin Mylan Pharma is indicated as adjunctive therapy in adults with partial seizures with or without secondary generalisation.

## Generalised Anxiety Disorder

Pregabalin Mylan Pharma is indicated for the treatment of Generalised Anxiety Disorder (GAD) in adults.

### **4.2 Posology and method of administration**

#### Posology

The dose range is 150 to 600 mg per day given in either two or three divided doses.

#### *Epilepsy*

Pregabalin treatment can be started with a dose of 150 mg per day given as two or three divided doses. Based on individual patient response and tolerability, the dose may be increased to 300 mg per day after 1 week. The maximum dose of 600 mg per day may be achieved after an additional week.

#### *Generalised Anxiety Disorder*

The dose range is 150 to 600 mg per day given as two or three divided doses. The need for treatment should be reassessed regularly.

Pregabalin treatment can be started with a dose of 150 mg per day. Based on individual patient response and tolerability, the dose may be increased to 300 mg per day after 1 week. Following an additional week the dose may be increased to 450 mg per day. The maximum dose of 600 mg per day may be achieved after an additional week.

#### *Discontinuation of pregabalin*

In accordance with current clinical practice, if pregabalin has to be discontinued, it is recommended this should be done gradually over a minimum of 1 week independent of the indication (see sections 4.4 and 4.8).

#### Special populations

##### *Renal impairment*

Pregabalin is eliminated from the systemic circulation primarily by renal excretion as unchanged drug. As pregabalin clearance is directly proportional to creatinine clearance (see section 5.2), dose reduction in patients with compromised renal function must be individualised according to creatinine clearance (CL<sub>cr</sub>), as indicated in Table 1 determined using the following formula:

$$CL_{cr}(\text{ml/min}) = \left[ \frac{1.23 \times [140 - \text{age (years)}] \times \text{weight (kg)}}{\text{serum creatinine } (\mu\text{mol/l)}} \right] (\times 0.85 \text{ for female patients})$$

Pregabalin is removed effectively from plasma by haemodialysis (50% of drug in 4 hours). For patients receiving haemodialysis, the pregabalin daily dose should be adjusted based on renal function. In addition to the daily dose, a supplementary dose should be given immediately following every 4 hour haemodialysis treatment (see Table 1).

Table 1. Pregabalin dose adjustment based on renal function

Creatinine clearance (CL <sub>cr</sub> ) (ml/min)	Total pregabalin daily dose *	Dose regimen
---	-------------------------------	--------------

	Starting dose (mg/day)	Maximum dose (mg/day)	
≥ 60	150	600	BID or TID
≥30 - <60	75	300	BID or TID
≥15 - <30	25-50	150	Once Daily or BID
< 15	25	75	Once Daily
<b>Supplementary dosage following haemodialysis (mg)</b>			
	25	100	Single dose <sup>+</sup>

TID = Three divided doses

BID = Two divided doses

\* Total daily dose (mg/day) should be divided as indicated by dose regimen to provide mg/dose

<sup>+</sup> Supplementary dose is a single additional dose

#### *Hepatic impairment*

No dose adjustment is required for patients with hepatic impairment (see section 5.2).

#### *Paediatric population*

The safety and efficacy of Pregabalin Mylan Pharma in children below the age of 12 years and in adolescents (12-17 years of age) have not been established. Currently available data are described in section 4.8, 5.1 and 5.2 but no recommendation on a posology can be made.

#### *Elderly*

Elderly patients may require a dose reduction of pregabalin due to a decreased renal function (see patients with renal impairment).

#### Method of administration

Pregabalin Mylan Pharma may be taken with or without food.

Pregabalin Mylan Pharma is for oral use only.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

### **4.4 Special warnings and precautions for use**

#### Diabetic patients

In accordance with current clinical practice, some diabetic patients who gain weight on pregabalin treatment may need to adjust hypoglycaemic medicinal products.

#### Hypersensitivity reactions

There have been reports in the post-marketing experience of hypersensitivity reactions, including cases of angioedema. Pregabalin should be discontinued immediately if symptoms of angioedema, such as facial, perioral, or upper airway swelling occur.

#### Dizziness, somnolence, loss of consciousness, confusion and mental impairment

Pregabalin treatment has been associated with dizziness and somnolence, which could increase the occurrence of accidental injury (fall) in the elderly population. There have also been post-marketing reports of loss of consciousness, confusion and mental impairment. Therefore, patients should be advised to exercise caution until they are familiar with the potential effects of the medicinal product.

#### Vision-related effects

In controlled trials, a higher proportion of patients treated with pregabalin reported blurred vision than did patients treated with placebo which resolved in a majority of cases with continued dosing. In the clinical studies where ophthalmologic testing was conducted, the incidence of visual acuity reduction and visual field changes was greater in pregabalin-treated patients than in placebo-treated patients; the incidence of fundoscopic changes was greater in placebo-treated patients (see section 5.1).

In the post-marketing experience, visual adverse reactions have also been reported, including loss of vision, visual blurring or other changes of visual acuity, many of which were transient. Discontinuation of pregabalin may result in resolution or improvement of these visual symptoms.

#### Renal failure

Cases of renal failure have been reported and in some cases discontinuation of pregabalin did show reversibility of this adverse reaction.

#### Withdrawal of concomitant anti-epileptic medicinal products

There are insufficient data for the withdrawal of concomitant anti-epileptic medicinal products, once seizure control with pregabalin in the add-on situation has been reached, in order to reach monotherapy on pregabalin.

#### Withdrawal symptoms

After discontinuation of short-term and long-term treatment with pregabalin, withdrawal symptoms have been observed in some patients. The following events have been mentioned: insomnia, headache, nausea, anxiety, diarrhoea, flu syndrome, nervousness, depression, pain, convulsion, hyperhidrosis and dizziness, suggestive of physical dependence. The patient should be informed about this at the start of the treatment.

Convulsions, including status epilepticus and grand mal convulsions, may occur during pregabalin use or shortly after discontinuing pregabalin.

Concerning discontinuation of long-term treatment of pregabalin, data suggest that the incidence and severity of withdrawal symptoms may be dose-related.

#### Congestive heart failure

There have been post-marketing reports of congestive heart failure in some patients receiving pregabalin. These reactions are mostly seen in elderly cardiovascular compromised patients during pregabalin treatment for a neuropathic indication. Pregabalin should be used with caution in these patients. Discontinuation of pregabalin may resolve the reaction.

#### Adverse reactions in patients with spinal cord injury

The incidence of adverse reactions in general, central nervous system adverse reactions and especially somnolence was increased in the treatment of central neuropathic pain due to spinal cord injury. This may be attributed to an additive effect due to concomitant medicinal products (e.g. anti-spasticity agents) needed for this condition.

#### Respiratory depression

There have been reports of severe respiratory depression in relation to pregabalin use. Patients with compromised respiratory function, respiratory or neurological disease, renal impairment, concomitant use of CNS depressants and the elderly may be at higher risk of experiencing this severe adverse reaction. Dose adjustments may be necessary in these patients (see section 4.2).

#### Suicidal ideation and behaviour

Suicidal ideation and behaviour have been reported in patients treated with anti-epileptic agents in several indications. A meta-analysis of randomised placebo controlled studies of anti-epileptic drugs has also shown a small increased risk of suicidal ideation and behaviour. The mechanism of this risk is not known and the available data do not exclude the possibility of an increased risk for pregabalin.

Therefore patients should be monitored for signs of suicidal ideation and behaviours and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of suicidal ideation or behaviour emerge.

#### Reduced lower gastrointestinal tract function

There are post-marketing reports of events related to reduced lower gastrointestinal tract function (e.g. intestinal obstruction, paralytic ileus, constipation) when pregabalin was co-administered with medications that have the potential to produce constipation, such as opioid analgesics. When pregabalin and opioids will be used in combination, measures to prevent constipation may be considered (especially in female patients and elderly).

#### Concomitant use with opioids

Caution is advised when prescribing pregabalin concomitantly with opioids due to risk of CNS depression (see section 4.5). In a case-control study of opioid users, those patients who took pregabalin concomitantly with an opioid had an increased risk for opioid-related death compared to opioid use alone (adjusted odds ratio [aOR], 1.68 [95% CI, 1.19 – 2.36]). This increased risk was observed at low doses of pregabalin ( $\leq 300$  mg, aOR 1.52 [95% CI, 1.04 – 2.22]) and there was a trend for a greater risk at high doses of pregabalin ( $> 300$  mg, aOR 2.51 [95% CI 1.24 – 5.06]).

#### Misuse, abuse potential or dependence

Cases of misuse, abuse and dependence have been reported. Caution should be exercised in patients with a history of substance abuse and the patient should be monitored for symptoms of pregabalin misuse, abuse or dependence (development of tolerance, dose escalation, drug-seeking behaviour have been reported).

#### Encephalopathy

Cases of encephalopathy have been reported, mostly in patients with underlying conditions that may precipitate encephalopathy.

#### Sodium content

This medicine contains less than 1 mmol sodium (23 mg) per capsule. That is to say essentially 'sodium-free'

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Since pregabalin is predominantly excreted unchanged in the urine, undergoes negligible metabolism in humans (<2% of a dose recovered in urine as metabolites), does not inhibit drug metabolism *in vitro*, and is not bound to plasma proteins, it is unlikely to produce, or be subject to, pharmacokinetic interactions.

##### *In vivo* studies and population pharmacokinetic analysis

Accordingly, in *in vivo* studies no clinically relevant pharmacokinetic interactions were observed between pregabalin and phenytoin, carbamazepine, valproic acid, lamotrigine, gabapentin, lorazepam, oxycodone or ethanol. Population pharmacokinetic analysis indicated that oral antidiabetics, diuretics, insulin, phenobarbital, tiagabine and topiramate had no clinically significant effect on pregabalin clearance.

##### Oral contraceptives, norethisterone and/or ethinyl oestradiol

Co-administration of pregabalin with the oral contraceptives norethisterone and/or ethinyl oestradiol does not influence the steady-state pharmacokinetics of either substance.

##### Central nervous system influencing medical products

Pregabalin may potentiate the effects of ethanol and lorazepam.

In the postmarketing experience, there are reports of respiratory failure, coma and deaths in patients taking pregabalin and opioids and/or other central nervous system (CNS) depressant medicinal products. Pregabalin appears to be additive in the impairment of cognitive and gross motor function caused by oxycodone.

##### Interactions and the elderly

No specific pharmacodynamic interaction studies were conducted in elderly volunteers. Interaction studies have only been performed in adults.

#### **4.6 Fertility, pregnancy and lactation**

##### Women of childbearing potential/Contraception in males and females

As the potential risk for humans is unknown, effective contraception must be used in women of child bearing potential.

##### Pregnancy

There are no adequate data from the use of pregabalin in pregnant women.

Studies in animals have shown reproductive toxicity (see section 5.3). The potential risk for humans is unknown.

Pregabalin Mylan Pharma should not be used during pregnancy unless clearly necessary (if the benefit to the mother clearly outweighs the potential risk to the foetus).



## Breast-feeding

Pregabalin is excreted into human milk (see section 5.2). The effect of pregabalin on newborns/infants is unknown. A decision must be made whether to discontinue breast-feeding or to discontinue pregabalin therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

## Fertility

There are no clinical data on the effects of pregabalin on female fertility.

In a clinical trial to assess the effect of pregabalin on sperm motility, healthy male subjects were exposed to pregabalin at a dose of 600 mg/day. After 3 months of treatment, there were no effects on sperm motility.

A fertility study in female rats has shown adverse reproductive effects. Fertility studies in male rats have shown adverse reproductive and developmental effects. The clinical relevance of these findings is unknown (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

Pregabalin Mylan Pharma may have minor or moderate influence on the ability to drive and use machines. Pregabalin Mylan Pharma may cause dizziness and somnolence and therefore may influence the ability to drive or use machines. Patients are advised not to drive, operate complex machinery or engage in other potentially hazardous activities until it is known whether this medicinal product affects their ability to perform these activities.

### **4.8 Undesirable effects**

#### Summary of the safety profile

The pregabalin clinical programme involved over 8900 patients exposed to pregabalin, of whom over 5600 were in double-blind placebo controlled trials. The most commonly reported adverse reactions were dizziness and somnolence. Adverse reactions were usually mild to moderate in intensity. In all controlled studies, the discontinuation rate due to adverse reactions was 12% for patients receiving pregabalin and 5% for patients receiving placebo. The most common adverse reactions resulting in discontinuation from pregabalin treatment groups were dizziness and somnolence.

#### Tabulated list of adverse reactions

In table 2 below all adverse reactions, which occurred at an incidence greater than placebo and in more than one patient, are listed by class and frequency (very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ), not known (cannot be estimated from the available data)). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

The adverse reactions listed may also be associated with the underlying disease and/or concomitant medicinal products.

The incidence of adverse reactions in general, CNS adverse reactions and especially somnolence was increased in the treatment of central neuropathic pain due to spinal cord injury (see section 4.4).

Additional reactions reported from post-marketing experience are included in italics in the list below.

**Table 2. Pregabalin Adverse Drug Reactions**

<b>System Organ Class</b>	<b>Adverse drug reactions</b>
<b>Infections and infestations</b>	
Common	Nasopharyngitis
<b>Blood and lymphatic system disorders</b>	
Uncommon	Neutropaenia
<b>Immune system disorders</b>	
Uncommon	<i>Hypersensitivity</i>
Rare	<i>Angioedema, allergic reaction</i>
<b>Metabolism and nutrition disorders</b>	
Common	Appetite increased
Uncommon	Anorexia, hypoglycaemia
<b>Psychiatric disorders</b>	
Common	Euphoric mood, confusion, irritability, disorientation, insomnia, libido decreased
Uncommon	Hallucination, panic attack, restlessness, agitation, depression, depressed mood, elevated mood, <i>aggression</i> , mood swings, depersonalisation, word finding difficulty, abnormal dreams, libido increased, anorgasmia, apathy
Rare	Disinhibition
<b>Nervous system disorders</b>	
Very common	Dizziness, somnolence, headache
Common	Ataxia, coordination abnormal, tremor, dysarthria, amnesia, memory impairment, disturbance in attention, paraesthesia, hypoaesthesia, sedation, balance disorder, lethargy
Uncommon	Syncope, stupor, myoclonus, <i>loss of consciousness</i> , psychomotor hyperactivity, dyskinesia, dizziness postural, intention tremor, nystagmus, cognitive disorder, <i>mental impairment</i> , speech disorder, hyporeflexia, hyperaesthesia, burning sensation, ageusia, <i>malaise</i>
Rare	

	<i>Convulsion, parosmia, hypokinesia, dysgraphia</i>
<b>Eye disorders</b>	
Common	Vision blurred, diplopia
Uncommon	Peripheral vision loss, visual disturbance, eye swelling, visual field defect, visual acuity reduced, eye pain, asthenopia, photopsia, dry eye, lacrimation increased, eye irritation
Rare	<i>Vision loss, keratitis, oscillopsia, altered visual depth perception, mydriasis, strabismus, visual brightness</i>
<b>Ear and labyrinth disorders</b>	
Common	Vertigo
Uncommon	Hyperacusis
<b>Cardiac disorders</b>	
Uncommon	Tachycardia, atrioventricular block first degree, sinus bradycardia, <i>congestive heart failure</i>
Rare	<i>QT prolongation, sinus tachycardia, sinus arrhythmia</i>
<b>Vascular disorders</b>	
Uncommon	Hypotension, hypertension, hot flushes, flushing, peripheral coldness
<b>Respiratory, thoracic and mediastinal disorders</b>	
Uncommon	Dyspnoea, epistaxis, cough, nasal congestion, rhinitis, snoring, nasal dryness
Rare	<i>Pulmonary oedema, throat tightness</i>
Not known	Respiratory depression
<b>Gastrointestinal disorders</b>	
Common	Vomiting, <i>nausea</i> , constipation, <i>diarrhoea</i> , flatulence, abdominal distension, dry mouth
Uncommon	Gastroesophageal reflux disease, salivary hypersecretion, hypoesthesia oral
Rare	Ascites, pancreatitis, <i>Swollen tongue</i> , dysphagia
<b>Hepatobiliary disorders</b>	

Uncommon	Elevated liver enzymes*
Rare	Jaundice
Very rare	Hepatic failure, hepatitis
<b>Skin and subcutaneous tissue disorders</b>	
Uncommon	Rash papular, urticaria, hyperhidrosis, <i>pruritus</i>
Rare	<i>Stevens Johnson syndrome</i> , cold sweat
<b>Musculoskeletal and connective tissue disorders</b>	
Common	Muscle cramp, arthralgia, back pain, pain in limb, cervical spasm
Uncommon	Joint swelling, myalgia, muscle twitching, neck pain, muscle stiffness
Rare	Rhabdomyolysis
<b>Renal and urinary disorders</b>	
Uncommon	Urinary incontinence, dysuria
Rare	Renal failure, oliguria, <i>urinary retention</i>
<b>Reproductive system and breast disorders</b>	
Common	Erectile dysfunction
Uncommon	Sexual dysfunction, ejaculation delayed, dysmenorrhoea, breast pain
Rare	Amenorrhoea, breast discharge, breast enlargement, <i>gynaecomastia</i>
<b>General disorders and administration site conditions</b>	
Common	Oedema peripheral, oedema, gait abnormal, fall, feeling drunk, feeling abnormal, fatigue
Uncommon	Generalised oedema, <i>face oedema</i> , chest tightness, pain, pyrexia, thirst, chills, asthenia
<b>Investigations</b>	
Common	Weight increased
Uncommon	Blood creatine phosphokinase increased, blood glucose increased, platelet count decreased, blood creatinine increased, blood potassium decreased, weight decreased

Rare	White blood cell count decreased
------	----------------------------------

\* Alanine aminotransferase increased (ALT) and aspartate aminotransferase increased (AST).

After discontinuation of short-term and long-term treatment with pregabalin withdrawal symptoms have been observed in some patients. The following reactions have been mentioned: insomnia, headache, nausea, anxiety, diarrhoea, flu syndrome, convulsions, nervousness, depression, pain, hyperhidrosis and dizziness, suggestive of physical dependence. The patient should be informed about this at the start of the treatment.

Concerning discontinuation of long-term treatment of pregabalin, data suggest that the incidence and severity of withdrawal symptoms may be dose-related.

#### Paediatric population

The pregabalin safety profile observed in five paediatric studies in patients with partial seizures with or without secondary generalization (12-week efficacy and safety study in patients 4 to 16 years of age, n=295; 14-day efficacy and safety study in patients 1 month to younger than 4 years of age, n=175; pharmacokinetic and tolerability study, n=65; and two 1 year open label follow on safety studies, n=54 and n=431) was similar to that observed in the adult studies of patients with epilepsy. The most common adverse events observed in the 12-week study with pregabalin treatment were somnolence, pyrexia, upper respiratory tract infection, increased appetite, weight increased, and nasopharyngitis. The most common adverse events observed in the 14-day study with pregabalin treatment were somnolence, upper respiratory tract infection, and pyrexia (see sections 4.2, 5.1 and 5.2).

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

### **4.9 Overdose**

#### Symptoms

In the post-marketing experience, the most commonly reported adverse reactions observed when pregabalin was taken in overdose included somnolence, confusional state, agitation, and restlessness. Seizures were also reported.

In rare occasions, cases of coma have been reported.

#### Management

Treatment of pregabalin overdose should include general supportive measures and may include haemodialysis if necessary (see section 4.2 Table 1).

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Anti-epileptics, other anti-epileptics, ATC code: N03AX16.

The active substance, pregabalin, is a gamma-aminobutyric acid analogue [(S)-3-(aminomethyl)-5-methylhexanoic acid].

#### Mechanism of action

Pregabalin binds to an auxiliary subunit ( $\alpha_2$ - $\delta$  protein) of voltage-gated calcium channels in the central nervous system.

#### Clinical efficacy and safety

##### *Epilepsy*

##### *Adjunctive Treatment*

Pregabalin has been studied in 3 controlled clinical trials of 12 week duration with either BID or TID dosing. Overall, the safety and efficacy profiles for BID and TID dosing regimens were similar.

A reduction in seizure frequency was observed by Week 1.

##### *Paediatric population*

The efficacy and safety of pregabalin as adjunctive treatment for epilepsy in paediatric patients below the age of 12 and adolescents has not been established. The adverse events observed in a pharmacokinetic and tolerability study that enrolled patients from 3 months to 16 years of age (n=65) with partial onset seizures were similar to those observed in adults. Results of a 12-week placebo-controlled study of 295 paediatric patients aged 4 to 16 years and a 14-day placebo-controlled study of 175 paediatric patients aged 1 month to younger than 4 years of age performed to evaluate the efficacy and safety of pregabalin as adjunctive therapy for the treatment of partial onset seizures and two 1 year open label safety studies in 54 and 431 paediatric patients respectively, from 3 months to 16 years of age with epilepsy indicate that the adverse events of pyrexia and upper respiratory infections were observed more frequently than in adult studies of patients with epilepsy (see sections 4.2, 4.8 and 5.2).

In the 12-week placebo-controlled study, paediatric patients (4 to 16 years of age) were assigned to pregabalin 2.5 mg/kg/day (maximum, 150 mg/day), pregabalin 10 mg/kg/day (maximum, 600 mg/day), or placebo. The percentage of subjects with at least a 50% reduction in partial onset seizures as compared to baseline was 40.6% of subjects treated with pregabalin 10 mg/kg/day (p=0.0068 versus placebo), 29.1% of subjects treated with pregabalin 2.5 mg/kg/day (p=0.2600 versus placebo) and 22.6% of those receiving placebo.

In the 14-day placebo-controlled study, paediatric patients (1 month to younger than 4 years of age) were assigned to pregabalin 7 mg/kg/day, pregabalin 14 mg/kg/day, or placebo. Median 24-hour seizure frequencies at baseline and at the final visit were 4.7 and 3.8 for pregabalin 7 mg/kg/day, 5.4 and 1.4 for pregabalin 14 mg/kg/day, and 2.9 and 2.3 for placebo, respectively. Pregabalin 14 mg/kg/day significantly reduced the log-transformed partial onset seizure frequency versus placebo (p=0.0223); pregabalin 7 mg/kg/day did not show improvement relative to placebo.

### *Monotherapy (newly diagnosed patients)*

Pregabalin has been studied in 1 controlled clinical trial of 56 week duration with BID dosing. Pregabalin did not achieve non-inferiority to lamotrigine based on the 6-month seizure freedom endpoint. Pregabalin and lamotrigine were similarly safe and well tolerated.

### *Generalised Anxiety Disorder*

Pregabalin has been studied in 6 controlled trials of 4-6 week duration, an elderly study of 8 week duration and a long-term relapse prevention study with a double blind relapse prevention phase of 6 months duration.

Relief of the symptoms of GAD as reflected by the Hamilton Anxiety Rating Scale (HAM-A) was observed by Week 1.

In controlled clinical trials (4-8 week duration) 52% of the pregabalin treated patients and 38% of the patients on placebo had at least a 50% improvement in HAM-A total score from baseline to endpoint.

In controlled trials, a higher proportion of patients treated with pregabalin reported blurred vision than did patients treated with placebo which resolved in a majority of cases with continued dosing.

Ophthalmologic testing (including visual acuity testing, formal visual field testing and dilated fundoscopic examination) was conducted in over 3600 patients within controlled clinical trials. In these patients, visual acuity was reduced in 6.5% of patients treated with pregabalin, and 4.8% of placebo-treated patients. Visual field changes were detected in 12.4% of pregabalin-treated, and 11.7% of placebo-treated patients. Fundoscopic changes were observed in 1.7% of pregabalin-treated and 2.1% of placebo-treated patients.

## **5.2 Pharmacokinetic properties**

Pregabalin steady-state pharmacokinetics are similar in healthy volunteers, patients with epilepsy receiving anti-epileptic drugs and patients with chronic pain.

### Absorption

Pregabalin is rapidly absorbed when administered in the fasted state, with peak plasma concentrations occurring within 1 hour following both single and multiple dose administration. Pregabalin oral bioavailability is estimated to be  $\geq 90\%$  and is independent of dose. Following repeated administration, steady state is achieved within 24 to 48 hours. The rate of pregabalin absorption is decreased when given with food resulting in a decrease in  $C_{max}$  by approximately 25-30% and a delay in  $t_{max}$  to approximately 2.5 hours. However, administration of pregabalin with food has no clinically significant effect on the extent of pregabalin absorption.

### Distribution

In preclinical studies, pregabalin has been shown to cross the blood brain barrier in mice, rats, and monkeys. Pregabalin has been shown to cross the placenta in rats and is present in the milk of lactating rats. In humans, the apparent volume of distribution of pregabalin following oral administration is approximately 0.56 l/kg. Pregabalin is not bound to plasma proteins.

### Biotransformation

Pregabalin undergoes negligible metabolism in humans. Following a dose of radiolabelled pregabalin, approximately 98% of the radioactivity recovered in the urine was unchanged pregabalin. The N-methylated derivative of pregabalin, the major metabolite of pregabalin found

in urine, accounted for 0.9% of the dose. In preclinical studies, there was no indication of racemisation of pregabalin S-enantiomer to the R-enantiomer.

#### Elimination

Pregabalin is eliminated from the systemic circulation primarily by renal excretion as unchanged drug. Pregabalin mean elimination half-life is 6.3 hours. Pregabalin plasma clearance and renal clearance are directly proportional to creatinine clearance (see section 5.2 Renal impairment).

Dose adjustment in patients with reduced renal function or undergoing haemodialysis is necessary (see section 4.2 Table 1).

#### Linearity/non-linearity

Pregabalin pharmacokinetics are linear over the recommended daily dose range. Inter-subject pharmacokinetic variability for pregabalin is low (<20%). Multiple dose pharmacokinetics are predictable from single-dose data. Therefore, there is no need for routine monitoring of plasma concentrations of pregabalin.

#### Gender

Clinical trials indicate that gender does not have a clinically significant influence on the plasma concentrations of pregabalin.

#### Renal impairment

Pregabalin clearance is directly proportional to creatinine clearance. In addition, pregabalin is effectively removed from plasma by haemodialysis (following a 4 hour haemodialysis treatment plasma pregabalin concentrations are reduced by approximately 50%). Because renal elimination is the major elimination pathway, dose reduction in patients with renal impairment and dose supplementation following haemodialysis is necessary (see section 4.2 Table 1).

#### Hepatic impairment

No specific pharmacokinetic studies were carried out in patients with impaired liver function. Since pregabalin does not undergo significant metabolism and is excreted predominantly as unchanged drug in the urine, impaired liver function would not be expected to significantly alter pregabalin plasma concentrations.

#### Paediatric population

Pregabalin pharmacokinetics were evaluated in paediatric patients with epilepsy (age groups: 1 to 23 months, 2 to 6 years, 7 to 11 years and 12 to 16 years) at dose levels of 2.5, 5, 10 and 15 mg/kg/day in a pharmacokinetic and tolerability study.

After oral administration of pregabalin in paediatric patients in the fasted state, in general, time to reach peak plasma concentration was similar across the entire age group and occurred 0.5 hours to 2 hours postdose.

Pregabalin  $C_{max}$  and AUC parameters increased in a linear manner with increasing dose within each age group. The AUC was lower by 30% in paediatric patients below a weight of 30 kg due to an increased body weight adjusted clearance of 43% for these patients in comparison to patients weighing  $\geq 30$  kg.



Pregabalin terminal half-life averaged about 3 to 4 hours in paediatric patients up to 6 years of age, and 4 to 6 hours in those 7 years of age and older.

Population pharmacokinetic analysis showed that creatinine clearance was a significant covariate of pregabalin oral clearance, body weight was a significant covariate of pregabalin apparent oral volume of distribution, and these relationships were similar in paediatric and adult patients.

Pregabalin pharmacokinetics in patients younger than 3 months old have not been studied (see sections 4.2, 4.8 and 5.1).

#### Elderly

Pregabalin clearance tends to decrease with increasing age. This decrease in pregabalin oral clearance is consistent with decreases in creatinine clearance associated with increasing age. Reduction of pregabalin dose may be required in patients who have age related compromised renal function (see section 4.2 Table 1).

#### Breast-feeding mothers

The pharmacokinetics of 150 mg pregabalin given every 12 hours (300 mg daily dose) was evaluated in 10 lactating women who were at least 12 weeks postpartum. Lactation had little to no influence on pregabalin pharmacokinetics. Pregabalin was excreted into breast milk with average steady-state concentrations approximately 76% of those in maternal plasma. The estimated infant dose from breast milk (assuming mean milk consumption of 150 ml/kg/day) of women receiving 300 mg/day or the maximum dose of 600 mg/day would be 0.31 or 0.62 mg/kg/day, respectively. These estimated doses are approximately 7% of the total daily maternal dose on a mg/kg basis.

### **5.3 Preclinical safety data**

In conventional safety pharmacology studies in animals, pregabalin was well-tolerated at clinically relevant doses. In repeated dose toxicity studies in rats and monkeys CNS effects were observed, including hypoactivity, hyperactivity and ataxia. An increased incidence of retinal atrophy commonly observed in aged albino rats was seen after long term exposure to pregabalin at exposures  $\geq 5$  times the mean human exposure at the maximum recommended clinical dose.

Pregabalin was not teratogenic in mice, rats or rabbits. Foetal toxicity in rats and rabbits occurred only at exposures sufficiently above human exposure. In prenatal/postnatal toxicity studies, pregabalin induced offspring developmental toxicity in rats at exposures  $> 2$  times the maximum recommended human exposure.

Adverse effects on fertility in male and female rats were only observed at exposures sufficiently in excess of therapeutic exposure. Adverse effects on male reproductive organs and sperm parameters were reversible and occurred only at exposures sufficiently in excess of therapeutic exposure or were associated with spontaneous degenerative processes in male reproductive organs in the rat. Therefore the effects were considered of little or no clinical relevance.

Pregabalin is not genotoxic based on results of a battery of *in vitro* and *in vivo* tests.

Two-year carcinogenicity studies with pregabalin were conducted in rats and mice. No tumours were observed in rats at exposures up to 24 times the mean human exposure at the maximum recommended clinical dose of 600 mg/day. In mice, no increased incidence of tumours was found at exposures similar to the mean human exposure, but an increased incidence of haemangiosarcoma was observed at higher exposures. The non-genotoxic mechanism of pregabalin-induced tumour formation in mice involves platelet changes and associated

endothelial cell proliferation. These platelet changes were not present in rats or in humans based on short term and limited long term clinical data. There is no evidence to suggest an associated risk to humans.

In juvenile rats the types of toxicity do not differ qualitatively from those observed in adult rats. However, juvenile rats are more sensitive. At therapeutic exposures, there was evidence of CNS clinical signs of hyperactivity and bruxism and some changes in growth (transient body weight gain suppression). Effects on the oestrus cycle were observed at 5-fold the human therapeutic exposure. Reduced acoustic startle response was observed in juvenile rats 1-2 weeks after exposure at >2 times the human therapeutic exposure. Nine weeks after exposure, this effect was no longer observable.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Capsules content

Hydroxypropylcellulose  
Maize starch  
Talc

#### Capsules shell

Iron oxide yellow (E172)  
Titanium dioxide (E171)  
Erythrosine (E127)  
Gelatin  
Sodium laurilsulfate  
Purified water

#### Printing ink

Shellac  
Iron oxide black (E172)  
Propylene glycolAmmonia solution, concentrated  
Potassium hydroxide  
Purified water

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years.

### **6.4 Special precautions for storage**

*Blister:* Store in the original package in order to protect from moisture.

*Bottle:* Keep the bottle tightly closed in order to protect from moisture.

## 6.5 Nature and contents of container

### Pregabalin Mylan Pharma 25 mg hard capsules

PVC/PVDC-Al blister pack containing 14, 21, 56, 84 and 100 hard capsules.  
PVC/PVDC-Al perforated unit dose blister pack containing 56 x 1, 84 x 1 and 100 x 1 hard capsules.

### Pregabalin Mylan Pharma 50 mg hard capsules

PVC/PVDC-Al blister pack containing 14, 21, 56, 84 and 100 hard capsules.  
PVC/PVDC-Al perforated unit dose blister pack containing 84 x 1 and 100 x 1 hard capsules.

### Pregabalin Mylan Pharma 75 mg hard capsules

PVC/PVDC-Al blister pack containing 14, 56 and 100 hard capsules.  
PVC/PVDC-Al perforated unit dose blister pack containing 14 x 1, 56 x 1 and 100 x 1 hard capsules.  
HDPE bottle pack containing 200 hard capsules.

### Pregabalin Mylan Pharma 100 mg hard capsules

PVC/PVDC-Al blister pack containing 21, 84 and 100 hard capsules.  
PVC/PVDC-Al perforated unit dose blister pack containing 84 x 1 and 100 x 1 hard capsule

### Pregabalin Mylan Pharma 150 mg hard capsules

PVC/PVDC-Al blister pack containing 14, 56 and 100 hard capsules.  
PVC/PVDC-Al perforated unit dose blister pack containing 14 x 1, 56 x 1 and 100 x 1 hard capsules  
HDPE bottle pack containing 200 hard capsules.

### Pregabalin Mylan Pharma 200 mg hard capsules

PVC/PVDC-Al blister pack containing 21, 84 and 100 hard capsules.  
PVC/PVDC-Al perforated unit dose blister pack containing 84 x 1 and 100 x 1 hard capsule

### Pregabalin Mylan Pharma 225 mg hard capsules

PVC/PVDC-Al blister pack containing 14, 56 and 100 hard capsules.  
PVC/PVDC-Al perforated unit dose blister pack containing 56 x 1 and 100 x 1 hard capsule.

### Pregabalin Mylan Pharma 300 mg hard capsules

PVC/PVDC-Al blister pack containing 14, 56 and 100 hard capsules.  
PVC/PVDC-Al perforated unit dose blister pack containing 56 x 1 and 100 x 1 hard capsules.  
HDPE bottle pack containing 200 hard capsules.

Not all pack sizes may be marketed.

## 6.6 Special precautions for disposal

No special requirements.

## 7. MARKETING AUTHORISATION HOLDER

Mylan S.A.S  
117 Allée des Parcs  
69800 Saint-Priest  
France

## 8. MARKETING AUTHORISATION NUMBER(S)

### Pregabalin Mylan Pharma 25 mg hard capsules

EU/1/15/998/001  
EU/1/15/998/002  
EU/1/15/998/003  
EU/1/15/998/004  
EU/1/15/998/005  
EU/1/15/998/006  
EU/1/15/998/007  
EU/1/15/998/008

### Pregabalin Mylan Pharma 50 mg hard capsules

EU/1/15/998/009  
EU/1/15/998/010  
EU/1/15/998/011  
EU/1/15/998/012  
EU/1/15/998/013  
EU/1/15/998/014  
EU/1/15/998/015

### Pregabalin Mylan Pharma 75 mg hard capsules

EU/1/15/998/016  
EU/1/15/998/017  
EU/1/15/998/018  
EU/1/15/998/019  
EU/1/15/998/020  
EU/1/15/998/021  
EU/1/15/998/022

### Pregabalin Mylan Pharma 100 mg hard capsules

EU/1/15/998/023  
EU/1/15/998/024  
EU/1/15/998/025  
EU/1/15/998/026  
EU/1/15/998/027

### Pregabalin Mylan Pharma 150 mg hard capsules

EU/1/15/998/028  
EU/1/15/998/029

EU/1/15/998/030  
EU/1/15/998/031  
EU/1/15/998/032  
EU/1/15/998/033  
EU/1/15/998/034

Pregabalin Mylan Pharma 200 mg hard capsules

EU/1/15/998/035  
EU/1/15/998/036  
EU/1/15/998/037  
EU/1/15/998/038  
EU/1/15/998/039

Pregabalin Mylan Pharma 225 mg hard capsules

EU/1/15/998/040  
EU/1/15/998/041  
EU/1/15/998/042  
EU/1/15/998/043  
EU/1/15/998/044

Pregabalin Mylan Pharma 300 mg hard capsules

EU/1/15/998/045  
EU/1/15/998/046  
EU/1/15/998/047  
EU/1/15/998/048  
EU/1/15/998/049  
EU/1/15/998/050

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 25 June 2015

Date of last renewal: DDMMYY

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

Medicinal Product no longer authorised

**ANNEX II**

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturers responsible for batch release

Mylan Hungary Kft  
Mylan utca 1, Komárom, 2900,  
Hungary

McDermott Laboratories Limited trading as Gerard Laboratories  
35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13  
Ireland

Mylan B.V.  
Dieselweg 25, 3752 LB Bunschoten  
The Netherlands

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to medical prescription.

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

Medicinal Product no longer authorised

**ANNEX III**

**LABELLING AND PACKAGE LEAFLET**



**A. LABELLING**  
Medicinal Product no longer authorised

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 25 mg hard capsules  
pregabalin

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each hard capsule contains 25 mg pregabalin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule.

14 hard capsules  
21 hard capsules  
56 hard capsules  
84 hard capsules  
100 hard capsules  
56 x 1 hard capsules  
84 x 1 hard capsules  
100 x 1 hard capsules

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S  
117 Allée des Parcs  
69800 Saint-Priest  
France

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/15/998/001  
EU/1/15/998/002  
EU/1/15/998/003  
EU/1/15/998/004  
EU/1/15/998/005  
EU/1/15/998/006  
EU/1/15/998/007  
EU/1/15/998/008

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Pregabalin Mylan Pharma 25 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC  
SN  
NN

Medicinal Product no longer authorised

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 25 mg hard capsules  
pregabalin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

Medicinal Product no longer authorised

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 50 mg hard capsules  
pregabalin

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each hard capsule contains 50 mg pregabalin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule.

14 hard capsules  
21 hard capsules  
56 hard capsules  
84 hard capsules  
100 hard capsules  
84 x 1 hard capsules  
100 x 1 hard capsules

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S  
117 Allée des Parcs  
69800 Saint-Priest  
France

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/15/998/009  
EU/1/15/998/010  
EU/1/15/998/011  
EU/1/15/998/012  
EU/1/15/998/013  
EU/1/15/998/014  
EU/1/15/998/015

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Pregabalin Mylan Pharma 50 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC  
SN  
NN

Medicinal Product no longer authorised



**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 50 mg hard capsules  
pregabalin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON FOR BLISTER AND BOTTLE**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 75 mg hard capsules  
pregabalin

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each hard capsule contains 75 mg pregabalin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule.

14 hard capsules  
56 hard capsules  
100 hard capsules  
200 hard capsules  
14 x 1 hard capsules  
56 x 1 hard capsules  
100 x 1 hard capsules

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

*Blister*: Store in the original package in order to protect from moisture.

*Bottle*: Keep the bottle tightly closed in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S  
117 Allée des Parcs  
69800 Saint-Priest  
France

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/15/998/016  
EU/1/15/998/017  
EU/1/15/998/018  
EU/1/15/998/022  
EU/1/15/998/019  
EU/1/15/998/020  
EU/1/15/998/021

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Pregabalin Mylan Pharma 75 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC  
SN  
NN

Medicinal Product no longer authorised

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 75 mg hard capsules  
pregabalin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

Medicinal Product no longer authorised

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**BOTTLE LABEL**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 75 mg hard capsules  
pregabalin

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each hard capsule contains 75 mg pregabalin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule.  
200 hard capsules

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

<For multilayer label only>  
'Peel here'

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Keep the bottle tightly closed in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S  
117 Allée des Parcs  
69800 Saint-Priest  
France

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/15/998/022

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 100 mg hard capsules  
pregabalin

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each hard capsule contains 100 mg pregabalin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule.

21 hard capsules  
84 hard capsules  
100 hard capsules  
84 x 1 hard capsules  
100 x 1 hard capsules

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP



**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S  
117 Allée des Parcs  
69800 Saint-Priest  
France

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/15/998/023  
EU/1/15/998/024  
EU/1/15/998/025  
EU/1/15/998/026  
EU/1/15/998/027

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Pregabalin Mylan Pharma 100 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC  
SN  
NN

Medicinal Product no longer authorised

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 100 mg hard capsules  
pregabalin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

Medicinal Product no longer authorised

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON FOR BLISTER AND BOTTLE**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 150 mg hard capsules  
pregabalin

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each hard capsule contains 150 mg pregabalin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule.

14 hard capsules  
56 hard capsules  
100 hard capsules  
200 hard capsules  
14 x 1 hard capsules  
56 x 1 hard capsules  
100 x 1 hard capsules

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

*Blister:* Store in the original package in order to protect from moisture.

*Bottle:* Keep the bottle tightly closed in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S  
117 Allée des Parcs  
69800 Saint-Priest  
France

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/15/998/028  
EU/1/15/998/029  
EU/1/15/998/030  
EU/1/15/998/034  
EU/1/15/998/031  
EU/1/15/998/032  
EU/1/15/998/033

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Pregabalin Mylan Pharma 150 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC  
SN  
NN

Medicinal Product no longer authorised

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 150 mg hard capsules  
pregabalin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

Medicinal Product no longer authorised

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**BOTTLE LABEL**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 150 mg hard capsules  
pregabalin

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each hard capsule contains 150 mg pregabalin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule.  
200 hard capsules

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

<For multilayer label only>

'Peel here'

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP



**9. SPECIAL STORAGE CONDITIONS**

Keep the bottle tightly closed in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S  
117 Allée des Parcs  
69800 Saint-Priest  
France

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/15/998/034

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 200 mg hard capsules  
pregabalin

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each hard capsule contains 200 mg pregabalin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule.

21 hard capsules  
84 hard capsules  
100 hard capsules  
84 x 1 hard capsules  
100 x 1 hard capsules

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S  
117 Allée des Parcs  
69800 Saint-Priest  
France

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/15/998/035  
EU/1/15/998/036  
EU/1/15/998/037  
EU/1/15/998/038  
EU/1/15/998/039

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Pregabalin Mylan Pharma 200 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC  
SN  
NN

Medicinal Product no longer authorised

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 200 mg hard capsules  
pregabalin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 225 mg hard capsules  
pregabalin

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each hard capsule contains 225 mg pregabalin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule.

14 hard capsules  
56 hard capsules  
100 hard capsules  
56 x 1 hard capsules  
100 x 1 hard capsules

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S  
117 Allée des Parcs  
69800 Saint-Priest  
France

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/15/998/040  
EU/1/15/998/041  
EU/1/15/998/042  
EU/1/15/998/043  
EU/1/15/998/044

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Pregabalin Mylan Pharma 225 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC  
SN  
NN

Medicinal Product no longer authorised



**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 225 mg hard capsules  
pregabalin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON FOR BLISTER AND BOTTLE**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 300 mg hard capsules  
pregabalin

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each hard capsule contains 300 mg pregabalin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule.

14 hard capsules  
56 hard capsules  
100 hard capsules  
200 hard capsules  
56 x 1 hard capsules  
100 x 1 hard capsules

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

*Blister*: Store in the original package in order to protect from moisture.

*Bottle*: Keep the bottle tightly closed in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S  
117 Allée des Pares  
69800 Saint-Priest  
France

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/15/998/045  
EU/1/15/998/046  
EU/1/15/998/047  
EU/1/15/998/050  
EU/1/15/998/048  
EU/1/15/998/049

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Pregabalin Mylan Pharma 300 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC  
SN  
NN

Medicinal Product no longer authorised

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 300 mg hard capsules  
pregabalin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

Medicinal Product no longer authorised

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**BOTTLE LABEL**

**1. NAME OF THE MEDICINAL PRODUCT**

Pregabalin Mylan Pharma 300 mg hard capsules  
pregabalin

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each hard capsule contains 300 mg pregabalin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule.  
200 hard capsules

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

<For multilayer label only>  
'Peel here'

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Keep the bottle tightly closed in order to protect from moisture.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan S.A.S  
117 Allée des Parcs  
69800 Saint-Priest  
France

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/15/998/050

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

Medicinal Product no longer authorised

**B. PACKAGE LEAFLET**



## Package leaflet: Information for the user

**Pregabalin Mylan Pharma 25 mg hard capsules**  
**Pregabalin Mylan Pharma 50 mg hard capsules**  
**Pregabalin Mylan Pharma 75 mg hard capsules**  
**Pregabalin Mylan Pharma 100 mg hard capsules**  
**Pregabalin Mylan Pharma 150 mg hard capsules**  
**Pregabalin Mylan Pharma 200 mg hard capsules**  
**Pregabalin Mylan Pharma 225 mg hard capsules**  
**Pregabalin Mylan Pharma 300 mg hard capsules**

pregabalin

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What Pregabalin Mylan Pharma is and what it is used for
2. What you need to know before you take Pregabalin Mylan Pharma
3. How to take Pregabalin Mylan Pharma
4. Possible side effects
5. How to store Pregabalin Mylan Pharma
6. Contents of the pack and other information

#### 1. What Pregabalin Mylan Pharma is and what it is used for

Pregabalin Mylan Pharma contains the active substance pregabalin which belongs to a group of medicines used to treat epilepsy and Generalised Anxiety Disorder (GAD) in adults.

**Epilepsy:** Pregabalin Mylan Pharma is used to treat a certain form of epilepsy (partial seizures with or without secondary generalisation) in adults. Your doctor will prescribe Pregabalin Mylan Pharma for you to help treat your epilepsy when your current treatment is not controlling your condition. You should take Pregabalin Mylan Pharma in addition to your current treatment. Pregabalin Mylan Pharma is not intended to be used alone, but should always be used in combination with other anti-epileptic treatment.

**Generalised Anxiety Disorder:** Pregabalin Mylan Pharma is used to treat Generalised Anxiety Disorder (GAD). The symptoms of GAD are prolonged excessive anxiety and worry that are difficult to control. GAD can also cause restlessness or feeling keyed up or on edge, being easily fatigued (tired), having difficulty concentrating or mind going blank, feeling irritable, having muscle tension or sleep disturbance. This is different to the stresses and strains of everyday life.

## 2. What you need to know before you take Pregabalin Mylan Pharma

### Do not take Pregabalin Mylan Pharma

If you are allergic to pregabalin or any of the other ingredients of this medicine (listed in section 6).

### Warnings and precautions

Talk to your doctor or pharmacist before taking Pregabalin Mylan Pharma.

- Some patients taking Pregabalin Mylan Pharma have reported symptoms suggesting an allergic reaction. These symptoms include swelling of the face, lips, tongue, and throat, as well as diffuse skin rash. Should you experience any of these reactions, you should contact your physician immediately.
- Pregabalin Mylan Pharma has been associated with dizziness and somnolence, which could increase the occurrence of accidental injury (fall) in elderly patients. Therefore, you should be careful until you are used to any effect the medicine might have.
- Pregabalin Mylan Pharma may cause blurring or loss of vision, or other changes in eyesight, many of which are temporary. You should immediately tell your doctor if you experience any changes in your vision.
- Some patients with diabetes who gain weight while taking pregabalin may need an alteration in their diabetic medicines.
- Certain side effects may be more common, such as sleepiness, because patients with spinal cord injury may be taking other medicines to treat, for example, pain or spasticity, that have similar side effects to Pregabalin and the severity of these effects may be increased when taken together.
- There have been reports of heart failure in some patients when taking Pregabalin Mylan Pharma; these patients were mostly elderly with cardiovascular conditions. **Before taking this medicine you should tell your doctor if you have a history of heart disease.**
- There have been reports of kidney failure in some patients when taking Pregabalin Mylan Pharma. If while taking Pregabalin Mylan Pharma you notice decreased urination, you should tell your doctor as stopping the medicine may improve this.
- A small number of people being treated with anti-epileptics such as Pregabalin Mylan Pharma have had thoughts of harming or killing themselves. If at any time you have these thoughts, immediately contact your doctor.
- When Pregabalin Mylan Pharma is taken with other medicines that may cause constipation (such as some types of pain medicines) it is possible that gastrointestinal problems may occur (e.g., constipation, blocked or paralysed bowel). Tell your doctor if you experience constipation, especially if you are prone to this problem.
- Before taking this medicine you should tell your doctor if you have a history of alcoholism or any drug abuse or dependence. Do not take more medicine than prescribed.
- There have been reports of convulsions when taking Pregabalin Mylan Pharma or shortly after stopping Pregabalin Mylan Pharma. If you experience a convulsion, contact your doctor immediately.

- There have been reports of reduction in brain function (encephalopathy) in some patients taking Pregabalin Mylan Pharma when they have other conditions. Tell your doctor if you have a history of any serious medical conditions, including liver or kidney disease.
- There have been reports of breathing difficulties. If you have nervous system disorders, respiratory disorders, renal impairment, or you are older than 65, your doctor may prescribe you a different dosing regimen. Contact your doctor if you experience trouble breathing or shallow breaths.

### **Children and adolescents**

The safety and efficacy in children and adolescents (under 18 years of age) has not been established and therefore, pregabalin should not be used in this age group.

### **Other medicines and Pregabalin Mylan Pharma**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Pregabalin Mylan Pharma and certain other medicines may influence each other (interaction). When taken with certain other medicines, which have sedative effects (including opioids), Pregabalin Mylan Pharma may potentiate these effects and could lead to respiratory failure, coma and death. The degree of dizziness, sleepiness and decreased concentration may be increased if Pregabalin Mylan Pharma is taken together with medicinal products containing:

- Oxycodone – (used as a pain-killer)
- Lorazepam – (used for treating anxiety)
- Alcohol

Pregabalin Mylan Pharma may be taken with oral contraceptives.

### **Pregabalin Mylan Pharma with food, drink and alcohol**

Pregabalin Mylan Pharma capsules may be taken with or without food.

It is advised not to drink alcohol while taking Pregabalin Mylan Pharma.

### **Pregnancy and breast-feeding**

Pregabalin Mylan Pharma should not be taken during pregnancy or when breast-feeding, unless you are told otherwise by your doctor. Effective contraception must be used by women of child-bearing potential. If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

### **Driving and using machines**

Pregabalin Mylan Pharma may produce dizziness, sleepiness and decreased concentration. You should not drive, operate complex machinery or engage in other potentially hazardous activities until you know whether this medicine affects your ability to perform these activities.

### **Sodium content**

This medicine contains less than 1 mmol sodium (23 mg) per capsule. That is to say essentially 'sodium-free'

## **3. How to take Pregabalin Mylan Pharma**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Your doctor will determine what dose is appropriate for you.

Pregabalin Mylan Pharma is for oral use only.

**Epilepsy or Generalised Anxiety Disorder:**

- Take the number of capsules as instructed by your doctor.
- The dose, which has been adjusted for you and your condition, will generally be between 150 mg and 600 mg each day.
- Your doctor will tell you to take Pregabalin Mylan Pharma either twice or three times a day. For twice a day take Pregabalin Mylan Pharma once in the morning and once in the evening, at about the same time each day. For three times a day take Pregabalin Mylan Pharma once in the morning, once in the afternoon and once in the evening, at about the same time each day.

If you have the impression that the effect of Pregabalin Mylan Pharma is too strong or too weak, talk to your doctor or pharmacist.

If you are an older patient (over 65 years of age), you should take Pregabalin Mylan Pharma normally except if you have problems with your kidneys.

Your doctor may prescribe a different dosing schedule and/or dose if you have problems with your kidneys.

Swallow the capsule whole with water.

Continue taking Pregabalin Mylan Pharma until your doctor tells you to stop.

**If you take more Pregabalin Mylan Pharma than you should**

Call your doctor or go to the nearest hospital emergency unit immediately. Take your box or bottle of Pregabalin Mylan Pharma capsules with you. You may feel sleepy, confused, agitated, or restless as a result of taking more Pregabalin Mylan Pharma than you should. Fits have also been reported.

**If you forget to take Pregabalin Mylan Pharma**

It is important to take your Pregabalin Mylan Pharma capsules regularly at the same time each day. If you forget to take a dose, take it as soon as you remember unless it is time for your next dose. In that case, just carry on with the next dose as normal. Do not take a double dose to make up for a forgotten dose.

**If you stop taking Pregabalin Mylan Pharma**

Do not stop taking Pregabalin Mylan Pharma unless your doctor tells you to. If your treatment is stopped it should be done gradually over a minimum of 1 week.

After stopping long and short-term Pregabalin Mylan Pharma treatment, you need to know that you may experience certain side effects. These include, trouble sleeping, headache, nausea, feeling anxious, diarrhoea, flu-like symptoms, convulsions, nervousness, depression, pain, sweating, and dizziness. These symptoms may occur more commonly or severely if you have been taking Pregabalin Mylan Pharma for a longer period of time.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Very common: may affect more than 1 in 10 people**

- Dizziness, drowsiness, headache

**Common: may affect up to 1 in 10 people**

- Increased appetite
- Feeling of elation, confusion, disorientation, decrease in sexual interest, irritability
- Disturbance in attention, clumsiness, memory impairment, loss of memory, tremor, difficulty with speaking, tingling feeling, numbness, sedation, lethargy, insomnia, fatigue, feeling abnormal
- Blurred vision, double vision
- Vertigo, problems with balance, fall
- Dry mouth, constipation, vomiting, flatulence, diarrhoea, nausea, swollen abdomen
- Difficulties with erection
- Swelling of the body including extremities
- Feeling drunk, abnormal style of walking
- Weight gain
- Muscle cramp, joint pain, back pain, pain in limb
- Sore throat

**Uncommon: may affect up to 1 in 100 people**

- Loss of appetite, weight loss, low blood sugar, high blood sugar
- Change in perception of self, restlessness, depression, agitation, mood swings, difficulty finding words, hallucinations, abnormal dreams, panic attacks, apathy, aggression, elevated mood, mental impairment, difficulty with thinking, increase in sexual interest, problems with sexual functioning including inability to achieve a sexual climax, delayed ejaculation
- Changes in eyesight, unusual eye movement, changes in vision including tunnel vision, flashes of light, jerky movements, reduced reflexes, increased activity, dizziness on standing, sensitive skin, loss of taste, burning sensation, tremor on movement, decreased consciousness, loss of consciousness, fainting, increased sensitivity to noise, feeling unwell
- Dry eyes, eye swelling, eye pain, weak eyes, watery eyes, eye irritation
- Heart rhythm disturbances, increased heart rate, low blood pressure, high blood pressure, changes in heart beat, heart failure
- Flushing, hot flushes
- Difficulty breathing, dry nose, nasal congestion
- Increased saliva production, heartburn, numb around mouth
- Sweating, rash, chills, fever
- Muscle twitching, joint swelling, muscle stiffness, pain including muscle pain, neck pain
- Breast pain
- Difficulty with or painful urination, incontinence
- Weakness, thirst, chest tightness
- Changes in blood and liver test results (blood creatinine phosphokinase increased, alanine aminotransferase increased, aspartate aminotransferase increased, platelet count decreased, neutropenia, increase in blood creatinine, decrease in blood potassium)
- Hypersensitivity, swollen face, itchiness, hives, runny nose, nose bleed, cough, snoring
- Painful menstrual periods
- Coldness of hands and feet

**Rare: may affect up to 1 in 1,000 people**

- Abnormal sense of smell, swinging vision, altered perception of depth, visual brightness, vision loss
- Dilated pupils, cross eyes
- Cold sweat, tightness of the throat, swollen tongue
- Inflammation of the pancreas
- Difficulty in swallowing
- Slow or reduced movement of the body

- Difficulty with writing properly
- Increased fluid in the abdomen
- Fluid in the lungs
- Convulsions
- Changes in the recording of electrical changes (ECG) in the heart which correspond to heart rhythm disturbances
- Muscle damage
- Breast discharge, abnormal breast growth, breast growth in males
- Interrupted menstrual periods
- Kidney failure, reduced urine volume, urinary retention
- Decrease in white blood cell count
- Inappropriate behaviour
- Allergic reactions (which may include difficulty breathing, inflammation of the eyes (keratitis) and a serious skin reaction characterized by rash, blisters, peeling skin and pain)
- Jaundice (yellowing of the skin and eyes).

**Very rare: may affect up to 1 in 10,000 people**

- Liver failure.
- Hepatitis (inflammation of the liver).

**If you experience swollen face or tongue or if your skin turns red and starts to blister or peel you should seek immediate medical advice.**

Certain side effects may be more common, such as sleepiness, because patients with spinal cord injury may be taking other medicines to treat, for example, pain or spasticity, that have similar side effects to Pregabalin and the severity of these effects may be increased when taken together.

The following side effects were reported in the postmarketing experience: trouble breathing, shallow breaths.

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Pregabalin Mylan Pharma**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

Blister: Store in the original package in order to protect from moisture.

Bottle: Keep the bottle tightly closed in order to protect from moisture.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Pregabalin Mylan Pharma contains**

The active substance is pregabalin. Each hard capsule contains either 25 mg, 50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 225 mg or 300 mg of pregabalin.

The other ingredients are: hydroxypropylcellulose, maize starch, talc, gelatin, titanium dioxide (E171), sodium laurilsulfate, purified water, shellac, black iron oxide (E172), propylene glycol, potassium hydroxide and concentrated ammonia solution, yellow iron oxide (E172) and erythrosine (E127).

Medicinal Product no longer authorised

## What Pregabalin Mylan Pharma looks like and contents of the pack

Hard capsule.

<p>Pregabalin Mylan Pharma 25 mg hard capsule</p>	<p>Light peach opaque cap and white opaque body, hard-shell gelatin capsule filled with white to off-white powder. The capsule is axially printed with <b>MYLAN</b> over <b>PB25</b> in black ink on cap and body.</p> <p>Available in blister packs containing 14, 21, 56, 84, 100 capsules and in perforated unit dose blister packs containing 56 x 1, 84 x 1, 100 x 1 capsules.</p>
<p>Pregabalin Mylan Pharma 50 mg hard capsule</p>	<p>Dark peach opaque cap and white opaque body, hard-shell gelatin capsule filled with white to off-white powder. The capsule is axially printed with <b>MYLAN</b> over <b>PB50</b> in black ink on cap and body.</p> <p>Available in blister packs containing 14, 21, 56, 84, 100 capsules and in perforated unit dose blister packs containing 84 x 1, 100 x 1 capsules.</p>
<p>Pregabalin Mylan Pharma 75 mg hard capsule</p>	<p>Light peach opaque cap and light peach opaque body, hard-shell gelatin capsule filled with white to off-white powder. The capsule is axially printed with <b>MYLAN</b> over <b>PB75</b> in black ink on cap and body.</p> <p>Available in blister packs containing 14, 56, 100 capsules, in perforated unit dose blister packs containing 14 x 1, 56 x 1, 100 x 1 capsules and in bottles containing 200 capsules.</p>
<p>Pregabalin Mylan Pharma 100 mg hard capsule</p>	<p>Dark peach opaque cap and dark peach opaque body, hard-shell gelatin capsule filled with white to off-white powder. The capsule is axially printed with <b>MYLAN</b> over <b>PB100</b> in black ink on cap and body.</p> <p>Available in blister packs containing 21, 84, 100 capsules and in perforated unit dose blister packs containing 84 x 1, 100 x 1 capsules.</p>
<p>Pregabalin Mylan Pharma 150 mg hard capsule</p>	<p>Light peach opaque cap and white opaque body, hard-shell gelatin capsule filled with white to off-white powder. The capsule is axially printed with <b>MYLAN</b> over <b>PB150</b> in black ink on cap and body.</p> <p>Available in blister packs containing 14, 56, 100 capsules, in perforated unit dose blister packs containing 14 x 1, 56 x 1, 100 x 1 capsules and in bottles containing 200 capsules.</p>



<p>Pregabalin Mylan Pharma 200 mg hard capsule</p>	<p>Light peach opaque cap and light peach opaque body, hard-shell gelatin capsule filled with white to off-white powder. The capsule is axially printed with <b>MYLAN</b> over <b>PB200</b> in black ink on cap and body.</p> <p>Available in blister packs containing 21, 84, 100 capsules and in perforated unit dose blister packs containing 84 x 1, 100 x 1 capsules.</p>
<p>Pregabalin Mylan Pharma 225 mg hard capsule</p>	<p>Dark peach opaque cap and dark peach opaque body, hard-shell gelatin capsule filled with white to off-white powder. The capsule is axially printed with <b>MYLAN</b> over <b>PB225</b> in black ink on cap and body.</p> <p>Available in blister packs containing 14, 56, 100 capsules and in perforated unit dose blister packs containing 56 x 1, 100 x 1 capsules.</p>
<p>Pregabalin Mylan Pharma 300 mg hard capsule</p>	<p>Light peach opaque cap and white opaque body, hard-shell gelatin capsule filled with white to off-white powder. The capsule is axially printed with <b>MYLAN</b> over <b>PB300</b> in black ink on cap and body.</p> <p>Available in blister packs containing 14, 56, 100 capsules, in perforated unit dose blister packs containing 56 x 1, 100 x 1 capsules and in bottles containing 200 capsules.</p>

Not all pack sizes may be marketed.

**Marketing Authorisation Holder**

Mylan S.A.S  
117 Allée des Parcs  
69800 Saint-Priest  
France

**Manufacturer**

Mylan Hungary Kft, Mylan utca 1, Komárom, 2900, Hungary

McDermott Laboratories Limited t/a Gerard Laboratories, 35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13, Ireland

Mylan B.V., Dieselweg 25, 3752 LB Bunschoten, The Netherlands

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

**België/Belgique/Belgien**

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00

**България**

Майлан ЕООД  
Тел.: +359 2 44 55 400

**Česká republika**

Mylan Healthcare CZ s.r.o.  
Tel: +420 222 004 400

**Danmark**

Mylan Denmark ApS  
Tel: +45 28 11 69 32

**Deutschland**

Mylan Healthcare GmbH  
Tel: +49 800 0700 800

**Eesti**

BGP Products Switzerland GmbH Eesti  
filiaal  
Tel: +372 6363 052

**Ελλάδα**

Generics Pharma Hellas ΕΠΕ  
Τηλ: +30 210 993 6410

**España**

Mylan Pharmaceuticals, S.L  
Tel: + 34 900 102 712

**France**

Mylan S.A.S  
Tel: +33 4 37 25 75 00

**Hrvatska**

Mylan Hrvatska d.o.o.  
Tel: +385 1 23 50 599

**Ireland**

Mylan Ireland limited  
Tel: +353 (0) 87 1694982

**Lietuva**

Mylan Healthcare UAB  
Tel: +370 5 205 1288

**Luxembourg/Luxemburg**

Mylan bvba/sprl  
Tel: + 32 (0)2 658 61 00  
(Belgique/Belgien)

**Magyarország**

Mylan EPD Kft  
Tel: + 36 1 465 2100

**Malta**

V.J. Salomone Pharma Ltd.  
Tel: + 356 21 22 01 74

**Nederland**

Mylan BV  
Tel: + 31 (0)20 426 3300

**Norge**

Mylan Healthcare Norge AS  
Tel: + 47 66 75 33 00

**Österreich**

Arcana Arzneimittel GmbH  
Tel: +43 1 416 2418

**Polska**

Mylan Healthcare Sp. z.o.o.  
Tel: + 48 22 546 64 00

**Portugal**

Mylan, Lda.  
Tel: + 351 21 412 72 56

**România**

BGP Products SRL  
Tel: +40 372 579 000

**Slovenija**

Mylan Healthcare d.o.o.  
Tel: + 386 1 23 63 180

**Ísland**

Icepharma hf  
Tel: +354 540 8000

**Italia**

Mylan Italia S.r.l.  
Tel: + 39 02 612 46921

**Κύπρος**

Varnavas Hadjipanayis Ltd  
Tηλ: +357 2220 7700

**Latvija**

Mylan Healthcare SIA  
Tel: +371 676 055 80

**Slovenská republika**

Mylan s.r.o.  
Tel: +421 2 32 199 100

**Suomi/Finland**

Mylan Finland OY  
Puh/Tel: +358 20 720 9555

**Sverige**

Mylan AB  
Tel: + 46 855 522 750

**United Kingdom**

Generics [UK] Ltd  
Tel: +44 1707 853000

**This leaflet was last revised in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.