

ANNEX I

**LIST OF THE NAMES, PHARMACEUTICAL FORMS, STRENGTHS OF THE MEDICINAL
PRODUCTS, ROUTE OF ADMINISTRATION AND MARKETING AUTHORISATION
HOLDERS IN THE MEMBER STATES**

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Austria | Pfizer Corporation Austria Ges.m.b.H. Floridsdorfer Hauptstraße 1 A - 1210 Wien Austria | Sortis 10 mg - Filmtabletten | 10mg | Film-coated tablet | Oral use |
| Austria | Pfizer Corporation Austria Ges.m.b.H. Floridsdorfer Hauptstraße 1 A - 1210 Wien Austria | Sortis 20 mg - Filmtabletten | 20mg | Film-coated tablet | Oral use |
| Austria | Pfizer Corporation Austria Ges.m.b.H. Floridsdorfer Hauptstraße 1 A - 1210 Wien Austria | Sortis 40 mg - Filmtabletten | 40mg | Film-coated tablet | Oral use |
| Austria | Pfizer Corporation Austria Ges.m.b.H. Floridsdorfer Hauptstraße 1 A - 1210 Wien Austria | Sortis 80 mg - Filmtabletten | 80mg | Film-coated tablet | Oral use |
| Belgium | Pfizer SA Boulevard de la Plaine 17 B-1050 Brussels, Belgium | Lipitor | 10mg | Film-coated tablet | Oral use |
| Belgium | Pfizer SA Boulevard de la Plaine 17 B-1050 Brussels, Belgium | Lipitor | 20mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Belgium | Pfizer SA Boulevard de la Plaine 17 B-1050 Brussels, Belgium | Lipitor | 40mg | Film-coated tablet | Oral use |
| Belgium | Pfizer SA Boulevard de la Plaine 17 B-1050 Brussels, Belgium | Lipitor | 80mg | Film-coated tablet | Oral use |
| Bulgaria | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 10mg | Film-coated tablet | Oral use |
| Bulgaria | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 20mg | Film-coated tablet | Oral use |
| Bulgaria | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 40mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Bulgaria | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 80mg | Film-coated tablet | Oral use |
| Cyprus | Pfizer Hellas A. E. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Lipitor | 10mg | Film-coated tablet | Oral use |
| Cyprus | Pfizer Hellas A. E. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Lipitor | 20mg | Film-coated tablet | Oral use |
| Cyprus | Pfizer Hellas A. E. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Lipitor | 40mg | Film-coated tablet | Oral use |
| Czech Republic | Pfizer, spol. s r.o., Stroupežnického 17, 150 00 Prague 5, Czech Republic | Sortis | 10mg | Film-coated tablet | Oral use |
| Czech Republic | Pfizer, spol. s r.o., Stroupežnického 17, 150 00 Prague 5, Czech Republic | Sortis | 20mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Czech Republic | Pfizer, spol. s r.o., Stroupežnického 17, 150 00 Prague 5, Czech Republic | Sortis | 40mg | Film-coated tablet | Oral use |
| Czech Republic | Pfizer, spol. s r.o., Stroupežnického 17, 150 00 Prague 5, Czech Republic | Sortis | 80mg | Film-coated tablet | Oral use |
| Denmark | Pfizer ApS, Lautrupvang 8, 2750 Ballerup, Denmark | Zarator | 10mg | Film-coated tablet | Oral use |
| Denmark | Pfizer ApS, Lautrupvang 8, 2750 Ballerup, Denmark | Zarator | 20mg | Film-coated tablet | Oral use |
| Denmark | Pfizer ApS, Lautrupvang 8, 2750 Ballerup, Denmark | Zarator | 40mg | Film-coated tablet | Oral use |
| Denmark | Pfizer ApS, Lautrupvang 8, 2750 Ballerup, Denmark | Zarator | 80mg | Film-coated tablet | Oral use |
| Estonia | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 10mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Estonia | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 20mg | Film-coated tablet | Oral use |
| Estonia | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 40mg | Film-coated tablet | Oral use |
| Estonia | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis 80 mg | 80mg | Film-coated tablet | Oral use |
| Finland | Pfizer Oy, Tietokuja 4, 00330 Helsinki, Finland | Lipitor | 10mg | Film-coated tablet | Oral use |
| Finland | Pfizer Oy, Tietokuja 4, 00330 Helsinki, Finland | Lipitor | 20mg | Film-coated tablet | Oral use |
| Finland | Pfizer Oy, Tietokuja 4, 00330 Helsinki, Finland | Lipitor | 40mg | Film-coated tablet | Oral use |
| Finland | Pfizer Oy, Tietokuja 4, 00330 Helsinki, Finland | Lipitor | 80mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--|---|---|------------------------|--|--|
| Finland | Pfizer Oy, Tietokuja 4, 00330 Helsinki, Finland | Orbeos | 10mg | Film-coated tablet | Oral use |
| Finland | Pfizer Oy, Tietokuja 4, 00330 Helsinki, Finland | Orbeos | 20mg | Film-coated tablet | Oral use |
| Finland | Pfizer Oy, Tietokuja 4, 00330 Helsinki, Finland | Orbeos | 40mg | Film-coated tablet | Oral use |
| Finland | Pfizer Oy, Tietokuja 4, 00330 Helsinki, Finland | Orbeos | 80mg | Film-coated tablet | Oral use |
| France | Pfizer Holding France 23-25 Avenue du Docteur Lannelongue 75668 Paris Cedex 14 France | Tahor | 10mg | Film-coated tablet | Oral use |
| France | Pfizer Holding France 23-25 Avenue du Docteur Lannelongue 75668 Paris Cedex 14 France | Tahor | 20mg | Film-coated tablet | Oral use |
| France | Pfizer Holding France 23-25 Avenue du Docteur Lannelongue 75668 Paris Cedex 14 France | Tahor | 40mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| France | Pfizer Holding France 23-25 Avenue du Docteur Lannelongue 75668 Paris Cedex 14 France | Tahor | 80mg | Film-coated tablet | Oral use |
| Germany | Pfizer Pharma GmbH, Linkstraße 10 10785 Berlin, Germany | Sortis 10 mg | 10mg | Film-coated tablet | Oral use |
| Germany | Pfizer Pharma GmbH, Linkstraße 10 10785 Berlin, Germany | Sortis 20 mg | 20mg | Film-coated tablet | Oral use |
| Germany | Pfizer Pharma GmbH, Linkstraße 10 10785 Berlin, Germany | Sortis 40 mg | 40mg | Film-coated tablet | Oral use |
| Germany | Pfizer Pharma GmbH, Linkstraße 10 10785 Berlin, Germany | Sortis 80 mg | 80mg | Film-coated tablet | Oral use |
| Germany | Pfizer Pharma GmbH, Linkstraße 10 10785 Berlin, Germany | Liprimar 10 mg | 10mg | Film-coated tablet | Oral use |
| Germany | Pfizer Pharma GmbH, Linkstraße 10 10785 Berlin, Germany | Liprimar 20 mg | 20mg | Film-coated tablet | Oral use |
| Germany | Pfizer Pharma GmbH, Linkstraße 10 10785 Berlin, Germany | Liprimar 40 mg | 40mg | Film-coated tablet | Oral use |
| Germany | Pfizer Pharma GmbH, Linkstraße 10 10785 Berlin, Germany | Liprimar 80 mg | 80mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Germany | Pfizer Pharma GmbH, Linkstraße 10 10785 Berlin, Germany | Atorvastatin 10 mg PD | 10mg | Film-coated tablet | Oral use |
| Germany | Pfizer Pharma GmbH, Linkstraße 10 10785 Berlin, Germany | Atorvastatin 20 mg PD | 20mg | Film-coated tablet | Oral use |
| Germany | Pfizer Pharma GmbH, Linkstraße 10 10785 Berlin, Germany | Atorvastatin 40 mg PD | 40mg | Film-coated tablet | Oral use |
| Germany | Pfizer Pharma GmbH, Linkstraße 10 10785 Berlin, Germany | Atorvastatin 80 mg PD | 80mg | Film-coated tablet | Oral use |
| Greece | Pfizer Hellas A. E. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Lipitor | 10mg | Film-coated tablet | Oral use |
| Greece | Pfizer Hellas A. E. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Lipitor | 20mg | Film-coated tablet | Oral use |
| Greece | Pfizer Hellas A. E. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Lipitor | 40mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Greece | Pfizer Hellas A. E. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Lipitor | 80mg | Film-coated tablet | Oral use |
| Greece | WIN MEDICA Ltd 41 Papdiamantopoulou Street 115 28 Ilisia Athens, Greece | Zarator | 10mg | Film-coated tablet | Oral use |
| Greece | WIN MEDICA Ltd 41 Papdiamantopoulou Street 115 28 Ilisia Athens, Greece | Zarator | 20mg | Film-coated tablet | Oral use |
| Greece | WIN MEDICA Ltd 41 Papdiamantopoulou Street 115 28 Ilisia Athens, Greece | Zarator | 40mg | Film-coated tablet | Oral use |
| Greece | Pfizer Hellas A. E. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Edovin | 10mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Greece | Pfizer Hellas A. E. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Edovin | 20mg | Film-coated tablet | Oral use |
| Greece | Pfizer Hellas A. E. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Edovin | 40mg | Film-coated tablet | Oral use |
| Hungary | Pfizer KFT, 1123 Budapest, Alkotás u. 53. MOM Park "F" Ép., Hungary | Sortis | 10mg | Film-coated tablet | Oral use |
| Hungary | Pfizer KFT, 1123 Budapest, Alkotás u. 53. MOM Park "F" Ép., Hungary | Sortis | 20mg | Film-coated tablet | Oral use |
| Hungary | Pfizer KFT, 1123 Budapest, Alkotás u. 53. MOM Park "F" Ép., Hungary | Sortis | 40mg | Film-coated tablet | Oral use |
| Hungary | Pfizer KFT, 1123 Budapest, Alkotás u. 53. MOM Park "F" Ép., Hungary | Sortis | 80mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|--|-------------------------------------|-----------------|--------------------------------------|--|
| Hungary | C.P. Pharma Gyógyszerkereskedel mi Kft. 2040 Budaörs Vasút u. 11. Hungary | Obradon | 10mg | Film-coated tablet | Oral use |
| Hungary | C.P. Pharma Gyógyszerkereskedel mi Kft. 2040 Budaörs Vasút u. 11. Hungary | Obradon | 20mg | Film-coated tablet | Oral use |
| Hungary | C.P. Pharma Gyógyszerkereskedel mi Kft. 2040 Budaörs Vasút u. 11. Hungary | Obradon | 40mg | Film-coated tablet | Oral use |
| Hungary | C.P. Pharma Gyógyszerkereskedel mi Kft. 2040 Budaörs Vasút u. 11. Hungary | Obradon | 80mg | Film-coated tablet | Oral use |
| Iceland | Pfizer ApS, Lautrupvang 8, 2750 Ballerup, Denmark | Zarator | 10mg | Film-coated tablet | Oral use |
| Iceland | Pfizer ApS, Lautrupvang 8, 2750 Ballerup, Denmark | Zarator | 20mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|--|-------------------------------------|-----------------|--------------------------------------|--|
| Iceland | Pfizer ApS, Lautrupvang 8, 2750 Ballerup, Denmark | Zarator | 40mg | Film-coated tablet | Oral use |
| Iceland | Pfizer ApS, Lautrupvang 8, 2750 Ballerup, Denmark | Zarator | 80mg | Film-coated tablet | Oral use |
| Ireland | Pfizer Ireland Pharmaceuticals, Pottery Road, Dun Laoghaire, Co Dublin Ireland | Lipitor | 10mg | Film-coated tablet | Oral use |
| Ireland | Pfizer Ireland Pharmaceuticals, Pottery Road, Dun Laoghaire, Co Dublin Ireland | Lipitor | 20mg | Film-coated tablet | Oral use |
| Ireland | Pfizer Ireland Pharmaceuticals, Pottery Road, Dun Laoghaire, Co Dublin Ireland | Lipitor | 40mg | Film-coated tablet | Oral use |
| Ireland | Pfizer Ireland Pharmaceuticals, Pottery Road, Dun Laoghaire, Co Dublin Ireland | Lipitor | 80mg | Film-coated tablet | Oral use |
| Italy | Pfizer Italia S.r.l. Via Isonzo, 71 04100 Latina - Italy | Xarator | 10mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Italy | Pfizer Italia S.r.l. Via Isonzo, 71 04100 Latina - Italy | Xarator | 20mg | Film-coated tablet | Oral use |
| Italy | Pfizer Italia S.r.l. Via Isonzo, 71 04100 Latina - Italy | Xarator | 40mg | Film-coated tablet | Oral use |
| Italy | Pfizer Italia S.r.l. Via Isonzo, 71 04100 Latina - Italy | Xarator | 80mg | Film-coated tablet | Oral use |
| Italy | Bioindustria Farmaceutici S.r.l Via Isonzo, 71 04100 Latina - Italy | Lipitor | 10mg | Film-coated tablet | Oral use |
| Italy | Bioindustria Farmaceutici S.r.l. Via Isonzo, 71 04100 Latina - Italy | Lipitor | 20mg | Film-coated tablet | Oral use |
| Italy | Bioindustria Farmaceutici S.r.l. Via Isonzo, 71 04100 Latina - Italy | Lipitor | 40mg | Film-coated tablet | Oral use |
| Italy | Bioindustria Farmaceutici S.r.l. Via Isonzo, 71 04100 Latina - Italy | Lipitor | 80mg | Film-coated tablet | Oral use |
| Italy | Pfizer Italia S.r.l. Via Isonzo, 71 04100 Latina - Italy | Torvast | 10mg | Film-coated tablet | Oral use |
| Italy | Pfizer Italia S.r.l. Via Isonzo, 71 04100 Latina - Italy | Torvast | 20mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Italy | Pfizer Italia S.r.l. Via Isonzo, 71 04100 Latina - Italy | Torvast | 40mg | Film-coated tablet | Oral use |
| Italy | Pfizer Italia S.r.l. Via Isonzo, 71 04100 Latina - Italy | Torvast | 80mg | Film-coated tablet | Oral use |
| Italy | Laboratori Guidotti S.p.A. Via Livornese, 897 - La Vettola 56010 Pisa - Italy | Totalip | 10mg | Film-coated tablet | Oral use |
| Italy | Laboratori Guidotti S.p.A. Via Livornese, 897 - La Vettola 56010 Pisa - Italy | Totalip | 20mg | Film-coated tablet | Oral use |
| Italy | Laboratori Guidotti S.p.A. Via Livornese, 897 - La Vettola 56010 Pisa - Italy | Totalip | 40mg | Film-coated tablet | Oral use |
| Italy | Laboratori Guidotti S.p.A. Via Livornese, 897 - La Vettola 56010 Pisa - Italy | Totalip | 80mg | Film-coated tablet | Oral use |
| Latvia | Pfizer Limited, Ramsgate Road, Sandwich, Kent CT13 9NJ, United Kingdom | Sortis | 10mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|--|-------------------------------------|-----------------|--------------------------------------|--|
| Latvia | Pfizer Limited, Ramsgate Road, Sandwich, Kent CT13 9NJ, United Kingdom | Sortis | 20mg | Film-coated tablet | Oral use |
| Latvia | Pfizer Limited, Ramsgate Road, Sandwich, Kent CT13 9NJ, United Kingdom | Sortis | 40mg | Film-coated tablet | Oral use |
| Latvia | Pfizer Limited, Ramsgate Road, Sandwich, Kent CT 13 9NJ, United Kingdom | Sortis | 80mg | Film-coated tablet | Oral use |
| Lithuania | Pfizer Limited, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 10mg | Film-coated tablet | Oral use |
| Lithuania | Pfizer Limited, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 20mg | Film-coated tablet | Oral use |
| Lithuania | Pfizer Limited, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 40mg | Film-coated tablet | Oral use |
| Lithuania | Pfizer Limited, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 80mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|--|-------------------------------------|-----------------|--------------------------------------|--|
| Luxembourg | Pfizer SA Boulevard de la Plaine 17 B-1050 Brussels, Belgium | Lipitor | 10mg | Film-coated tablet | Oral use |
| Luxembourg | Pfizer SA Boulevard de la Plaine 17 B-1050 Brussels, Belgium | Lipitor | 20mg | Film-coated tablet | Oral use |
| Luxembourg | Pfizer SA Boulevard de la Plaine 17 B-1050 Brussels, Belgium | Lipitor | 40mg | Film-coated tablet | Oral use |
| Luxembourg | Pfizer SA Boulevard de la Plaine 17 B-1050 Brussels, Belgium | Lipitor | 80mg | Film-coated tablet | Oral use |
| Malta | Pfizer Hellas S.A. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Lipitor | 10mg | Film-coated tablet | Oral use |
| Malta | Pfizer Hellas S.A. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Lipitor | 20mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|--|-------------------------------------|-----------------|--------------------------------------|--|
| Malta | Pfizer Hellas S.A. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Lipitor | 40mg | Film-coated tablet | Oral use |
| Malta | Pfizer Hellas S.A. 243, Messoghion Ave., 154 51 Neo Psychiko, Athens, Greece | Lipitor | 80mg | Film-coated tablet | Oral use |
| Netherlands | Pfizer bv Rivium Westlaan 142 2909 LD Capelle a/d IJssel The Netherlands | Lipitor | 10mg | Film-coated tablet | Oral use |
| Netherlands | Pfizer bv Rivium Westlaan 142 2909 LD Capelle a/d IJssel The Netherlands | Lipitor | 20mg | Film-coated tablet | Oral use |
| Netherlands | Pfizer bv Rivium Westlaan 142 2909 LD Capelle a/d IJssel The Netherlands | Lipitor | 40mg | Film-coated tablet | Oral use |
| Netherlands | Pfizer bv Rivium Westlaan 142 2909 LD Capelle a/d IJssel The Netherlands | Lipitor | 80mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Norway | Pfizer AS Pb. 3 1324 Lysaker Norway | Lipitor | 10mg | Film-coated tablet | Oral use |
| Norway | Pfizer AS Pb. 3 1324 Lysaker Norway | Lipitor | 20mg | Film-coated tablet | Oral use |
| Norway | Pfizer AS Pb. 3 1324 Lysaker Norway | Lipitor | 40mg | Film-coated tablet | Oral use |
| Norway | Pfizer AS Pb. 3 1324 Lysaker Norway | Lipitor | 80mg | Film-coated tablet | Oral use |
| Poland | Parke-Davis GmbH, Pfizerstrasse 1, 76 139 Karlsruhe, Germany | Sortis 10 | 10mg | Film-coated tablet | Oral use |
| Poland | Parke-Davis GmbH, Pfizerstrasse 1, 76 139 Karlsruhe, Germany | Sortis 20 | 20mg | Film-coated tablet | Oral use |
| Poland | Parke-Davis GmbH, Pfizerstrasse 1, 76 139 Karlsruhe, Germany | Sortis 40 | 40mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Poland | Pfizer Polska Sp.z o.o. ul. Rzymowskiego 28 02-697 Warszawa Poland | Sortis 80 | 80 mg | Film-coated tablet | Oral use |
| Portugal | Laboratorios Pfizer, Lda., Lagoas Park, Edificio 10, 2740-271 Porto Salvo, Portugal | Zarator | 10mg | Film-coated tablet | Oral use |
| Portugal | Laboratorios Pfizer, Lda., Lagoas Park, Edificio 10, 2740-271 Porto Salvo, Portugal | Zarator | 20mg | Film-coated tablet | Oral use |
| Portugal | Laboratorios Pfizer, Lda., Lagoas Park, Edificio 10, 2740-271 Porto Salvo, Portugal | Zarator | 40mg | Film-coated tablet | Oral use |
| Portugal | Laboratorios Pfizer, Lda., Lagoas Park, Edificio 10, 2740-271 Porto Salvo, Portugal | Zarator | 80 mg | Film-coated tablet | Oral use |
| Portugal | Parke-Davis - Produtos Farmaceuticos, Lda., Lagoas Park, Edificio 10, 2740-271 Porto Salvo, Portugal | Atorvastatina Parke-Davis | 10mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Portugal | Parke-Davis - Produtos Farmaceuticos, Lda., Lagoas Park, Edificio 10, 2740-271 Porto Salvo, Portugal | Atorvastatina Parke-Davis | 20mg | Film-coated tablet | Oral use |
| Portugal | Parke-Davis - Produtos Farmaceuticos, Lda., Lagoas Park, Edificio 10, 2740-271 Porto Salvo, Portugal | Atorvastatina Parke-Davis | 40mg | Film-coated tablet | Oral use |
| Portugal | Parke-Davis - Produtos Farmaceuticos, Lda., Lagoas Park, Edificio 10, 2740-271 Porto Salvo, Portugal | Atorvastatina Parke-Davis | 80 mg | Film-coated tablet | Oral use |
| Portugal | Farmogene - Produtos Farmacêuticos, Lda, Lagoas Park, Edificio 10, 2740-271 Porto Salvo, Portugal | Texzor | 10mg | Film-coated tablet | Oral use |
| Portugal | Farmogene - Produtos Farmacêuticos, Lda, Lagoas Park, Edificio 10, 2740-271 Porto Salvo, Portugal | Texzor | 20mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|--|-------------------------------------|-----------------|--------------------------------------|--|
| Portugal | Farmogene - Produtos Farmacêuticos, Lda, Lagoas Park, Edificio 10, 2740-271 Porto Salvo, Portugal | Texzor | 40mg | Film-coated tablet | Oral use |
| Portugal | Farmogene - Produtos Farmacêuticos, Lda., Lagoas Park, Edificio 10, 2740-271 Porto Salvo, Portugal | Texzor | 80 mg | Film-coated tablet | Oral use |
| Romania | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 10mg | Film-coated tablet | Oral use |
| Romania | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 20mg | Film-coated tablet | Oral use |
| Romania | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 40mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Romania | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 80 mg | Film-coated tablet | Oral use |
| Slovak Republic | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 10mg | Film-coated tablet | Oral use |
| Slovak Republic | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 20mg | Film-coated tablet | Oral use |
| Slovak Republic | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 40mg | Film-coated tablet | Oral use |
| Slovak Republic | Pfizer Europe MA EEIG, Ramsgate Road, Sandwich, Kent, CT13 9NJ, United Kingdom | Sortis | 80 mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|---------------------------------------|-----------------|--------------------------------------|--|
| Slovenia | Pfizer Luxembourg SARL, 51, Avenue J. F. Kennedy, L-1855 Luxembourg, Luxembourg | Sortis 10 mg filmsko obložene tablete | 10mg | Film-coated tablet | Oral use |
| Slovenia | Pfizer Luxembourg SARL, 51, Avenue J. F. Kennedy, L-1855 Luxembourg, Luxembourg | Sortis 20 mg filmsko obložene tablete | 20mg | Film-coated tablet | Oral use |
| Slovenia | Pfizer Luxembourg SARL, 51, Avenue J. F. Kennedy, L-1855 Luxembourg, Luxembourg | Sortis 40 mg filmsko obložene tablete | 40mg | Film-coated tablet | Oral use |
| Slovenia | Pfizer Luxembourg SARL, 51, Avenue J. F. Kennedy, L-1855 Luxembourg, Luxembourg | Sortis 80 mg filmsko obložene tablete | 80 mg | Film-coated tablet | Oral use |
| Spain | Parke Davis. S.L. Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Zarator | 10mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Spain | Parke Davis. S.L. Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Zarator | 20mg | Film-coated tablet | Oral use |
| Spain | Parke Davis. S.L. Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Zarator | 40mg | Film-coated tablet | Oral use |
| Spain | Parke Davis. S.L. Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Zarator | 80 mg | Film-coated tablet | Oral use |
| Spain | Pfizer, S.A. Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Cardyl | 10mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|--|-------------------------------------|-----------------|--------------------------------------|--|
| Spain | Pfizer, S.A. Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Cardyl | 20mg | Film-coated tablet | Oral use |
| Spain | Pfizer, S.A. Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Cardyl | 40mg | Film-coated tablet | Oral use |
| Spain | Pfizer, S.A. Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Cardyl | 80 mg | Film-coated tablet | Oral use |
| Spain | Nostrum Farma, S.A., Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Atorvastatina Nostrum | 10mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|---|-------------------------------------|-----------------|--------------------------------------|--|
| Spain | Nostrum Farma, S.A., Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Atorvastatina Nostrum | 20mg | Film-coated tablet | Oral use |
| Spain | Nostrum Farma, S.A., Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Atorvastatina Nostrum | 40mg | Film-coated tablet | Oral use |
| Spain | Nostrum Farma, S.A., Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Atorvastatina Nostrum | 80 mg | Film-coated tablet | Oral use |
| Spain | PHARMACIA GRUPO PFIZER, S.L. Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Atorvastatina Pharmacia | 10mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|--|-------------------------------------|-----------------|--------------------------------------|--|
| Spain | PHARMACIA GRUPO PFIZER, S.L. .Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Atorvastatina Pharmacia | 20mg | Film-coated tablet | Oral use |
| Spain | PHARMACIA GRUPO PFIZER, S.L. Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Atorvastatina Pharmacia | 40mg | Film-coated tablet | Oral use |
| Spain | PHARMACIA GRUPO PFIZER, S.L. Avda. de Europa 20B Parque Empresarial la Moraleja 28108 Alcobendas, (Madrid) Spain | Atorvastatina Pharmacia | 80 mg | Film-coated tablet | Oral use |
| Spain | Almirall S.A. General Mitre, 151 08022 - Barcelona Spain | Prevenor | 10mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|--|-------------------------------------|-----------------|--------------------------------------|--|
| Spain | Almirall S.A. General Mitre, 151 08022 - Barcelona Spain | Prevencor | 20mg | Film-coated tablet | Oral use |
| Spain | Almirall S.A. General Mitre, 151 08022 - Barcelona Spain | Prevencor | 40mg | Film-coated tablet | Oral use |
| Spain | Almirall S.A. General Mitre, 151 08022 - Barcelona Spain | Prevencor | 80 mg | Film-coated tablet | Oral use |
| Sweden | Pfizer AB 191 90 Sollentuna Sweden | Lipitor | 10mg | Film-coated tablet | Oral use |
| Sweden | Pfizer AB 191 90 Sollentuna Sweden | Lipitor | 20mg | Film-coated tablet | Oral use |
| Sweden | Pfizer AB 191 90 Sollentuna Sweden | Lipitor | 40mg | Film-coated tablet | Oral use |
| Sweden | Pfizer AB 191 90 Sollentuna Sweden | Lipitor | 80 mg | Film-coated tablet | Oral use |
| United Kingdom | Pfizer Ireland Pharmaceuticals, Pottery Road, Dun Laoghaire, Co Dublin Ireland | Lipitor | 10mg | Film-coated tablet | Oral use |

| <u>Member State</u> <u>EU/EEA</u> | <u>Marketing</u> <u>Authorisation Holder</u> | <u>Invented name</u> <u>Name</u> | <u>Strength</u> | <u>Pharmaceutical</u> <u>Form</u> | <u>Route of</u> <u>administration</u> |
|--------------------------------------|--|-------------------------------------|-----------------|--------------------------------------|--|
| United Kingdom | Pfizer Ireland Pharmaceuticals, Pottery Road, Dun Laoghaire, Co Dublin Ireland | Lipitor | 20mg | Film-coated tablet | Oral use |
| United Kingdom | Pfizer Ireland Pharmaceuticals, Pottery Road, Dun Laoghaire, Co Dublin Ireland | Lipitor | 40mg | Film-coated tablet | Oral use |
| United Kingdom | Pfizer Ireland Pharmaceuticals, Pottery Road, Dun Laoghaire, Co Dublin Ireland | Lipitor | 80 mg | Film-coated tablet | Oral use |

ANNEX II

AMENDMENTS TO THE SUMMARY OF PRODUCT CHARACTERISTICS AND PACKAGE LEAFLET

**These amendments to the SPC, labelling and package leaflet are valid at the time of the
Commission Decision.**

**After the Commission Decision the Member State Competent Authorities will update the
product information as required.**

**AMENDMENTS TO BE INCLUDED IN THE RELEVANT SECTIONS OF THE SUMMARY
OF PRODUCT CHARACTERISTICS**

1. NAME OF THE MEDICINAL PRODUCT

[See Annex I - To be completed nationally]
{PRODUCT NAME} 10 mg film-coated tablet

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

{As currently approved}.

3. PHARMACEUTICAL FORM

{As currently approved}.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Hypercholesterolaemia

{PRODUCT NAME} is indicated as an adjunct to diet for reduction of elevated total cholesterol, LDL-cholesterol, apolipoprotein B, and triglycerides in **patients adults, adolescents and children aged 10 years or older** with primary hypercholesterolaemia including familial hypercholesterolaemia (heterozygous variant) or combined (mixed) hyperlipidaemia (Corresponding to Types IIa and IIb of the Fredrickson classification) when response to diet and other nonpharmacological measures is inadequate.

{PRODUCT NAME} is also indicated to reduce total-C and LDL-C in **patients adults** with homozygous familial hypercholesterolaemia as an adjunct to other lipid-lowering treatments (e.g. LDL apheresis) or if such treatments are unavailable.

Prevention of cardiovascular disease

Prevention of cardiovascular events in adult patients estimated to have a high risk for a first cardiovascular event (see Section 5.1), as an adjunct to correction of other risk factors.

4.2 Posology and method of administration

{As currently approved}.

Paediatric use

~~Pediatric use should only be carried out by specialists.~~

~~Experience in pediatrics is limited to a small number of patients (age 4–17 years) with severe dyslipidemias, such as homozygous familial hypercholesterolemia. The recommended starting dose in this population is 10 mg of atorvastatin per day. The dose may be increased to 80 mg daily, according to the response and tolerability. Developmental safety data in this population have not been evaluated.~~

Hypercholesterolaemia:

Paediatric use should only be carried out by physicians experienced in the treatment of paediatric hyperlipidaemia and patients should be re-evaluated on a regular basis to assess progress.

For patients aged 10 years and above, the recommended starting dose of atorvastatin is 10 mg per day with titration up to 20 mg per day. Titration should be conducted according to the individual response and tolerability in paediatric patients. Safety information for paediatric patients treated with doses above 20 mg, corresponding to about 0.5 mg/kg, is limited.

There is limited experience in children between 6-10 years of age (see section 5.1). Atorvastatin is not indicated in the treatment of patients below the age of 10 years.

Other pharmaceutical forms/strengths may be more appropriate for this population.

4.3 Contraindications

{As currently approved}.

4.4 Special warnings and precautions for use

{As currently approved}.

Paediatric use

Developmental safety in the paediatric population has not been established (see section 4.8).

4.5 Interaction with other medicinal products and other forms of interaction

{As currently approved}.

Paediatric population

Drug-drug interaction studies have only been performed in adults. The extent of interactions in the paediatric population is not known. The above mentioned interactions for adults and the warnings in section 4.4 should be taken into account for the paediatric population.

4.6 Pregnancy and lactation

{As currently approved}.

4.7 Effects on ability to drive and use machines

{As currently approved}.

4.8 Undesirable effects

{As currently approved}.

Paediatric Population

The clinical safety database includes safety data for 249 paediatric patients who received atorvastatin, among which 7 patients were < 6 years old, 14 patients were in the age range of 6 to 9, and 228 patients were in the age range of 10 to 17.

Nervous system disorders

Common: Headache

Gastrointestinal disorders

Common: Abdominal pain

Investigations

Common: Alanine aminotransferase increased, blood creatine phosphokinase increased

Based on the data available, frequency, type and severity of adverse reactions in children are expected to be the same as in adults. There is currently limited experience with respect to long-term safety in the paediatric population.

4.9 Overdose

{As currently approved}.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

{As currently approved}.

Paediatric Population

Heterozygous Familial Hypercholesterolaemia in Paediatric Patients aged 6-17 years old

An 8-week, open-label study to evaluate pharmacokinetics, pharmacodynamics, and safety and tolerability of atorvastatin was conducted in children and adolescents with genetically confirmed heterozygous familial hypercholesterolemia and baseline LDL-C ≥ 4 mmol/L. A total of 39 children and adolescents, 6 to 17 years of age, were enrolled. Cohort A included 15 children, 6 to 12 years of age and at Tanner Stage 1. Cohort B included 24 children, 10 to 17 years of age and at Tanner Stage ≥ 2 .

The initial dose of atorvastatin was 5 mg daily of a chewable tablet in Cohort A and 10 mg daily of a tablet formulation in Cohort B. The atorvastatin dose was permitted to be doubled if a subject had not attained target LDL-C of < 3.35 mmol/L at Week 4 and if atorvastatin was well tolerated.

Mean values for LDL-C, TC, VLDL-C, and Apo B decreased by Week 2 among all subjects. For subjects whose dose was doubled, additional decreases were observed as early as 2 weeks, at the first assessment, after dose escalation. The mean percent decreases in lipid parameters were similar for both cohorts, regardless of whether subjects remained at their initial dose or doubled their initial dose. At Week 8, on average, the percent change from baseline in LDL-C and TC was approximately 40% and 30%, respectively, over the range of exposures.

Heterozygous Familial Hypercholesterolaemia in Paediatric Patients aged 10-17 years old

In a double-blind, placebo controlled study followed by an open-label phase, 187 boys and postmenarchal girls 10-17 years of age (mean age 14.1 years) with heterozygous familial hypercholesterolaemia (FH) or severe hypercholesterolaemia were randomised to atorvastatin (n=140) or placebo (n=47) for 26 weeks and then all received atorvastatin for 26 weeks.. The dosage of atorvastatin (once daily) was 10 mg for the first 4 weeks and up-titrated to 20 mg if the LDL-C level was > 3.36 mmol/l. Atorvastatin significantly decreased plasma levels of total-C,

LDL-C, triglycerides, and apolipoprotein B during the 26 week double-blind phase. The mean achieved LDL-C value was 3.38 mmol/l (range: 1.81-6.26 mmol/l) in the atorvastatin group compared to 5.91 mmol/l (range: 3.93-9.96 mmol/l) in the placebo group during the 26-week double-blind phase.

An additional paediatric study of atorvastatin versus colestipol in patients with hypercholesterolaemia aged 10-18 years demonstrated that atorvastatin (N=25) caused a significant reduction in LDL-C at week 26 ($p<0.05$) compared with colestipol (N=31).

A compassionate use study in patients with severe hypercholesterolaemia (including homozygous hypercholesterolaemia) included 46 paediatric patients treated with atorvastatin titrated according to response (some subjects received 80 mg atorvastatin per day). The study lasted 3 years: LDL-cholesterol was lowered by 36%.

The long-term efficacy of atorvastatin therapy in childhood to reduce morbidity and mortality in adulthood has not been established.

The European Medicines Agency has waived the obligation to submit the results of studies with atorvastatin in children aged 0 to less than 6 years in the treatment of heterozygous hypercholesterolaemia and in children aged 0 to less than 18 years in the treatment of homozygous familial hypercholesterolaemia, combined (mixed) hypercholesterolaemia, primary hypercholesterolaemia and in the prevention of cardiovascular events (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

{As currently approved}.

Special populations

{As currently approved}.

- Paediatric: Pharmacokinetic data in the paediatric population are not available. In an open-label, 8-week study, Tanner Stage 1 (N=15) and Tanner Stage ≥ 2 (N=24) paediatric patients (ages 6-17 years) with heterozygous familial hypercholesterolemia and baseline LDL-C ≥ 4 mmol/L were treated with 5 or 10 mg of chewable or 10 or 20 mg of film-coated atorvastatin tablets once daily, respectively. Body weight was the only significant covariate in atorvastatin population PK model. Apparent oral clearance of atorvastatin in paediatric subjects appeared similar to adults when scaled allometrically by body weight. Consistent decreases in LDL-C and TC were observed over the range of atorvastatin and o-hydroxyatorvastatin exposures.
- {As currently approved}.

5.3 Preclinical safety data

{As currently approved}.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

{As currently approved}.

6.2 Incompatibilities

{As currently approved}.

6.3 Shelf life

{As currently approved}.

6.4 Special precautions for storage

{As currently approved}.

6.5 Nature and contents of container

{As currently approved}.

6.6 Special precautions for disposal

{As currently approved}.

7. MARKETING AUTHORISATION HOLDER

[To be completed nationally]

[See Annex I - To be completed nationally]

{Name and address}

<{tel}>

<{fax}>

<{e-mail}>

8. MARKETING AUTHORISATION NUMBER(S)

[To be completed nationally]

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

[To be completed nationally]

10. DATE OF REVISION OF THE TEXT

1. NAME OF THE MEDICINAL PRODUCT

[See Annex I - To be completed nationally]
{PRODUCT NAME} 20 mg film-coated tablet

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

{As currently approved}.

3. PHARMACEUTICAL FORM

{As currently approved}.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Hypercholesterolaemia

{PRODUCT NAME} is indicated as an adjunct to diet for reduction of elevated total cholesterol, LDL-cholesterol, apolipoprotein B, and triglycerides in **patients adults, adolescents and children aged 10 years or older** with primary hypercholesterolaemia including familial hypercholesterolaemia (heterozygous variant) or combined (mixed) hyperlipidaemia (Corresponding to Types IIa and IIb of the Fredrickson classification) when response to diet and other nonpharmacological measures is inadequate.

{PRODUCT NAME} is also indicated to reduce total-C and LDL-C in **patients adults** with homozygous familial hypercholesterolaemia as an adjunct to other lipid-lowering treatments (e.g. LDL apheresis) or if such treatments are unavailable.

Prevention of cardiovascular disease

Prevention of cardiovascular events in adult patients estimated to have a high risk for a first cardiovascular event (see Section 5.1), as an adjunct to correction of other risk factors.

4.2 Posology and method of administration

{As currently approved}.

Paediatric use

~~Paediatric use should only be carried out by specialists.~~

~~Experience in pediatrics is limited to a small number of patients (age 4–17 years) with severe dyslipidemias, such as homozygous familial hypercholesterolemia. The recommended starting dose in this population is 10 mg of atorvastatin per day. The dose may be increased to 80 mg daily, according to the response and tolerability. Developmental safety data in this population have not been evaluated.~~

Hypercholesterolaemia:

Paediatric use should only be carried out by physicians experienced in the treatment of paediatric hyperlipidaemia and patients should be re-evaluated on a regular basis to assess progress.

For patients aged 10 years and above, the recommended starting dose of atorvastatin is 10 mg per day with titration up to 20 mg per day. Titration should be conducted according to the individual response and tolerability in paediatric patients. Safety information for paediatric patients treated with doses above 20 mg, corresponding to about 0.5 mg/kg, is limited.

There is limited experience in children between 6-10 years of age (see section 5.1). Atorvastatin is not indicated in the treatment of patients below the age of 10 years.

Other pharmaceutical forms/strengths may be more appropriate for this population.

4.3 Contraindications

{As currently approved}.

4.4 Special warnings and precautions for use

{As currently approved}.

Paediatric use

Developmental safety in the paediatric population has not been established (see section 4.8).

4.5 Interaction with other medicinal products and other forms of interaction

{As currently approved}.

Paediatric population

Drug-drug interaction studies have only been performed in adults. The extent of interactions in the paediatric population is not known. The above mentioned interactions for adults and the warnings in section 4.4 should be taken into account for the paediatric population.

4.6 Pregnancy and lactation

{As currently approved}.

4.7 Effects on ability to drive and use machines

{As currently approved}.

4.8 Undesirable effects

{As currently approved}.

Paediatric Population

The clinical safety database includes safety data for 249 paediatric patients who received atorvastatin, among which 7 patients were < 6 years old, 14 patients were in the age range of 6 to 9, and 228 patients were in the age range of 10 to 17.

Nervous system disorders
Common: Headache

Gastrointestinal disorders
Common: Abdominal pain

Investigations

Common: Alanine aminotransferase increased, blood creatine phosphokinase increased

Based on the data available, frequency, type and severity of adverse reactions in children are expected to be the same as in adults. There is currently limited experience with respect to long-term safety in the paediatric population.

4.9 Overdose

{As currently approved}.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

{As currently approved}.

Paediatric Population

Heterozygous Familial Hypercholesterolaemia in Paediatric Patients aged 6-17 years old

An 8-week, open-label study to evaluate pharmacokinetics, pharmacodynamics, and safety and tolerability of atorvastatin was conducted in children and adolescents with genetically confirmed heterozygous familial hypercholesterolemia and baseline LDL-C ≥ 4 mmol/L. A total of 39 children and adolescents, 6 to 17 years of age, were enrolled. Cohort A included 15 children, 6 to 12 years of age and at Tanner Stage 1. Cohort B included 24 children, 10 to 17 years of age and at Tanner Stage ≥ 2 .

The initial dose of atorvastatin was 5 mg daily of a chewable tablet in Cohort A and 10 mg daily of a tablet formulation in Cohort B. The atorvastatin dose was permitted to be doubled if a subject had not attained target LDL-C of < 3.35 mmol/L at Week 4 and if atorvastatin was well tolerated.

Mean values for LDL-C, TC, VLDL-C, and Apo B decreased by Week 2 among all subjects. For subjects whose dose was doubled, additional decreases were observed as early as 2 weeks, at the first assessment, after dose escalation. The mean percent decreases in lipid parameters were similar for both cohorts, regardless of whether subjects remained at their initial dose or doubled their initial dose. At Week 8, on average, the percent change from baseline in LDL-C and TC was approximately 40% and 30%, respectively, over the range of exposures.

Heterozygous Familial Hypercholesterolaemia in Paediatric Patients aged 10-17 years old

In a double-blind, placebo controlled study followed by an open-label phase, 187 boys and postmenarchal girls 10-17 years of age (mean age 14.1 years) with heterozygous familial hypercholesterolaemia (FH) or severe hypercholesterolaemia were randomised to atorvastatin (n=140) or placebo (n=47) for 26 weeks and then all received atorvastatin for 26 weeks. The dosage of atorvastatin (once daily) was 10 mg for the first 4 weeks and up-titrated to 20 mg if the LDL-C level was > 3.36 mmol/l. Atorvastatin significantly decreased plasma levels of total-C, LDL-C, triglycerides, and apolipoprotein B during the 26 week double-blind phase. The mean achieved LDL-C value was 3.38 mmol/l (range: 1.81-6.26 mmol/l) in the atorvastatin group compared to 5.91 mmol/l (range: 3.93-9.96 mmol/l) in the placebo group during the 26-week double-blind phase.

An additional paediatric study of atorvastatin versus colestipol in patients with hypercholesterolaemia aged 10-18 years demonstrated that atorvastatin (N=25) caused a significant reduction in LDL-C at week 26 ($p<0.05$) compared with colestipol (N=31).

A compassionate use study in patients with severe hypercholesterolaemia (including homozygous hypercholesterolaemia) included 46 paediatric patients treated with atorvastatin titrated according to response (some subjects received 80 mg atorvastatin per day). The study lasted 3 years: LDL-cholesterol was lowered by 36%.

The long-term efficacy of atorvastatin therapy in childhood to reduce morbidity and mortality in adulthood has not been established.

The European Medicines Agency has waived the obligation to submit the results of studies with atorvastatin in children aged 0 to less than 6 years in the treatment of heterozygous hypercholesterolaemia and in children aged 0 to less than 18 years in the treatment of homozygous familial hypercholesterolaemia, combined (mixed) hypercholesterolaemia, primary hypercholesterolaemia and in the prevention of cardiovascular events (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

{As currently approved}.

Special populations

{As currently approved}.

- ~~Paediatric: Pharmacokinetic data in the paediatric population are not available.~~ **In an open-label, 8-week study, Tanner Stage 1 (N=15) and Tanner Stage ≥ 2 (N=24) paediatric patients (ages 6-17 years) with heterozygous familial hypercholesterolemia and baseline LDL-C ≥ 4 mmol/L were treated with 5 or 10 mg of chewable or 10 or 20 mg of film-coated atorvastatin tablets once daily, respectively. Body weight was the only significant covariate in atorvastatin population PK model. Apparent oral clearance of atorvastatin in paediatric subjects appeared similar to adults when scaled allometrically by body weight. Consistent decreases in LDL-C and TC were observed over the range of atorvastatin and o-hydroxyatorvastatin exposures.**
- {As currently approved}.

5.3 Preclinical safety data

{As currently approved}.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

{As currently approved}.

6.2 Incompatibilities

{As currently approved}.

6.3 Shelf life

{As currently approved}.

6.4 Special precautions for storage

{As currently approved}.

6.5 Nature and contents of container

{As currently approved}.

6.6 Special precautions for disposal

{As currently approved}.

7. MARKETING AUTHORISATION HOLDER

[To be completed nationally]

[See Annex I - To be completed nationally]

{Name and address}

<{tel}>

<{fax}>

<{e-mail}>

8. MARKETING AUTHORISATION NUMBER(S)

[To be completed nationally]

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

[To be completed nationally]

10. DATE OF REVISION OF THE TEXT

1. NAME OF THE MEDICINAL PRODUCT

[See Annex I - To be completed nationally]
{PRODUCT NAME} 40 mg film-coated tablet

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

{As currently approved}.

3. PHARMACEUTICAL FORM

{As currently approved}.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Hypercholesterolaemia

{PRODUCT NAME} is indicated as an adjunct to diet for reduction of elevated total cholesterol, LDL-cholesterol, apolipoprotein B, and triglycerides in **patients adults, adolescents and children aged 10 years or older** with primary hypercholesterolaemia including familial hypercholesterolaemia (heterozygous variant) or combined (mixed) hyperlipidaemia (Corresponding to Types IIa and IIb of the Fredrickson classification) when response to diet and other nonpharmacological measures is inadequate.

{PRODUCT NAME} is also indicated to reduce total-C and LDL-C in **patients adults** with homozygous familial hypercholesterolaemia as an adjunct to other lipid-lowering treatments (e.g. LDL apheresis) or if such treatments are unavailable.

Prevention of cardiovascular disease

Prevention of cardiovascular events in adult patients estimated to have a high risk for a first cardiovascular event (see Section 5.1), as an adjunct to correction of other risk factors.

4.2 Posology and method of administration

{As currently approved}.

Paediatric use

~~Paediatric use should only be carried out by specialists.~~

~~Experience in pediatrics is limited to a small number of patients (age 4–17 years) with severe dyslipidemias, such as homozygous familial hypercholesterolemia. The recommended starting dose in this population is 10 mg of atorvastatin per day. The dose may be increased to 80 mg daily, according to the response and tolerability. Developmental safety data in this population have not been evaluated.~~

Hypercholesterolaemia:

Paediatric use should only be carried out by physicians experienced in the treatment of paediatric hyperlipidaemia and patients should be re-evaluated on a regular basis to assess progress.

For patients aged 10 years and above, the recommended starting dose of atorvastatin is 10 mg per day with titration up to 20 mg per day. Titration should be conducted according to the individual response and tolerability in paediatric patients. Safety information for paediatric patients treated with doses above 20 mg, corresponding to about 0.5 mg/kg, is limited.

There is limited experience in children between 6-10 years of age (see section 5.1). Atorvastatin is not indicated in the treatment of patients below the age of 10 years.

Other pharmaceutical forms/strengths may be more appropriate for this population.

4.3 Contraindications

{As currently approved}.

4.4 Special warnings and precautions for use

{As currently approved}.

Paediatric use

Developmental safety in the paediatric population has not been established (see section 4.8).

4.5 Interaction with other medicinal products and other forms of interaction

{As currently approved}.

Paediatric population

Drug-drug interaction studies have only been performed in adults. The extent of interactions in the paediatric population is not known. The above mentioned interactions for adults and the warnings in section 4.4 should be taken into account for the paediatric population.

4.6 Pregnancy and lactation

{As currently approved}.

4.7 Effects on ability to drive and use machines

{As currently approved}.

4.8 Undesirable effects

{As currently approved}.

Paediatric Population

The clinical safety database includes safety data for 249 paediatric patients who received atorvastatin, among which 7 patients were < 6 years old, 14 patients were in the age range of 6 to 9, and 228 patients were in the age range of 10 to 17.

Nervous system disorders

Common: Headache

Gastrointestinal disorders
Common: Abdominal pain

Investigations

Common: Alanine aminotransferase increased, blood creatine phosphokinase increased

Based on the data available, frequency, type and severity of adverse reactions in children are expected to be the same as in adults. There is currently limited experience with respect to long-term safety in the paediatric population.

4.9 Overdose

{As currently approved}.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

{As currently approved}.

Paediatric Population

Heterozygous Familial Hypercholesterolaemia in Paediatric Patients aged 6-17 years old

An 8-week, open-label study to evaluate pharmacokinetics, pharmacodynamics, and safety and tolerability of atorvastatin was conducted in children and adolescents with genetically confirmed heterozygous familial hypercholesterolemia and baseline LDL-C ≥ 4 mmol/L. A total of 39 children and adolescents, 6 to 17 years of age, were enrolled. Cohort A included 15 children, 6 to 12 years of age and at Tanner Stage 1. Cohort B included 24 children, 10 to 17 years of age and at Tanner Stage ≥ 2 .

The initial dose of atorvastatin was 5 mg daily of a chewable tablet in Cohort A and 10 mg daily of a tablet formulation in Cohort B. The atorvastatin dose was permitted to be doubled if a subject had not attained target LDL-C of < 3.35 mmol/L at Week 4 and if atorvastatin was well tolerated.

Mean values for LDL-C, TC, VLDL-C, and Apo B decreased by Week 2 among all subjects. For subjects whose dose was doubled, additional decreases were observed as early as 2 weeks, at the first assessment, after dose escalation. The mean percent decreases in lipid parameters were similar for both cohorts, regardless of whether subjects remained at their initial dose or doubled their initial dose. At Week 8, on average, the percent change from baseline in LDL-C and TC was approximately 40% and 30%, respectively, over the range of exposures.

Heterozygous Familial Hypercholesterolaemia in Paediatric Patients aged 10-17 years old

In a double-blind, placebo controlled study followed by an open-label phase, 187 boys and postmenarchal girls 10-17 years of age (mean age 14.1 years) with heterozygous familial hypercholesterolaemia (FH) or severe hypercholesterolaemia were randomised to atorvastatin (n=140) or placebo (n=47) for 26 weeks and then all received atorvastatin for 26 weeks.. The dosage of atorvastatin (once daily) was 10 mg for the first 4 weeks and up-titrated to 20 mg if the LDL-C level was > 3.36 mmol/l. Atorvastatin significantly decreased plasma levels of total-C, LDL-C, triglycerides, and apolipoprotein B during the 26 week double-blind phase. The mean achieved LDL-C value was 3.38 mmol/l (range: 1.81-6.26 mmol/l) in the atorvastatin group compared to 5.91 mmol/l (range: 3.93-9.96 mmol/l) in the placebo group during the 26-week double-blind phase.

An additional paediatric study of atorvastatin versus colestipol in patients with hypercholesterolaemia aged 10-18 years demonstrated that atorvastatin (N=25) caused a significant reduction in LDL-C at week 26 ($p<0.05$) compared with colestipol (N=31).

A compassionate use study in patients with severe hypercholesterolaemia (including homozygous hypercholesterolaemia) included 46 paediatric patients treated with atorvastatin titrated according to response (some subjects received 80 mg atorvastatin per day). The study lasted 3 years: LDL-cholesterol was lowered by 36%.

The long-term efficacy of atorvastatin therapy in childhood to reduce morbidity and mortality in adulthood has not been established.

The European Medicines Agency has waived the obligation to submit the results of studies with atorvastatin in children aged 0 to less than 6 years in the treatment of heterozygous hypercholesterolaemia and in children aged 0 to less than 18 years in the treatment of homozygous familial hypercholesterolaemia, combined (mixed) hypercholesterolaemia, primary hypercholesterolaemia and in the prevention of cardiovascular events (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

{As currently approved}.

Special populations

{As currently approved}.

- ~~Paediatric: Pharmacokinetic data in the paediatric population are not available.~~ **In an open-label, 8-week study, Tanner Stage 1 (N=15) and Tanner Stage ≥ 2 (N=24) paediatric patients (ages 6-17 years) with heterozygous familial hypercholesterolemia and baseline LDL-C ≥ 4 mmol/L were treated with 5 or 10 mg of chewable or 10 or 20 mg of film-coated atorvastatin tablets once daily, respectively. Body weight was the only significant covariate in atorvastatin population PK model. Apparent oral clearance of atorvastatin in paediatric subjects appeared similar to adults when scaled allometrically by body weight. Consistent decreases in LDL-C and TC were observed over the range of atorvastatin and o-hydroxyatorvastatin exposures.**
- {As currently approved}.

5.3 Preclinical safety data

{As currently approved}.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

{As currently approved}.

6.2 Incompatibilities

{As currently approved}.

6.3 Shelf life

{As currently approved}.

6.4 Special precautions for storage

{As currently approved}.

6.5 Nature and contents of container

{As currently approved}.

6.6 Special precautions for disposal

{As currently approved}.

7. MARKETING AUTHORISATION HOLDER

[To be completed nationally]

[See Annex I - To be completed nationally]

{Name and address}

<{tel}>

<{fax}>

<{e-mail}>

8. MARKETING AUTHORISATION NUMBER(S)

[To be completed nationally]

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

[To be completed nationally]

10. DATE OF REVISION OF THE TEXT

1. NAME OF THE MEDICINAL PRODUCT

[See Annex I - To be completed nationally]
{PRODUCT NAME} 80 mg film-coated tablet

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

{As currently approved}.

3. PHARMACEUTICAL FORM

{As currently approved}.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Hypercholesterolaemia

{PRODUCT NAME} is indicated as an adjunct to diet for reduction of elevated total cholesterol, LDL-cholesterol, apolipoprotein B, and triglycerides in **patients adults, adolescents and children aged 10 years or older** with primary hypercholesterolaemia including familial hypercholesterolaemia (heterozygous variant) or combined (mixed) hyperlipidaemia (Corresponding to Types IIa and IIb of the Fredrickson classification) when response to diet and other nonpharmacological measures is inadequate.

{PRODUCT NAME} is also indicated to reduce total-C and LDL-C in **patients adults** with homozygous familial hypercholesterolaemia as an adjunct to other lipid-lowering treatments (e.g. LDL apheresis) or if such treatments are unavailable.

Prevention of cardiovascular disease

Prevention of cardiovascular events in adult patients estimated to have a high risk for a first cardiovascular event (see Section 5.1), as an adjunct to correction of other risk factors.

4.2 Posology and method of administration

{As currently approved}.

Paediatric use

~~Paediatric use should only be carried out by specialists.~~

~~Experience in pediatrics is limited to a small number of patients (age 4–17 years) with severe dyslipidemias, such as homozygous familial hypercholesterolemia. The recommended starting dose in this population is 10 mg of atorvastatin per day. The dose may be increased to 80 mg daily, according to the response and tolerability. Developmental safety data in this population have not been evaluated.~~

Hypercholesterolaemia:

Paediatric use should only be carried out by physicians experienced in the treatment of paediatric hyperlipidaemia and patients should be re-evaluated on a regular basis to assess progress.

For patients aged 10 years and above, the recommended starting dose of atorvastatin is 10 mg per day with titration up to 20 mg per day. Titration should be conducted according to the individual response and tolerability in paediatric patients. Safety information for paediatric patients treated with doses above 20 mg, corresponding to about 0.5 mg/kg, is limited.

There is limited experience in children between 6-10 years of age (see section 5.1). Atorvastatin is not indicated in the treatment of patients below the age of 10 years.

Other pharmaceutical forms/strengths may be more appropriate for this population.

4.3 Contraindications

{As currently approved}.

4.4 Special warnings and precautions for use

{As currently approved}.

Paediatric use

Developmental safety in the paediatric population has not been established (see section 4.8).

4.5 Interaction with other medicinal products and other forms of interaction

{As currently approved}.

Paediatric population

Drug-drug interaction studies have only been performed in adults. The extent of interactions in the paediatric population is not known. The above mentioned interactions for adults and the warnings in section 4.4 should be taken into account for the paediatric population.

4.6 Pregnancy and lactation

{As currently approved}.

4.7 Effects on ability to drive and use machines

{As currently approved}.

4.8 Undesirable effects

{As currently approved}.

Paediatric Population

The clinical safety database includes safety data for 249 paediatric patients who received atorvastatin, among which 7 patients were < 6 years old, 14 patients were in the age range of 6 to 9, and 228 patients were in the age range of 10 to 17.

Nervous system disorders

Common: Headache

Gastrointestinal disorders
Common: Abdominal pain

Investigations

Common: Alanine aminotransferase increased, blood creatine phosphokinase increased

Based on the data available, frequency, type and severity of adverse reactions in children are expected to be the same as in adults. There is currently limited experience with respect to long-term safety in the paediatric population.

4.9 Overdose

{As currently approved}.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

{As currently approved}.

Paediatric Population

Heterozygous Familial Hypercholesterolaemia in Paediatric Patients aged 6-17 years old

An 8-week, open-label study to evaluate pharmacokinetics, pharmacodynamics, and safety and tolerability of atorvastatin was conducted in children and adolescents with genetically confirmed heterozygous familial hypercholesterolemia and baseline LDL-C ≥ 4 mmol/L. A total of 39 children and adolescents, 6 to 17 years of age, were enrolled. Cohort A included 15 children, 6 to 12 years of age and at Tanner Stage 1. Cohort B included 24 children, 10 to 17 years of age and at Tanner Stage ≥ 2 .

The initial dose of atorvastatin was 5 mg daily of a chewable tablet in Cohort A and 10 mg daily of a tablet formulation in Cohort B. The atorvastatin dose was permitted to be doubled if a subject had not attained target LDL-C of < 3.35 mmol/L at Week 4 and if atorvastatin was well tolerated.

Mean values for LDL-C, TC, VLDL-C, and Apo B decreased by Week 2 among all subjects. For subjects whose dose was doubled, additional decreases were observed as early as 2 weeks, at the first assessment, after dose escalation. The mean percent decreases in lipid parameters were similar for both cohorts, regardless of whether subjects remained at their initial dose or doubled their initial dose. At Week 8, on average, the percent change from baseline in LDL-C and TC was approximately 40% and 30%, respectively, over the range of exposures.

Heterozygous Familial Hypercholesterolaemia in Paediatric Patients aged 10-17 years old

In a double-blind, placebo controlled study followed by an open-label phase, 187 boys and postmenarchal girls 10-17 years of age (mean age 14.1 years) with heterozygous familial hypercholesterolaemia (FH) or severe hypercholesterolaemia were randomised to atorvastatin (n=140) or placebo (n=47) for 26 weeks and then all received atorvastatin for 26 weeks. The dosage of atorvastatin (once daily) was 10 mg for the first 4 weeks and up-titrated to 20 mg if the LDL-C level was > 3.36 mmol/l. Atorvastatin significantly decreased plasma levels of total-C, LDL-C, triglycerides, and apolipoprotein B during the 26 week double-blind phase. The mean achieved LDL-C value was 3.38 mmol/l (range: 1.81-6.26 mmol/l) in the atorvastatin group compared to 5.91 mmol/l (range: 3.93-9.96 mmol/l) in the placebo group during the 26-week double-blind phase.

An additional paediatric study of atorvastatin versus colestipol in patients with hypercholesterolaemia aged 10-18 years demonstrated that atorvastatin (N=25) caused a significant reduction in LDL-C at week 26 ($p<0.05$) compared with colestipol (N=31).

A compassionate use study in patients with severe hypercholesterolaemia (including homozygous hypercholesterolaemia) included 46 paediatric patients treated with atorvastatin titrated according to response (some subjects received 80 mg atorvastatin per day). The study lasted 3 years: LDL-cholesterol was lowered by 36%.

The long-term efficacy of atorvastatin therapy in childhood to reduce morbidity and mortality in adulthood has not been established.

The European Medicines Agency has waived the obligation to submit the results of studies with atorvastatin in children aged 0 to less than 6 years in the treatment of heterozygous hypercholesterolaemia and in children aged 0 to less than 18 years in the treatment of homozygous familial hypercholesterolaemia, combined (mixed) hypercholesterolaemia, primary hypercholesterolaemia and in the prevention of cardiovascular events (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

{As currently approved}.

Special populations

{As currently approved}.

- Paediatric: ~~Pharmacokinetic data in the paediatric population are not available.~~ **In an open-label, 8-week study, Tanner Stage 1 (N=15) and Tanner Stage ≥ 2 (N=24) paediatric patients (ages 6-17 years) with heterozygous familial hypercholesterolemia and baseline LDL-C ≥ 4 mmol/L were treated with 5 or 10 mg of chewable or 10 or 20 mg of film-coated atorvastatin tablets once daily, respectively. Body weight was the only significant covariate in atorvastatin population PK model. Apparent oral clearance of atorvastatin in paediatric subjects appeared similar to adults when scaled allometrically by body weight. Consistent decreases in LDL-C and TC were observed over the range of atorvastatin and o-hydroxyatorvastatin exposures.**
- {As currently approved}.

5.3 Preclinical safety data

{As currently approved}.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

{As currently approved}.

6.2 Incompatibilities

{As currently approved}.

6.3 Shelf life

{As currently approved}.

6.4 Special precautions for storage

{As currently approved}.

6.5 Nature and contents of container

{As currently approved}.

6.6 Special precautions for disposal

{As currently approved}.

7. MARKETING AUTHORISATION HOLDER

[To be completed nationally]

[See Annex I - To be completed nationally]

{Name and address}

<{tel}>

<{fax}>

<{e-mail}>

8. MARKETING AUTHORISATION NUMBER(S)

[To be completed nationally]

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

[To be completed nationally]

10. DATE OF REVISION OF THE TEXT

**AMENDMENTS TO BE INCLUDED IN THE RELEVANT SECTIONS OF THE PACKAGE
LEAFLET**

PACKAGE LEAFLET: INFORMATION FOR THE USER

{PRODUCT NAME} 10 mg film-coated tablets

Atorvastatin calcium

[See Annex I - To be completed nationally]

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What {PRODUCT NAME} is and what it is used for
2. Before you take {PRODUCT NAME}
3. How to take {PRODUCT NAME}
4. Possible side effects
5. How to store {PRODUCT NAME}
6. Further information

1. WHAT {PRODUCT NAME} IS AND WHAT IT IS USED FOR

{As currently approved}.

2. BEFORE YOU TAKE {PRODUCT NAME}

{As currently approved}.

3. HOW TO TAKE {PRODUCT NAME}

The usual starting dose of {PRODUCT NAME} is 10 mg once a day **in adults and children aged 10 years or older**. This may be increased if necessary by your doctor until you are taking the amount you need. Your doctor will adapt the dosage at intervals of 4 weeks or more. The maximum dose of {PRODUCT NAME} is 80 mg once daily **for adults and 20 mg once daily for children**.

{As currently approved}.

4. POSSIBLE SIDE EFFECTS

{As currently approved}.

5. HOW TO STORE {PRODUCT NAME}

{As currently approved}.

6. FURTHER INFORMATION

What {PRODUCT NAME} contains

{As currently approved}.

This medicine is available as 5 mg, 10 mg, 20 mg and 40 mg chewable tablets and 10 mg, 20 mg, 40 mg and 80 mg film-coated tablets.

Marketing Authorisation Holder and Manufacturer

{Name and address}

<{tel}>

<{fax}>

<{e-mail}>

[See Annex I - To be completed nationally]

This medicinal product is authorised in the Member States of the EEA under the following names:

{As currently approved}.

This leaflet was last approved in {MM/YYYY}.

[To be completed nationally]

PACKAGE LEAFLET: INFORMATION FOR THE USER

{PRODUCT NAME} 20 mg film-coated tablets

Atorvastatin calcium

[See Annex I - To be completed nationally]

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What {PRODUCT NAME} is and what it is used for
2. Before you take {PRODUCT NAME}
3. How to take {PRODUCT NAME}
4. Possible side effects
5. How to store {PRODUCT NAME}
6. Further information

1. WHAT {PRODUCT NAME} IS AND WHAT IT IS USED FOR

{As currently approved}.

2. BEFORE YOU TAKE {PRODUCT NAME}

{As currently approved}.

3. HOW TO TAKE {PRODUCT NAME}

The usual starting dose of {PRODUCT NAME} is 10 mg once a day **in adults and children aged 10 years or older**. This may be increased if necessary by your doctor until you are taking the amount you need. Your doctor will adapt the dosage at intervals of 4 weeks or more. The maximum dose of {PRODUCT NAME} is 80 mg once daily **for adults and 20 mg once daily for children**.

{As currently approved}.

4. POSSIBLE SIDE EFFECTS

{As currently approved}.

5. HOW TO STORE {PRODUCT NAME}

{As currently approved}.

6. FURTHER INFORMATION

What {PRODUCT NAME} contains

{As currently approved}.

This medicine is available as 5 mg, 10 mg, 20 mg and 40 mg chewable tablets and 10 mg, 20 mg, 40 mg and 80 mg film-coated tablets.

Marketing Authorisation Holder and Manufacturer

{Name and address}

<{tel}>

<{fax}>

<{e-mail}>

[See Annex I - To be completed nationally]

This medicinal product is authorised in the Member States of the EEA under the following names:

{As currently approved}.

This leaflet was last approved in {MM/YYYY}.

[To be completed nationally]

PACKAGE LEAFLET: INFORMATION FOR THE USER

{PRODUCT NAME} 40 mg film-coated tablets

Atorvastatin calcium

[See Annex I - To be completed nationally]

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What {PRODUCT NAME} is and what it is used for
2. Before you take {PRODUCT NAME}
3. How to take {PRODUCT NAME}
4. Possible side effects
5. How to store {PRODUCT NAME}
6. Further information

1. WHAT {PRODUCT NAME} IS AND WHAT IT IS USED FOR

{As currently approved}.

2. BEFORE YOU TAKE {PRODUCT NAME}

{As currently approved}.

3. HOW TO TAKE {PRODUCT NAME}

The usual starting dose of {PRODUCT NAME} is 10 mg once a day **in adults and children aged 10 years or older**. This may be increased if necessary by your doctor until you are taking the amount you need. Your doctor will adapt the dosage at intervals of 4 weeks or more. The maximum dose of {PRODUCT NAME} is 80 mg once daily **for adults and 20 mg once daily for children**.

{As currently approved}.

4. POSSIBLE SIDE EFFECTS

{As currently approved}.

5. HOW TO STORE {PRODUCT NAME}

{As currently approved}.

6. FURTHER INFORMATION

What {PRODUCT NAME} contains

{As currently approved}.

This medicine is available as 5 mg, 10 mg, 20 mg and 40 mg chewable tablets and 10 mg, 20 mg, 40 mg and 80 mg film-coated tablets.

Marketing Authorisation Holder and Manufacturer

{Name and address}

<{tel}>

<{fax}>

<{e-mail}>

[See Annex I - To be completed nationally]

This medicinal product is authorised in the Member States of the EEA under the following names:

{As currently approved}.

This leaflet was last approved in {MM/YYYY}.

[To be completed nationally]

PACKAGE LEAFLET: INFORMATION FOR THE USER

{PRODUCT NAME} 80 mg film-coated tablets

Atorvastatin calcium

[See Annex I - To be completed nationally]

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What {PRODUCT NAME} is and what it is used for
2. Before you take {PRODUCT NAME}
3. How to take {PRODUCT NAME}
4. Possible side effects
5. How to store {PRODUCT NAME}
6. Further information

1. WHAT {PRODUCT NAME} IS AND WHAT IT IS USED FOR

{As currently approved}.

2. BEFORE YOU TAKE {PRODUCT NAME}

{As currently approved}.

3. HOW TO TAKE {PRODUCT NAME}

The usual starting dose of {PRODUCT NAME} is 10 mg once a day **in adults and children aged 10 years or older**. This may be increased if necessary by your doctor until you are taking the amount you need. Your doctor will adapt the dosage at intervals of 4 weeks or more. The maximum dose of {PRODUCT NAME} is 80 mg once daily **for adults and 20 mg once daily for children**.

{As currently approved}.

4. POSSIBLE SIDE EFFECTS

{As currently approved}.

5. HOW TO STORE {PRODUCT NAME}

{As currently approved}.

6. FURTHER INFORMATION

What {PRODUCT NAME} contains

{As currently approved}.

This medicine is available as 5 mg, 10 mg, 20 mg and 40 mg chewable tablets and 10 mg, 20 mg, 40 mg and 80 mg film-coated tablets.

Marketing Authorisation Holder and Manufacturer

{Name and address}

<{tel}>

<{fax}>

<{e-mail}>

[See Annex I - To be completed nationally]

This medicinal product is authorised in the Member States of the EEA under the following names:

{As currently approved}.

This leaflet was last approved in {MM/YYYY}.

[To be completed nationally]

ANNEX III
CONDITIONS OF THE MARKETING AUTHORISATION

The National Health Authorities shall ensure the following conditions are fulfilled by the Marketing Authorisation Holder:

- Submit a Risk Management Plan (or its update) for Sortis and associated names at National level, taking into account the new paediatric data and the CHMP recommendations. The Risk Management Plan should include the ongoing study A2581173 (3-year study of the safety and follow-up study of efficacy of atorvastatin treatment of children and adolescents aged 6 years to less than 18 years with heterozygous familial hypercholesterolaemia).
- Restart the cycle of PSUR submission for Sortis and associated names as follows:
 - Six-monthly PSURs until two full years of experience with the paediatric indication in the EU has been gained
 - Yearly PSURs for the following two years
 - Thereafter submission at 3-yearly intervals

The PSURs should focus on the use in the paediatric population.