

## **I. SUMMARY OF THE RISK MANAGEMENT PLAN FOR BRETARIS GENUAIR**

This is a summary of the risk management plan (RMP) for Bretaris Genuair. The RMP details important risks of Bretaris Genuair and how more information will be obtained about Bretaris Genuair's risks and uncertainties (missing information).

Bretaris Genuair's Summary of Product Characteristics (SmPC) and its package leaflet give essential information to healthcare professionals and patients on how Bretaris Genuair should be used.

Important new concerns or changes to the current ones will be included in updates of Bretaris Genuair's RMP.

### **I.1 THE MEDICINE AND WHAT IT IS USED FOR**

Bretaris Genuair is authorised for COPD (see SmPC for the full indication). It contains acclidinium bromide as the active substance and it is given by inhalation.

Further information about the evaluation of Bretaris Genuair benefits can be found in Bretaris Genuair's EPAR, including in its plain-language summary, available on the EMA website, under the medicine's webpage:

<https://www.ema.europa.eu/en/medicines/human/EPAR/bretaris-genuair>

### **I.2 RISKS ASSOCIATED WITH THE MEDICINE AND ACTIVITIES TO MINIMISE OR FURTHER CHARACTERISE THE RISKS**

Important risks of acclidinium, together with measures to minimise such risks and the proposed studies for learning more about acclidinium's risks, are outlined below.

Measures to minimise the risks identified for medicinal products can be:

- Specific information, such as warnings, precautions, and advice on correct use, in the package leaflet addressed to patients and healthcare professionals;
- Important advice on the medicine's packaging;
- The authorised pack size — the amount of medicine in a pack is chosen so to ensure that the medicine is used correctly;
- The medicine's legal status — the way a medicine is supplied to the patient (e.g. with or without prescription) can help to minimise its risks.

Together, these measures constitute routine risk minimisation measures.

In addition to these measures, information about adverse reactions is collected continuously and regularly analysed so that immediate action can be taken as necessary. These measures constitute routine pharmacovigilance activities.

If important information that may affect the safe use of acridinium is not yet available, it is listed under missing information below.

### **I.2.1 List of important risks and missing information**

Important risks of acridinium are risks that need special risk management activities to further investigate or minimise the risk, so that the medicinal product can be safely administered. Important risks can be regarded as identified or potential. Identified risks are concerns for which there is sufficient proof of a link with the use of acridinium. Potential risks are concerns for which an association with the use of this medicine is possible based on available data, but this association has not been established yet and needs further evaluation. Missing information refers to information on the safety of the medicinal product that is currently missing and needs to be collected (e.g. on the long-term use of the medicine);

**Table I-1 List of important potential risks and missing information**

Important potential risks	<ul style="list-style-type: none"><li>• Cardiac events (cardiac failure, myocardial infarction)</li><li>• Cerebrovascular events (stroke, transient ischaemic attack)</li><li>• Mortality</li></ul>
Missing Information	<ul style="list-style-type: none"><li>• Patients with the following concomitant illnesses: newly diagnosed or unstable arrhythmias, recent myocardial infarction, unstable angina or heart failure</li></ul>

### **I.2.2 Summary of important risks**

**Table I-2 Important potential risk: Cardiac events**

Evidence for linking the risk to the medicine	Acridinium increases the heart rate in some patients, which might increase the risk of cardiac failure and myocardial infarction.
Risk factors and risk groups	Anticipated patient groups at risk of cardiac events include elderly and those with a history of cardiac disease, hypertension, dyslipidaemia, hyperglycaemia or diabetes mellitus, and body mass index >30 kg/m <sup>2</sup> .
Risk minimisation measures	Routine risk minimisation measures: SmPC Section 4.4 and PIL Section 2
Additional pharmacovigilance activities	Additional pharmacovigilance activities: Post-authorisation safety study D6560R00004 to assess the effect of Bretaris Genuair on mortality, cardiac and cerebrovascular events.

**Table I-3            Important potential risk: Cerebrovascular events**

Evidence for linking the risk to the medicine	If acclidinium increases the risk of myocardial infarction or atrial fibrillation, the risk of blood clots forming in the left atrium or left ventricle also increases, which in turn increases the risk of stroke.
Risk factors and risk groups	COPD patients have a moderate increase in the risk of strokes compared with non-COPD patients. Known risk factors include hypertension, dyslipidaemia, diabetes, atrial fibrillation, arterial stenosis, smoking, obesity, poor diet and physical inactivity.
Risk minimisation measures	Routine risk minimisation measures: None.
Additional pharmacovigilance activities	Additional pharmacovigilance activities: PASS study D6560R00004

**Table I-4            Important potential risk: Mortality**

Evidence for linking the risk to the medicine	Although a risk of increased mortality has been identified for other similar compounds, there is currently no evidence of an increase in mortality with Bretaris Genuair
Risk factors and risk groups	Studies have shown risk of mortality in patients with COPD to be greater in men and greatest in elderly adults aged 75 years and older.
Risk minimisation measures	Routine risk minimisation measures: None.
Additional pharmacovigilance activities	Additional pharmacovigilance activities: PASS study D6560R00004

**Table I-5            Missing information: Patients with the following concomitant illnesses: newly diagnosed or unstable arrhythmias, recent myocardial infarction, unstable angina or heart failure**

Risk minimisation measures	Routine risk minimisation measures: SmPC section 4.4, PIL section 2. Additional pharmacovigilance activities: D6560R00004/PASS (cardiac components),
----------------------------	--

### **I.2.3            Post-authorisation development plan**

#### **I.2.3.1           Studies which are conditions of the marketing authorisation**

Study short name: D6560R00004

Rationale and study objectives: To evaluate the cardiovascular safety concerns and all-cause mortality of acclidinium, acclidinium/formoterol and other bronchodilators used in patients with COPD.