

26 June 2014
EMA/CHMP/365686/2014
Committee for Medicinal Products for Human Use (CHMP)

Summary of opinion¹ (post authorisation)

Avastin bevacizumab

On 26 June 2014, the Committee for Medicinal Products for Human Use (CHMP) adopted a positive opinion recommending a variation to the terms of the marketing authorisation for the medicinal product Avastin. The marketing authorisation holder for this medicinal product is Roche Registration Ltd. They may request a re-examination of the CHMP opinion, provided that they notify the European Medicines Agency in writing of their intention within 15 days of receipt of the opinion.

The CHMP adopted a new indication as follows:

“Bevacizumab in combination with paclitaxel, topotecan, or pegylated liposomal doxorubicin is indicated for the treatment of adult patients with platinum-resistant recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who received no more than two prior chemotherapy regimens and who have not received prior therapy with bevacizumab or other VEGF inhibitors or VEGF receptor–targeted agents (see Section 5.1).”.

Detailed conditions for the use of this product will be described in the updated summary of product characteristics (SmPC), which will be published in the revised European public assessment report (EPAR), and will be available in all official European Union languages after the variation to the marketing authorisation has been granted by the European Commission.

For information, the full indication(s) for Avastin will be as follows²:

Bevacizumab in combination with fluoropyrimidine-based chemotherapy is indicated for treatment of adult patients with metastatic carcinoma of the colon or rectum.

Bevacizumab in combination with paclitaxel is indicated for first-line treatment of adult patients with metastatic breast cancer. For further information as to human epidermal growth factor receptor 2 (HER2) status, please refer to section 5.1.

Bevacizumab in combination with capecitabine is indicated for first-line treatment of adult patients with metastatic breast cancer in whom treatment with other chemotherapy options including taxanes or

¹ Summaries of positive opinion are published without prejudice to the Commission decision, which will normally be issued within 44 days (Type II variations) and 67 days (Annex II applications) from adoption of the opinion.

² The text in bold represents the new or the amended indication.

anthracyclines is not considered appropriate. Patients who have received taxane and anthracycline-containing regimens in the adjuvant setting within the last 12 months should be excluded from treatment with Avastin in combination with capecitabine. For further information as to HER2 status, please refer to section 5.1.

Bevacizumab, in addition to platinum-based chemotherapy, is indicated for first-line treatment of adult patients with unresectable advanced, metastatic or recurrent non-small cell lung cancer other than predominantly squamous cell histology.

Bevacizumab in combination with interferon alfa-2a is indicated for first line treatment of adult patients with advanced and/or metastatic renal cell cancer.

Bevacizumab, in combination with carboplatin and paclitaxel is indicated for the front-line treatment of adult patients with advanced (International Federation of Gynecology and Obstetrics (FIGO) stages III B, III C and IV) epithelial ovarian, fallopian tube, or primary peritoneal cancer.

Bevacizumab, in combination with carboplatin and gemcitabine, is indicated for treatment of adult patients with first recurrence of platinum-sensitive epithelial ovarian, fallopian tube or primary peritoneal cancer who have not received prior therapy with bevacizumab or other VEGF inhibitors or VEGF receptor–targeted agents”.

Bevacizumab in combination with paclitaxel, topotecan, or pegylated liposomal doxorubicin is indicated for the treatment of adult patients with platinum-resistant recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who received no more than two prior chemotherapy regimens and who have not received prior therapy with bevacizumab or other VEGF inhibitors or VEGF receptor–targeted agents (see Section 5.1).