| EU (MA) number | Invented name | Strength | Pharmaceutical form | Target species | Route of administration | Packaging | Content | Package size | Withdrawal period |
|-----------------|---------------|-----------|------------------------|---------------------------|--|--------------|---------|-----------------|--|
| EU/2/20/258/001 | Increxxa | 100 mg/ml | Solution for injection | Cattle, pigs and sheep | Subcutaneous use (cattle), intramuscular use (pigs, sheep) | Vial (glass) | 20 ml | 1 vial | Meat and offal: cattle: 22 days; pigs: 13 days; sheep: 16 days; |
| EU/2/20/258/002 | Increxxa | 100 mg/ml | Solution for injection | Cattle, pigs and sheep | Subcutaneous use (cattle), intramuscular use (pigs, sheep) | Vial (glass) | 50 ml | 1 vial | Meat and offal: cattle: 22 days; pigs: 13 days; sheep: 16 days; |
| EU/2/20/258/003 | Increxxa | 100 mg/ml | Solution for injection | Cattle, pigs and sheep | Subcutaneous use (cattle), intramuscular use (pigs, sheep) | Vial (glass) | 100 ml | 1 vial | Meat and offal: cattle: 22 days; pigs: 13 days; sheep: 16 days; |
| EU/2/20/258/004 | Increxxa | 100 mg/ml | Solution for injection | Cattle, pigs and sheep | Subcutaneous use (cattle), intramuscular use (pigs, sheep) | Vial (glass) | 250 ml | 1 vial | Meat and offal: cattle: 22 days; pigs: 13 days; sheep: 16 days; |
| EU/2/20/258/005 | Increxxa | 100 mg/ml | Solution for injection | Cattle, | Subcutaneous use | Vial (glass) | 500 ml | 1 vial | Meat and offal: cattle: 22 days; |
| EU/2/20/258/006 | Increxxa | 25 mg/ml | Solution for injection | Pigs | Intramuscular use | Vial (glass) | 50 ml | 1 vial | Meat and offal: pigs: 13 days; |
| EU/2/20/258/007 | Increxxa | 25 mg/ml | Solution for injection | Pigs | Intramuscular use | Vial (glass) | 100 ml | 1 vial | Meat and offal: pigs: 13 days; |
| EU/2/20/258/008 | Increxxa | 25 mg/ml | Solution for injection | Pigs | Intramuscular use | Vial (glass) | 250 ml | 1 vial | Meat and offal: pigs: 13 days; |