

MA (EU) Number	Invented name	Strength	Pharmaceutical form	Target species	Route of administration	Packaging	Content	Package size	Withdrawal period
EU/2/09/098/001	Melovem	5 mg/ml	Solution for injection	Cattle, pigs	Subcutaneous use, intramuscular use	Injection vial (type I glass)	100 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs)
EU/2/09/098/002	Melovem	20 mg/ml	Solution for injection	Cattle, horses and pigs	Intramuscular use, intravenous use, subcutaneous use	Injection vial (type I glass)	50 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs, horses) Milk: 5 days (cattle)
EU/2/09/098/003	Melovem	20 mg/ml	Solution for injection	Cattle, horses and pigs	Intramuscular use, intravenous use, subcutaneous use	Injection vial (type I glass)	100 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs, horses) Milk: 5 days (cattle)
EU/2/09/098/004	Melovem	20 mg/ml	Solution for injection	Cattle, horses and pigs	Intramuscular use, intravenous use, subcutaneous use	Injection vial (type I glass)	250 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs, horses) Milk: 5 days (cattle)
EU/2/09/098/005	Melovem	30 mg/ml	Solution for injection	Cattle, pigs	Intramuscular use, subcutaneous use	Injection vial (type I glass)	50 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs) Milk: 5 days (cattle)
EU/2/09/098/006	Melovem	30 mg/ml	Solution for injection	Cattle, pigs	Intramuscular use, subcutaneous use	Injection vial (type I glass)	100 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs) Milk: 5 days (cattle)
EU/2/09/098/007	Melovem	30 mg/ml	Solution for injection	Cattle, pigs	Intramuscular use, subcutaneous use	Injection vial (type I glass)	250 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs) Milk: 5 days (cattle)