

<b>EU (MA) number</b>	<b>Invented name</b>	<b>Strength</b>	<b>Pharmaceutical form</b>	<b>Target species</b>	<b>Route of administration</b>	<b>Packaging</b>	<b>Content</b>	<b>Package size</b>	<b>Withdrawal period</b>
EU/2/97/004/001	Metacam	5 mg/ml	Solution for injection	Cattle and pigs	Subcutaneous or intravenous use (cattle), Intramuscular use (pigs)	Vial (glass)	100 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs)
EU/2/97/004/003	Metacam	1.5 mg/ml	Oral suspension	Dogs	Oral use	Bottle (Polyethylene)	10 ml	1 bottle	Not applicable
EU/2/97/004/004	Metacam	1.5 mg/ml	Oral suspension	Dogs	Oral use	Bottle (Polyethylene)	32 ml	1 bottle	Not applicable
EU/2/97/004/005	Metacam	1.5 mg/ml	Oral suspension	Dogs	Oral use	Bottle (Polyethylene)	100 ml	1 bottle	Not applicable
EU/2/97/004/006	Metacam	5 mg/ml	Solution for injection	Dogs and cats	Subcutaneous or intravenous use (dogs), subcutaneous use (cats)	Vial (glass)	10 ml	1 vial	Not applicable
EU/2/97/004/007	Metacam	20 mg/ml	Solution for injection	Cattle, pigs and horses	Subcutaneous or intravenous use (cattle), intramuscular use (pigs), intravenous use (horses)	Vial (glass)	50 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs/horses) Milk: 5 days
EU/2/97/004/008	Metacam	20 mg/ml	Solution for injection	Cattle, pigs and horses	Subcutaneous or intravenous use (cattle), intramuscular use (pigs), intravenous use (horses)	Vial (glass)	100 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs/horses) Milk: 5 days
EU/2/97/004/009	Metacam	15 mg/ml	Oral suspension	Horses	Oral use	Bottle (Polyethylene)	100 ml	1 bottle	Meat and offal : 3 days

<b>EU (MA) number</b>	<b>Invented name</b>	<b>Strength</b>	<b>Pharmaceutical form</b>	<b>Target species</b>	<b>Route of administration</b>	<b>Packaging</b>	<b>Content</b>	<b>Package size</b>	<b>Withdrawal period</b>
EU/2/97/004/010	Metacam	5 mg/ml	Solution for injection	Cattle and pigs	Subcutaneous or intravenous use (cattle), intramuscular use (pigs)	Vial (glass)	100 ml	12 vials	Meat and offal: 15 days (cattle), 5 days (pigs)
EU/2/97/004/011	Metacam	5 mg/ml	Solution for injection	Dogs and cats	Subcutaneous or intravenous use (dogs), subcutaneous use (cats)	Vial (glass)	20 ml	1 vial	Not applicable
EU/2/97/004/012	Metacam	0.5 mg/ml	Oral suspension	Dogs	Oral use	Bottle (Polyethylene)	15 ml	1 bottle	Not applicable
EU/2/97/004/013	Metacam	0.5 mg/ml	Oral suspension	Dogs	Oral use	Bottle (Polyethylene)	30 ml	1 bottle	Not applicable
EU/2/97/004/014	Metacam	20 mg/ml	Solution for injection	Cattle, pigs and horses	Subcutaneous or intravenous use (cattle), intramuscular use (pigs), intravenous use (horses)	Vial (glass)	50 ml	12 vials	Meat and offal: 15 days (cattle), 5 days (pigs/horses) Milk: 5 days
EU/2/97/004/015	Metacam	20 mg/ml	Solution for injection	Cattle, pigs and horses	Subcutaneous or intravenous use (cattle), intramuscular use (pigs), intravenous use (horses)	Vial (glass)	100 ml	12 vials	Meat and offal: 15 days (cattle), 5 days (pigs/horses) Milk: 5 days
EU/2/97/004/026	Metacam	0.5 mg/ml	Oral suspension	Cats, guinea pigs	Oral use	Polyethylene bottle	15 ml	1 bottle	Not applicable

<b>EU (MA) number</b>	<b>Invented name</b>	<b>Strength</b>	<b>Pharmaceutical form</b>	<b>Target species</b>	<b>Route of administration</b>	<b>Packaging</b>	<b>Content</b>	<b>Package size</b>	<b>Withdrawal period</b>
EU/2/97/004/027	Metacam	20 mg/ml	Solution for injection	Cattle, pigs and horses	Subcutaneous or intravenous use (cattle), intramuscular use (pigs), intravenous use (horses)	Vial (glass)	20 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs/horses) Milk: 5 days
EU/2/97/004/028	Metacam	20 mg/ml	Solution for injection	Cattle, pigs and horses	Subcutaneous or intravenous use (cattle), intramuscular use (pigs), intravenous use (horses)	Vial (glass)	20 ml	12 vials	Meat and offal: 15 days (cattle), 5 days (pigs/horses) Milk: 5 days
EU/2/97/004/029	Metacam	1.5 mg/ml	Oral suspension	Dogs	Oral use	Bottle (Polyethylene)	180 ml	1 bottle	Not applicable
EU/2/97/004/030	Metacam	15 mg/ml	Oral suspension	Horses	Oral use	Bottle (Polyethylene)	250 ml	1 bottle	Meat and offal: 3 days
EU/2/97/004/031	Metacam	20 mg/ml	Solution for injection	Cattle, pigs and horses	Subcutaneous or intravenous use (cattle), intramuscular use (pigs), intravenous use (horses)	Vial (glass)	250 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs/horses) Milk: 5 days
EU/2/97/004/032	Metacam	20 mg/ml	Solution for injection	Cattle, pigs and horses	Subcutaneous or intravenous use (cattle), intramuscular use (pigs), intravenous use (horses)	Vial (glass)	250 ml	6 vials	Meat and offal: 15 days (cattle), 5 days (pigs/horses) Milk: 5 days
EU/2/97/004/033	Metacam	0.5 mg/ml	Oral suspension	Cats, guinea pigs	Oral use	Polyethylene bottle	10 ml	1 bottle	Not applicable

<b>EU (MA) number</b>	<b>Invented name</b>	<b>Strength</b>	<b>Pharmaceutical form</b>	<b>Target species</b>	<b>Route of administration</b>	<b>Packaging</b>	<b>Content</b>	<b>Package size</b>	<b>Withdrawal period</b>
EU/2/97/004/034	Metacam	0.5 mg/ml	Oral suspension	Cats. guinea pigs	Oral use	Polyethylene bottle	3 ml	1 bottle	Not applicable
EU/2/97/004/035	Metacam	5 mg/ml	Solution for injection	Cattle and pigs	Subcutaneous or intravenous use (cattle), intramuscular use (pigs)	Vial (glass)	20 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs)
EU/2/97/004/036	Metacam	5 mg/ml	Solution for injection	Cattle and pigs	Subcutaneous or intravenous use (cattle), intramuscular use (pigs)	Vial (glass)	20 ml	12 vials	Meat and offal: 15 days (cattle), 5 days (pigs)
EU/2/97/004/037	Metacam	5 mg/ml	Solution for injection	Cattle and pigs	Subcutaneous or intravenous use (cattle), intramuscular use (pigs)	Vial (glass)	50 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (pigs)
EU/2/97/004/038	Metacam	5 mg/ml	Solution for injection	Cattle and pigs	Subcutaneous or intravenous use (cattle), intramuscular use (pigs)	Vial (glass)	50 ml	12 vials	Meat and offal: 15 days (cattle), 5 days (pigs)
EU/2/97/004/039	Metacam	2 mg/ml	Solution for injection	Cats	Subcutaneous use	Vial (glass)	10 ml	1 vial	Not applicable
EU/2/97/004/040	Metacam	2 mg/ml	Solution for injection	Cats	Subcutaneous use	Vial (glass)	20 ml	1 vial	Not applicable
EU/2/97/004/041	Metacam	15 mg/ml	Oral suspension	Pigs	Oral use	Bottle (Polyethylene)	100 ml	1 bottle	Meat and offal: 5 days
EU/2/97/004/042	Metacam	15 mg/ml	Oral suspension	Pigs	Oral use	Bottle (Polyethylene)	250 ml	1 bottle	Meat and offal: 5 days
EU/2/97/004/043	Metacam	1 mg	Chewable tablets	Dogs	Oral use	Blisters (Alu/Alu)	7 tablets	1 blister (7 tablets)	Not applicable

<b>EU (MA) number</b>	<b>Invented name</b>	<b>Strength</b>	<b>Pharmaceutical form</b>	<b>Target species</b>	<b>Route of administration</b>	<b>Packaging</b>	<b>Content</b>	<b>Package size</b>	<b>Withdrawal period</b>
EU/2/97/004/044	Metacam	1 mg	Chewable tablets	Dogs	Oral use	Blisters (Alu/Alu)	7 tablets	12 blisters (84 tablets)	Not applicable
EU/2/97/004/045	Metacam	1 mg	Chewable tablets	Dogs	Oral use	Blisters (Alu/Alu)	7 tablets	36 blisters (252 tablets)	Not applicable
EU/2/97/004/046	Metacam	2.5 mg	Chewable tablets	Dogs	Oral use	Blisters (Alu/Alu)	7 tablets	1 blister (7 tablets)	Not applicable
EU/2/97/004/047	Metacam	2.5 mg	Chewable tablets	Dogs	Oral use	Blisters (Alu/Alu)	7 tablets	12 blisters (84 tablets)	Not applicable
EU/2/97/004/048	Metacam	2.5 mg	Chewable tablets	Dogs	Oral use	Blisters (Alu/Alu)	7 tablets	36 blisters (252 tablets)	Not applicable
EU/2/97/004/049	Metacam	0.5 mg/ml	Oral suspension	Cats, guinea pigs	Oral use	Polyethylene bottle	30 ml	1 bottle	Not applicable
EU/2/97/004/050	Metacam	40 mg/ml	Solution for injection	Cattle, and horses	Intravenous use (cattle and horses), subcutaneous use (cattle)	Vial (glass)	50 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (horses) Milk: 5 days (cattle)
EU/2/97/004/051	Metacam	40 mg/ml	Solution for injection	Cattle, and horses	Intravenous use (cattle and horses), subcutaneous use (cattle)	Vial (glass)	100 ml	1 vial	Meat and offal: 15 days (cattle), 5 days (horses) Milk: 5 days (cattle)
EU/2/97/004/052	Metacam	40 mg/ml	Solution for injection	Cattle, and horses	Intravenous use (cattle and horses), subcutaneous use (cattle)	Vial (glass)	50 ml	12 vials	Meat and offal: 15 days (cattle), 5 days (horses) Milk: 5 days (cattle)
EU/2/97/004/053	Metacam	40 mg/ml	Solution for injection	Cattle, and horses	Intravenous use (cattle and horses), subcutaneous use (cattle)	Vial (glass)	100 ml	12 vials	Meat and offal: 15 days (cattle), 5 days (horses) Milk: 5 days (cattle)