



London, 16 July 2009
Doc. Ref.: EMEA/HMPC/577784/2008

**COMMITTEE ON HERBAL MEDICINAL PRODUCTS
(HMPC)**

DRAFT

**COMMUNITY HERBAL MONOGRAPH ON *ECHINACEA PURPUREA* (L.) MOENCH.,
RADIX**

DISCUSSION IN WORKING PARTY ON COMMUNITY MONOGRAPHS AND COMMUNITY LIST (MLWP)	January 2009 May 2009 July 2009
ADOPTION BY HMPC FOR RELEASE FOR CONSULTATION	16 July 2009
END OF CONSULTATION (DEADLINE FOR COMMENTS)	15 December 2009
REDISCUSSION IN WORKING PARTY ON COMMUNITY MONOGRAPHS AND COMMUNITY LIST (MLWP)	
ADOPTION BY HMPC	

Comments should be provided using this [template](#) to hmpc.secretariat@emea.europa.eu
Fax: +44 20 75 23 70 51

KEYWORDS	Herbal medicinal products; HMPC; Community herbal monographs; Traditional use; <i>Echinacea purpurea</i> (L.) Moench.; Echinaceae purpureae radix; purple coneflower root
-----------------	---

**COMMUNITY HERBAL MONOGRAPH ON *ECHINACEA PURPUREA* (L.) MOENCH.,
RADIX**

1. NAME OF THE MEDICINAL PRODUCT

To be specified for the individual finished product.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION^{1,2}

<u>Well-established use</u>	<u>Traditional use</u>
	<p>With regard to the registration application of Article 16d(1) of Directive 2001/83/EC as amended</p> <p><i>Echinacea purpurea</i> (L.) Moench., radix (purple coneflower root)</p> <p>i) Herbal substance Not applicable.</p> <p>ii) Herbal preparations Dry extract (6.5:1), extraction solvent ethanol 45% (v/v).</p>

3. PHARMACEUTICAL FORM

<u>Well-established use</u>	<u>Traditional use</u>
	<p>Herbal preparations in solid dosage forms for oral and oromucosal use.</p> <p>The pharmaceutical form should be described by the European Pharmacopoeia full standard term.</p>

4. CLINICAL PARTICULARS

4.1. Therapeutic indications

<u>Well-established use</u>	<u>Traditional use</u>
	Traditional herbal medicinal product for

¹ The material complies with the Eur. Ph. monograph (ref.: 01/2008:1824)

² The declaration of the active substance(s) for an individual finished product should be in accordance with relevant herbal quality guidance

	<p>supportive treatment of common cold.</p> <p>The product is a traditional herbal medicinal product for use in specified indication exclusively based upon long-standing use.</p>
--	--

4.2. Posology and method of administration

<p><u>Well-established use</u></p>	<p><u>Traditional use</u></p> <p>Posology</p> <p><i>Adolescents, adults, elderly</i></p> <p><u>Oral and oromucosal use:</u></p> <p>1 chewable tablet containing 40 mg extract (6.5:1) and corresponding to 260 mg of herbal substance, every second hour (maximum 9 tablets a day).</p> <p>The use in children under 12 years of age is contraindicated (see section 4.3 'Contraindications').</p> <p>Duration of use</p> <p>The therapy should start at first signs of common cold.</p> <p>If the symptoms persist longer than 10 days during the use of the medicinal product, a doctor or a qualified health care practitioner should be consulted.</p> <p>Method of administration</p> <p>Oral and oromucosal use.</p>
------------------------------------	---

4.3. Contraindications

<p><u>Well-established use</u></p>	<p><u>Traditional use</u></p> <p>Hypersensitivity to the active substance or to plants of the Asteraceae (Compositae) family.</p> <p>Progressive systemic diseases such as tuberculosis, diseases of the white blood cells system, collagenoses, multiple sclerosis, AIDS, HIV infections and other immune diseases.</p> <p>Children under 12 years of age.</p>
------------------------------------	---

4.4. Special warnings and precautions for use

<u>Well-established use</u>	<u>Traditional use</u> If the symptoms worsen or high fever occurs during the use of the medicinal product, a doctor or a qualified health care practitioner should be consulted. There is a possible risk of allergic reactions in sensitive individuals. Those patients should consult their doctor before using <i>Echinacea</i> . There is a possible risk of anaphylactic reactions in atopic patients. Atopic patients should consult their doctor before using <i>Echinacea</i> .
-----------------------------	---

4.5. Interactions with other medicinal products and other forms of interaction

<u>Well-established use</u>	<u>Traditional use</u> None reported.
-----------------------------	--

4.6. Pregnancy and lactation

<u>Well-established use</u>	<u>Traditional use</u> Safety during pregnancy and lactation has not been established. In the absence of sufficient data, the use during pregnancy and lactation is not recommended.
-----------------------------	--

4.7. Effects on ability to drive and use machines

<u>Well-established use</u>	<u>Traditional use</u> No studies on the effects on the ability to drive and use machines have been performed.
-----------------------------	---

4.8. Undesirable effects

<u>Well-established use</u>	<u>Traditional use</u> Hypersensitive reactions (skin reactions). The frequency is not known. If other adverse reactions not mentioned above occur, a doctor or a qualified health care practitioner should be consulted.
-----------------------------	---

4.9. Overdose

<u>Well-established use</u>	<u>Traditional use</u>
	No case of overdose has been reported.

5. PHARMACOLOGICAL PROPERTIES

5.1. Pharmacodynamic properties

<u>Well-established use</u>	<u>Traditional use</u>
	Not required as per Article 16c(1)(a)(iii) of Directive 2001/83/EC as amended.

5.2. Pharmacokinetic properties

<u>Well-established use</u>	<u>Traditional use</u>
	Not required as per Article 16c(1)(a)(iii) of Directive 2001/83/EC as amended.

5.3. Preclinical safety data

<u>Well-established use</u>	<u>Traditional use</u>
	Not required as per Article 16c(1)(a)(iii) of Directive 2001/83/EC as amended, unless necessary for the safe use of the product. Tests on reproductive toxicity, genotoxicity and carcinogenicity have not been performed.

6. PHARMACEUTICAL PARTICULARS

<u>Well-established use</u>	<u>Traditional use</u>
	Not applicable.

7. DATE OF COMPILATION/LAST REVISION

16 July 2009