European Medicines Agency
Evaluation of Medicines for Human Use

London, 8 March 2007
Doc. Ref: EMEA/HMPC/513578/2006

COMMITTEE ON HERBAL MEDICINAL PRODUCTS (HMPC)

DRAFT

LIST OF REFERENCES FOR ASSESSMENT OF *RHAMNUS PURSHIANUS* D.C.

.

The EMEA acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

German monograph of Rhamni purshianae cortex (Amerikanische Faulbaumrinde), BanzNr. 133 vom 21.07.1993

Bundesinstitut für Arzneimittel und Medizinprodukte. Bekanntmachung über die Zulassung und Registrierung von Arzneimitteln – Abwehr von Arzneimittelrisiken, Stufe II – Humanarzneimitteln zur innerlichen Anwendung, ausgenommen nach HAB 1 hergestellte homöopathische Arzneimittel, welche Drogen, Drogenzubereitungen oder isolierte Inhaltsstoffe der Anthranoid- (Hydroxyanthracenderivat-) haltigen Pflanzengattungen Andira, Cassia, Rhamnus, Rheum oder Aloe enthalten (German pharmacovigilance actions for anthranoide-containing laxatives) vom 21. Juni 1996, Bundesanzeiger vom 05.07.1996 Nr. 123, Seite 7581

ESCOP monograph “RHAMNI PURSHIANI CORTEX” (Cascara), 2nd edition 2003, Thieme Verlag Stuttgart, New York

Tarpila S et al. Efficacy of ground flaxseed on constipation in patients with irritable bowel syndrome. Current Topics in Nutraceutical Research 2004; 2: 119-25

Goodman & Gilman’s. The Pharmacological Basis of Therapeutics. Eleventh edition. Section VI / Drugs affecting gastrointestinal function. Stimulant (Irritant) Laxatives

HagerROM 2003, Springer Verlag, Heidelberg: Rhamni purshiani cortex (Cascararinde)

Fairbairn JW et al. Cascarosides A and B. Journal of pharmaceutical sciences 1977; Vol 66 (9): 1300-3

Griffini A et al. Isolation and characterization of pure cascarosides A, B, C, and D. Planta Medica 1992; Vol 58 (7): A593-A594

Manitto P et al. Studies on cascara, Part 2. Structures of cascarosides E and F. Journal of Natural Products 1995; Vol 58 (3): 419-23

Van Os FHL. Anthraquinones derivatives in vegetable laxatives. Pharmacology 1976; 14 (suppl 1): 7-17

7 Westferry Circus, Canary Wharf, London, E14 4HB, UK
Tel. (44-20) 74 18 84 00 Fax (44-20) 75 23 70 51
E-mail: mail@emea.europa.eu http://www.emea.europa.eu

Submission of a review of data from the national toxicology program and relevant to the status of cascara sagrada ingredients as over-the counter drug active ingredients by the American herbal products association, December 19, 2002, Docket No. 78N-036L

Lang W. Pharmacokinetic-metabolic studies with 14C-aloe emodin after oral administration to male and female rats. Pharmacology 1993; 47 (Suppl 1): 110-9

Liang JW et al. Emodin pharmacokinetics in rabbits. Planta medica 1995; Vol 61 (5): 406-8

Capasso F et al. Effect of indomethacin on aloin and 1,8 dioxianthraquinone-induced production of prostaglandins in rat isolated colon. Prostaglandins 1983; 26 (4): 557-62

Cohen MM. The effect of cathartics on prostaglandin synthesis by rat gastro-intestinal tract. Prostaglandins, leukotrienes, and medicine 1982; Vol 8 (4): 389-97

Izzo AA et al. NG-nitro-L-arginine methyl ester reduces senna- and cascara-induced diarrhoea and fluid secretion in the rat. European journal of pharmacology 1996; 301 (1-3): 137-42

Izzo AA et al. The role of constitutive and inducible nitric oxide synthase in senna- and cascara-induced diarrhoea in the rat. European journal of pharmacology 1997; 323 (1): 93-7

Dreessen M, Lemli J. Studies in the field of drugs containing anthraquinone deriviatives. XXXVI. The metabolism of cascarosides by intestinal bacteria. Pharm. Acta Helv 1988; 63 Nr. 9-10: 287-9

De Witte P, Lemli L. The metabolism of anthranoid laxatives. Hepato-gastroenterology 1990; 37 (6): 601-5

Müller SO et al. Biotransformation of the anthraquinones emodin and chrysophanol by cytochrome P450 enzymes. Bioactivation to genotoxic metabolites. Drug metabolism and disposition: the biological fate of chemicals 1998; 26 (6): 540-6

NTP Technical Report on the Toxicology and Carcinogenesis Studies of Emodin in F344/N Rats and B6C3F$_1$ Mice. National Toxicology Programm, June 2001, NTP TR 493, NIH Publication No. 01-3952

Masuda T et al. 2-Hydroxyemodin, an active metabolite of emodin in the hepatic microsomes of rats. Mutation Research 1985; 149: 327-332

Mengs U et al. Lack of emodin genotoxicity in the mouse micronucleus assay. Mutation Research 1997; 393: 289-93

Jahnke GD et al. Developmental toxicity evaluation of emodin in rats in mice. Birth Defects Research Part B – Developtmental and Reproductive Toxicology 2004; 71 (2): 89-101

Heidemann a et al. Genotoxicity of aloeemodin in vitro and in vivo. Mutation Research 1996; 367: 123-3

Schörkhuber M et al. Effect of anthraquinone-laxatives on the prolifaration and urokinase secretion of normal, premalignant and malignant colonic epithelial cells. European journal of cancer: official journal for European Organisation for Research and Treatment of Cancer (EORTC) (and) European Association for Cancer Research (EACR) 1998; 34 (7): 1091-8

Siegers CP et al. Sennosides and aloin do not promote dimethylhydrazine-induced colorectal tumors in mice. Pharmacology 1993; 47, No.1: 205-8

Mereto E et al. Evaluation of the potential carcinogenic activity of senna and cascara glycosides for the rat colon. Cancer letters 1996; 101 (1): 79-83

Borelli F et al. Effect of bisacodyl and cascara on growth of aberrant crypt foci and malignant tumors in the rat colon. Life sciences 2001; 69 (16): 1871-7

Vyth A and Kamp PE. Detection of anthraquinone laxatives in the urine. Pharm Weekbl SCI ED 1979; No.2: 456-59

Ewe K. Obstipation und Laxanzien. Dtsch. Apoth. Ztg. 1994; 134: 825-33

Gabler F. Obstipation und Laxanzien. PTA heute 8,1994, 5, 380-8

Tavares IA et al. Effects of anthraquinone derivatives on PAF release by human gastrointestinal mucosa in vitro. Phytotherapy Research 1996; 10 (Suppl.1): S20-S21

Izzo AA et al. Recent findings on the mode of action of laxatives: the role of platelet activating factor and nitric oxide. Trends in pharmacological sciences 1998; 19 (10): 403-5

Koch A. Aloe als Laxans: ein neues Argument für die Überprüfung der Droge (Aloe as a laxative: a new argument for the investigation of the drug). Dtsch Apoth Ztg 1995; 135 (12): 29-32

Mueller SO et al. Occurence of emodin, chrysophanol and physcion in vegetables, herbs and liquors. Genotoxicity and anti-genotoxicity of the anthraquinones and of the whole plants. Food and Chemical Toxicology 1999; 37: 481-91

Che QM et al. Isolation of a human intestinal bacterium capable of transforming barabaloin to aloe-emodin anthrone. Planta medica 1991; 57 (1): 15-9

Yagi T, Yamauchi K. Synergistic effects of anthraquinones on the purgative activity of rhein anthrone in mice. Journal of Pharmacy and Pharmacology 1999; 51 (1): 93-5

Stern FH. Constipation – an omnipresent symptom: effect of a preparation containing prune concentrate and cascarin. Journal of the American Geriatrics Society 1966; 14 (11): 1153-5

Mauracher R et al. Gekreuzter Doppelblindvergleich von gereinigtem Glucofrangulin mit zwei anderen Laxantien. Comparison of glucofrangulin in purified form with two other laxatives. Praxis 1976; 65 (21): 655-8

Borgia M et al. A double-blind double-controlled multicenter study of the therapeutic efficacy of a well-known combination of medicinal herbs. Clin Ter 1985; 114 (5): 401-9

Foert E et al. Cascara stimulates gallbladder contraction: a randomised placebo-controlled double-blind trial. Gastroenterology 1994; 106 (4 Suppl): A338

Wicke L. Zur Darmreinigung vor Irrigoskopien. Bowel cleansing before irrigoscopy. Wiener medizinische Wochenschrift 1981; 131 (5): 133-4

Novetsky GJ et al. Cleansing the colon in gallium-67 scintigraphy: a prospective comparison of regimens. AJR American journal of roentgenology 1981; 137 (5): 979-81

Fork FT. Colon cleansing regimens. Gastrointest Radiol 1982; 7: 383-9

Hangartner PJ et al. Comparison of three colon cleansing methods: evaluation of a randomised clinincal trail with 300 ambulatory patients. Endoscopy 1989; 21: 272-5

Phillip J et al. Vorbereitung zur Koloskopie mit “Golytely” – eine sichere Methode? Vergleichende histologische und klinische Untersuchung zwischen Lavage und salinischem Laxans. Preparation for colonoscopy using Golytely – a sure method? Comparative histological and clinical study between lavage and saline laxatives. Medizinische Klinik 1990; 85 (7): 415-20

Børkje B et al. Effectiveness and acceptability of three bowel cleansing regimes. Scandinavian journal of gastroenterology 1991; 26 (2): 162-6

D’Angelo L: Effects of cascarosides and their metabolites on colonic intestinal muscle. Acta Toxicologica et Therapeutica 1993; 14 (3): 193-7

Williams CL et al. A summary of conference recommendations on dietary fiber in childhood: Conference on Dietary Fiber in Childhood, New York (New York, USA), May 24, 1994. Pediatrics 1995; 96: 1023-28

HagerROM 2003, Springer Verlag, Heidelberg: Frangulae cortex (Faulbaumrinde)

Faber P et al. Relevance of rhein excretion into breast milk. Pharmacology 1988; 36 (Suppl 1) 212-20

Heidemann A et al. The Genotoxicity Status of Senna. Pharmacology 1993; 47, Suppl. 1: 178-186

Sitzmann FC et al. Lokale Nebenwirkungen an der Haut durch Anthraglykoside. Pädiatr Prax 1979; 21: 355-8

Beuers U et al. Hepatitis after chronic abuse of senna. Lancet (England) 1991; Vo.: 337: 372-373 (Senna reference 91)

Vanderperren B et al. Acute liver failure with renal impairment related to the abuse of senna anthraquinone glycosides. Ann Pharmacotherapy 2005; 39: 1353-7 (Senna reference 122)

Müller-Lissner S. Myths and misconceptions about chronic constipation. Am J Gastroenterol 2005; 100: 232-42 (corresponding to senna reference 111)

Smith B. Effect of irritant purgatives on the myentric plexus in man and mouse. Gut 1968; 9: 139-45 (corresponding to senna reference 112)

Riemann JF et al. The fine structure of colonic submucosal nerves in patients with chronic laxative abuse. Scand J Gastroenterol 1980; 15: 761-8 (corresponding to senna reference 113)

Riemann JF et al. Ultrastructural changes in the gut autonomic nervous system following laxative abuse and in other conditions. Scand J Gastroenterol 1982; 71: 11-24 (corresponding to senna reference 114)

Berkelhammer C et al. Low-volume oral colonoscopy bowel preparation: sodium phosphate and magnesium citrate. Gastrointest Endosc 2002; 56: 89-94 (corresponding to senna reference 115)

Meisel JL et al. Human rectal mucosa: proctoscopic and morphological changes caused by laxatives. Gastroenterology 1977; 72: 1274-9 (corresponding to senna reference 116)

Pockros PJ et al. Golytely lavage versus a standard colonoscopy preparation. Effect on normal colonic mucosal histology. Gastroenterology 1985; 88: 545-8 (corresponding to senna reference 117)

Riecken EO et al. The effect of an anthraquinone laxative on colonic nerve tissue: A controlled trial in constipated women. Z Gastroenterol 1990; 28: 660-4 (corresponding to senna reference 118)

Danan G and Benichou C. Causality assessment of adverse reactions to drugs – I. A novel method based on the conclusions of international consensus meetings: application to drug-induced liver injuries. J Clin Epidemiology 1993; Vol 46 (11):1323-30

Benichou C et al. Causality assessment of adverse reactions to drugs – II. An original model for validation of drug causality assessment methods: case reports with positive rechallenge. J Clin Epidemiology 1993; Vol.46 (11): 1331-36

Nadir A et al. Cascara sagrada-induced intrahepatic cholestasis causing portal hypertension: case report and review of herbal hepatotoxicity. American Journal of Gastroenterology 2000; 95 (1): 3634-7
Adesunloye BA Acute renal failure due to herbal remedy CKLS. The American Journal of Medicine 2003; 115: 506-7

Madaus G. Lehrbuch der Biologischen Heilmittel, Band I, Georg Thieme Verlag, Leipzig 1938

The British Pharmaceutical Codex, published by direction of the Council of the Pharmaceutical Society of Great Britain,1911.

The Eclectic Materia Medica, Pharmacology and Therapeutics, by Harvey Wickes Felter, M.D., 1922

Hagers Handbuch der Pharmazeutischen Praxis, Dr. G. Frerichs, G. Arends, Dr. H. Zörnig (Herausgeber), 2. Band: 558 – 63, Berlin, Verlag von Julius Springer 1927

Martindale – The Extra Pharmacopoiea – 25th ed 1967. RG Todd (editor). London. The pharmaceutical press

Dragendorff G. Die Heilpflanzen der verschiedenen Völker und Zeiten. Werner Fritsch, München 1967
The Dispensatory of the United States of America, edited by Joseph P. Remington, Horatio C. Wood and others, 1918

Giavina-Bianchi Jr PF et al. Occupational respiratory allergic disease induced by Passiflora alata and Rhamnus purshiana. Annals of Allergy, Asthma & Immunology 1997; 79: 449-54

Schmidt L. Vergleichende Pharmakologie und Toxikologie der Laxantien. Arch Exper Path Pharmakol 1955; 226: 207-18

Konsensuspapier Expertenforum: Obstipation und Laxanzien. Praxis 1999; 88: 1269-74 (corresponding to senna reference 104)

De Witte P; Lemli J: Metabolism of 14C-rhein and 14C-rhein anthrone in rats. Pharmacology 1988; 36 (Suppl 1): 152-7

Westendorf J et al. Genotoxicity of naturally occurring hydroxyanthraquinones. Mutat Res 1990; 240: 1-12

Helmholz H et al. Genotoxizität der Faulbaumrinde. Genotoxicity of buckthorn bark. Pharmazeutische Zeitung 1993; 138 (Oct 28): 48-50

Steinegger. Pharmakognosie – Phytopharmazie, Springer Verlag 7. Auflage

Siegers C-P et al. Anthranoid laxative abuse – A risk for colorectal cancer? Gut 1993; 34 (8): 1099-101 (corresponding to Frangula 41)

Kune GA et al. The role of chronic constipation, diarrhea, and laxative use in the etiology of large-bowel cancer. Data from the Melbourne Colorectal Cancer Study. Diseases of the colon and rectum 1988; 31 (7): 507 – 12 (corresponding to Frangula 42)

Kune GA. Laxative use not a risk for colorectal cancer: data from the Melbourne Colorectal Cancer Study. Zeitschrift für Gastroenterologie 1993; 31 (2): 140-3 (corresponding to Frangula 43)

Nusko G et al. Retrospective study on laxative use and melanosis coli as risk factors for colorectal neoplasma. Pharmacology 1993; 47 Suppl 1: 234-41 (corresponding to Frangula 44)

Sonnenberg A, Muller AD. Constipation and cathartics as risk factors of colorectal cancer: A meta-analysis. Pharmacology 1993; 47 (Suppl.): 224-33 (corresponding to Frangula 45)

Loew D et al. Anthranoid laxatives: cause of colorectal cancer? Deutsche Apotheker-Zeitung 1994; 134 (Aug 25): 30, 32-33 (corresponding to Frangula 46)

Jacobs EJ et White E. Constipation, laxative use, and colon cancer among middle-aged adults. Epidemiology 1998; 9: 385-91 (corresponding to senna reference 109)

Nusko G et al. Anthranoide laxative use is not a risk factor for colorectal neoplasia: results of a prospective case control study. Gut 2000; 46: 651-55 (corresponding to Frangula 47)

Roberts MC et al. Constipation, laxative use, and colon cancer in a North Carolina population. Am J Gastroenterol 2003; 98: 857-64 (corresponding to senna reference 110)

Nilsson SE et al. Laxative treatment elevates plasma homocysteine: A study on a population-based Swedish sample of old people. European Journal of Clinical Pharmacology 2004; 60 (1): 45-49 (corresponding to Frangula 49)

Jae Sik Joo et al. Alterations in colonic anatomy induced by chronic stimulant laxatives: The cathartic colon revisited. Journal of Clinical Gastroenterology 1998; 26 (4): 283-6 (corresponding to Frangula 50)

Van Gorkom BA et al. Review article: anthranoid laxatives and their potential carcinogenic effects. Alimentary pharmacology & therapeutics 1999; 13 (4): 443-52 (corresponding to Frangula 51)

Willems M et al. Anthranoid self-medication causing rapid development of melanosis coli. The netherlands journal of medicine 2003; 61 (1): 22-4 (corresponding to Frangula 48)