



EUROPEAN MEDICINES AGENCY
SCIENCE MEDICINES HEALTH

27 January 2011
EMA/HMPC/560733/2010
Committee on Herbal Medicinal Products (HMPC)

List of references supporting the assessment of *Chamaemelum nobile* (L.) All., flos

Draft

The Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Abramson W, Basch E, Cheung L, Dacey C, Giese N, Hashmi S, Santos A, Seamon E, Ulbricht C, Varghese M, Weissner W, Woods J. Chamomile (*Matricaria recutita*, *Chamaemelum nobile*). In: Ulbricht C, Basch E editors. Natural Standard Professional Database, Foods, Herbs & Supplements. Natural Standard Inc. 2010. Available at:
<http://www.naturalstandard.com/naturalstandard/monographs/monoframeset.asp?monograph=/monographs/herbssupplements/aux1-chamomile.asp&patientVersion=/monographs/herbssupplements/patient-chamomile.asp>
Accessed 01/07/2010.

Abou-Zied EN, Rizk AM. Phytochemical investigation of *Anthemis nobilis* growing in Egypt. *Qual Plant Mater Veg* 1973, 22:141-144.

Antonelli A, Fabbri C. Study on Roman chamomile (*Chamaemelum nobile* L. All.) oil. *JEOR* 1998, 10:571-574.

Augustin B, Javorka S, Giovannini R, Rom P. Magyar gyógynövények [Hungarian Herbal Drugs] I. Általános és leíró rész (I. General and describing part) Földművelésügyi Minisztérium, Budapest, 1948, 299-300. [Hungarian]

Bail S, Buchbauer G, Jirovetz L, Denkova Z, Slavchev A, Stoyanova A, Schmidt E, Geissler M. Antimicrobial Activities of Roman Chamomile Oil From France and Its Main Compounds. *JEOR* 2009, 21:283-286.

Balbaa SI, Zaki AY, El-Zalabani SM. The volatile oil of *Anthemis nobilis* L. growing in Egypt. *Egypt J Pharmaceut Sci* 1975, 16:161-173.

Bandoniene D, Pukalskas A, Venskutonis PR, Gruzdiene D. Preliminary screening of antioxidant activity of some plant extracts in rapeseed oil. *Food Res Int* 2000, 33:785-791.



- Barbetti P, Casinovi CG. [Sesquiterpene lactones of pharmacological importance: structural and functional properties. II. Germacranolides]. *Ann Ist Super Sanita* 1981, 17:283-322.
- Barnes J, Anderson LA, Phillipson JD. Herbal Medicines. 3rd ed. Pharmaceutical Press, London 2002, 156-158.
- Benesova V, Herout V, Sorm F. Terpenes. CLXX. Isolation and structure of nobilin, a sesquiterpenic lactone with a ten-membered ring. *Collect Czech Chem Commun* 1964, 29:3096-3101.
- Benesova V, Samek Z, Herout V, Sorm F. The structure of nobilin. *Tetrahedron Lett* 1970, 57:5017-5020.
- Bisset NG, editor. Herbal Drugs and Phytopharmaceuticals (Translated from: Wichtl M, editor, Teedrogen. Ein Handbuch für die Praxis auf wissenschaftlicher Grundlage. 2nd ed). Medpharm Scientific Publishers GmbH, Stuttgart 1994. 140-142.
- Blumenthal M, Busse WR, Goldberg A, Gruenwald J, et al., editors. The Complete German Commission E Monographs; Therapeutic Guide to Herbal Medicines. American Botanical Council, Austin Texas 1998, 320-321.
- Bohlmann F, Bornowski H, Schönowsky H. Polyacetylenverbindungen, XXXIX Über heterocyclisch substituierte Acetylenverbindungen aus dem Tribus Anthemideae L. *Chem Ber* 1962, 95:1733-1741.
- Bohlmann F, Burkhardt T, Zdero C. Naturally Occuring Acetylenes. Academic Press, London, New York 1973.
- Bohlmann F, Zdero C. Polyacetylenverbindungen, IC. Über ein Thiophenlacton aus Chamaemelum nobile L. *Chem Ber* 1966, 99:1226-1228.
- Bossuyt L, Dooms-Goossens A. Contact sensitivity to nettles and camomile in 'alternative' remedies. *Contact Dermatitis* 1994, 31:131-132.
- Bradley PR, ed. British herbal compendium, Vol. 1. British Herbal Medicine Association, Guildford 1992, 191-193.
- British Herbal Pharmacopoeia 1974 (BHP, 1974) Sections One, Two & Three. British Herbal Medicine Association, London 1971.
- Bruneton J. Pharmacognosy, Phytochemistry, Medicinal Plants. 2nd ed. Lavoisier Publishing Inc., Paris 1999, 335-337.
- Chao SC, Young DG, Oberg CJ, Nakaoka K. Inhibition of methicillin-resistant Staphylococcus aureus (MRSA) by essential oils. *Flav Fragr J* 2008, 23:444-449.
- Chao SC, Young DG, Oberg CJ. Screening for inhibitory activity of essential oils on selected bacteria, fungi and viruses. *JEOR* 2000, 12:639-649.
- Chaumont JP. Polyphenols (flavonoids, phenolic acids, coumarins) of the capitula of the Roman camomile (Anthemis nobilis). *Plant Med Phytoter* 1969, 3:167-174.
- Chialva F, Gabri G, Liddle PAP, Ulian F. Qualitative Evaluation of Aromatic Herbs by Direct Headspace GC Analysis - Applications of the Method and Comparison with the Traditional Analysis of Essential Oils. *J High Resolut Chrom Chrom Comm* 1982, 5:182-188.
- Choi WI, Lee SG, Park HM, Ahn YJ. Toxicity of Plant Essential Oils to Tetranychus urticae (Acari: Tetranychidae) and Phytoseiulus persimilis (Acari: Phytoseiidae). *J Econ Entomol* 2004, 97:553-558.
- Duarte MCT, Figueira GM, Sartoratto A, Rehder VLG, Delarmelina C. Anti-Candida activity of Brazilian medicinal plants. *J Ethnopharmacol* 2005, 97:305-311.

Eddouks M, Lemhadri A, Zeggwagh NA, Michel JB. Potent hypoglycaemic activity of the aqueous extract of *Chamaemelum nobile* in normal and streptozotocin-induced diabetic rats. *Diabetes Res Clin Pr* 2005, 67:189-195.

European Pharmacopoeia 6th ed. Chamomile flower, Roman – *Chamomillae romanae flos*. Council of Europe. 1/2008: 0380; 1487-1488.

Evans WC. Trease and Evan's Pharmacognosy. 13th ed. Bailliere Tindall, London 1989, 471-473.

Farnsworth NR, Bingel AS, Cordell GA, Crane FA, Fong HH. Potential value of plants as sources of new antifertility agents I. *J Pharm Sci* 1975, 64:535-598.

Fauconnier ML, Jaziri M, Homes J, Shimomura K, Marlier M. *Anthemis nobilis* L. (Roman chamomile): In vitro culture, micropropagation, and the production of essential oils. In: Bajaj YPS, editor. *Biotechnology in Agriculture and Forestry*. Vol. 37. Springer-Verlag, Berlin 1996, 16-37.

FDA, 2010. Listing of Food Additive Status Part I. Available at:

<http://www.fda.gov/Food/FoodIngredientsPackaging/FoodAdditives/FoodAdditiveListings/ucm091048.htm>

Accessed 21/10/2010.

Franke R, Schilcher H, editors. *Chamomile – Industrial Profiles*. CRC Press, Boca Raton 2005.

Giordano-Labadie F, Schwarze HP, Bazex J. Allergic contact dermatitis from camomile used in phytotherapy. *Contact Dermatitis* 2000, 42:247.

Grabarczyk B, Drozd B, Hladon B, Wojciechowska J. Sesquiterpene lactones. Part XV. New cytostatic active sesquiterpene lactone from herb of *Anthemis nobilis* L. *Pol J Pharmacol Pharm* 1977, 29:419-423.

Hänsel R, Keller K, Rimpler H, Schneider G, editors. *Hagers Handbuch der Pharmazeutischen Praxis. Drogen A-D*. Vol. 4. 5th ed. Vol 5. Springer-Verlag, Berlin 1993, 808-817.

Hausen BM, Vieluf IK, editors. *Allergiepflanzen – Handbuch und Atlas*. Nikol Verlagsgesellschaft mbH. and Co. KG, Hamburg 1997, 90-91.

Herisset A, Chaumont JP, Paris RR. Phenolic acids and coumarins from Roman camomile (*Anthemis nobilis*), simple variety. *Plant Med Phytoter* 1974, 8:306-313.

Herisset A, Chaumont JP, Paris RR. Flavonoids of Roman camomille (*Anthemis nobilis*), simple variety. *Plant Med Phytoter* 1973, 7:234-240.

Herisset A, Chaumont JP, Paris RR. Roman camomile (*Anthemis nobilis*) polyphenols. *Plant Med Phytoter* 1970, 4:189-200.

Herisset A, Paris RR, Chaumont JP. Flavonoids of Roman camomile (*Anthemis nobilis*). *Plant Med Phytoter* 1971, 5:234-239.

Hiller K, Melzig MF. *Lexikon der Arzneipflanzen und Drogen*. Erster Band A bis K. Spektrum Akademischer Verlag GmbH, Heidelberg, Berlin 1999, 179-180.

Holub M, Samek Z. Terpenes 235. Isolation and Structure of 3-Epinobilin, 1.10-Epoxyneobilin and 3-Dehydronobilin - Other Sesquiterpenic Lactones from Flowers of *Anthemis-Nobilis* I Revision of Structure of Nobilin and Eucannabinolide. *Collect Czech Chem Commun* 1977, 42:1053-1064.

Inouye S, Uchida K, Abe S. Vapor activity of 72 essential oils against a Trichophyton mentagrophytes. *J Infect Chemother* 2006, 12:210-216.

- Klimes I, Lamparsky D. Unsaturated components in the essential oil of *Anthemis nobilis* L. (Roman chamomile). *Perfum & Flavor* 1984, 9:1-13.
- König GM, Wright AD, Keller WJ, Judd RL, Bates S, Day C. Hypoglycaemic activity of an HMG-containing flavonoid glucoside, chamaemeloside, from *Chamaemelum nobile*. *Planta Med* 1998, 64:612-614.
- Laurelle R. [Roman chamomile; technics for its culture and drying.]. *J Pharm Belg* 1959, 14:232-241.
- Leung AY. Encyclopedia of common natural ingredients used in foods, drugs and cosmetics. John Wiley, New York 1980.
- Lukacs M. [The phytochemical investigation of *Anthemis nobilis* L.] PhD Thesis, Szent-Györgyi Albert Medical University Szeged 1990.
- Ma CM, Winsor L, Daneshtalab M. Quantification of spiroether isomers and herniarin of different parts of *Matricaria matricarioides* and flowers of *Chamaemelum nobile*. *Phytochem Anal* 2007, 18:42-49.
- Maddocks-Jennings W. Critical incident: idiosyncratic allergic reactions to essential oils. *Complement Ther Nurs Midwifery* 2004, 10:58-60.
- Magro A, Carolino M, Bastos M, Mexia A. Efficacy of plant extracts against stored-products fungi. *Rev Iberoam Micol* 2006, 23:176-178.
- Mayer R, Rucker G. Peroxides as plant constituents, III: 1 β -Hydroperoxyisobornoline, a new hydroperoxide from the flowers of *Anthemis nobilis*. *Arch Pharm (Weinheim)* 1987, 320:318-322.
- McGeorge BC, Steele MC. Allergic contact dermatitis of the nipple from Roman chamomile ointment. *Contact Dermatitis* 1991, 24:139-140.
- Melegari M, Albasini A, Pecorari P, Vampa G, Rinaldi M, Rossi TBA. Chemical characteristics and pharmacological properties of the essential oils of *Anthemis nobilis*. *Fitoterapia* 1988, 59:449-455.
- Nano GM, Sacco T, Frattini C. Ricerche botaniche e chimiche su *Anthemis nobilis* L. ed alcune sue cultivar. *Essenze Deriv Agrum* 1976, 46:171-175.
- Newall CA, Anderson LA, Phillipson JD. Herbal Medicines – A Guide for Health-Care Professionals. Pharmaceutical Press, London 1996, 72-73.
- Omidbaigi R, Sefidkon F, Kazemi F. Influence of drying methods on the essential oil content and composition of Roman chamomile. *Flav Fragr J* 2004, 19:196-198.
- Omidbaigi R, Sefidkon F, Kazemi F. Roman chamomile oil: comparison between hydrodistillation and supercritical fluid extraction. *JEOBP* 2003, 6:191-194.
- Opdyke DJL. Monographs on fragrance raw materials: Chamomile oil German and Roman. *Food Cosmet Toxicol* 1974, 12(Suppl.):1423-1425.
- Paulsen E. Contact sensitization from Compositae-containing herbal remedies and cosmetics. *Contact Dermatitis* 2002, 47:189-198.
- PDR for Herbal Medicines, 4th ed. Thomson, Montvale 2000.
- Pereira F, Santos R, Pereira A. Contact dermatitis from chamomile tea. *Contact Dermatitis* 1997, 36:307.
- Piccaglia R, Marotti M, Giovanelli E, Deans SG, Eaglesham E. Antibacterial and antioxidant properties of Mediterranean aromatic plants. *Ind Crop Prod* 1993, 2:47-50.

Pietta P, Mauri P, Bruno A, Rava A, Manera E, Ceva P. Identification of Flavonoids from Ginkgo-Biloba L, Anthemis-Nobilis I and Equisetum-Arvense I by High-Performance Liquid-Chromatography with Diode-Array UV Detection. *J Chrom A* 1991, 553:223-231.

Podsedeck A, Sikora M, Kalembe D. The determination of antioxidant activity of essential oils. *SOFW Journal* 2009, 135:4-5.

Povilaityte V, Venskutonis PR. Antioxidative activity of purple peril (*Perilla frutescens* L.), Moldavian dragonhead (*Dracocephalum moldavica* L.), and Roman chamomile (*Anthemis nobilis* L.) extracts in rapeseed oil. *JAOCs* 2000, 77:951-956.

Rápóti J, Romváry V. Gyógyító növények. 4th ed. Medicina Kiadó, Budapest 1974, 219.

Rossi T, Melegari M, Bianchi A, Albasini A, Vampa G. Sedative, anti-inflammatory and anti-diuretic effects induced in rats by essential oils of varieties of *Anthemis nobilis*: a comparative study. *Pharmacol Res Commun* 1988, 5:71-74.

Rucker G, Mayer R, Lee KR. [Peroxides as plant constituents. 6. Hydroperoxides from the blossoms of Roman chamomile, *Anthemis nobilis* L]. *Arch Pharm (Weinheim)* 1989, 322:821-826.

Samek Z, Holub M, Grabarczyk H, Drozd B, Herout V. Terpenes 236. Structure of Hydroxyisobornil - Cytostatically Active Sesquiterpenic Lactone from Leaves of *Anthemis-Nobilis* I. *Collect Czech Chem Commun* 1977, 42:1065-1068.

Schrader A, Eckey H, Rohr M. Examination of the activity of irradiation-lowering substances on the human skin exemplified with various chamomile extracts. *SOFW Journal* 1997, 123:3-11.

Tognolini M, Barocelli E, Ballabeni V, Bruni R, Bianchi A, Chiavarini M, Impicciatore M. Comparative screening of plant essential oils: phenylpropanoid moiety as basic core for antiplatelet activity. *Life Sci* 2006, 78:1419-1432.

Tschan GM, König GM, Wright AD, Sticher O. Chamaemeloside, a new flavonoid glycoside from *Chamaemelum nobile*. *Phytochemistry* 1996, 41:643-646.

Venskutonis PR, Gruzdiene DTD, Tirzitis G. Assessment of antioxidant activity of plant extracts by different methods. In: Baser, KHC, Franz, G, Canigueral, S, Demirci, F et al. editors. The IIIrd World Congress on Medicinal and Aromatic Plants. *Acta Hort* (Proceedings of WOCMAP III: Perspectives in Natural Product Chemistry) 2005, 677:99-107.

Wichtl M, editor. Herbal Drugs and Phytopharmaceuticals (Translated from: Wichtl M, editor, Teedrogen und Phytopharmaka. Ein Handbuch für die Praxis auf wissenschaftlicher Grundlage. 4th ed). CRC Press, Boca Raton 2004, 127-129.

Wilkinson S, Aldridge J, Salmon I, Cain E, Wilson B. An evaluation of aromatherapy massage in palliative care. *Pall Med* 1999, 13:409-417.

Witherup KM, McLaughlin JL, Judd RL, Ziegler MH, Medon PJ, Keller WJ. Identification of 3-Hydroxy-3-Methylglutaric Acid (HMG) as a Hypoglycemic Principle of Spanish Moss (*Tillandsia Usneoides*). *J Nat Prod* 1995, 58:1285-1290.

Zeggwagh NA, Moufid A, Michel JB, Eddouks M. Hypotensive effect of *Chamaemelum nobile* aqueous extract in spontaneously hypertensive rats. *Clin Exp Hypertens* 2009, 31:440-450.

Zwaving JH. Echte en roomse kamille, hun verschil en overeenkomst in samestelling en toepassing. *Pharm Weekbl* 1982, 117:157-165.

Zyczyńska-Baloniak I, Matuszelewska H, Dudzińska J. Neutral constituents of the flowers of *Anthemis nobilis* L. *Acta Pol Pharm* 1971, 28:625-634.