



EUROPEAN MEDICINES AGENCY
SCIENCE MEDICINES HEALTH

25 November 2010
EMA/HMPC/369803/2009
Committee on Herbal Medicinal Products (HMPC)

List of references supporting the assessment of *Chelidonium majus* L., herba

Draft

The Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Amoros M, Fauconnier B, Girre L. Propriétés antivirales de quelques extraits de plantes indigènes. *Annals Pharm Françaises* 1977; 35: 371-376

Ansari K, Dhawan A, Subhash K, Khanna, Das M. Protective effect of bioantioxidants on argemone oil/sanguinarine alkaloid induced genotoxicity in mice. *Cancer Letters* 2006; 244:109-118

Barnes J, Anderson L, Phillipson D. *Herbal Medicines: A Guide for Healthcare Professionals*. Pharmaceutical Press, London 2007, 136-145

Basini G, Santini S, Bussolati S, Grasselli F. The Plant Alkaloid Sanguinarine is a Potential Inhibitor of Follicular Angiogenesis. *Journal of Reproduction and Development* 2007; 53(3):573-579

Benninger J, Schneider HT, Schuppan D, Kirchner T, Hahn EG. Acute hepatitis induced by greater celandine (*Chelidonium majus*). *Gastroenterology* 1999; 117: 1234-1237

Bichler B. Fallbericht aus der Praxis: Hepatitis unklarer Genese. *Phytotherapie* 2009; 6: 19-20

Boegge SC, Kesper S, Verspohl EJ, Nahrstedt A. Reduction of ACh-induced contraction of rat isolated ileum by coptisine, (+)-caffeoylmalic acid, *Chelidonium majus*, and *Corydalis lutea* extracts. *Planta Medica* 1996; 62(2):173-4

Boulware RT, Southard GL, Yankell SL. Sanguinaria extract, a new agent for the control of volatile sulphur compounds in the oral cavity. *Journal of the Society of Cosmetic Chemists* 1985; 36(4):297-302

Bruneton J. *Pharmacognosie en phytochimie: plantes médicinales*, 3e ed TEC et DOC. Lavoisier, Paris 1999, 917

Caolo MA, Stermitz FR. Benzophenanthridinium salt equilibria. *Heterocycles* 1979; 12(1):11-5



- Chen CM, Chang HC. Determination of berberine in plasma, urine and bile by high-performance liquid chromatography. *Journal of Chromatography B* 1995; 665:117-123
- Chung HS, An HJ, Jeong HJ, Won JH, Hong SH, Kim HM. Water extract isolated from *Chelidonium majus* enhances nitric oxide and tumour necrosis factor- α production via nuclear factor- κ B activation in mouse peritoneal macrophages. *Journal of Pharmacy and Pharmacology* 2004; 56(1):129-34
- Colombo ML, Bosisio E. Pharmacological activities of *Chelidonium majus* L (Papaveraceae). *Pharmacological Research* 1996; 33(2):127-34
- Crijns APG, De SMet PAGM, Van den Heuvel M, Schot BW, Haagsma EB. Acute hepatitis na gebruik van een plantaardig preparaat met stinkende gouwe (*Chelidonium majus*). *Ned Tijdschr Geneesk* 2002; 146(3):124-127
- Delfosse M. Drogues végétales et plantes médicinales. *APB* 1998, 131-132
- Dvorak Z, Sovadinova I, Blaha L, Giesy J, Ulrichova J. Quaternary benzo [c]phenanthridine alkaloids sanguinarine and chelerythrine do not affect transcriptional activity of aryl hydrocarbon receptor: Analyses in rat hepatoma cell line H4IIE.luc. *Food and Chemical Toxicology* 2006; 44:1466-1473
- ESCOP Monographs 2nd ed. European Scientific Cooperative on Phytotherapy, editor. Thieme, Stuttgart 2003, 74-78
- Etxenagusia MA, Anda M, Gonzalez-Mahave I, Fernandez E, Fernandez de Corres L. Contact dermatitis from *Chelidonium majus* (greater celandine). *Contact Dermatitis* 2000; 43:47
- European Pharmacopoeia 6th ed. Council of Europe. 2008: 1861
- Faddejewa MD, Belyaeva TN, Rosanov YM, Sedova VM, Sokolovskaya EL. Studies on the complex-formation with DNA and the effect on DNA hydrolysis, RNA-synthesis and cellular membrane ATPase systems of some antitumoragents including alkaloids. *Studia Biophysica* 1984; 104(1-3):267-9
- Fik E, et al. New plant glycoprotein against methicillin resistant *Staphylococci* and *Enterococci*. *Acta microbiologica polonica* 1997; 46:325-327
- Frankos VH, Brusick DJ, Johnson EM. et al. Safety of sanguinaria extract as used in commercial toothpaste and oral rinse products. *J Can Dent Associat* 1990; 56(7):41-47
- Granger I, Serradeillegat C, Augereau JM, Gleye J. Benzophenanthridine alkaloids isolated from *eschscholtzia californica* cell suspension cultures interact with Vasopressin (V1) receptors. *Planta Medica* 1992; 58(1):35-8
- Häberlein H, et al. *Chelidonium majus* L.: components with in vitro affinity for the GABA_A receptor. Positive cooperation of alkaloids. *Planta Med* 1996; 62:227-231
- Hänsel R, Keller K, Rimpler H, Schneider G. Hagers Handbuch der Pharmazeutischen Praxis. Drogen E-O, Springer Verlag, Berlin 1992
- Hardeman E, Van Overbeke L, Ilegems S, Ferrante M. Acute hepatitis induced by greater celandine (*Chelidonium majus*). *Acta Gastro-Enterologica belgica* 2008; 71(2)281-282
- Hiller KO, Ghorbani M, Schilcher H. Antispasmodic and relaxant activity of chelidonine, protopine, coptisine, and *Chelidonium majus* extracts on isolated guinea-pig ileum. *Planta Medica* 1998; 64(8):758-60
- Hladon B, et al. Cytotoxic activity of some *Chelidonium majus* alkaloids on human and animal tumor cell cultures in vitro. *Ann Pharm (Poznan)* 1978; 13:61-8

- Li HL, Zhang WD, Zhang C, Liu RH, Wang XW, Wang XL, Zhu JB, Chen CL. Bioavailability and pharmacokinetics of four active alkaloids of traditional Chinese medicine Yanhuanglian in rats following intravenous and oral administration. *Journal of Pharmaceutical and Biomedical Analysis* 2006; 41:1342-1346
- Ishii H, Ichikawa Y, Kawanabe E, Ishikawa M, Ishikawa T, Kuretani K, Inomata M and Hoshi A. Studies on the chemical constituents of rutaceous plants. LX. Development of a versatile method for Syntheses of the antitumor benzophenanthridine alkaloids. Efficient syntheses and antitumour activities of Nitidine and related nonphenolic benzophenanthridine alkaloids. *Chem Pharm Bull* 1985; 33:4139-51
- Kaminsky V, Lin K, Filyak Y, Stoika R. Differential effect of sanguinarine, chelerythrine and chelidonine on DNA damage and cell viability in primary mouse spleen cells and mouse leukemic cells. *Cell Biology International* 2007; 32:271-277
- Kardos J, Blasko G, Simonyi M. Enhancement of gamma-aminobutyric acid receptor binding by protopine-type alkaloids. *Arzneimittel-Forschung/Drug Research* 1986; 36-1(6):939-40
- Kéry A, et al. Antiviral alkaloid in *Chelidonium majus* L. *Acta Pharm Hung* 1987; 57:19-25
- Kim Y, Shin M, Chung J, Kim E, Koo G, Lee C, Kim C. Modulation of *Chelidonium herba* on GABA activated chloride current in rat PAG neurons. *Am J Chin Med* 2001; 29:265-279
- Kim DJ, Ahn B, Han BS, Tsuda H. Potential preventive effects of *Chelidonium majus* L (Papaveraceae) herb extract on glandular stomach tumor development in rats treated with N-methyl-N'-nitro-N nitrosoguanidine (MNNG) and hypertonic sodium chloride. *Cancer Letters* 1997; 112(2):203-8
- Kim HK, Farnsworth NR, Blomster RN, Fong HHS. Biological and phytochemical evaluation of plants. V. Isolation of 2 cytotoxic alkaloids from *Chelidonium majus*. *Journal of Pharmaceutical Sciences* 1969; 58(3):372
- Kokoska L, Polesny Z, Rada V, Nepovim A, Vanek T. Screening of some Siberian medicinal plants for antimicrobial activity. *Journal of Ethnopharmacology* 2002; 82(1):51-53
- Kosina P, Walterova D, Ulrichova J, Lichnovsky V, Stiborova M, Rydlova H, Vicar J, Krecman V, Brabec M, Simanek V. Sanguinarine and chelerythrine: assessment of safety on pigs in ninety days feeding experiment. *Food and Chemical Toxicology* 2004; 42:85-91
- Krey AK, Hahn FE. Berberine – complex with DNA. *Science* 1969; 166(3906):755
- Leclerc H. *Précis de Phytothérapie: thérapeutique par les plantes françaises*, Masson Editeurs, Paris 1954, 268-270
- Lenfeld J, Kroutil M, Marsalek E, Slavik J, Preininger V, Simanek V. Isolation, chemistry and biology of alkaloids from plants of the papaveraceae: anti-inflammatory activity of quaternary benzophenanthridine alkaloids from *Chelidonium majus*. *Planta Medica* 1981; 43(2):161-5
- Lozyuk LV. Antiviral properties of some compounds of plant origin. *Mikrobiol Zh (Kiev)* 1977; 39:343-348
- Ma WG, Fukushi Y, Tahara S, Osawa T. Fungitoxic alkaloids from Hokkaido Papaveraceae. *Fitoterapia* 2000; 71(5):527-34
- Mahajan VM, et al. Antimycotic activity of Berberine sulphate: an alkaloid from an Indian medicinal herb. *Sabouraudia* 1982; 20:79-81
- Maiti M, Nandi R, Chaudhuri K. Sanguinarine – a monofunctional intercalating alkaloid *Febs Letters* 1982; 142(2):280-4

- Malikova J, Zdarilova A, Hlobilkova A. Effects of sanguinarine and chelerythrine on the cell cycle and apoptosis. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2006; 150(1):5-12
- Matos OC, Baeta J, Silva MJ, Ricardo CPP. Sensitivity of *Fusarium* strains to *Chelidonium majus* L. extracts. *Journal of Ethnopharmacology* 1999; 66(2):151-8
- Mitscher LA, Park YH, Clark D, Clark GW 3rd, Hammesfahr PD, Wu WN, Beal JL. Antimicrobial agents from higher plants. An investigation of *Hunnemannia fumariaefolia* pseudoalcoholates of sanguinarine and chelerythrine. *Lloydia (J Nat Prod)* 1978; 41:145-150
- Moffat AC, Osselton MD, Widdop B, Galichet LY. Clarke's Analysis of drugs and poisons. The Pharmaceutical Press, London 2004, 692-693
- Moro PA, Casseti F, Giugliano G, Falce M, Mazzanti G, Mennitti-Ippolito F, Raschetti R, Santuccio C. Hepatitis from Greater celandine (*Chelidonium majus* L.): review of literature and report of a new case. *J Ethnopharmacology* 2009; 124:328-332
- Pitea M and Margineanu C. Correlations between chemical structure and antibacterial activity of berberine. *Clujul med* 1972; 45:465-9
- Psotova J, Klejdus B, Vecera R, Kosina P, Kuban V, Vicar J, Simanek V, Ulrichova J. A liquid chromatographic-mass spectrometric evidence of dihydrosanguinarine as a first metabolite of sanguinarine transformation in rat. *Journal of Chromatography B* 2006; 830:165–172
- Ritter R, et al. Clinical trial on standardised celandine extract in patients with functional epigastric complaints: results of a placebo-controlled double-blind trial. *Complement Ther Med* 1993; 1:189-193
- Rogelj B, Popovic T, Ritonja A, Strukelj B, Brzin J. Chelidocystatin, a novel phytocystatin from *Chelidonium majus*. *Phytochemistry* 1998; 49(6):1645-9
- Runge D, Rudolf H, Ullrich A, Ludwig K. Hepatotoxic assessment of *Chelidonium majus* L. and its alkaloids in primary hepatocyte cultures of different species. *Zeitschrift für Phytotherapie* 2009; 30 (Suppl. 1)
- Saglam H, Arar G. Cytotoxic activity and quality control determinations on *Chelidonium majus*. *Fitoterapia* 2003; 74:127-129
- Sethi ML. Enzyme inhibition 6. inhibition of reverse transcriptase activity by protoberberine alkaloids and structure activity relations. *Journal of Pharmaceutical Sciences* 1983; 72(5):538-41
- Sethi ML. Screening of benzophenanthridine alkaloids for their inhibition of reverse-transcriptase activity and preliminary report on the structure activity relationships. *Canadian Journal of Pharmaceutical Sciences* 1981; 16(1):29-34
- Sethi ML. Enzyme inhibition 7. comparison of inhibition of reverse transcriptase and antileukemic activities exhibited by protoberberine and benzophenanthridine alkaloids and structure activity relationships. *Phytochemistry* 1985; 24(3):447-54
- Shin M-C. Modulation of *Chelidonium herba* on glycine-activated and glutamate-activated chloride current in rat PAG neurons. *Clin chim acta* 2002; 337:93-101
- Smekal E, Kubova N, Kleinwachter V. DNA – chelerythrine interaction and binding isotherms. *Studia Biophysica* 1985; 105(2):77-84
- Smekal E, Kubova N, Kleinwachter V. Interaction of benzophenanthridine alkaloid sanguinarine with DNA. *Studia Biophysica* 1984; 101:125-32

- Smekal E, Kubova N. Alkaloids of protoberberinen and benzophenanthridine groups as intercalating drugs with DNA. *Studia Biophysica* 1984; 101:121-3
- Sokoloff B. Oncostatic and oncolytic factors present in certain plants. *Oncology* 1968; 22(1):49-60
- Song JY, Yang HO, Pyo SN, Jung IS, Yi SY, Yun YS. Immunomodulatory activity of protein-bound polysaccharide extracted from *Chelidonium majus*. *Archives of Pharmacal Research* 2002; 25(2):158-64
- Song JY, Yang HO, Shim JY, et al. Radiation protective effect of an extract from *Chelidonium majus*. *International Journal of Hematology* 2003; 78(3):226-32
- Southard G, et al. Benzophenanthridine alkaloids containing solutions for treating periodontal diseases. US Patent 4,683,133, 1987
- Stiborová M, Simánek V, Frei E, Hobza P, Ulrichová J. DNA adduct formation from quaternary benzo[c]phenanthridine alkaloids sanguinarine and chelerythrine as revealed by the 32P-postlabeling technique. *Chem Biol Interact* 2002; 140(3):231-42
- Stiborova M, Vostalova J, Zdarilova A, Ulrichova J, Hudecek J, Tschirner K, Simanel V. *Macleaya cordata* extract and Sangrovit® genotoxicity assessment in vivo. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2008; 152(1):35-39
- Stickel F, Seitz HK, Hahn EG, Schuppan D. Hepatotoxizität von Arzneimittel pflanzlichen Ursprungs. *Z Gastroenterol* 2001; 39:225-237
- Tan GT, Pezzuto JM, Kinghorn AD, Hughes SH. Evaluation of natural products as inhibitors of human immunodeficiency virus type 1 (HIV-1) reverse transcriptase. *J Nat Prod* 1991; 54:143-154
- Tomè F, Colombo M. Distribution of alkaloids in *Chelidonium majus* and factors affecting their accumulation. *Phytochemistry* 1995; 40(1):37-39
- Tsurui Yakuhin Kogyo KK. Anticaries agents. Jpn. Kokai tokkyo JP Patent 8,285,319, 1982
- Vahensieck U, et al. The effect of *Chelidonium majus* herb extract on choleresis in the isolated perfused rat liver. *Plant med* 1995; 61:267-270
- Van Hellemont J. *Fytotherapeutisch compendium*, Belgian Pharmaceutical Society, Brussels, 144-146
- Vavreckova C, Gawlik I, Muller K. Benzophenanthridine alkaloids of *Chelidonium majus* .1. Inhibition of 5- and 12-lipoxygenase by a non-redox mechanism. *Planta Medica* 1996a; 62(5):397-401
- Vavreckova C, Gawlik I, Muller K. Benzophenanthridine alkaloids of *Chelidonium majus* .2. Potent inhibitory action against the growth of human keratinocytes. *Planta Medica* 1996b; 62(6):491-4
- Wegener T, Wagner H. The active components and the pharmacological multi-target principle of STW 5 (Iberogast). *Phytomedicine* 2006; 13:20-35
- Weiss RF, Fintelmann V. *Lehrbuch der phytotherapie*. Hippocrates Verlag, Stuttgart 1999, 131-132
- Zdarilova A, Vrzal R, Rypka M, Ulrichova J, Dvorak Z. Investigation of sanguinarine and chelerythrine effects on CYP1A1 expression and activity in human hepatoma cells. *Food and Chemical Toxicology* 2006; 44:242-249
- Zhu B, Ahrens FA. Effect of Berberine on intestinal secretion mediated by *Escherichia-coli* heat stable enterotoxin in jejunum of pigs. *American Journal of Veterinary Research* 1982; 43(9):1594-8

Zuo GY, Meng FY, Hao XY, Zhang YL, Wang GCH, Xu GL. Antibacterial Alkaloids from *Chelidonium majus* Linn (Papaveracea) against clinical isolates of methicillin-resistant *Staphylococcus aureus*. J Pharm Pharmaceut Sci 2008; 11(4):90-94

References not cited

Blumenthal M, Busse W, Goldberg A, Gruenwald J, Hall T, Riggins C, Rister R. The Complete German Commission E Monographs. American Botanical Council, Austin Texas 1998, 105-106

Capasso F, Gaginella T, Lieb E. Phytotherapie: A quick reference to herbal medicine, Springer Verlag, Berlin Heidelberg New York 2003, 307

Cohen HG, Seifen EE, Straub KD, Tiefenback C, Stermitz FR. Structural specificity of NaK ATPase inhibition by sanguinarine, an isoquinoline benzophenanthridine alkaloid. Biochemical Pharmacology 1978; 27(21):2555-8

Gansauge F, Ramadani M, Pressmar J, Gansauge S, Muehling B, Stecker K, Cammerer G, Leder G, Beger H NSC-631570 (Ukrain) in the palliative treatment of pancreatic cancer: results of a phase II trial. Langenbeck's Arch Surg 2002; 386:570-574

Horvath J, et al. Antiviral effect of *Chelidonium* extracts. Proc Int Congr Chemother 13th 1983; 9:124/106-124/110

Kelber O, Wittwer A, Lapke C, Kroll U, Weiser D, Okpanyi SN, Heilmann J. Ex vivo/in vitro absorption of STW 5 (Iberogast) and its extract components. Phytomedicine 2006; 13:107-113

Mackraj I, Govender T, Gathiram P. Sanguinarine. Cardiovascular Therapeutics 2008; 26:75-83

Madisch A, et al. Ein phytotherapeutikum und seine modifizierte rezeptur bei functioneller dyspepsie. Ergebnisse einer doppelblinden plazebokontrollierten vergleichstudie. Z Gastroenterol 2001; 39:511-517

Niu C, et al. Review of the studies on *Chelidonium majus* L. Zhongguo Yaoxue Zazhi 1994; 29:138-140

Sárközi A, Then M, Szentmihályi K. Mineral element content of greater celandine (*Chelidonium majus* L.) Acta Alimentaria 2005; 34:113-120

Stickel F, Schuppan D. Herbal medicine in the treatment of liver diseases. Digestive and Liver Disease 2007; 39:293-304