



EUROPEAN MEDICINES AGENCY
SCIENCE MEDICINES HEALTH

22 November 2016
EMA/HMPC/424585/2016
Committee on Herbal Medicinal Products (HMPC)

List of references supporting the assessment of *Echinacea purpurea* (L.) Moench, radix

Draft

The European Medicines Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Abouelella AM, Shahein YE, Tawfik SS, Zahran AM (2007). Phytotherapeutic effects of *Echinacea purpurea* in gamma-irradiated mice. *J Vet Sci* 8: 341-351

Agnew LL, Guffogg SP, Matthias A, Lehmann RP, Bone KM, Watson K (2005). *Echinacea* intake induces an immune response through altered expression of leucocyte hsp70, increased white cell counts and improved erythrocyte antioxidant defences. *J Clin Pharm Ther* 30: 363-369

Ardjomand-Woelkart K, Kollroser M, Magnes C, Sinner F, Frye RF, Derendorf H *et al.* (2011). Absolute/relative bioavailability and metabolism of dodeca-2E,4E,8Z,10E/Z-tetraenoic acid isobutylamides (tetraenes) after intravenous and oral single doses to rats. *Med* 77(16): 1794-1799

Barbour EK, Abou Assi CA, Shaib H, Hamadeh S, Murtada M, Mahmoud G *et al.* (2015). Evaluation of a *Salmonella* Enteritidis vaccine and related ELISA for respective induction and assessment of acquired immunity to the vaccine and/or *Echinacea purpurea* in Awassi Ewes Original Research Article. *Vaccine* 33(19): 2228-2231

Barcz E, Sommer E, Nartowska J, Balan B, Chorostowska-Wynimko J, Skopińska-Różewska E (2007). Influence of *Echinacea purpurea* intake during pregnancy on fetal growth and tissue angiogenic activity. *Folia Histochem Cytobiol* 45(1): 35-39

Barnes J, Anderson LA, Phillipson JD (2007). *Echinacea*. In: *Herbal Medicines*, 3rd ed., Pharmaceutical Press, London, UK: 217-236

Barnes J, Anderson, LA, Gibbons S, Phillipson JD (2005). *Echinacea* species (*Echinacea angustifolia* (DC.) Hell., *Echinacea pallida* (Nutt.) Nutt., *Echinacea purpurea* (L.) Moench): A review of their chemistry, pharmacology and clinical properties. *J Pharm Pharmacol* 57: 929-954

Barrett B (2003). Medicinal properties of *Echinacea*: a critical review. *Phytomedicine* 10: 66-86



- Barrett B, Brown R, Rakel D, Mundt M, Bone K, Barlow S *et al.* (2010). Echinacea for treating the common cold: a randomized trial. *Ann Intern Med* 153(12): 769-777
- Barrett B, Brown RL, Locken K, Maberry R, Bobula JA, D'Alessio D (2002). Treatment of the common cold with unrefined Echinacea. A randomized, double-blind, placebo-controlled trial. *Ann Intern Med* 137: 939-946
- Barrett B, Vohmann M, Calabrese C (1999). Echinacea for upper respiratory tract infection. *J Farm Pract* 48: 628-635
- Barth A, Hovhannisyanyan A, Jamalyan K, Narimanyan M (2015). Antitussive effect of a fixed combination of *Justicia adhatoda*, *Echinacea purpurea* and *Eleutherococcus senticosus* extracts in patients with acute upper respiratory tract infection: A comparative, randomized, double-blind, placebo-controlled study. *Phytomedicine* 22(13): 1195-1200
- Bauer R, Jurcic K, Puhlmann J, Wagner H (1988). Immunologische In-vivo- und In-vitro-Untersuchungen mit Echinacea-Extrakten. *Arzneim Forsch* 38: 276-281
- Bauer R, Liersch R (1993). Echinacea. In: Hagers Buch der Pharmazeutischen Praxis, 5th ed., Vol. 5: Drogen E-O, Springer-Verlag, Berlin, Germany: 1-34.
- Bauer R, Remiger P (1989). TLC and HPLC Analysis of Alkamides in Echinacea Drugs. *Planta Med* 55: 367-371
- Bauer R, Remiger P, Jurcic K, Wagner H (1989). Beeinflussung der Phagozytose-Aktivität durch Echinacea-Extrakte. *Z Phytother* 10: 43-48
- Benson JM, Pokorny AJ, Rhule A, Wenner CA, Kandhi V, Cech NB *et al.* (2010). Echinacea purpurea extracts modulate murine dendritic cell fate and function. *Food Chem Toxicol* 48(5): 1170-1177
- Beuscher N, Bodinet I, Willigmann D, Egert (1995). Immunomodulierende Eigenschaften von Wurzelextrakten verschiedenen Echinacea-Arten. *Z Phytother* 16: 157-166
- Beuscher N, Schet KH, Bodinet C, Egert D (1990). Modulation of host resistance by polymer substances from *Baptisia tinctoria* and *Echinacea purpurea*. In: Immunotherapeutic prospects of infectious diseases (Masihi KN, Lange W, eds.), Springer-Verlag, Berlin-Heidelberg, Germany: 59-63
- Binns SE, Hudson J, Merali S, Arnason JT (2002). Antiviral activity of characterized extracts from *Echinacea* spp. (Heliantheae: Asteraceae) against herpes simplex virus (HSV-1). *Planta Med* 68: 780-783
- Binns SE, Purgina B, Bergeron C, Smith ML, Ball L, Baum BR, Arnason JT (2000). Light-mediated antifungal activity of Echinacea extracts. *Planta Med* 66: 241-244
- Blumenthal M, Goldberg A, Brinckmann J (2000). Echinacea purpurea root. In: Herbal Medicine - Expanded Commission E monographs, American Botanical Council, Austin, TX, USA: 98-102
- Bodinet C, Beuscher N (1991). Antiviral and Immunological Activity of Glycoproteins from *Echinacea purpurea* Radix. *Planta Med* 57(2): 33-34
- Bodinet C, Freudenstein J (1999). Effect of an orally applied aqueous-ethanolic extract of a mixture of *Thuja occidentalis* herba, *Baptisia tinctoriae* radix, *Echinacea purpureae* radix and *Echinacea pallidae* radix on antibody response against sheep red blood cells in mice. *Planta Med* 65: 695-699
- Bodinet K (1999). Immunopharmakologische Untersuchungen an einem pflanzlichen Immunomodulator [Dissertation]. Ernst-Moritz-Arndt-Universität, Greifswald, Germany
- Bossaer JB, Odle BL (2012). Probable etoposide interaction with Echinacea. *J Diet Suppl* 9(2): 90-95

- Bradley P (2006). Echinacea. In: British Herbal Compendium, Vol. 2., British Herbal Medicine Association, Bournemouth, UK: 129-141
- Bräunig B, Dorn M, Limburg E, Knick E, Bausendorf (1992). Echinaceae purpureae radix: zur stärkung der körpereigenen Abwehr bei grippalen Infekten. Z Phytother 13: 7-13
- Brinkeborn RM, Shah DV, Degenring FH (1999). Echinaforce® and other Echinacea fresh plant preparations in the treatment of the common cold. A randomized, placebo-controlled, double-blind clinical trial. Phytomedicine 6: 1-6.
- Brousseau M, Miller SC (2005). Enhancement of natural killer cells and increased survival of aging mice fed daily Echinacea root extract from youth. Biogerontology 6: 157-163.
- Budzinski JW, Foster BC, Vandenhoeck S, Arnason JT (2000). An in vitro evaluation of human cytochrome P450 3A4 inhibition by selected commercial herbal extracts and tinctures. Phytomedicine 7: 273-282
- Cech NB, Kandhi V, Davis JM, Hamilton A, Eads D, Laster SM (2010). Echinacea and its alkylamides: effects on the influenza A-induced secretion of cytokines, chemokines, and PGE₂ from RAW 264.7 macrophage-like cells. Int Immunopharmacol 10(10): 1268-1278
- Chen Y, Fu T, Tao T, Yang J, Chang Y, Wang M, Kim L, Qu L, Cassady J, Scalzo R, Wang X (2005). Macrophage Activating Effects of New Alkamides from the Roots of Echinacea Species. J Nat Prod 68: 773-776
- Chicca A, Adinolfi B, Martinotti E, Fogli S, Breschi MC, Pellati F, Benvenuti S, Nieri P (2007). Cytotoxic effects of Echinacea root hexanic extracts on human cancer cell lines. J Ethnopharmacol 110: 148-153
- Chicca A, Pellati F, Adinolfi B, Matthias A, Massarelli I, Benvenuti S, Martinotti E, Bianucci AM, Bone K, Lehmann R, Nieri P (2008). Cytotoxic activity of polyacetylenes and polyenes isolated from roots of Echinacea pallida. Br J Pharmacol 153: 879-885
- Clifford LJ, Nair MG, Rana J, Dewitt DL (2002). Bioactivity of alkamides isolated from Echinacea purpurea (L.) Moench. Phytomedicine 9: 249-253
- Currier NL, Miller SC (2000). Natural killer cells from aging mice treated with extracts from Echinacea purpurea are quantitatively and functionally rejuvenated. Exp Gerontol 35: 627-639
- Currier NL, Miller SC (2001). Echinacea purpurea and melatonin augment natural-killer cells in leukemic mice and prolong life span. J Altern Complement Med 7: 241-251
- Currier NL, Miller SC (2002). The effect of immunization with killed tumor cells, with/without feeding of Echinacea purpurea in an erythroleukemic mouse model. J Altern Complement Med 8: 49-58
- Dalby-Brown L, Barsett H, Landbo AK, Meyer AS, Mølgaard P (2005). Synergistic antioxidative effects of alkamides, caffeic acid derivatives, and polysaccharide fractions from Echinacea purpurea on in vitro oxidation of human low-density lipoproteins. J Agric Food Chem 53: 9413-9423
- Di Carlo G, Nuzzo I, Capasso R, Sanges MR, Galdiero E, Capasso F, Carratelli CR (2003). Modulation of apoptosis in mice treated with Echinacea and St. John's wort. Pharmacol Res 48: 273-277
- El-Ashmawy NE, El-Zamarany EA, Salem ML, El-Bahrawy HA, Al-Ashmawy GM (2015). In vitro and in vivo studies of the immunomodulatory effect of Echinacea purpurea on dendritic cells. J Gen Eng Biotech 13(2): 185-192
- European Pharmacopoeia (2008). Purple Coneflower Root. In: European Pharmacopoeia, 6.0., Vol. 2, Council of Europe. 01/2008: 1824

European Scientific Cooperative On Phytotherapy (ESCOP) (2003). *Echinaceae purpureae radix*. In: ESCOP Monographs, 2nd ed., Thieme, New York, NY, USA: 137-141

European Scientific Cooperative On Phytotherapy (ESCOP) (2009). *Echinaceae purpureae radix*. In: ESCOP Monographs, 2nd ed., Suppl. 2009, Thieme, New York, NY, USA: 102-109

Fast DJ, Balles JA, Scholten JD, Mulder T, Rana J (2015). *Echinacea purpurea* root extract inhibits TNF release in response to Pam3Csk4 in a phosphatidylinositol-3-kinase dependent manner. *Cell Immunol* 297(2): 94-99

Freeman C, Spelman K (2008). A critical evaluation of drug interactions with *Echinacea* spp. *Mol Nutr Food Res* 52: 789-798

Gallo M, Sarkar M, Au W, Pietrzak K, Comas B, Smith M, Jaeger TV, Einarson A, Koren G (2000). Pregnancy outcome following gestational exposure to *Echinacea*: a prospective controlled study. *Arch Intern Med* 160(20): 3141-3143

Gertsch J, Schoop R, Kuenzle U, Suter A (2004). *Echinacea* alkylamides modulate TNF-alpha gene expression via cannabinoid receptor CB2 and multiple signal transduction pathways. *FEBS Lett* 577: 563-569

Goel V, Chang C, Slama J, Barton R, Bauer R, Gahler R, Basu T (2002a). *Echinacea* stimulates macrophage function in the lung and spleen of normal rats. *J Nutr Biochem* 13: 487-492

Goel V, Chang C, Slama JV, Barton R, Bauer R, Gahler R, Basu TK (2002b). Alkylamides of *Echinacea purpurea* stimulate alveolar macrophage function in normal rats. *Int Immunopharmacol* 2: 381-387

Goel V, Lovlin R, Barton R, Lyon MR, Bauer R, Lee TD, Basu TK (2004). Efficacy of a standardized *Echinacea* preparation (Echinilin) for the treatment of the common cold: a randomized, double-blind, placebo-controlled trial. *J Clin Pharm Ther* 29: 75-83

Goel V, Lovlin R, Chang C, Slama JV, Barton R, Gahler R, Bauer R, Goonewardene L, Basu TK (2005). A proprietary extract from the *Echinacea* plant (*Echinacea purpurea*) enhances systemic immune response during a common cold. *Phytother Res* 19: 689-694

Gorski JC, Huang SM, Pinto A, Hamman MA, Hilligoss JK, Zaheer NA, Desai M, Miller M, Hall SD (2004). The effect of *Echinacea* (*Echinacea purpurea* root) on cytochrome P450 activity in vivo. *Clin Pharmacol Ther* 75: 89-100

Gryzlak BM, Wallace RB, Zimmerman MB, Nisly NL (2007). National surveillance of herbal dietary supplement exposures: the poison control center experience. *Pharmacoepidemiol Drug Saf* 16: 947-957

Haller CA, Anderson IB, Kim SY, Blanc PD (2002). An evaluation of selected herbal reference texts and comparison to published reports of adverse herbal events. *Adverse Drug React Toxicol Rev* 21: 143-150

Hellum BH, Nilsen OG (2007). The in vitro inhibitory potential of trade herbal products on human CYP2D6-mediated metabolism and the influence of ethanol. *Basic Clin Pharmacol Toxicol* 101: 350-358

Henneicke-von Zepelin H, Hentschel C, Schnitker J, Kohnen R, Köhler G, Wüstenberg P (1999). Efficacy and safety of a fixed combination phytomedicine in the treatment of the common cold (acute viral respiratory tract infection): results of a randomised, double blind, placebo controlled, multicentre study. *Curr Med Res Opin* 15: 214-227

- Hohmann J, Rédei D, Forgo P, Szabó P, Freund TF, Haller J. *et al.* (2011). Alkamides and a neolignan from *Echinacea purpurea* roots and the interaction of alkamides with G-protein-coupled cannabinoid receptors. *Phytochemistry* 72(14-15): 1848-1853
- Hou CC, Chen CH, Yang NS, Chen YP, Lo CP, Wang SY *et al.* (2010). Comparative metabolomics approach coupled with cell- and gene-based assays for species classification and anti-inflammatory bioactivity validation of *Echinacea* plants. *J Nutr Biochem* 21(11): 1045-1059
- Hou CC, Huang CC, Shyur LF (2011). *Echinacea* alkamides prevent lipopolysaccharide/D-galactosamine-induced acute hepatic injury through JNK pathway-mediated HO-1 expression. *J Agric Food Chem* 59(22): 11966-11974
- Hu C, Kitts DD (2000). Studies on the antioxidant activity of *Echinacea* root extract. *J Agric Food Chem* 48: 1466-72
- Huntley AL, Thompson Coon J, Ernst E (2005). The safety of herbal medicinal products derived from *Echinacea* species: a systematic review. *Drug Saf* 28: 387-400
- Jager H, Meinel L, Dietz B, Lapke C, Bauer R, Merkle HP, Heilmann J (2002). Transport of Alkamides from *Echinacea* Species through Caco-2 Monolayers. *Planta Med* 68: 469-471
- Jawad M, Schoop R, Suter A, Klein P, Eccles R (2012). Safety and Efficacy Profile of *Echinacea purpurea* to Prevent Common Cold Episodes: A Randomized, Double-Blind, Placebo-Controlled Trial. *Evid Based Compl Altern Med*: 841315
- Jurcic K, Melchart D, Holzmann M, Martin P, Bauer R, Doenecke A, Wagner H (1989). Zwei Probandenstudien zur Stimulierung der Granulozytenphagozytose durch *Echinacea*-Extrakt-haltige Präparate. *Z Phytother* 10: 67-70
- Karsch-Völk M, Barrett B, Kiefer D, Bauer R, Ardjomand-Woelkart K, Linde K (2014). *Echinacea* for preventing and treating the common cold. *Cochrane Database of Systematic Reviews* 2: 1-91
- Kotowska D, El-Houri RB, Borkowski K, Petersen RK, Fretté XC, Wolber G *et al.* (2014). Isomeric C12-alkamides from the roots of *Echinacea purpurea* improve basal and insulin-dependent glucose uptake in 3T3-L1 adipocytes. *Planta Med* 80(18): 1712-1720
- LaLone CA, Hammer KD, Wu L, Bae J, Leyva N, Liu Y, Solco AK, Kraus GA, Murphy PA, Wurtele ES, Kim OK, Seo KI, Widrlechner MP, Birt DF (2007). *Echinacea* species and alkamides inhibit prostaglandin E(2) production in RAW264.7 mouse macrophage cells. *J Agric Food Chem* 55: 7314-7322
- Linde K, Barret B, Bauer R, Melchart D, Woelkart K (2006). *Echinacea* for preventing and treating the common cold. *Cochrane Database of Systematic Reviews*, Issue 1, In: *The Cochrane Library* 2008, Issue 3, John Wiley & Sons, Ltd., Chichester, UK: 1-88
- Lindenmuth GF, Lindenmuth EB (2000). The efficacy of *Echinacea* compound herbal tea preparation on the severity and duration of upper respiratory and flu symptoms: a randomized, double-blind placebo-controlled study. *J Alt Complem Med* 6: 327-334
- Maffei Facino R, Carini M, Aldini G, Saibene L, Pietta P, Mauri P (1995). Echinacoside and caffeoyl conjugates protect collagen from free radical-induced degradation: a potential use of *Echinacea* extracts in the prevention of skin photodamage. *Planta Med* 61: 510-514
- Matthias A, Addison RS, Agnew LL, Bone KM, Watson K, Lehmann RP (2007). Comparison of *Echinacea* alkylamide pharmacokinetics between liquid and tablet preparations. *Phytomedicine* 14: 587-590
- Matthias A, Addison RS, Penman KG, Dickinson RG, Bone KM, Lehmann RP (2005b). *Echinacea* alkamide disposition and pharmacokinetics in humans after tablet ingestion. *Life Sci* 77: 2018-2029

- Matthias A, Banbury L, Bone KM, Leach DN, Lehmann RP (2008). Echinacea alkylamides modulate induced immune responses in T-cells. *Fitoterapia* 79: 53-58
- Matthias A, Blanchfield JT, Penman KG, Toth I, Lang CS, De Voss JJ, Lehmann RP (2004). Permeability studies of alkylamides and caffeic acid conjugates from Echinacea using a Caco-2 cell monolayer model. *J Clin Pharm Ther* 29: 7-13
- Matthias A, Gillam EM, Penman KG, Matovic NJ, Bone KM, De Voss JJ, Lehmann RP (2005a). Cytochrome P450 enzyme-mediated degradation of Echinacea alkylamides in human liver microsomes. *Chem Biol Interact* 155: 62-70
- Mazza G, Cottrell T (1999). Volatile components of roots, stems, leaves, and flowers of Echinacea species. *J Agric Food Chem* 47: 3081-3085
- McCann DA, Solco A, Liu Y, Macaluso F, Murphy PA, Kohut ML, Senchina DS (2007). Cytokine- and interferon-modulating properties of Echinacea spp. root tinctures stored at -20°C for 2 years. *J Interferon Cytokine Res* 27: 425-436
- Melchart D, Linde K, Fischer P, Kaesmayr J (2004). Echinacea for preventing and treating the common cold (Cochrane review). In: *The Cochrane Library, Issue 2*, John Wiley & Sons, Chichester, UK
- Melchart D, Linde K, Worku F, Bauer R, Wagner H (1994). Immunomodulation with Echinacea. A systematic review of controlled clinical trials. *Phytomed* 1: 245-254
- Melchart D, Linde K, Worku F, Sarkady L, Holzmann M, Jurcic K, Wagner H (1995). Results of five randomized studies on the immunomodulatory activity of preparations of Echinacea. *J Altern Complement Med* 1: 145-160
- Melchart D, Walther E, Linde K, Brandmaier R, Lersch C (1998). Echinacea root extracts for the prevention of upper respiratory tract infections: a double-blind, placebo-controlled randomized trial. *Arch Fam Med* 7: 541-545
- Merali S, Binns S, Paulin-Levasseur M, Ficker C, Smith M, Baum B, Brovelli E, Arnason JT (2003). Antifungal and Anti-inflammatory Activity of the Genus Echinacea. *Pharm Biol* 41: 412-420
- Moltó J, Valle M, Miranda C, Cedeño S, Negredo E, Barbanoj MJ *et al.* (2011). Herb-drug interaction between Echinacea purpurea and darunavir-ritonavir in HIV-infected patients. *Antimicrob Agents Chemother* 55(1): 326-330
- Moltó J, Valle M, Miranda C, Cedeño S, Negredo E, Clotet B (2012). Herb-drug interaction between Echinacea purpurea and etravirine in HIV-infected patients. *Antimicrob Agents Chemother* 56(10): 5328-5331
- Müller-Jakic B, Breu W, Pröbstle A, Redl K, Greger H, Bauer R (1994). In vitro inhibition of cyclooxygenase and 5-lipoxygenase by alkamides from Echinacea and Achillea species. *Planta Med* 60: 37-40
- Mullins RJ (1998). Echinacea-associated anaphylaxis. *Med J Austr* 168: 170-171
- Mullins RJ, Heddle R (2002). Adverse reactions associated with Echinacea: the Australian experience. *Ann Allergy Asthma Immunol* 88: 42-51
- PDR (2007). Echinacea. In: Gruenwald J, Brendler T, Jaenicke C (editors). *PDR for herbal medicines*, 4th Edition, Thomson Healthcare Inc., Montvale, NJ, USA: 266-276
- Pellati F, Benvenuti S, Magro L, Melegari M, Soragni F (2004). Analysis of phenolic compounds and radical scavenging activity of Echinacea spp. *J Pharm Biomed Anal* 35: 289-301

Perri D, Dugoua JJ, Mills E, Koren G (2006). Safety and efficacy of Echinacea (*Echinacea angustifolia*, *E. purpurea* and *E. pallida*) during pregnancy and lactation. *Can J Clin Pharmacol* 13(3): 262-267

Pugh ND, Jackson CR, Pasco DS (2013). Total bacterial load within *Echinacea purpurea*, determined using a new PCR-based quantification method, is correlated with LPS levels and in vitro macrophage activity. *Planta Med* 79(1): 9-14

Raduner S, Majewska A, Chen JZ, Xie XQ, Hamon J, Faller B, Altmann KH, Gertsch J (2006). Alkylamides from *Echinacea* are a new class of cannabinomimetics. Cannabinoid type 2 receptor-dependent and -independent immunomodulatory effects. *J Biol Chem* 281: 14192-14206

Raso GM, Pacilio M, Di Carlo G, Esposito E, Pinto L, Meli R (2002). *In-vivo* and *in-vitro* anti-inflammatory effect of *Echinacea purpurea* and *Hypericum perforatum*. *J Pharm Pharmacol* 54: 1379-1383

Rauš K, Pleschka S, Klein P, Schoop R, Fisher P (2015). Effect of an Echinacea-Based Hot Drink Versus Oseltamivir in Influenza Treatment: A Randomized, Double-Blind, Double-Dummy, Multicenter, Noninferiority Clinical Trial. *Curr Therap Res* 77: 66-72

Rininger JA, Kickner S, Chigurupati P, McLean A, Franck Z (2000). Immunopharmacological activity of Echinacea preparations following simulated digestion on murine macrophages and human peripheral blood mononuclear cells. *J Leukoc Biol* 68: 503-10

Rizzello CG, Coda R, Macías DS, Pinto D, Marzani B, Filannino P *et al.* (2013). Lactic acid fermentation as a tool to enhance the functional features of Echinacea spp. *Microb Cell Fact* 12: 44

Romiti N, Pellati F, Nieri P, Benvenuti S, Adinolfi B, Chieli E (2008). P-Glycoprotein inhibitory activity of lipophilic constituents of *Echinacea pallida* roots in a human proximal tubular cell line. *Planta Med* 74: 264-266

Schoop R, Klein P, Suter A, Johnston SL (2006). *Echinacea* in the prevention of induced rhinovirus colds: a meta-analysis. *Clin Ther* 28: 174-183

Senchina DS, Flagel LE, Wendel JF, Kohut ML (2008). Phenetic comparison of seven *Echinacea* species based on immunomodulatory characteristics. *Econ Bot* 60: 205-211

Senchina DS, McCann DA, Asp JM, Johnson JA, Cunnick JE, Kaiser MS, Kohut ML (2005). Changes in immunomodulatory properties of *Echinacea* spp. root infusions and tinctures stored at 4 degrees C for four days. *Clin Chim Acta* 355: 67-82

Senchina DS, Wu L, Flinn GN, Konopka del N, McCoy JA, Widrlechner MP, Wurtele ES, Kohut ML (2006). Year-and-a-half old, dried *Echinacea* roots retain cytokine-modulating capabilities in an *in vitro* human older adult model of influenza vaccination. *Planta Med* 72: 1207-1215

Shah SA, Sander S, White CM, Rinaldi M, Coleman CI (2007). Evaluation of *Echinacea* for the prevention and treatment of the common cold: a meta-analysis. *Lancet Infect Dis* 7: 473-480

Sharma M, Arnason JT, Burt A, Hudson JB (2006). *Echinacea* extracts modulate the pattern of chemokine and cytokine secretion in rhinovirus-infected and uninfected epithelial cells. *Phytother Res* 20: 147-152

Shin DM, Choi KM, Lee YS, Kim W, Shin KO, Oh S, Jung JC *et al.* (2014). *Echinacea purpurea* root extract enhances the adipocyte differentiation of 3T3-L1 cells. *Arch Pharm Res* 37(6): 803-812

Skaidickas D, Kondrotas A, Baltrusaitis K (2004). The effect of *Echinacea purpurea* extract on sexual glands of male rats. *Medicina (Kaunas)* 40: 1211-1218

- Skaudickas D, Kondrotas AJ, Baltrusaitis K, Vaitiekaitis G (2003). Effect of *Echinacea* (*Echinacea purpurea* (L.) Moench) preparations on experimental prostate gland. *Medicina (Kaunas)* 39: 761-766
- Skwarek T, Tynecka Z, Glowniak K, Lutostanska E (1996). *Echinacea* L. – Inducer of interferons. *Herba Polonica* 42: 110-117
- Sloley BD, Urichuk LJ, Tywin C, Coutts RT, Pang PK, Shan JJ (2001). Comparison of chemical components and antioxidants capacity of different *Echinacea* species. *J Pharm Pharmacol* 53: 849-857
- South EH, Exon JH (2001). Multiple immune functions in rats fed *Echinacea* extracts. *Immunopharmacol Immunotoxicol* 23: 411-421
- Speroni E, Govoni P, Guizzardi S, Renzulli C, Guerra MC (2002). Anti-inflammatory and cicatrizing activity of *Echinacea pallida* Nutt. root extract. *J Ethnopharmacol* 79: 265-272
- Stimpel M, Proksch A, Wagner H, Lohmann-Matthes ML (1984). Macrophage activation and induction of macrophage cytotoxicity by purified polysaccharide fractions from the plant *Echinacea purpurea*. *Infect Immun* 46: 845-849
- Sun LZ, Currier NL, Miller SC (1999). The American coneflower: a prophylactic role involving nonspecific immunity. *J Altern Complement Med* 5: 437-446
- Thygesen L, Thulin J, Mortensen A, Skibsted LH, Molgaard P (2007). Antioxidant activity of cichoric acid and alkamides from *Echinacea purpurea*, alone and in combination. *Food Chem* 101: 74-81
- Tiralongo E, Lea RA, Wee SS, Hanna MM, Griffiths LR (2012). Randomised, double blind, placebo-controlled trial of echinacea supplementation in air travellers. *E- Based Complem Altern Med* 417267
- Todd DA, Gullede TV, Britton ER, Oberhofer M, Leyte-Lugo M, Moody AN *et al.* (2015). Ethanolic *Echinacea purpurea* Extracts Contain a Mixture of Cytokine-Suppressive and Cytokine-Inducing Compounds, Including Some That Originate from Endophytic Bacteria. *PLoS One* 10(5): e0124276
- Uluşık D, Keskin E (2012). Effects of ginseng and echinacea on cytokine mRNA expression in rats. *ScieWorld J* 2012: 942025
- Wacker A, Hilbig W (1978). Virushemmung mit *Echinacea purpurea*. *Planta Med* 33: 89-102
- Wagner H, Breu W, Willer F, Wierer M, Remiger P, Schwenker G (1989). *In vitro* inhibition of arachidonate metabolism by some alkamides and prenylated phenols. *Planta Med* 55: 566-567
- Wagner H, Jurcic K (2002). Immunological studies of Revitonil, a phytopharmaceutical containing *Echinacea purpurea* and *Glycyrrhiza glabra* root extract. *Phytomedicine* 9: 390-397
- Wagner H, Proksch A, Riess-Maurer I, Vollmar A, Odenthal S, Stuppner H, Jurcic K, Le Turdu M, Fang JN (1984). Immunostimulating action of polysaccharides (heteroglycans) from higher plants. *Arzneimittelforsch* 35: 1069-1075
- Westendorf J (1982). Carito® – *in-vitro* Untersuchungen zum Nachweiss spasmolytischer und kontraktiler Einflüsse. *Therapiewoche* 32: 6291-6297
- WHO (1999). Radix echinaceae. In: WHO monographs on selected medicinal plants, Vol. 1, WHO, Geneva, Switzerland: 128-135
- Wolters Kluwer Health (2004). *Echinacea*. In: The Review of Natural Products, Wolters Kluwer Health, St. Louis, MO, USA
- Wu L, Rowe EW, Jeftinija K, Jeftinija S, Rizshsky L, Nikolau BJ *et al.* (2010). Echinacea-induced cytosolic Ca²⁺ elevation in HEK293. *BMC Complem Altern Med* 10: 72

Zhai Z, Haney D, Wu L, Solco A, Murphy PA, Wurtele ES, Kohut ML, Cunnick JE (2007b). Alcohol extracts of *Echinacea* inhibit production of nitric oxide and tumor necrosis factor-alpha by macrophages *in vitro*. *Food Agric Immunol* 18: 221-236

Zhai Z, Liu Y, Wu L, Senchina DS, Wurtele ES, Murphy PA, Kohut ML, Cunnick JE (2007a). Enhancement of innate and adaptive immune functions by multiple *Echinacea* species. *J Med Food* 10: 423-434

Zoutewelle G, Van Wijk R (1990). Effects of *Echinacea purpurea* extracts on fibroblast populated collagen lattice contraction. *PhytotherRes* 4: 77-81