

28 January 2015
EMA/HMPC/678996/2013
Committee on Herbal Medicinal Products (HMPC)

List of references supporting the assessment of *Vaccinium myrtillus L., fructus*

Draft

The Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Aktaş C, Şenkal V, Sarıkaya S, Kartı S. Bilberry potentiates warfarin effect? *Turk J Geriatr* 2011, 14(1): 79-81

Alfieri R, Sole P. Influence des anthocyanosides administrés par voie oro-perlinguale sur l'adapto-electroretinogramme (AERG) en lumière rouge chez l'Homme. [Influence of anthocyanosides, in oral-perlingual administration, on the adapto-electroretinogram (AERG) in red light in humans.] *C R Seances Soc Biol Fil* 1966, 160(8): 1590–1593

Allegra C, Pollari G, Criscuolo A, Bonifacio M. Antocianosidi e sistema microvasculotessutale. *Minerva Angiol* 1982, 7: 39–44

Anderson KG, Anderson A, Connor CG. Potential use of bilberry for dry eye relief. *Optometry* 2011, 82(6): 380

Anthocyanins. Available at: <http://www.inchem.org/documents/jecfa/jecmono/v17je05.htm>. Accessed 25/07/2012

Anthony JP, Fyfe L, Stewart D, McDougall GJ, Smith HV. The effect of blueberry extracts on *Giardia duodenalis* viability and spontaneous excystation of *Cryptosporidium parvum* oocysts, *in vitro*. *Methods* 2007, 42(4): 339–348

Anthony JP, Fyfe L, Stewart D, McDougall GJ. Differential effectiveness of berry polyphenols as anti-giardial agents. *Parasitology* 2011, 138(9): 1110–1116

Aura AM. Microbial metabolism of dietary phenolic compounds in the colon. *Phytochem Rev* 2008, 7(3): 407–429

Aura AM, Martin-Lopez P, O'Leary KA, Williamson G, Oksman-Caldentey KM, Poutanen K, et al. *In vitro* metabolism of anthocyanins by human gut microflora. *Eur J Nutr* 2005, 44(3): 133–142

Barnes J, Anderson LA, Philipson JD. Bilberry. In: *Herbal Medicines: a guide for healthcare professionals*, Pharmaceutical Press, London 2002, 73–77



Bell DR, Gochenaur K. Direct vasoactive and vasoprotective properties of anthocyanin-rich extracts. *J Appl Physiol* 2006, 100(4):1164-1170

Belleoud L, Leluan D, Boyer Y. Etude des effets des glucosides d'anthocyane sur la vision nocturne du personnel navigant. *Rev Med Aeronaut Spat* 1967, 6:5-10

Bertuglia S, Malandrino S, Colantuoni A. Effect of *Vaccinium myrtillus* anthocyanosides on ischaemia reperfusion injury in hamster cheek pouch microcirculation. *Pharmacol Res* 1995, 31(3-4):183-187

Bettini V, Guerra B, Martino R, Ton P, Tegazzini V. Contractile responses of isolated rat stomach to stimulation of postganglionic cholinergic fibers in the presence of *Vaccinium myrtillus* anthocyanosides. *Fitoterapia* 1986, 57:211-216

Bettini V, Mayellaro F, Ton P, Zanella P. Effects of *Vaccinium myrtillus* anthocyanosides on vascular smooth muscle. *Fitoterapia* 1984a, 55:265-272

Bettini V, Mayellaro F, Ton P, Zogno M. Interactions between *Vaccinium myrtillus* anthocyanosides and serotonin on splenic artery smooth muscle. *Fitoterapia* 1984b, 55:201-208

Bettini V, Mayellaro F, Patron E, Ton P, Terribile Wiel VM. Inhibition by *Vaccinium myrtillus* anthocyanosides of barium-induced contractions in segments of internal thoracic vein. *Fitoterapia* 1984c, 55:323-327

Bettini V, Mayellaro F, Pilla I, Terribile Wiel VM. Mechanical responses of isolated coronary arteries to barium in the presence of *Vaccinium myrtillus* anthocyanosides. *Fitoterapia* 1985a, 56:3-10

Bettini V, Aragno R, Bettini MB, Bettini V, Fiori A, Martino R, et al. Study of the mechanism whereby anthocyanosides potentiate the effect of catecholamines on coronary vessels. *Fitoterapia* 1985b, 56:67-72

Bettini V, Aragno R, Bettini M, Braggion G, Calore L, Concolato MT, et al. Vasodilator and inhibitory effects of *Vaccinium myrtillus* anthocyanosides on the contractile responses of coronary artery segments to acetylcholine: role of the prostacyclins and of the endothelium-derived relaxing factor. *Fitoterapia* 1991, 62:15-28

Bettini V, Aragno R, Bettini MB, Braggion G, Calore L, Morimando L, et al. Facilitating influence of *Vaccinium myrtillus* anthocyanosides on the acetylcholine-induced relaxation of isolated coronary arteries: role of the endothelium-derived relaxing factor. *Fitoterapia* 1993, 64:45-57

Bhargava SK. Anti-spermatogenic activity of malvidin chloride in langur monkeys (*Presbytis entellus entellus* Dufresne). *Int J Androl* 1990, 13(3):207-215

Bilberry Fruit Extract. SUMMARY OF DATA FOR CHEMICAL SELECTION 84082-34-8. Available at: http://ntp-server.niehs.nih.gov/ntp/htdocs/Chem_Background/ExSumPdf/Bilberry_508.pdf. Accessed 20/10/2013

Biedermann L, Mwinyi J, Scharl M, Frei P, Zeitz J, Kullak-Ublick GA, et al. Bilberry ingestion improves disease activity in mild to moderate ulcerative colitis - an open pilot study. *J Crohns Colitis* 2013, 7(4):271-279

Bomser J, Madhavi DL, Singletary K, Smith MA. *In vitro* anticancer activity of fruit extracts from *Vaccinium* species. *Planta Med* 1996, 62(3):212-216

Boniface R, Robert AM. Einfluss von Anthocyanen auf den Bindegewebsmetabolismus beim Menschen. [Effect of anthocyanins on human connective tissue metabolism in the human.] *Klin Monbl Augenheilkd* 1996, 209(6):368-372

Bornsek SM, Ziberna L, Polak T, Vanzo A, Ulrich NP, Abram V, et al. Bilberry and blueberry anthocyanins act as powerful intracellular antioxidants in mammalian cells. *Food Chem* 2012, 134(4):1878-1884

Bottecchia D, Bettini V, Martino R, Camerra G. Preliminary report on the inhibitory effect of *Vaccinium myrtillus* anthocyanosides on platelet aggregation and clot retraction. *Fitoterapia* 1987, 58:3-8

Brenneisen VR, Steinegger E. Zur Analytik der Polyphenole der Früchte von *Vaccinium myrtillus* L. (Ericaceae) " Pharm. Acta Helv 1981a, 56: 180-185

Brenneisen VR, Steinegger E. Quantitativer Vergleich der Polyphenole in früchten von *Vaccinium myrtillus* L. unter schiedlichen Reifegrades. *Pharm Acta Helv* 1981b, 56:341-343

Brown EM, Gill CIR, McDougall GJ, Stewart D. Mechanism underlying the anti-proliferative effects of berry components in *in vitro* models of colon cancer. *Curr Pharm Biotech* 2012, 13:200-209

Burdulis D, Ivanauskas L, Dirse V, Kazlauskas S, Razukas A. Study of diversity of anthocyanin composition in bilberry (*Vaccinium myrtillus* L.) fruits. *Medicina (Kaunas)* 2007, 43(12):971-977

Cahyana Y, Gordon MH. Interaction of anthocyanins with human serum albumin: Influence of pH and chemical structure on binding. *Food Chem* 2013, 141:2278-2285

Camire M E. Phytochemicals in the *Vaccinium* Family Bilberries, Blueberries, and Cranberries. In: MS Meskin, et al., editors. *Phytochemicals in Nutrition and Health*. CRC Press 2002, 19-40

Campbell FM, Nicol PF, Moar KM, Cruickshank M, Hoggard N. Lower levels of damaged protein biomarkers in the plasma of overweight type 2 diabetic men following supplementation with a standardised bilberry extract. *Proc Nutr Soc* 2012, 71:E130

Canter PH, Ernst E. Anthocyanosides of *Vaccinium myrtillus* (bilberry) for night vision - a systematic review of placebo-controlled trials. *Surv Ophthalmol* 2004, 49: 38-50

Capasso F, Gaginella TS, Grandolini G, Izzo AA. Bilberry. In: *Phytotherapy*, Springer Verlag, Berlin 2003, 125-126

Cerutti R, Ferruzzi BE, Balbi E, Santelli F, Sbrignadello C. Utilità degli antocianosidi del mirtillo nella profilassi degli effetti collaterali minori nella contraccezione con dispositivi intrauterine medicate al rame. [Value of *Vaccinium myrtillus* anthocyanosides in the prophylaxis of minor side effects with copper intrauterine device contraception.] *Ginecol Clin* 1984, 3-4: 244–249

Clifford MN. Anthocyanins – nature, occurrence and dietary burden. *J Sci Food Agric* 2000, 80:1063-1072

Cluzel C, Bastide P, Wegman R, Tronche P. Activités enzymatiques de la rétine et anthocyanosides extraits de *Vaccinium myrtillus* (lactate-deshydrogenase, α -hydroxybutyrate- deshydrogenase, 6-phosphogluconate- deshydrogenase, α -glycerophosphate-deshydrogenase, 5-nucleotidase, phosphoglucosidomerase). *Biochem Pharmacol* 1970, 19:2295-302

Cluzel C, Bastide P, Tronche P. Activités phosphoglucomutase et glucose-6-phosphatasique de la rétine et anthocyanosides extraits de *Vaccinium myrtillus* (étude *in vitro* et *in vivo*). *CR Soc Biol* 1969, 163:147-50

Coget PJ, Merlen JF. Étude clinique d'un nouvel agent de protection vasculaire le Difrarel 20, composé d'anthocyanosides extraits du *Vaccinium myrtillus*. *Phlébologie* 1968, 21:221-228

Commission E Monograph. Myrtilli fructus. Bundesanzeiger No. 76 of 23 April 1987, amended in Bundesanzeiger No. 50 of 13 March 1990

Contestabile M, Appolloni R, Suppressa F, D'Alba E, Pecorelli B. Trattamento prolungato con antocianosidi del mirtillo ad alto dosaggio: risposte elettrofisiologiche nel paziente miope. [Prolonged treatment with blueberry anthocyanosides administered in high doses: electrophysiological response in myopic patients.] *Boll Ocul* 1991, 70:1157-1169

Cristoni A, Magistretti MJ. Antiulcer and healing activity of *Vaccinium myrtillus* anthocyanosides. *Farmaco Prat* 1987, 42:29-43

Dall'Asta M, Calani L, Tedeschi M, Jechiu L, Brighenti F, Del Rio D. Identification of microbial metabolites derived from *in vitro* fecal fermentation of different polyphenolic food sources. *Nutrition* 2012, 28:197-203

Delgado-Vargas F., Jiménez AR, Paredes-López O. Natural Pigments: Carotenoids, Anthocyanins, and Betalains — Characteristics, Biosynthesis, Processing, and Stability. *Crit Rev Food Sci Nutr* 2003, 40: 173–289

Djuv A, Nilsen OG, Steinsbekk A The co-use of conventional drugs and herbs among patients in Norwegian general practice: a cross-sectional study. *BMC Complement Altern Med* 2013, 30, 13:295

Ek S, Kartimo H, Mattila S, Tolonen A. Characterization of phenolic compounds from lingonberry (*Vaccinium vitis-idaea*). *J Agric Food Chem* 2006, 54:9834–9842

ESCOP Monographs 2nd ed. Myrtilli fructus – Bilberry fruit. European Scientific Cooperative on Phytotherapy, editor. Thieme, Stuttgart 2003, 345-350

Esseelen M, Fritz J, Hutter M, Teller N, Baechler S, Boettler U, et al. Anthocyanin-rich extracts suppress the DNA-damaging effects of topoisomerase poisons in human colon cancer cells. *Mol Nutr Food Res* 2011, 55(Suppl 1):S143-153

European Pharmacopoeia 8th ed. Bilberry fruit, dried - Myrtilli fructus siccus. Council of Europe. 01/2008:1588

European Pharmacopoeia 8th ed. Bilberry fruit, fresh - Myrtilli fructus recens. Council of Europe. 01/2008:1602

European Pharmacopoeia 8th ed. Bilberry fructus dry extract prepared from fresh bilberry fruit – Myrtilli fructus recentis extractum siccum raffinatum et normatum. Council of Europe. 07/2008:2394

Felgines C, Talavera S, Texier O, Gil-Izquierdo A, Lamaison J-L, Remesy C. Blackberry Anthocyanins Are Mainly Recovered from Urine as Methylated and Glucuronidated Conjugates in Humans. *J Agric Food Chem* 2005, 53:7721-7727

Felgines C, Talavera S, Texier O, Besson C, Fogliano V, Lamaison JL, et al. Absorption and metabolism of red orange juice anthocyanins in rats. *Br J Nutr* 2006, 95:898-904

Felgines C, Texier O, Besson C, Vitaglione P, Lamaison J, Fogliano V, et al. Influence of glucose on cyanidin 3-glucoside absorption in rats. *Mol Nutr Food Res* 2008, 52:959–964

Fleschhut J, Kratzer F, Rechkemmer G, Kulling ES. Stability and biotransformation of various dietary anthocyanins *in vitro*. *Eur J Nutr* 2006, 45:7-18

Fossen T, Cabrita L, Andersen EM. Colour and stability of pure anthocyanins influenced by pH including the alkaline region. *Food Chem* 1998, 63:435-440

Friedrich H, Schönert J. Untersuchungen über einige Inhaltsstoffe der Blätter und Früchte von *Vaccinium myrtillus*. *Planta Med* 1973, 24:90-100

Frohne D. Heilpflanzenlexikon. Ein Leitfaden auf wissenschaftlicher Grundlage. *Vaccinium myrtillus* Stuttgart, 2006 [Polish Translation]

Fuchikami H, Satoh H, Tsujimoto M, Ohdo S, Ohtani H, Sawada Y. Effects of herbal extracts on the function of human organic anion-transporting polypeptide OATP-B. *Drug Metab Dispos* 2006, 34:577-582

Galvano F, Vitaglione P, Li Volti G, Di Giacomo C, Gazzolo D, Vanella L, et al. Protocatechuic acid: the missing human cyanidins' metabolite. *Mol Nutr Food Res* 2008, 52:386-387, author reply 388

Garcia V, Diban N, Gorri D, Keiski R, Urtiaga A, Ortiz I. Separation and concentration of bilberry impact aroma compound from dilute model solution by pervaporation. *J Chem Technol Biotechnol* 2008, 83:973-982

Gardner Z, McGuffin M. Botanical Safety Handbook. *Vaccinium myrtillus* L. American Herbal Products Association, Taylor and Francis 2013

Gatta L. *Vaccinium myrtillus* anthocyanosides in the treatment of venous stasis: controlled clinical study on sixty patients. *Fitoterapia* 1988, 59:19-26

Gloria E, Perla A. Effetto degli antocianosidi sulla soglia visiva assoluta [Effect of anthocyanosides on the absolute visual threshold.] *Ann Oftalmol Clin Ocul* 1966, 92:595-607

Govindaraghavan S. Pharmacopoeial HPLC identification methods are not sufficient to detect adulterations in commercial bilberry (*Vaccinium myrtillus*) extracts. Anthocyanin profile provides additional clues. *Fitoterapia*. 2014 Sep 20. pii: S0367-326X(14)00255-X. doi: 10.1016/j.fitote.2014.09.007. [Epub ahead of print]

Grismondi G. Contributo al trattamento delle flebopatie da stasi in gravidanza. [Treatment of pregnancy induced phlebopathies.] *Minerva Gin* 1981, 33:221-230

Gruenwald J, Brendler T, Jaenicke C, editors. PDR for Herbal Medicines. 4th ed. Medical Economics Co., Inc., Montvale 2000

Hansel R, Keller K, Rimpler H, Schneider G, editors. Hagers Handbuch der Pharmazeutischen Praxis. Drogen P-Z. Vol 6. Berlin: Springer, 1994

Heinonen IM, Meyer AS, Frankel EN. Antioxidant activity of berry phenolics on human low-density lipoprotein and liposome oxidation, *J Agric Food Chem* 1998, 46:4107-4112

Heinonen M. Antioxidant activity and antimicrobial effect of berry phenolics--a Finnish perspective. *Mol Nutr Food Res* 2007, 51:684-691

Hou DX, Yanagita T, Uto T, Masuzaki S, Fujii M. Anthocyanidins inhibit cyclooxygenase-2 expression in LPS-evoked macrophages: structure-activity relationship and molecular mechanisms involved. *Biochem Pharmacol* 2005, 70:417-25

Huttunen S, Toivanen M, Arkko S, Ruponen M, Tikkanen-Kaukanen C. Inhibition activity of wild berry juice fractions against *Streptococcus pneumoniae* binding to human bronchial cells. *Phytother Res* 2011, 25:122-127

Ichiyanagi T, Hatano Y, Matsugo S, Konishi T. Structural dependence of HPLC separation pattern of anthocyanins from Bilberry (*Vaccinium myrtillus* L.). *Chem Pharm Bull* 2004a, 52:628-630

Ichiyanagi T, Kashiwada Y, Ikeshiro Y, Shida, Y, Hatano, Y Shida Y, et al. Complete assignment of bilberry (*Vaccinium myrtillus* L.) anthocyanins separated by capillary zone electrophoresis. *Chem Pharm Bull* 2004b, 52:226–229

Jayle GE, Aubert L. Action des glucosides d'anthocyanes sur la vision scotopique et mesopique du sujet normal. *Therapie* 1964, 19:171–175

Jayle GE, Aubry M, Gavini H, Braccini G, de la Baume C. Etude concernant l'action sur la vision nocturne des anthocyanosides extrait du *Vaccinium myrtillus*. *Ann Ocul* 1965, 198:556–62

Johnson J, Maher P, Hanneken A. The flavonoid, eriodictyol, induces long-term protection in ARPE-19 cells through its effects on Nrf2 activation and phase 2 gene expression. *Invest Ophthalmol Vis Sci* 2009, 50:2398-2406

Kahle K, Kraus M, Scheppach W, Ackermann M, Ridder F, Richling E. Studies on apple and blueberry fruit constituents: Do the polyphenols reach the colon after ingestion? *Mol Nutr Food Res* 2006, 50: 418–423

Kalt W, McDonald J E, Ricker R D and Lu X. Anthocyanin content and profile within and among blueberry species. *Can J Plant Sci* 1999, 79:617–623

Kalt W, Hanneken A, Milbury P, Tremblay F. Recent research on polyphenolics in vision and eye health. *J Agric Food Chem* 2010, 58:4001-4007

Karlsen A, Paur I, Bøhn SK, Sakhi AK, Borge GI, Serafini M, et al. Bilberry juice modulates plasma concentration of NF-kappaB related inflammatory markers in subjects at increased risk of CVD. *Eur J Nutr* 2010, 49:345-355

Katsume N, Iwashita K, Tsushida T, Yamaki K, Kobori M. Induction of apoptosis in cancer cells by Bilberry (*Vaccinium myrtillus*) and the anthocyanins. *J Agric Food Chem* 2003, 51:68-75

Kay CD. Aspects of anthocyanin absorption, metabolism and pharmacokinetics in humans. *Nutr Rev* 2006, 19:137–146

Kay CD, Mazza GJ, Holub BJ. Anthocyanins exist in the circulation primarily as metabolites in adult men. *J Nutr* 2005, 135:2582-2588

Kähkönen MP, Hopia AI, Heinonen M. Berry phenolics and their antioxidant activity. *J Agric Food Chem* 2001, 49:4076-4082

Kähkönen MP, Heinämäki J, Ollilainen V, Heinonen M. Berry anthocyanins: Isolation, identification and antioxidant activities. *J Sci Food Agric* 2003, 83:1403-1411

Kemperman RA, Bolca S, Roger LC, Vaughan EE. Novel approaches for analysing gut microbes and dietary polyphenols: challenges and opportunities. *Microbiology* 2010, 156:3224-3231

Keppler K., Humpf H-U. Metabolism of anthocyanins and their phenolic degradation products by the intestinal microflora. *Bioorg Med Chem* 2005, 13:5195-5205

Kim ES, Yu SY, Kwon SJ, Kwon OW, Kim SY, Kim TW, et al. Clinical Evaluation of Patients with non proliferative diabetic retinopathy following medication of anthocyanoside: Multicenter Study. *Korean Ophthalmol Soc* 2008, 49:1629-1633

Kolehmainen M, Mykkänen O, Kirjavainen PV, Leppänen T, Moilanen E, Adriaens M, et al-. Bilberries reduce low-grade inflammation in individuals with features of metabolic syndrome. *Mol Nutr Food Res* 2012, 56:1501-1510

Kong JM, Chia LS, Goh NK, Chia TF, Brouillard R. Analysis and biological activities of anthocyanins. *Phytochemistry* 2003, 64: 923-933

Kramer JH. Anthocyanosides of *Vaccinium myrtillus* (Bilberry) for Night Vision—A Systematic Review of Placebo-controlled Trials. *Surv Ophthalmol* 2004, 49: 618

Lamy S, Lafleur R, Bédard V, Moghrabi A, Barrette S, Gingras D, et al. Anthocyanidins inhibit migration of glioblastoma cells: structure-activity relationship and involvement of the plasminolytic system. *J Cell Biochem* 2007, 100: 100-111

Laplaud PM, Lelubre A, Chapman MJ. Antioxidant action of *Vaccinium myrtillus* extract human low density lipoproteins *in vitro*: initial observations. *Fundam Clin Pharmacol* 1997, 11: 35-40

Lätti AK, Riihinen KR, Kainulainen PS. Analysis of anthocyanin variation in wild population of bilberry (*Vaccinium myrtillus* L.) in Finland. *J Agric Food Chem* 2008, 56: 190-196

Levy Y, Glovinsky Y. The effect of anthocyanosides on night vision. *Eye* 1998, 12: 967-969

Lietti A, Forni G. Studies on *Vaccinium myrtillus* anthocyanosides: II. Aspects of anthocyanins pharmacokinetics in the rat. *Arzneimittelforschung* 1976, 26: 832-835

Magnasco A, Zingirian M. Influenza degli antocianosidi sulla soglia retinica differenziale mesopica [Influence of anthocyanosides on the mesopic differential threshold of the retina.] *Ann Oftalmol Clin Ocul* 1966, 92: 188-193

Maher P, Hanneken A. Flavonoids Protect Retinal Ganglion Cells from Oxidative Stress-Induced Death. *IOVS* 2005, 46: 4796-4803

Maher P, Hanneken A. Flavonoids protect retinal ganglion cells from ischemia *in vitro*. *Exp Eye Res* 2008, 86: 366-374

Malaveille C, Hautefeuille A, Pignatelli B, Talaska G, Vineis P, Bartsch H. Dietary phenolics as anti-mutagens and inhibitors of tobacco-related DNA adduction in the urothelium of smokers. *Carcinogenesis* 1996, 17: 2193-2200

Manach C, Williamson G, Morand C, Scalbert A, Rémesy C. Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. *Am J Clin Nutr* 2005, 81(Suppl 1): 230-242.

Martín-Aragón S, Basabe B, Benedí J, Villar A. Antioxidant Action of *Vaccinium myrtillus* L. *Phytother Res* 1997, 12: 104-106

Martín-Aragón S, Basabe B, Benedí J, Villar A. *In vitro* and *in vivo* Antioxidant Properties of *Vaccinium myrtillus*. *Pharm Biol* 1999, 37: 109-113

Martin-Aragon S, Basabe B, Benedi J, Villar A. Antioxidant action of *Vaccinium myrtillus* L. *Phytother Res* 1998, 12: 104-106

Matsumoto H, Nakamura Y, Tachibanaki S, Kawamura S, Hirayama M. Stimulatory Effect of Cyanidin 3-Glycosides on the Regeneration of Rhodopsin. *J Agric Food Chem* 2003, 51: 3560-3563

Matuschek MC, Hendriks WH, McGhie TK, Reynolds GW. The jejunum is the main site of absorption for anthocyanins in mice. *J Nutr Biochem* 2006, 17: 31-36

Mayser HM, Wilhelm H. Effects of anthocyanosides on contrast vision (abstract). *Invest Ophthalmol Vis Sci* 2001, 42(Suppl): 63

Mazza G, Kay CD, Cottrell T, Holub BJ. Absorption of anthocyanins from blueberries and serum antioxidant status in human subjects. *J Agric Food Chem* 2002, 50: 7731–7737

Määttä-Riihinen KR, Kamal-Eldin A, Mattila PH, González-Paramás AM, Törrönen AR. Distribution and contents of phenolic compounds in eighteen Scandinavian berry species. *J Agric Food Chem* 2004, 52: 4477–4486

McGhie T, Walton MC. The bioavailability and absorption of anthocyanins: Towards a better understanding. *Molecular Nutr Food Res* 2007, 51: 702–713

Mian E. Anthocyanosides and microvessel walls: new findings on the mechanism of action of their protective effect in syndromes due to abnormal capillary fragility. *Minerva Med* 1977, 68: 3565–3581

Milbury PE, Graf B, Curran-Celentano JM, Blumberg JM. Bilberry (*Vaccinium myrtillus*) anthocyanins modulate heme oxygenase-1 and glutathione S transferase-pi expression in ARPE-19 cells. *Invest Ophthalmol Vis Sci* 2007, 48: 2343–2343

Mills S, Bone K. Bilberry fruit (*Vaccinium myrtillus*). In: Principles and Practice of Phytotherapy. Churchill Livingstone, Edinburgh 2000, 297–301

Morazzoni P, Bombardelli E. Review: *Vaccinium myrtillus* L. *Fitoterapia* 1996, 67: 3–29

Morazzoni P, Magistretti M. Activity of Myrtocyan®, an anthocyanoside complex from *Vaccinium myrtillus* (VMA), on platelet aggregation and adhesiveness. *Fitoterapia* 1990, 61: 13–22

Morazzoni P, Livio S, Scilingo A, Malandrino S. *Vaccinium myrtillus* anthocyanosides pharmacokinetics in rats. *Arzneimittelforschung Drug Res* 1991, 41: 128–131

McGhie T, Walton MC. The bioavailability and absorption of anthocyanins: Towards a better understanding. *Mol Nutr Food Res* 2007, 51: 702–713

Muth ER, Laurent JM, Jasper P. The effect of bilberry nutritional supplementation on night visual acuity and contrast sensitivity. *Alt Med Rev* 2000, 5(2): 164–173

Nielsen ILF, Dragsted LO, Ravn-Haren G, Freese R, Rasmussen SE. Absorption and excretion of black currant anthocyanins in humans and Watanabe heritable hyperlipidemic rabbits. *J Agric Food Chem* 2003, 51: 2813–2820

Nohynek LJ, Alakomi HL, Kähkönen MP, Heinonen M, Helander IM, Oksman-Caldentey KM, et al. Berry phenolics: antimicrobial properties and mechanisms of action against severe human pathogens. *Nutr Cancer* 2006, 54: 18–32

Nurmi T, Mursu J, Heinonen M, Nurmi A, Hiltunen R, Voutilainen S. Metabolism of berry anthocyanins to phenolic acids in humans. *J Agric Food Chem* 2009, 57: 2274–2281

Ogawa K, Oyagi A, Tank J, Kobayashi S, Hara H. The protective effect and action mechanism of *Vaccinium myrtillus* L. on gastric ulcer in mice. *Phytother Res* 2011, 25: 1160–1165

Paoletti A, Gallo E, Benemei S, Vietri M, Lapi F, Volpi R, et al. Interactions between Natural Health Products and Oral Anticoagulants: Spontaneous Reports in the Italian Surveillance System of Natural Health Products. *Evid Based Complement Alternat Med* 2011, 612150: 1–5

Passamonti S, Vrhovsek U, Vanzo A, Mattivi F. The stomach as a site for anthocyanins from food. *FEBS Lett* 2003, 544: 210–213

Patras A, Brunton NP, O'Donnell C, Tiwari BK. Effects of thermal processing on thermal stability in foods, mechanisms of kinetics and degradation. *Trends Food Sci Techn* 2010, 21: 3–11

Pennarola R, Roco P, Matarazzo G, De Martino F, Mormile A, Labate L. L'azione terapeutica degli antocianosidi nelle alterazioni microcircolatorie da polinevrite da collanti. [The therapeutic action of the anthocyanosides in microcirculatory changes due to adhesive-induced polyneuritis.] *Gazz Med Ital* 1980, 139: 485–491

Perossini M, Chiellini S, Guidi G, Siravo D. Diabetic and hypertensive retinopathy therapy with *Vaccinium myrtillus* anthocyanosides (Tegens) double-blind placebo-controlled clinical trial. *Ann Ottalmol Clin Ocul* 1987, 113: 1173–1190

Peris J, Pascual B, Garcia L, Muro N, Surra G, Vinuesa T. Drug herb interactions with oral anticoagulants. *Pharm World Sci* 2008, 30: 677–678 Abstr. DI-81, No 5. Published in: Reactions Weekly 2008, 1229:5

Prior RL, Cao G, Martin A, Sofic E, McEwen J, O'Brien C, et al. Antioxidant Capacity As Influenced by Total Phenolic and Anthocyanin Content, Maturity, and Variety of *Vaccinium* Species. *J Agric Food Chem* 1998, 46: 2686–2693

Prior RL, Cao G, Martin A, Sofic E, McEwen J, O'Brien C, et al. Antioxidant capacity as influenced by total phenolic and anthocyanin content, maturity, and variety of *Vaccinium* species. *J Agric Food Chem* 1998, 46: 2686–93

Prior RL, Cao G. Antioxidant Phytochemicals in Fruits and Vegetables: Diet and Health Implications. *Hort Science* 2000, 35: 588–592

Pulliero G, Montin S, Bettini, Martino R, Mogno C, Lo Castro G. *Ex vivo* study of the inhibitory effects of *Vaccinium myrtillus* anthocyanosides on human platelet aggregation. *Fitoterapia* 1989, 60: 69–75

Puupponen-Pimiä R, Nohynek L, Alakomi HL, Oksman-Caldentey KM. Bioactive berry compounds – novel tools against human pathogens. *Appl Microbiol Biotechn* 2005a, 67: 8–18

Puupponen-Pimiä R, Nohynek L, Hartmann-Schmidlin S, Kähkönen M, Heinonen M, Maatta-Riihinen K, et al. Berry phenolics selectively inhibit the growth of intestinal pathogens. *J Appl Microbiol* 2005b, 98: 991–1000

Puupponen-Pimiä R, Nohynek L, Alakomi HL, Oksman-Caldentey KM. The action of berry phenolics against human intestinal pathogens. *Biofactors* 2005c, 23: 243–251

Rauha JP, Remes S, Heinonen M, Hopia A, Kähkönen M, Kujala T, et al. Antimicrobial effects of Finnish plant extracts containing flavonoids and other phenolic compounds. *Int J Food Microbiol* 2000, 56: 3–12

Repossi P, Malagola R, De Cadilhac C. The role of anthocyanosides on vascular permeability in diabetic retinopathy. *Ann Ottalmol Clin Ocul* 1987, 113: 357–361

Riihinen K, Ryyränen A, Toivanen M, Könönen E, Törrönen R, Tikkanen-Kaukanen C. Antiaggregation potential of berry fractions against pairs of *Streptococcus mutans* with *Fusobacterium nucleatum* or *Actinomyces naeslundii*. *Phytother Res* 2011, 25: 81–87

Rivas-Gonzalo JC. Analysis of anthocyanins. In: Santos-Buelga C, Williamson G, editors. *Methods in Polyphenol Analysis*. The Royal Society of Chemistry Cambridge, 2003, 338–358

Rotblatt M, Ziment I. Evidence-Based Herbal Medicine. Bilberry (*Vaccinium myrtillus*). In: Hanley and Belfus, Philadelphia 2002: 92–97

Sakakibara H, Ogawa T, Koyanagi A, Kobayashi S, Goda T, Kumazawa S, et al. Distribution and excretion of bilberry anthocyanins [corrected] in mice. *J Agric Food Chem* 2009, 57: 7681–7686

Sala D, Rolando M, Rossi P, Pisarello L. Effetto degli antocianosidi sulle "performances" visive alle basse luminanze. [Effect of anthocyanosides on visual performances at low illumination.] *Minerva Oftalmol* 1979, 21:283–285

Scharrer A, Ober M. [Anthocyanosides in the treatment of retinopathies.] *Klin Monatsbl Augenheilkd* 1981, 178:386–389 [German]

Schilcher H. Phytotherapie in der Kinderheilkunde. Wissenschaftliche Verlagsgesellschaft, Stuttgart 1992 [Polish Edition PZWL, Warszawa 1998]

Schimmer O, Krüger A, Paulini H, Haefele F. An evaluation of 55 commercial plant extracts in the Ames mutagenicity test. *Pharmazie* 1994, 49:448-451

Selma MV, Espín JC, Tomás-Barberán FA. Interaction between phenolics and gut microbiota: role in human health. *J Agric Food Chem* 2009, 57:6485-6501

Shim SH, Kim JM, Choi CY, Kim CY, Park KH. *Ginkgo biloba* extract and bilberry anthocyanins improve visual function in patients with normal tension glaucoma. *J Med Food* 2012, 15:818-823

Song J, Li Y, Ge J, Duan Y, Sze SC, Tong Y, et al. Protective effect of bilberry (*Vaccinium myrtillus* L.) extracts on cultured human corneal limbal epithelial cells (HCLEC). *Phytother Res* 2010, 24:520-524

Steinert RE, Ditscheid B, Netzel M, Jahreis G. Absorption of black currant anthocyanins by monolayers of human intestinal epithelial Caco-2 cells mounted in ussing type chambers. *J Agric Food Chem* 2008, 56:4995-5001

Stoner GD, Wang L-S, Casto BC. Laboratory and clinical studies of cancer chemoprevention by antioxidants in berries. *Carcinogenesis* 2008, 29:1665–1674

Su Z. Anthocyanins and Flavonoids of *Vaccinium* L. *Pharm Crops* 2012, 3:7-37

Svobodová A, Rambousková J, Walterová D, Vostalová J. Bilberry extract reduces UVA-induced oxidative stress in HaCaT keratinocytes: a pilot study. *Biofactors* 2008, 33:249-266

Svobodová A, Zdarilová A, Vostálová J. *Lonicera caerulea* and *Vaccinium myrtillus* fruit polyphenols protect HaCaT keratinocytes against UVB-induced phototoxic stress and DNA damage. *J Dermatol Sci* 2009, 56:196-204

Szakiel A, Pączkowski C, Huttunen S. Triterpenoid content of berries and leaves of bilberry *Vaccinium myrtillus* from Finland and Poland. *J Agric Food Chem* 2012, 60:11839-11849

Talavéra S, Felgines C, Texier O, Besson C, Lamaison JL, Rémesy C. Anthocyanins are efficiently absorbed from the stomach in anesthetized rats. *J Nutr* 2003, 133:4178-4182

Talavera S, Felgines C, Texier O, Besson C, Manach C, Lamaison JL, et al. Anthocyanins are efficiently absorbed from the small intestine in rats. *J Nutr* 2004, 134:2275-2279

Tangen O, Berman HJ, Marfey R. Gel Filtration. A New Technique for Separation of Blood Platelets from Plasma. *Thromb Diath Haemorrh* 1971, 25:268-278

Tirupula KC, Balem F, Yanamala N, Klein-Seetharaman J. pH-dependent interaction of rhodopsin with cyanidin-3-glucoside. 2. Functional aspects. *Photochem Photobiol* 2009, 85:463-70

Toivanen M, Ryynänen A, Huttunen S, Duricová J, Riihinens K, Törrönen R, et al. Binding of *Neisseria meningitidis* pili to berry polyphenolic fractions. *J Agric Food Chem* 2009, 57:3120-3127

Toivanen M, Huttunen S, Lapinjoki S, Tikkainen-Kaukanen C. Inhibition of adhesion of *Neisseria meningitidis* to human epithelial cells by berry juice polyphenolic fractions. *Phytother Res* 2011, 25:828-832

Triebel S, Trieu HL, Richling E. Modulation of inflammatory gene expression by a bilberry (*Vaccinium myrtillus* L.) extract and single anthocyanins considering their limited stability under cell culture conditions. *J Agric Food Chem* 2012, 60:8902-8910

Trumbeckaité S, Burdulis D, Raudonė L, Liobikas J, Toleikis A, Janulis V. Direct effects of *Vaccinium myrtillus* L. fruit extracts on rat heart mitochondrial functions. *Phytother Res* 2013, 27:499-506

Upton R, Graff A, Länger R, Sudberg S, Sudberg E, Miller T, et al. Bilberry Fruit, *Vaccinium myrtillus* L. American Herbal Pharmacopoeia and Therapeutic Compendium: Standards of Analysis, Quality Control, and Therapeutics. (2001). Eds. R. Upton, A Graff, C Petrone. Santa Cruz, CA: American Herbal Pharmacopoeia

Valentová K, Ulrichová J, Cvak L, Šimánek V. Cytoprotective effect of a bilberry extract against oxidative damage of rat hepatocytes. *Food Chem* 2007, 101:912-917

Vannini L, Samuelli R, Coffano M, Tibaldi L. Studio del comportamento pupillare in seguito a somministrazioni di antocianosidi.[Study of the papillary reflex after anthocyanoside administration.] *Bollett Ocul* 1986, 65(Suppl 6):569-577

Viljanen K, Kylli P, Kivistö R, Heinonen M. Inhibition of protein and lipid oxidation in liposomes by berry phenolics. *J Agric Food Chem* 2004, 52:7419-7424

Vitaglione P, Donnarumma G, Napolitano A, Galvano F, Gallo A, Scalfi L, et al. Protocatechuic acid is the major human metabolite of cyanidin-glucosides. *J Nutr* 2007, 137:2043-2048

Wang LS, Stoner GD. Anthocyanins and their role in cancer prevention. *Cancer Lett* 2008, 269:281-290

Wichtl M. Herbal Drugs and Phytopharmaceuticals. Medpharm GmbH Scientific Publisher, CRC Press, Stuttgart 1994

Williamson G, Clifford MN. Colonic metabolites of berry polyphenols: the missing link to biological activity? *Br J Nutr* 2010, 104(Suppl 3):48-66

Williamson G, Manach C. Bioavailability and bioefficacy of polyphenols in humans. II. Review of 93 intervention studies. *Am J Clin Nutr* 2005, 81(Suppl 1):243-255.

Yamamoto M, Yamaura K, Ishiwatari M, Shimada M, Kado S, Seki H, et al. Degradation Index for Quality Evaluation of Commercial Dietary Supplements of Bilberry Extract. *J Food Sci* 2013, 78:477-483

Yamamoto M, Yamaura K, Ishiwatari M, Ueno K. Difficulty for consumers in choosing commercial bilberry supplements by relying only on product label information. *Pharmacognosy Res* 2013, 5:212-215

Yanamala N, Tirupula KC, Balem F, Klein-Seetharaman J. pH-dependent Interaction of Rhodopsin with Cyanidin-3-glucoside. 1. Structural Aspects. *Photochem Photobiol* 2009, 85:454-462

Yue X, Xu Z. Changes of anthocyanins, anthocyanidins, and antioxidant activity in bilberry extract during dry heating. *J Food Sci* 2008, 73:494-499

Zadok D, Levy Y, Glovinsky Y. The effect of anthocyanosides in a multiple oral dose on night vision. *Eye* 1999, 13: 734-736

Zanetti-Ripamonti G. Piante Medicinali Nostre. Mirtillo (*Vaccinium myrtillus L.*). Istituto Editoriale Ticinese, Lugano-Bellinzona 1940

Zhang Z, Kou X, Fugal K, McLaughlin J. Comparison of HPLC methods for determination of anthocyanins and anthocyanidins in bilberry extracts. *J Agric Food Chem* 2004, 52: 688-691

Ziberna L, Lunder M, Tramer F, Drevenšek G, Passamonti S. The endothelial plasma membrane transporter bilitranslocase mediates rat aortic vasodilation induced by anthocyanins. *Nutr Metab Cardiovasc Dis* 2013, 23: 68-74