

30 May 2017
EMA/HMPC/186849/2017
Committee on Herbal Medicinal Products (HMPC)

List of references supporting the assessment of *Artemisia absinthium L.*, herba

Final

The European Medicines Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Abass K, Reponen P, Mattila S, Pelkonen O. Metabolism of α-thujone in human liver preparations *in vitro*. *Xenobiotica* 2011, 41(2):101-11. doi: 10.3109/00498254.2010.528066. Epub 2010 Nov 18.

AB-DDR. Pharmacopoeia of the German Democratic Republic. Berlin, Akademie-Verlag, 1988

Algieri F, Rodriguez-Nogales A, Rodriguez-Cabezas ME, Risco S, Ocete MA, Galvez J. Botanical Drugs as an Emerging Strategy in Inflammatory Bowel Disease: A Review. *Mediators of Inflamm* 2015, Article ID 179616, 14 pages

Ariño A, Arberas I, Renobales G et al. Seasonal variations in wormwood (*Artemisia absinthium L.*) essential oil composition. *J Essent Oil Res* 1999a, 11: 619-622

Ariño A, Arberas I, Renobales G et al. Essential oil of *Artemisia absinthium L.* from the Spanish Pyrenees. *J Essent Oil Res* 1999b, 11: 182-184

Ariño A, Arberas I, Renobales G et al. Influence of extraction method and storage conditions on the volatile oil of wormwood (*Artemisia absinthium L.*). *Eur Food Res Technol* 1999c, 209(2): 126-129

Arnold WN. Absinthe. *Sci Am* 1989, 260(6): 112-117

Baumann IC. Über die Wirkung von *Chelidonium*, *Curcuma*, Absinth und *Carduus Marianus* auf die Galle- und Pankreassekretion bei Hepatopathien. *Med Mschr* 1975, 29:173-180

Baumann IC, Glatzel H, Muth HW . Untersuchungen der Wirkung von Wermut (*Artemisia absinthium L.*) auf die Gallen- und Pankreasaff-Sekretion des Menschen.

Z Allgemeinmed 1975, 51(17): 784-791

Bayerl C, Jung EG . Allergic contact stomatitis from aristochol, a phytotherapeutic cholagogue. *Contact Derm* 1996, 34(3): 222-223

Blaschek W, Ebel S, Hackenthal E, et al., editors. *Absinthii herba*. HagerROM 2006: Hagers Handbuch der Drogen und Arzneistoffe. Version 5.0. Springer Medizin Verlag, Heidelberg 2006



Blumberger W, Glatzel H. Über die Speichelwirksamkeit von Bitterstoffen. *Planta Med* 1966, 14: 52-60

Bonkovsky HL, Cable EE, Cable JW et al. Porphyrogenic properties of the terpenes, camphor, pinene and thujone (With a note on historic implications for absinthe and the illness of Vincent van Gogh). *Biochem Pharmacol* 1992, 43: 2359-2368

British Herbal Pharmacopoeia. British Herbal Medicine Association, Bournemouth 1983

Canadanovic-Brunet JM, Djilas SM, Cetkovic GS et al. Free radical scavenging activity of wormwood (*Artemisia absinthium* L.) extracts. *J Sci Food Agric* 2005, 85: 265-272

Carnat AP, Madesclaire M, Chavignon O et al. Cis-Chrysanthenol, a main component in essential oil of *Artemisia absinthium* L. growing in Auvergne (Massif Central), France. *J Essent Oil Res* 1992, 4: 487-490

Centini F, Laurini GP. Una intossicazione da olio di salvia. *Zacchia* 1987, 60: 263-274

Chialva F, Liddle PAP, Godlia G . Chemotaxonomy of wormwood (*Artemisia absinthium* L.). (I. Composition of the essential oil of several chemotypes). *Z Lebensm Unters Forsch* 1983, 176: 363-366

Chiasson H, Belanger A, Bostanian N et al. Acaricidal properties of *Artemisia absinthium* and *Tanacetum vulgare* (Asteraceae) essential oils obtained by three methods of extraction. *J Econ Entomol* 2001, 94(1): 167-171

Cobb S . A case of epilepsy with a general discussion of pathology. *Med Clin N Am* 1922, 5: 1403-1420

Council Directive of 22 June 1988 on the approximation of the laws of the Member States relating to flavourings for use in foodstuffs and to source materials for their production (88/388/EEC) (OJ L 184, 15.7.198, p. 61-66)

Deutscher Arzneimittel-Codex (DAC). Ergänzungsbuch zum Arzneibuch. Hrsg. ABDA – Bundesvereinigung Deutscher Apothekerverbände. Eschborn; Govi-Verlag Pharmazeutischer Verlag GmbH and Stuttgart; Deutscher Apotheker Verlag, 2007

Del Castillo J, Anderson M, Rubottom GM. Marijuana, absinthe and the central nervous system. *Nature* 1975, 253(5490): 365-366

Dettling A, Grass H, Schuff A et al. Absinthe: attention performance and mood under the influence of thujone. *J Stud Alcohol* 2004, 65: 573-581

EI Makrini NI, Hassam B. Artemisia absinthium: burning plant! *Pan Afr Med J* 2016, 23:10

EMA/HMPC/732886/2010. Public statement on the use of herbal medicinal products containing thujone. Committee on Herbal Medicinal Products (HMPC)

European Pharmacopoeia 6th ed. Wormwood – Absinthii herba. Council of Europe 01/2008:1380

Firouzi R, Azadbakht M, Nabinedjad A. Anti-listerial activity of essential oils of some plants. *J Appl Anim Res* 1998, 14: 75-80

De Freitas MV, Netto Rde C, da Costa Huss JC et al. Influence of aqueous crude extracts of medicinal plants on the osmotic stability of human erythrocytes. *Toxicol In Vitro* 2008, 22: 219-224

Gambelunghe C, Melai P. Absinthe: enjoying a new popularity among young people? *Forens Sci Int* 2002, 130: 183-186

Gebhardt R. Stimulation of acid secretion by extracts of *Gentiana lutea* L. in cultured cells from rat gastric mucosa. *Pharm & Pharmacol Lett* 1997, 7(2-3): 106-108

- Ghédira K, Goetz P. *Artemisia absinthium* L.: absinthe (Asteraceae) *Phytothérapie* 2016, 14:125-129
- German Pharmacopoeia (1872). Berlin, Verlag der königlichen Geheimen Ober-Hofdruckerei
- Gessner O. Gift- und Arzneipflanzen von Mitteleuropa. Hrsg. G. Orzechowski, 3. Aufl., Heidelberg, Carl-Winter-Universitätsverlag. 1974
- Gilani AH, Janbaz KH. Preventive and curative effects of *Artemisia absinthium* on acetaminophen and CCl₄-induced hepatotoxicity. *Gen Pharmac* 1995, 26(2): 309-315
- Guarrera PM. Traditional phytotherapy in Central Italy (Marche, Abruzzo, and Latium). *Fitoterapia* 2005, 76: 1-25
- Haffner & Schultz. Normdosen der gebräuchlichen Arzneimittel. Hrsg. Schmid W & Schultz O-E, 5. Aufl., Stuttgart, Wissenschaftliche Verlagsgesellschaft mbH, 1969
- Hänsel R, Sticher O. Pharmakognosie – Phytopharmazie. 8. Aufl.; Heidelberg; Springer Medizin Verlag, 2007
- Hausen BM, Vieluf IK. Allergiepflanzen – Pflanzenallergene. 2. Aufl., Landsberg/München, ecomed Verlagsgesellschaft mbH, 1997
- Hernandez H, Mendiola J, Torres D et al. Effect of aqueous extracts of *Artemisia* on the in vitro culture of *Plasmodium falciparum*. *Fitoterapia* 1990, 61(6): 540-541
- Höld KM, Sirisoma NS, Ikeda T et al. α-Thujone (the active component of absinthe): γ-Aminobutyric acid type A receptor modulation and metabolic detoxification. *PNAS* 2000a, 97(8): 3826-3831
- Höld KM, Sirisoma NS, Sparks SE et al. (2000b). α-Thujone and absinthe: structural and metabolic aspects of neurotoxic action. TOXI 023, 219th ACS National meeting, San Francisco, CA, 26-30th March 2000
- Höld KM, Sirisoma NS, Casida JE. Detoxification of α- and β-thujones (the active ingredients of absinthe): site specificity and species differences in cytochrome P450 oxidation in vitro and in vivo. *Chem Res Toxicol* 2001, 14: 589-595
- Hose S. Der Wermut – *Artemisia absinthium* L. Arzneipflanze für Kranke und "Kultige". *Z Phytother* 2002, 23: 187-194
- Ikram M, Shafi N, Mir I et al. 24zeta-Ethylcholesta-7,22-Dien-3beta-ol: a possibly antipyretic constituent of *Artemisia absinthium*. *Planta Med* 1987, 53(4): 389
- Ishida T, Toyota M, Asakawa Y. Terpenoid biotransformation in mammals. V. Metabolism of (+)-citronellal, (±)-7-hydroxycitronellal, citral, (-)-perillaldehyde, (-)-myrtenal, cuminaldehyde, thujone and (±)-carvone in rabbits. *Xenobiotica* 1989, 19(8): 843-855
- Jaenson TGT, Palsson K, Borg-Karlson AK. Evaluation of extracts and oils of tick-repellent plants from Sweden. *Med Veter Entomol* 2005, 19: 345-352
- Jagusch M. Die Pankreasenzymstimulierende Wirkung von *Artemisia absinthium* im Vergleich zur Wirkung einer alkoholischen Placeboslösung. Inaugural-Dissertation Bayerische Julius-Maximilians-Universität, Würzburg, 1988
- Jiang Y, He X, Ortiz de Montellano PR. Radical intermediates in the catalytic oxidation of hydrocarbons by bacterial and human cytochrome P450 enzymes. *Biochem* 2006, 45(2): 533-542
- Juteau F, Jerkovic I, Masotti V et al. Composition and Antimicrobial Activity of the Essential Oil of *Artemisia absinthium* from Croatia and France. *Planta Med* 2003, 69: 158-161

Kalemba JG, Kurowska A, Majda T et al. Analysis of essential oils in aspects of their influence to insects. *Zeszyty Nauk Politechnik Lodz* 1993, 589: 5-14

Karwowska K, Stegman J, Duszkiewicz-Reinhard W et al. Studies on isolation and chemical composition of biologically active compounds of wormwood (*Artemisia absinthium*). Part II. *Ann Warsaw Agricult Univ – SGGW Horticult* 1997, 18:105-108

Kaul VK, Nigam SS, Dhar KL . Antimicrobial activities of the essential oils of *Artemisia absinthium* Linn, *Artemisia vestita* Wall, and *Artemisia vulgaris* Linn. *Ind J Pharm* 1976, 38(1): 21-22

Kaul VK, Nigam SS, Banerjee AK. Insecticidal activity of some essential oils. *Ind J Pharm* 1978, 40(1):22

Khattak SG, Gilani SN, Ikram M. Antipyretic studies on some indigenous Pakistani medicinal plants. *J Ethnopharmacol* 1985, 14:45-51

Kistler B. Die galletreibende Wirkung der Einzeldroge Wermut gegenüber Placebo bei intraduodenaler Applikation am Menschen im Doppelblindversuch. Inaugural-Dissertation Bayerische Julius-Maximilians-Universität, Würzburg, 1988

Kocaoglu C, Ozel A. Persistent metabolic acidosis and severe diarrhoea due to *Artemisia absinthium* poisoning. *Pak Med Assoc* 2014, 64(9):1081-1083

Kordali S, Kotan R, Mavi A et al. Determination of the chemical composition and antioxidant activity of the essential oil of *Artemisia dranunculus* and of the antifungal and antibacterial activities of Turkish *Artemisia absinthium*, *A. dranunculus*, *A. santonicum*, and *Artemisia spicigera* essential oils. *J Agric Food Chem* 2005, 53: 9452-9458

Kordali S, Aslan I, Çalmaşur O et al. Toxicity of essential oils isolated from three *Artemisia* species and some of their major components to granary weevil, *Sitophilus granaries* (L.) (Coleoptera: Curculionidae). *Ind Crops Prod* 2006, 23: 162-170

Krebs S, Omer TN, Omer B. Wormwood (*Artemisia absinthium*) suppresses tumour necrosis factor alpha and accelerates healing in patients with Crohn's disease – a controlled clinical trial. *Phytomed* 2010, 5: 305-309

Kreitmair H. *Artemisia absinthium* L. – der echte Wermut. *Pharmazie* 1951, 6: 27-28

Kröner LU, Padosch SA, Lachenmeier DW et al. Untersuchungen zur forensischen Relevanz thujonhaltiger Spirituosen unter besonderer Berücksichtigung toxikologisch-analytischer Aspekte. *Blutalkohol* 2005, 42: 263-271

Lee HG, Kim H, Oh WK et al. Tetramethoxy Hydroxyflavone p7F downregulates inflammatory mediators via the inhibition of Nuclear Faktor κB. *Ann NY Acad Sci* 2004, 1030: 555-568

Lewin L. Gifte und Vergiftungen. 4. Aufl., Berlin, Verlag von Georg Stilke, 1929

Li Y, Ohizumi Y. Search for constituents with neurotrophic factor-potentiating activity from the medicinal plants of Paraguay and Thailand. *Yakugaku Zasshi* 2004, 124(7): 417-424

Madaus G (1976). Lehrbuch der biologischen Heilmittel. Hildesheim, Georg Olms Verlag (reprint of Madaus G. (1938) Lehrbuch der biologischen Heilmittel. Leipzig, Georg Thieme Verlag)

Margaria R . Acute and sub-acute toxicity study on thujone. Unpublished report of Istituto di Fisiologia, Università di Milano (cited from the SCF (2002) Opinion of the Scientific Committee on Food on Thujone, 1963 (available at http://ec.europa.eu/food/fs/sc/scf/out162_en.pdf)

Martindale. The Extra Pharmacopoeia. 29th edition. The Pharmaceutical Press London, 1989

Mendiola J, Bosa M, Perez N et al. Extracts of Artemisia abrotanum and Artemisia absinthium inhibit growth of Naegleria fowleri in vitro. *Transact Royal Soc Trop Med Hyg* 1991, 85:78-79

Meschler JP, Howlett AC. Thujone exhibits low affinity for cannabinoid receptors but fails to evoke cannabimimetic responses. *Pharmacol Biochem Behav* 1999, 62(3):473-480

Milett Y, Jouglard J, Steinmetz MD et al. Toxicity of some essential plant oils. Clinical and experimental study. *Clin Toxicol* 1981, 18(12): 1485-1498

Muto T, Watanabe T, Okamura M, et al. Thirteen-week repeated dose toxicity study of wormwood (Artemisia absinthium) extract in rats. *J Toxicol Sci* 2003, 28(5): 471-478

Ng SC, Lam YT, Tsoi KKF, et al. Systematic review: the efficacy of herbal therapy in inflammatory bowel disease. *Aliment Pharmacol Ther* 2013, 38:854-863

Niesel S. Untersuchungen zum Freisetzungerverhalten und zur Stabilität ausgewählter wertbestimmender Pflanzeninhaltsstoffe unter besonderer Berücksichtigung moderner phytochemischer Analysenverfahren. Inaugural-Dissertation Freie Universität, Berlin, 1992

National Toxicology Program (NTP). Testing status of alpha-thujone and alpha/beta-thujone. 2003 (available at http://ntp-apps.niehs.nih.gov/ntp_tox/index.cfm) National Toxicology Program (NTP). Technical Report No. 570: Toxicology and Carcinogenesis Studies of alpha,beta-Thujone (CAS No. 76231-76-0) in F344/N Rats and B6C3F1 Mice (Gavage studies), 2010 (available at <http://ntp.niehs.nih.gov/go/36136>)

Omer B, Krebs S, Omer H et al. Steroid-sparing effect of wormwood (Artemisia absinthium) in Crohn's Disease: A double-blind placebo-controlled study. *Phytomed* 2007, 14: 87-95

Opdyke DL. Monographs on fragrance raw materials. *Food Cosmet Toxicol* 1975, 13: 683-923

Pinto-Sconamiglio W . Effetti del tujone sull'attività spontanea e sul comportamento condizionato del ratto. *Bull Chim Farm* 1967, 107: 780-791

Qureshi RA, Ahmad M, Yousaf Z et al. Taxonomic study and medicinal importance of three species of the genus Artemisia LINN. *Pakistan J Forest* 2002, 52(1): 57-64

Ramos A, Visozo A, Piloto J et al. Screening of antimutagenicity via antioxidant activity in Cuban medicinal plants. *J Ethnopharmacol* 2003, 87: 241-246

Rao VSN, Menezes AMS, Gadelha MGT. Antifertility screening of some indigenous plants of Brasil. *Fitoterapia* 1987, 59(1): 17-20

Rice KC, Wilson RS. (-)-3-Isothujone, a small non-nitrogenous molecule with antinociceptive activity in mice. *J Med Chem* 1976, 19(8): 1054-1057

Roth L, Daunderer M, Kormann K. Giftpflanzen – Pflanzengifte. 4. Aufl., Hamburg, Nikol Verlagsgesellschaft mbH & Co KG, 1994

Rozengurt E. Taste receptors in the gastrointestinal tract. I. Bitter taste receptors and α-gustducin in the mammalian gut. *Am J Physiol Gastroint Liver Physiol* 2006, 291: G171-177

SCF (2002). Opinion of the Scientific Committee on Food on Thujone (available at: http://ec.europa.eu/food/fs/sc/scf/out162_en.pdf)

Schulz V, Hänsel R. Rationale Phytotherapie. 5. Aufl., Berlin Heidelberg New York, Springer Verlag, 2004

Shafi N, Khan GA, Ghauri EG. Antiulcer effect of *Artemisia absinthium* L. in rats. *Pak J Sci Ind Res* 2004, 47(2): 130-134

Stanković N, Mihajlov-Krstev T, Zlatković B, et al. Antibacterial and Antioxidant Activity of Traditional Medicinal Plants from the Balkan Peninsula. *NJAS - Wageningen J Life Sci* 2016a, 78:21-28

Stanković N, Mihajlov-Krstev T, Zlatković B, et al. Comparative Study of Composition, Antioxidant, and Antimicrobial Activities of Essential Oils of Selected Aromatic Plants from Balkan Peninsula. *Planta Med* 2016b, 82:650-661

Sternini C. Taste receptors in the gastrointestinal tract. IV. Functional implications of bitter taste receptors in gastrointestinal chemosensing. *Am J Physiol Gastrointest Liver Physiol* 2007, 292(2): G457-61.

Surber W. Etude de toxicité sous-chronique de la thujone sur rats. Rapport Final. Institut Battelle, Geneve, 1962

Tahir M, Siddiqui MMH, Khan AB. Effect of Afsanteen (*Artemisia absinthium* Linn.) in acute intestinal amoebiasis. *Hamdrad Med* 1997, 40(3): 24-27

Tegtmeier M, Harnischfeger G. Methods for the reduction of thujone content in pharmaceutical preparations of *Artemisia*, *Salvia* and *Thuja*. *Eur J Pharm Biopharm* 1994, 40(5): 337-340

Teuscher E. Pharmakognosie. 3. Aufl., Berlin, Akademie-Verlag, 1989

Tosi B, Bonora A, Dall'Olio G et al. (1991). Screening for toxic thiophene compounds from crude drugs of the family Compositae used in Northern Italy. *Phytother Res* 5(2): 59-62

Wake G, Court J, Pickering A et al. CNS acetylcholine receptor activity in European medicinal plants traditionally used to improve failing memory. *J Ethnopharmacol* 2000, 69: 105-114

Wegener T . Anwendung eines Trockenextraktes aus *Gentiana lutea* radix bei dyspeptischem Symptomenkomplex. *Z Phytother* 1998, 19: 163-164

Weisbord SD, Soule JB, Kimmel PL . Poison on line – acute renal failure caused by oil of wormwood purchased through the internet. *New Eng Med* 1997, 337:825-527

Wichtl M . Teedrogen und Phytotherapeutika. 4. Aufl., Stuttgart, Wissenschaftliche Verlagsgesellschaft mbH, 2002Zafar MM, Hamdard ME, Hameed A. Screening of *Artemisia absinthium* for antimalarial effects on *Plasmodium berghei* in mice: a preliminary report. *J Ethnopharmacol* 1990, 30: 223-226