



London, 4 September 2008
Doc. Ref. EMEA/HMPC/260098/2007

COMMITTEE ON HERBAL MEDICINAL PRODUCTS (HMPC)

LIST OF REFERENCES SUPPORTING THE ASSESSMENT REPORT ON:

Avenae herba and Avenae fructus
***Avena sativa* L.**
(oat herb and oat fruit)

The EMEA acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Alexandrescu DT., Vaillant JG., Dasanu CA. Effect of treatment with a colloidal oatmeal lotion on the acneform eruption induced by epidermal growth factor receptor and multiple tyrosine-kinase inhibitors. *Clinical and Experimental Dermatology* 2007; 32: 71-4.

Anand CL. Effects of *Avena sativa* on Cigarette smoking. *Nature* 1971; 233: 496.

Anonymus (1944) United States Patent Office; Patent 2,355,028 Oat product patented Aug. 1, 1944.

Anonymous. *The British Herbal Pharmacopoeia*, London, 1976; 27.

Anonymus. *The US Pharmacopoeia* 30: Colloidal Oatmeal. Monograph, 1990.

Anonymous (2007). http://www.e-gezondheid.be/nl/gezondheid_tijdschrift/gezondheid_homeopathie_per_product/Avena_sativa_complex-10837-949-art.htm

Anonymus (2008) <http://www.vitaliashop.de/Gesundheit/Aufbau/Hafer-Heilpflanzensaft::800.html>

Aries MF., Vaissiere C., Fabre B., Charveron M., Gall Y. Immunomodulatory activity of *Avena rhealba* : interest in skin inflammatory disorders. *J. Invest. Dermatol.* 1999a; 113 (3): 329.

Aries MF., Vassiere C., Ceruh L., Chaveron M., Gall Y. *Avena rhealba* activity in cutaneous wound healing process. *J. Invest. Dermatol.* 1999b; 112 (1): 132.

Aries MF., Vaissiere C., Giraud E., Fabre B., Charveron M., Gall Y. Continuous dermogenic inflammation : induction of endothelial NO synthase (eNOS) expression by substance P, modulation by *Avena rhealba*. *J. Invest. Dermatol.* 2000; 115 (3): Annual Congress of the Society for Dermatological Research (Lyon 17-19 May 2000): abstract not submitted.

Aries MF., Vaissiere C., Fabre B., Charveron M., Gall Y. *Avena rhealba* inhibits arachidonic acid cascade, CPLA2 and COX expression in human keratinocytes. Interest in cutaneous inflammatory disorders. *J. Invest. Dermatol.* 2003; 121 (1): JID abstract 0046.

Aries MF., Vaissiere C., Pinelli E., Pipy B., Charveron M. Avena Rhealba® inhibits A23187-stimulated arachidonic acid mobilisation, eicosanoid release, and cPLA2 expression in human keratinocytes : potential in cutaneous inflammatory processes. *Biol. Pharm. Bull.* 2005; 28(4): 601-6.

Barnes PJ., Taylor PW. Gamma tocopherol in barley germ. *Phytochemistry* 1979; 20 (7): 1753-4.

Berg A., Konig D., Deibert P., Grathwohl D., Baumstark MW., Franz IW. Effect of an oat bran enriched diet on the atherogenic lipid profile in patients with an increased coronary heart disease risk. A controlled randomized lifestyle intervention study. *Ann. Nutr. Metab.* 2003; 47: 306-11.

Bognar A. Determination of vitamin A in food using HPLC: results of the collaborative study of the working group 'vitamin analysis' within the framework of §35 LMBG. *Z. Lebensmittel-Unters. Forschung* 1986; 182 (6): 492-7.

Boisnic S., Branchet-Gumila MC., Coutanceau C. Inhibitory effect of oatmeal extract oligomer on vasoactive intestinal peptide-induced inflammation in surviving skin. *Int. J. Tissue React.* 2003; 25: 41-6.

Boisnic S., Branchet MC., Ermosilla V. Healing effect of a spray containing Rhealba oat colloidal extract in an in vitro reconstitution model of skin. *Int. J. Tissue Reactions* 2005; 27: 83-9.

Brinegar AC., Peterson DM. Separation and characterisation of oat globulin polypeptides. *Arch. Biochem. Biophysics* 1982; 219: 71-9.

Bringelsson S., Dimberg LH., Kamal-Eldin A. Effects of commercial processing on levels of antioxydants in oats (*Avena sativa* L.). *J. Agric. Food Chem.* 2002; 50: 1890-6.

Bratt K., Sunnerheim K., Bryngelsson S. et al. Avenanthramides in oats (*Avena sativa* L.) and structure-antioxydant activity relationships. *J. Agric. Food Chem.* 2003; 51: 594-600.

Bye C., Fowle ASE., Letley E., Wilkinson S. Lack of effect of *Avena sativa* on cigarette smoking. *Nature* 1974; 252: 580-1.

Camplone G., Arcangeli F., Bonifazi E., Gelmetti C., Menni S., Mulas P., Paradisi M., Patrizi A. The use of colloidal oatmeal products in the care of children with mild atopic dermatitis. *Eur. J. Pediatric Dermatology* 2004; 14: 157-60.

Chen C-Y., Milbury P., Kwak H-K., Collins W., Samuel P., Blumberg JB. Avenanthramides and phenolic acids from oats are bioavailable and act synergistically with vitamin C to enhance hamster and human LDL resistance to oxidation. *J. Nutr.* 2004; 134: 1459-66.

Czerwinski J., Bartnikowska E., Leontowicz H. et al. Oat (*Avena sativa* L.) and amaranth (*Amaranthus hypochondriacus*) meals positively affect lipid profile in rats fed cholesterol-containing diets. *J. Nutritional Biochem.* 2004; 15: 622-9.

Daniels DG., Martin HF. Antioxydants in oats: mono-esters of caffeic and ferulic acids. *J. Sci. Fd. Agric.* 1967; 18: 589-95.

Delaney B., Nicolosi RJ., Wilson TA. et al. Beta-glucan fractions from barley and oats are similarly antiatherogenic in hypercholesterolemic Syrian golden hamsters. *J. Nutr.* 2003; 133: 468-95.

Dick LA. Colloidal emollient baths in pediatric dermatoses. *Arch. of Pediatrics* 1958; 75: 506-8.

Dodoneus Rembertus. Cruydt-Boeck. Plantijn-Moretus, Antwerpen, 1608; 898.

Dor R., Shanahan DJ. Oats and coeliac disease. *Gut* 2002; 51: 755-8.

Drisch P. Studie zur Senkung erhöhter Harnsäurespiegel mit einem Phytotherapeutikum. *Natura-med* 1988; 12 Sonderndruck.

El Bakali AE., Kühne EA., Grosshans E., Manciet JR., Pons-Guiraud A., Morel P., Dubertret L., Dargassies J., Dupuy P. Potentiel allergénique des extraits d'avoine dans la dermatite atopique. Document Pierre Fabre October 2000.

Fabre B. *Avena sativa*, demande d'inscription en usage topique. Pierre Fabre Innovation Développement. Concept paper 2004.

Franks AG. Dermatological uses of baths. *The Am. Practitioner and Digest of Treatments* 1958; 12: 1998-2000.

Frølich W., Nyman M. Minerals, phytate and dietary fibre in different fractions of oat-grain. *J. Cereal Science* 1988; 7: 73-82.

Grimalt R., Mengeaud V., Cambazard F. The steroid-sparing effect of an emollient therapy in infants with atopic dermatitis: a randomized controlled study. *Dermatology* 2007; 214: 61-7.

Hänsel R., Keller K., Rimpler H., Schneider G. Hager's Handbuch der Pharmazeutischen Praxis. Drogen A-D. Springer-Verlag, Berlin 1992: 437-46.

Heinoven M., Ollilainen V., Linkola E., Varo P., Koivistoinen P. Carotenoids and retinoids in Finnish foods: cereal and bakery products. *Cereal Chem.* 1989; 66 (4): 270-3.

Högborg I., Laurin P., Fälth-Magnusson K. Et al. Oats to children with newly diagnosed coeliac disease: a randomized double blind study. *Gut* 2004; 53: 649-54.

Jahn-Deesbach W. New studies on the thiamin (vitamin B1) content of grain. *Getreide, Mehl, Brot* 1979; 33 (10): 258-64.

Janatuinen EK., Kemppainen TA., Julkunen RJK. et al. No harm from five year ingestion of oats in coeliac disease. *Gut* 2002; 50: 332-5.

Karmally W., Montez MG., Palmas W. et al. Cholesterol-lowering benefits of oat-containing cereal in Hispanic Americans. *J. Am. Diet. Assoc.* 2005; 105: 967-70.

Katz DL., Nawaz H., Boukhalil et al. Effects of oat and wheat cereals on endothelial response. *Preventive Medicine* 2001; 33: 476-86.

Katz DL., Evans MA., Chan W. et al. Oats, antioxydants and endothelial function in overweight, dyslipidemic adults. *J. Am. Coll. Nutr.* 2004; 23: 397-403.

Kerckhoffs DAJM., Hornstra G., Mensink RP. Cholesterol-lowering effect of beta-glucan from oat bran in mildly hypercholesterolemic subjects may decrease when beta-glucan is incorporated into bread and cookies. *Am. J. Clin. Nutr.* 2003; 78: 221-7.

Krug E. Senkung des Harnsäurespiegels durch ein kombiniertes Phytotherapeutikum. *Acta Medica Empirica* 1985; 34 (6): 407-9.

Lasztity R. Oats proteins. I. Introduction and structure of the kernels. In: *The chemistry of cereal proteins*. CRC Press Inc. Boca Raton 1984, pp. 157-64.

Leclerc H. *Le Livre des Plantes Médicinales et Vénéneuses de France*. Ed. P. Lechevalier, Paris, 1947 ; 178-9.

- Leone F. Treatment of diaper rash. *The Connecticut State Medical Journal* 1957; 9: 815.
- List P.H. & Hörhammer L. *Hagers handbuch der Pharmazeutischen Praxis*. Springer Verlag, Berlin 1972 ; 342-4.
- Liu L., Zubik L., Collins FW., Marko M., Meydani M. The antiatherogenic potential of oat phenolic compounds. *Atherosclerosis* 2004; 175: 39-49.
- Lundin KEA., Nilsen EM., Scott HG. Et al. Oats induced villous atrophy in coeliac disease. *Gut* 2003; 52: 1649-52.
- Madaus G. *Lehrbuch der biologischen Heilmittel*. Georg Thieme Verlag , Leipzig 1938.
- Martinez-Tome M., Murcia MA., Frega N. et al. Evaluation of antioxidant capacity of cereal brans. *J. Agric. Food Chem.* 2004; 52: 4690-9.
- Matheson JD., Clauton J., Muller MJ. The reduction of itch during burn wound healing. *J. Burn Care & Rehabilitation* 2001; 22: 76-81.
- Mizuno A. Clinical evaluation of A-Derma Exomega Cream for atopic dry skin. *Skin Research* 2005; 4: 581-7.
- MonteBovi AJ. The colloidal demulcents. I. Physical and chemical properties. *Am. J. Pharmacy* 1954; 126: 1.
- Nie L., Wise ML., Peterson DM., Meydani M. Avenanthramide, a polyphenol from oats, inhibits vascular smooth muscle cell proliferation and enhances nitric oxide production. *Atherosclerosis* 2006; 186 (2): 260-6.
- Pacifico A., De Angelis L., Fargnoli MC., De Felice C., Chimenti S., Peris K. Clinical trial on Aveeno skin relief moisturizing lotion in patients with itching accompanied by skin lesions and xerosis. *J. Applied Research* 2005; 5: 325-30.
- Peraaho M., Kaukinen E., Mustalahti K. Et al. Effect of an oats-containing gluten-free diet on symptoms and quality of life in coeliac disease. A randomized study. *Scand. J. Gastroenterol.* 2004 ; 39 : 27-31.
- Pins JJ., Geleva D., Keenan JM., Frazel C., O'Connor PJ., Cherney LM. Do whole-grain oat cereals reduce the need for antihypertensive medications and improve blood pressure control? *J. Fam. Pract.* 2002; 51: 353-9.
- Rancé F., Dargassies J., Dupuy P., Schmitt AM., Guérin L., Dutau G. Faut-il contre-indiquer l'utilisation des emollients à base d'avoine chez l'enfant atopique? *Rev. Fr. Allergol. Immunol. Clin.* 2001 ; 41 : 477-83.
- Robitaille J., Fontaine-Bisson B., Couture P., Tchernof A., Vohl MC. Effect of an oat bran-rich supplement on the metabolic profile of overweight premenopausal women. *Ann. Nutr. Metab.* 2005; 49: 141-8.
- Saltzman E., Das SK., Lichtenstein AH., et al. An oat-containing hypocaloric diet reduces systolic blood pressure and improves lipid profile beyond effects of weight loss in men and women. *J. Nutr.* 2001; 131: 1465-70.
- Schneider E. *Avena sativa – Hafer als Heilpflanze* 1985; 6 (5): 165-7.
- Schneider E. Lösliche Silikate im grünen Hafer. *Zeitschrift für Phytotherapie* 1990: 129-32.

Smith GC. The treatment of various dermatoses associated with dry skin. J. of the South Carolina Medical Association 1958; LIV: 8.

Storsrud S., Hulthen LR., Lenner RA. Beneficial effects of oats in the gluten free diet of adults with special references to nutrient status, symptoms and subjective experiences. Br. J. Nutr. 2003; 90: 101-7.

Thompson T. Oats and the gluten-free diet. J. Am. Diet Assoc. 2003; 103: 376-9.

Torphy TJ., Zhou HL., Cieslinski LB. Stimulation of beta adrenoceptors in a human monocyte cell line (U937) up-regulates cyclic AMP-specific phosphodiesterase activity. J. Pharmacol. Exp. Ther. 1992; 263: 1195-1205.

van Elteren G. (2008) <http://plantaardigheden.nl/plant/beschr/gonnve/haver.htm> (last access 11 June 2008)

Van Hellemont J. Fytotherapeutisch Compendium. Belgian Pharmaceutical Society, Brussels 1985; 90.

Varjonen E., Vainio E., Kalimo K., Juntunen-Backman K., Savolainen J. Skin-prick test and RAST responses to cereals in children with atopic dermatitis. Characterization of IgE-binding components in wheat and oats by an immunoblotting method. Clinical and Experimental Allergy 1995; 25: 1100-7.

Vié K., Cours-Darne S., Vienne M., Boyer F., Fabre B., Dupuy P. Modulating effects of oatmeal extracts in the sodium lauryl sulfate skin irritancy model. Skin Pharmacol. Appl. Skin Physiology 2002; 15: 120-4.

Wenzig E., Kunert O., Ferreira D., Schmid M., Schuhly W., Bauer R., Hiermann A. Flavonolignans from *Avena sativa*. J. Nat. Prod. 2005; 68: 289-92.

Yun C-H., Estrada A., Van Kessel A., Park B-C., Laarveld B. B-Glucan, extracted from oat, enhances disease resistance against bacterial and parasitic infections. FEMS Immunology and Medical Microbiology 2003; 35: 67-75.

Zhou M., Panchuk-Voloshina N. A one-step fluorometric method for the continuous measurement of monoamine oxidase activity. Anal. Biochem. 1997; 253: 169-174.