



**This document was valid from 16 July 2009 until May 2016.
It is now superseded by a [new version](#) adopted by the HMPC on
30 May 2017 and published on the EMA website.**

COMMITTEE ON HERBAL MEDICINAL PRODUCTS (HMPC)

FINAL

LIST OF REFERENCES SUPPORTING THE ASSESSMENT REPORT ON

**Absinthii herba
Artemisia absinthium L., herba
(wormwood)**

The EMEA acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

1. AB-DDR (1988). Pharmacopoeia of the German Democratic Republic. Berlin, Akademie-Verlag
2. Ariño A, Arberas I, Renobales G et al. (1999a). Seasonal variations in wormwood (*Artemisia absinthium* L.) essential oil composition. J Essent Oil Res 11: 619-622
3. Ariño A, Arberas I, Renobales G et al. (1999b). Essential oil of *Artemisia absinthium* L. from the Spanish Pyrenees. J Essent Oil Res 11: 182-184
4. Ariño A, Arberas I, Renobales G et al. (1999c). Influence of extraction method and storage conditions on the volatile oil of wormwood (*Artemisia absinthium* L.). Eur Food Res Technol 209(2): 126-129
5. Arnold WN (1989). Absinthe. Sci Am 260(6): 112-117
6. Baumann IC (1975). Über die Wirkung von *Chelidonium*, *Curcuma*, Absinth und *Carduus marianus* auf die Galle- und Pankreassekretion bei Hepatopathien. Med Mschr 29:173-180
7. Baumann IC, Glatzel H, Muth HW (1975). Untersuchungen der Wirkung von Wermut (*Artemisia absinthium* L.) auf die Gallen- und Pankreassaft-Sekretion des Menschen. Z Allgemeinmed 51(17): 784-791
8. Bayerl C & Jung EG (1996). Allergic contact stomatitis from aristochol, a phytotherapeutic cholagogue. Contact Derm 34(3): 222-223
9. Bielenberg J (2002). Die "Grüne Fee". ÖAZ Aktuell, issue 12/2002; www.oecz.at

10. Blumberger W & Glatzel H (1966). Über die Speichelwirksamkeit von Bitterstoffen. *Planta Med* 14: 52-60
11. Bonkovsky HL, Cable EE, Cable JW et al. (1992). Porphyrigenic properties of the terpenes, camphor, pinene and thujone (With a note on historic implications for absinthe and the illness of Vincent van Gogh). *Biochem Pharmacol* 43: 2359-2368
12. British Herbal Pharmacopoeia. British Herbal Medicine Association, Bournemouth 1983
13. Canadanovic-Brunet JM, Djilas SM, Cetkovic GS et al. (2005). Free radical scavenging activity of wormwood (*Artemisia absinthium* L.) extracts. *J Sci Food Agric* 85: 265-272
14. Carnat AP, Madesclaire M, Chavignon O et al. (1992). Cis-Chrysanthenol, a main component in essential oil of *Artemisia absinthium* L. growing in Auvergne (Massif Central), France. *J Essent Oil Res* 4: 487-490
15. Centini F & Laurini GP (1987). Una intossicazione da olio di salvia. *Zacchia* 60: 263-274
16. Chialva F, Liddle PAP, Godlia G (1983). Chemotaxonomy of wormwood (*Artemisia absinthium* L.). (I. Composition of the essential oil of several chemotypes). *Z Lebensm Unters Forsch* 176: 363-366
17. Chiasson H, Belanger A, Bostanian N et al. (2001). Acaricidal properties of *Artemisia absinthium* and *Tanacetum vulgare* (Asteraceae) essential oils obtained by three methods of extraction. *J Econ Entomol* 94(1): 167-171
18. Cobb S (1922). A case of epilepsy with a general discussion of pathology. *Med Clin N Am* 5: 1403-1420
19. Council Directive of 22 June 1988 on the approximation of the laws of the Member States relating to flavourings for use in foodstuffs and to source materials for their production (88/388/EEC) (OJ L 184, 15.7.198, p. 61-66)
20. Council of Europe (2007). Natural source of flavourings. Report No. 2. Council of Europe Publishing, Strasbourg
21. DAC (2007). Deutscher Arzneimittel-Codex. Ergänzungsbuch zum Arzneibuch. Hrsg. ABDA – Bundesvereinigung Deutscher Apothekerverbände. Eschborn; Govi-Verlag Pharmazeutischer Verlag GmbH and Stuttgart; Deutscher Apotheker Verlag
22. Del Castillo J, Anderson M, Rubottom GM (1975). Marijuana, absinthe and the central nervous system. *Nature* 253(5490): 365-366
23. Dettling A, Grass H, Schuff A et al. (2004). Absinthe: attention performance and mood under the influence of thujone. *J Stud Alcohol* 65: 573-581
24. European Pharmacopoeia 6th ed. Wormwood – Absinthii herba. Council of Europe 01/2008:1380
25. Firouzi R, Azadbakht M, Nabinedjad A (1998). Anti-listerial activity of essential oils of some plants. *J Appl Anim Res* 14: 75-80
26. de Freitas MV, Netto Rde C, da Costa Huss JC et al. (2008). Influence of aqueous crude extracts of medicinal plants on the osmotic stability of human erythrocytes. *Toxicol In Vitro* 22: 219-224

27. Gambelunghe C & Melai P (2002). Absinthe: enjoying a new popularity among young people? *Forens Sci Int* 130: 183-186
28. Gebhardt R (1997). Stimulation of acid secretion by extracts of *Gentiana lutea* L. in cultured cells from rat gastric mucosa. *Pharm & Pharmacol Lett* 7(2-3): 106-108
29. German Pharmacopoeia (1872). Berlin, Verlag der königlichen Geheimen Ober-Hofdruckerei
30. Gessner O (1974). Gift- und Arzneipflanzen von Mitteleuropa. Hrsg. G. Orzechowski, 3. Aufl., Heidelberg, Carl-Winter-Universitätsverlag
31. Gilani AH & Janbaz KH (1995). Preventive and curative effects of *Artemisia absinthium* on acetaminophen and CCl₄-induced hepatotoxicity. *Gen Pharmac* 26(2): 309-315
32. Guarrera PM (2005). Traditional phytotherapy in Central Italy (Marche, Abruzzo, and Latium). *Fitoterapia* 76: 1-25
33. Haffner & Schultz (1969). Normdosen der gebräuchlichen Arzneimittel. Hrsg. Schmid W & Schultz O-E, 5. Aufl., Stuttgart, Wissenschaftliche Verlagsgesellschaft mbH
34. Absinthii herba. In Blaschek W, Ebel S, Hackenthal E, Holzgrabe U, Keller K, Reichling J, Schulz V, editors (2006). *HagerROM 2006: Hagers Handbuch der Drogen und Arzneistoffe*. Springer Medizin Verlag, Heidelberg
35. Hänsel R & Sticher O (2007). *Pharmakognosie – Phytopharmazie*. 8. Aufl.; Heidelberg; Springer Medizin Verlag
36. Hausen BM & Vieluf IK (1997). *Allergiepflanzen – Pflanzenallergene*. 2. Aufl., Landsberg/München, ecomed Verlagsgesellschaft mbH
37. Hernandez H, Mendiola J, Torres D et al. (1990). Effect of aqueous extracts of *Artemisia* on the in vitro culture of *Plasmodium falciparum*. *Fitoterapia* 61(6): 540-541
38. Höld KM, Sirisoma NS, Ikeda T et al. (2000a). α -Thujone (the active component of absinthe): γ -Aminobutyric acid type A receptor modulation and metabolic detoxification. *PNAS* 97(8): 3826-3831
39. Höld KM, Sirisoma NS, Sparks SE et al. (2000b). α -Thujone and absinthe: structural and metabolic aspects of neurotoxic action. *TOXI 023*, 219th ACS National meeting, San Francisco, CA, 26-30th March 2000
40. Höld KM, Sirisoma NS, Casida JE (2001). Detoxification of α - and β -thujones (the active ingredients of absinthe): site specificity and species differences in cytochrome P450 oxidation in vitro and in vivo. *Chem Res Toxicol* 14: 589-595
41. Hose S (2002). Der Wermut – *Artemisia absinthium* L. Arzneipflanze für Kranke und “Kultige”. *Z Phytother* 23: 187-194
42. Ikram M, Shafi N, Mir I et al. (1987). 24zeta-Ethylcholesta-7,22-Dien-3beta-ol: a possibly antipyretic constituent of *Artemisia absinthium*. *Planta Med* 53(4): 389
43. Ishida T, Toyota M, Asakawa Y (1989). Terpenoid biotransformation in mammals. V. Metabolism of (+)-citronellal, (\pm)-7-hydroxycitronellal, citral, (-)-perillaldehyde, (-)-myrtenal, cuminaldehyde, thujone and (\pm)-carvone in rabbits. *Xenobiotica* 19(8): 843-855
44. Jaenson TGT, Palsson K, Borg-Karlson AK (2005). Evaluation of extracts and oils of tick-repellent plants from Sweden. *Med Veter Entomol* 19: 345-352

45. Jagusch M (1988). Die Pankreasenzymstimulierende Wirkung von *Artemisia absinthium* im Vergleich zur Wirkung einer alkoholischen Placebolösung. Inaugural-Dissertation Bayerische Julius-Maximilians-Universität, Würzburg
46. Juteau F, Jerkovic I, Masotti V et al. (2003). Composition and Antimicrobial Activity of the Essential Oil of *Artemisia absinthium* from Croatia and France. *Planta Med* 69: 158-161
47. Kalembe JG, Kurowska A, Majda T et al. (1993). Analysis of essential oils in aspects of their influence to insects. *Zeszyty. Nauk. Politechnik. Lodz.* 589: 5-14
48. Karwowska K, Stegman J, Duszkievicz-Reinhard W et al. (1997). Studies on isolation and chemical composition of biologically active compounds of wormwood (*Artemisia absinthium*). Part II. *Ann Warsaw Agricult Univ – SGGW Horticult.* 18:105-108
49. Kaul VK, Nigam SS, Dhar KL (1976). Antimicrobial activities of the essential oils of *Artemisia absinthium* Linn, *Artemisia vestita* Wall, and *Artemisia vulgaris* Linn. *Ind J Pharm* 38(1): 21-22
50. Kaul VK, Nigam SS, Banerjee AK (1978). Insecticidal activity of some essential oils. *Ind J Pharm* 40(1): 22
51. Keith HM & Stavrayk GW (1953). Experimental convulsions induced by administration of thujone. A pharmacologic study of the influence of the autonomic nervous system on these convulsions. *Arch Neurol Psychiatr* 34:1022
52. Khattak SG, Gilani SN, Ikram M (1985). Antipyretic studies on some indigenous Pakistani medicinal plants. *J Ethnopharmacol* 14: 45-51
53. Kim JO, Kim YS, Lee JH et al (1993). Antimutagenic effect of the major volatile compounds identified from mugwort (*Artemisia asiatica nakai*) leaves. *J Korean Soc Food Nutr* 21(3): 308-313
54. Kistler B (1988). Die galletreibende Wirkung der Einzeldroge Wermut gegenüber Placebo bei intraduodenaler Applikation am Menschen im Doppelblindversuch. Inaugural-Dissertation Bayerische Julius-Maximilians-Universität, Würzburg
55. Kordali S, Kotan R, Mavi A et al. (2005). Determination of the chemical composition and antioxidant activity of the essential oil of *Artemisia dranunculus* and of the antifungal and antibacterial activities of Turkish *Artemisia absinthium*, *A. dranunculus*, *A. santonicum*, and *Artemisia spicigera* essential oils. *J Agric Food Chem* 53: 9452-9458
56. Kordali S, Aslan I, Çalmaşur O et al. (2006). Toxicity of essential oils isolated from three *Artemisia* species and some of their major components to granary weevil, *Sitophilus granaries* (L.) (Coleoptera: Curculionidae). *Ind Crops Prod* 23: 162-170
57. Kreitmair H (1951). *Artemisia absinthium* L. – der echte Wermut. *Pharmazie* 6: 27-28
58. Kröner LU, Padosch SA, Lachenmeier DW et al. (2005). Untersuchungen zur forensischen Relevanz thujonhaltiger Spirituosen unter besonderer Berücksichtigung toxikologisch-analytischer Aspekte. *Blutalkohol* 42: 263-271
59. Lee HG, Kim H, Oh WK et al. (2004). Tetramethoxy Hydroxyflavone p7F downregulates inflammatory mediators via the inhibition of Nuclear Faktor κB. *Ann NY Acad Sci* 1030: 555-568
60. Lewin L (1929). *Gifte und Vergiftungen*. 4. Aufl., Berlin, Verlag von Georg Stilke

61. Li Y & Ohizumi Y (2004). Search for constituents with neurotrophic factor-potentiating activity from the medicinal plants of Paraguay and Thailand. *Yakugaku Zasshi* 124(7): 417-424
62. Madaus G (1976). *Lehrbuch der biologischen Heilmittel*. Hildesheim, Georg Olms Verlag (reprint of Madaus G. (1938) *Lehrbuch der biologischen Heilmittel*. Leipzig, Georg Thieme Verlag)
63. Margaria R (1963). Acute and sub-acute toxicity study on thujone. Unpublished report of the Instituto de Fisiologia, Universita di Milano (cited from the SCF (2002) Opinion of the Scientific Committee on Food on Thujone (http://ec.europa.eu/food/fs/sc/scf/out162_en.pdf))
64. Martindale. *The Extra Pharmacopoeia*. 29th edition. The Pharmaceutical Press London, 1989
65. Mendiola J, Bosa M, Perez N et al. (1991). Extracts of *Artemisia abrotanum* and *Artemisia absinthium* inhibit growth of *Naegleria fowleri* in vitro. *Transact Royal Soc Trop Med Hyg* 85: 78-79
66. Meschler JP & Howlett AC (1999). Thujone exhibits low affinity for cannabinoid receptors but fails to evoke cannabimimetic responses. *Pharmacol Biochem Behav* 62(3): 473-480
67. Milett Y, Jouglard J, Steinmetz MD et al. (1981). Toxicity of some essential plant oils. Clinical and experimental study. *Clin Toxicol* 18(12): 1485-1498
68. Muto T, Watanabe T, Okamura M et al. (2003). Thirteen-week repeated dose toxicity study of wormwood (*Artemisia absinthium*) extract in rats. *J Toxicol Sci* 28(5): 471-478
69. Niesel S (1992). Untersuchungen zum Freisetzungsverhalten und zur Stabilität ausgewählter wertbestimmender Pflanzeninhaltsstoffe unter besonderer Berücksichtigung moderner phytochemischer Analysenverfahren. Inaugural-Dissertation Freie Universität, Berlin
70. NTP (2003). National Toxicology Program. Testing status of alpha-thujone and alpha/beta-thujone. http://ntp-apps.niehs.nih.gov/ntp_tox/index.cfm
71. Omer B, Krebs S, Omer H et al. (2007). Steroid-sparing effect of wormwood (*Artemisia absinthium*) in Crohn's Disease: A double-blind placebo-controlled study. *Phytomedicine* 14: 87-95
72. Opdyke DL (1975). Monographs on fragrance raw materials. *Food Cosmet Toxicol* 13: 683-923
73. Pinto-Sconamiglio W (1967). Effetti del tujone sull'attività spontanea e sul comportamento condizionato del ratto. *Bull Chim Farm* 107: 780-791
74. Qureshi RA, Ahmad M, Yousaf Z et al. (2002). Taxonomic study and medicinal importance of three species of the genus *Artemisia* LINN. *Pakistan J Forest* 52(1): 57-64
75. Ramos A, Visozo A, Piloto J et al. (2003). Screening of antimutagenicity via antioxidant activity in Cuban medicinal plants. *J Ethnopharmacol* 87: 241-246
76. Rao VSN, Menezes AMS, Gadelha MGT (1987). Antifertility screening of some indigenous plants of Brasil. *Fitoterapia* 59(1): 17-20
77. Rice KC & Wilson RS (1976). (-)-3-Isotujone, a small non-nitrogenous molecule with antinociceptive activity in mice. *J Med Chem* 19(8): 1054-1057
78. Roth L, Daunderer M, Kormann K (1994). *Giftpflanzen – Pflanzengifte*. 4. Aufl., Hamburg, Nikol Verlagsgesellschaft mbH & Co KG

79. Rozenfurt E (2006). Taste receptors in the gastrointestinal tract. I. Bitter taste receptors and α -gustducin in the mammalian gut. *Am J Physiol Gastroint Liver Physiol* 291: G171-177
80. SCF (2002). Opinion of the Scientific Committee on Food on Thujone (http://ec.europa.eu/food/fs/sc/scf/out162_en.pdf)
81. Schulz V & Hänsel R (2004) *Rationale Phytotherapie*. 5. Aufl., Berlin Heidelberg New York, Springer Verlag
82. Shafi N, Khan GA, Ghauri EG (2004). Antiulcer effect of *Artemisia absinthium* L. in rats. *Pak J Sci Ind Res* 47(2): 130-134
83. Surber W (1962). Etude de toxicité sous-chronique de la thujone sur rats. Rapport Final. Institut Battelle, Geneve
84. Tahir M, Siddiqui MMH, Khan AB (1997). Effect of Afsanteen (*Artemisia absinthium* Linn.) in acute intestinal amoebiasis. *Hamdrad Med* 40(3): 24-27
85. Tegtmeier M & Harnischfeger G (1994). Methods for the reduction of thujone content in pharmaceutical preparations of *Artemisia*, *Salvia* and *Thuja*. *Eur J Pharm Biopharm* 40(5): 337-340
86. Teuscher E (1989). *Pharmakognosie*. 3. Aufl., Berlin, Akademie-Verlag
87. Tosi B, Bonora A, Dall'Olio G et al. (1991). Screening for toxic thiophene compounds from crude drugs of the family Compositae used in Northern Italy. *Phytother Res* 5(2): 59-62
88. Wake G, Court J, Pickering A et al. (2000). CNS acetylcholine receptor activity in European medicinal plants traditionally used to improve failing memory. *J Ethnopharmacol* 69: 105-114
89. Wegener T (1998). Anwendung eines Trockenextraktes aus *Gentiana lutea* radix bei dyspeptischem Symptomenkomplex. *Z Phytother* 19: 163-164
90. Weisbord SD, Soule JB, Kimmel PL (1997). Poison on line – acute renal failure caused by oil of wormwood purchased through the internet. *New Eng Med* 337:825-527
91. Weiss RF & Fintelmann V (1999). *Lehrbuch der Phytotherapie*. 9. Aufl., Stuttgart, Hippokrates Verlag
92. Wichtl M (2002). *Teedrogen und Phytotherapeutika*. 4. Aufl., Stuttgart, Wissenschaftliche Verlagsgesellschaft mbH
93. Zafar MM, Hamdard ME, Hameed A (1990). Screening of *Artemisia absinthium* for antimalarial effects on *Plasmodium berghei* in mice: a preliminary report. *J Ethnopharmacol* 30: 223-226