



**This document was valid from July 2008 until February 2016.
It is now superseded by a [new version](#) adopted by the HMPC on
2 February 2016 and published on the EMA website.**

Equiseti herba
Equisetum arvense L., herba
(horsetail)

The EMEA acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

(I) References cited in the Assessment Report

Agustín-Ubide MP; Martínez-Cócerca C; Alonso-Llamazares A; Robledo T; Lombardero M; Dominguez J; Carballo MA: Diagnostic approach to anaphylaxis by carrot, related vegetables and horsetail (*Equisetum arvense*) in a homemaker. *Allergy*, 2004; 59 (7):786-7.

Amarowicz R; Pegg RB; Rahimi-Moghaddam P; Barl B; Weil JA: Free-radical scavenging capacity and antioxidant activity of selected plant species from the Canadian prairies. *Food Chemistry*, 2004; 84 (4):551-62.

Aswal BS; Bhakuni DS; Goel AK; Kar K; Mehrotra BN: Screening of Indian Plants for Biological Activity: Part XI. *Indian Journal of Experimental Biology*, 1984; 22 (9):487-504.

Belkin M; Fitzgerald DB; Felix MD: Tumor-damaging capacity of plant materials: II. Plants used as diuretics: *Journal of the National Cancer Institute*, 1952; 13 (3):741-4.

Bézanger-Beauquesne L; Pinkas M; Torck M; Trotin F: *Plantes médicinales des régions tempérées*. 2. Ed. Maloine SA (ed.), 1990; 18.

Bézanger-Beauquesne L; Pinkas M; Torck M: *Les plantes dans la thérapeutique moderne*. Ed. Maloine SA (ed.), 1986; 199-200.

Blumenthal M; Busse WR; Goldberg A; Gruenwald J; Hall T; Riggins W; Rister R; Klein S: *The Complete German Commission E Monographs; Therapeutic Guide to Herbal Medicines*; Ed. The American Botanical Council, Austin, Texas, 1998; 150-1.

Bradley PR: *British Herbal Compendium-Volume 1*; Ed. British Herbal Medicine Association, 1992; 92-4.

Breitwieser K: Versuche über die Brauchbarkeit von Herba Equiseti als Diureticum. *Arch. Pharm.* (1939) 277, 53-61.

British Herbal Medicine Association (BHMA): *A guide to traditional herbal medicines*; Ed. British Herbal Medicine Association, 2003; 134.

British Herbal Medicine Association (BHMA): A guide to traditional herbal medicines; Ed. British Herbal Medicine Association, 1996; 76.

Cetto AA; Wiedenfeld H; Revilla MC; Sergio IA: Hypoglycemic effect of Equisetum myriochaetum aerial parts on streptozotocin diabetic rats. Journal of Ethnopharmacology, 2000; 72 (1/2):129-33.

Conseil de l'Europe: Preparations de plantes utilisees en tant Que matieres premieres dans les produits cosmetiques. 1re edition ; Conseil de l'Europe, 1989; 142-3.

Cow D: Einige Studien über Diurese. Arch. für exper. Path. (1912) 69, 393-411.

Dos Santos Junior JG; Blanco MM; Do Monte FH; Russi M; Lanziotti VM; Leal LK; Cunha GM: Sedative and anticonvulsant effects of hydroalcoholic extract of Equisetum arvense. Fitoterapia, 2005; 76 (6):508-13.

Dos Santos Junior JG; do Monte FHM; Blanco MM; Lanziotti VMdNB; Maia FD; Leal LKdA: Cognitive enhancement in aged rats after chronic administration of Equisetum arvense L. with demonstrated antioxidant properties in vitro. Pharmacology Biochemistry and Behavior, 2005; 81 (3):593-600.

Duke JA: Handbook of medicinal herbs; CRC Press, 1985; 178-9, 515, 555.

Duke JA: Handbook of medicinal herbs; CRC Press, 2002; 391-2.

Equisetum. In: British Herbal Pharmacopoeia Part One, British Herbal Medicinal Association, 1976, 75.

Equisetum. In: Ainley Wade, Martindale The Extra Pharmacopoeia, 27 th Ed, Pharmaceutical Society of Great Britain 1977, 1754.

Equiseti herba, Schachtelhalmkraut, Monographie der Kommission E. In: Bundesanzeiger Nr. 173 vom 18.09.1986.

Equiseti herba. Pharmacopoea Helvetica, Verlag Eidgenössische Drucksachen- und Materialzentrale Bern 1971

Equisetum Stem-Equiseti herba.(01/2008: 1825 corrected 6.0) In European Pharmacopoeia, 6.0 Vol 2 Council of Europe, 2008, 1794-5.

Equiseti herba Schachtelhalmkraut. Ergänzungsbuch zum Deutschen Arzneibuch 6. Ausgabe, DAV Stuttgart, 1953, 244-245.

Equiseti herba. Medicaments à base plantes: Les Cahiers de l'Agence No.3 Agence du Médicament [Agency instructions Nr. 3] (notice: former version: Bulletin officiel Nr.. 90/22 1990) 1998, 43, 68, 70, 77.

Fabre B; Geay B; Beaufils P: Thiaminase activity in Equisetum arvense and its extracts. Plantes Medicinales Phytothérapie, 1993; 26 (3):190-7.

Frohne D, Pfänder HJ: A colour atlas of poisonous plants: Ed. A Wolfe Science Book, 1984; 103-4. AR

Garnier G, Bezanger-Beauquesne L, Debraux G: Ressources medicinales de la flore francaise; Ed. Vigot Freres, 1961; 1:90-1.

Graefe EU; Veit M: Urinary metabolites of flavonoids and hydroxycinnamic acids in humans after application of a crude extract from *Equisetum arvense*. *Phytomedicine*, 1999; 6 (4):239-46.

Goun EA; Petrichenko VM; Solodnikov SU; Suhinina TV; Kline MA; Cunningham G; Nguyen C; Miles H: Anticancer and antithrombin activity of Russian plants: *Journal of Ethnopharmacology*; 2002; 81 (3):337-42.

Grases F; Melero G; Costa-Bauza A; Prieto R; March JG: Urolithiasis and phytotherapy. *International Urology and Nephrology*, 1994; 26 (5):507-11.

Guerin JC; Reveillere HP: Antifungal Activity of Plant Extracts in Therapy. I. Study of 41 Extracts against 9 Fungal Species. (Fr.). *Annales Pharmaceutiques Françaises*, 1984; 42 (6):553-9.

Hänsel R., Keller K., Rimpel H., Schneider G.: *Hagers Handbuch der Pharmazeutischen Praxis* 5. Ed.; Springer Verlag, Berlin, 1993, Vol. 5: Drogen E-O, 64-9.

Hamon NW; Awang DV: Horsetail: *Canadian Pharmaceutical Journal*, 1992; 125:399-401.

Heisey RM; Gorham BK: Antimicrobial Effects of Plant Extracts on *Streptococcus mutans*, *Candida albicans*, *Trichophyton rubrum* and Other Micro-Organisms. *Letters in Applied Microbiology*; 1992; 14 (4):136-9.

Henderson JA; Evans EV; McIntosh RA: The antithiamine action of *Equisetum*. *Journal of the American Veterinary Medical Association*, 1952; 120 (903):375-378.

Herre E: Die diuretische Wirkung der in Naturheilkunde und Volksmedizin angewandten harntreibenden Mittel. *Arch. exp. Path. Pharm.* (1937) 184, 710-15.

Hesse A; Vahlensieck W: Harntee 400 for the Prophylaxis of Urolithiasis Recurrence. [Ger.] *Therapiewoche*, 1985; 35 (16):1975-80.

Hiermann A: *Equisetum*. In: Blaschek W, Ebel S, Hackenthal E, Holzgrabe U, Keller K, Reichling J (eds.): *Hagers Handbuch der Drogen und Arzneistoffe*. 2003. Hager ROM 2003; Springer electronic media

Hiermann A: *Equisetum*. In: Blaschek W, Ebel S, Hackenthal E, Holzgrabe U, Keller K, Reichling J (eds.): *Hagers Handbuch der Drogen und Arzneistoffe*. 2006. Hager ROM 2006; Springer electronic media

Hoppe HA: *Equisetum hyemale*. In: Hoppe HA: *Drogenkunde*, Walter de Gruyter Verlag Berlin, 8. Ed. 1975; 174.

Husson GP; Vilagines R; Delaveau P: Research into the antiviral properties of a few natural extracts: *Annales Pharmaceutiques Françaises*, 1986; 44 (1):41-8.

Jaretsky R.; Breitwieser K; Neuwald F: Diureseversuche mit Kieselsäure und Kieselsäuredrogen an Ratten. *Arch. Pharm.* (1938) 276, 552-59.

Jaretsky R: Entgegnung zu vorstehenden Ausführungen Vollmers "Über die diuretische Wirkung der *Radix Ononidis* und der *Herba Equiseti*" *Arch. Pharm.* (1940) 278, 44-7.

Jean-Blain C; Grisvard M: *Plantes veneneuses-toxicologie*, La Maison Rustique, 1973; 20-1.

Joksic G; Stankovic M; Novak A: Antibacterial medicinal plants *Equiseti herba* and *Ononidis radix* modulate micronucleus formation in human lymphocytes in vitro. *Journal of Environmental Pathology, Toxicology and Oncology*, 2003; 22 (1):41-8.

Katalinic V; Milos M; Kulisic T; Jukic M: Screening of 70 medicinal plant extracts for antioxidant capacity and total phenols. *Food Chemistry*, 2006; 94 (4):550-7.

Kloucek P; Polesny Z; Svobodova B; Vlkova E; Kokoska L: Antibacterial screening of some Peruvian medicinal plants used in Callería District. *Journal of Ethnopharmacology*, 2005; 99 (2):309-12.

Kreitmair H: Pharmakologische Versuche mit einigen einheimischen Pflanzen. *Merck's Jahresbericht*, (1936) 50, 103-110.

Kreitmair H: Equisetum arvense-der Ackerschachtelhalm. *Pharmazie* (1953) 3, 298-300.

LaGow B (ed.): Horsetail. *PDR for Herbal Medicines* (3rd edition): Ed. Thomson PDR, 2004; 450.

Länger: Eine südamerikanische Heilpflanze, wissenschaftlich erforscht; *ÖAZ Aktuell* (4) 2002

Lemus I; Garcia R; Erazo S; Pena R; Parada M; Fuenzalida M: Diuretic activity of a Equisetum bogotense tea (Platero herb): evaluation in healthy volunteers. *Journal of Ethnopharmacology*, 1996; 54:55-8.

Lovkova MY; Shelepova OV; Sokolova SM; Sabirova NS; Rabinovich AM: Selenium in medicinal plants of the Russian flora. *Biology Bulletin of the Russian Academy of Sciences*, 1993; 20 (6):669-673 paper translated from *Izvestiya Rossiiskoi Akademii Nauk, Seriya Biologicheskaya* 1993; 6:833-8.

Madaus G: Equisetum arvense et Equisetum hiemale. In: Madaus G: *Lehrbuch der biologischen Heilmittel*. Georg Thieme Verlag Leipzig, 1938; 1:1267-78.

Madaus G: Equisetum arvense et Equisetum hyemale. In: Madaus G: *Lehrbuch der biologischen Heilmittel*, Georg Olms Verlag Hildesheim, 1976 (ISBN 3-4870-589); 1267-77.

Maeda H; Miyamoto K; Sano T: Occurrence of dermatitis in rats fed a cholesterol diet containing field horsetail (Equisetum arvense L.). *Journal of Nutritional Science and Vitaminology*, 1997; 43 (5):553-63.

Mills S; Bone K: *The essential guide to herbal safety*. Ed. Elsevier, Churchill Livingstone, 2005; 476-8.

do Monte FHM; dos Santos Júnior JG; Russi M; Lanziotti VMNB; Leal LKAM; de Cunha GMA: Antinociceptive and anti-inflammatory properties of the hydroalcoholic extract of stems from Equisetum arvense L. in mice. *Pharmacological Research*, 2004; 49 (3):239-43.

Nagai T; Myoda T; Nagashima T: Antioxidative activities of water extract and ethanol extract from field horsetail (tsukushi) Equisetum arvense L.. *Food Chemistry*, 2005; 91 (3):389-94.

Oh H; Kim DH; Cho JH; Kim YC: Hepatoprotective and free radical scavenging activities of phenolic petrosins and flavonoids isolated from Equisetum arvense. *Journal of Ethnopharmacology*, 2004; 95(2-3):421-4.

Parihar P: Antifungal efficacy of various pteridophytic plant parts extracts: a study in vitro. *Advances in Plant Sciences*, 2002; 15 (1):35-8.

Paris RR, Moyses H: *Precis de matiere medicale*. (2eme edition): Ed. Masson, 1976; 1:376-8.

Perazella: Drug induced hyperkalemia – the reply. *The American Journal of Medicine* 2002, Volume 112, 334.

Pérez Gutiérrez RM; Yescas Laguna G; Walkowski A: Diuretic activity of Mexican equisetum. *Journal of Ethnopharmacology*, 1985; 14 (2/3):269-72.

Pohl RW: Toxicity of ferns and Equisetum: *American Fern Journal*, 1955; 45 (3):95-7.

Rapp WF: The toxicity of Equisetum: *American Fern Journal*, 1954; 44 (4):148-54.

Revilla MC; Andrade-Cetto A; Islas S; Wiedenfeld H. Titel: Hypoglycemic effect of Equisetum myriochaetum aerial parts on type 2 diabetic patients. *Journal of ethnopharmacology* 2002; VOL: 81 (1); p. 117-20.

Robert S: Phytotherapie et Affections du Systeme Uro-Genital. [Phytotherapy and affections of the urogenital system] *Journal de Pharmacie de Belgique*, 1986; 41 (5):40-5.

Ryu JH; Ahn H; Kim JY; Kim YK: Inhibitory activity of plant extracts on nitric oxide synthesis in LPS-activated macrophages. *Phytotherapy research: PTR*, 2003; 17 (5):485-9.

Saint-Paul A: La composition elementaire de quelques Equisetum medicaux: *Plantes medicinales et phytotherapie*, 1980; 14(2):73-82.

Schachtelhalmkraut. In: Braun R Standardzulassungen für Fertigarzneimittel, DAV Stuttgart Govi Verlag GmbH Frankfurt 2004.

Schauss AG; Steels E: Preliminary evidence of the safety and efficacy of UroLogic (TM), an herbal formulation for the treatment of urinary incontinence and overactive bladder. *Experimental Biology 2006 Meeting*, 2006; San Francisco, CA, USA; American Association of Anatomists

Schneider K; Kubelka W: Zur Hamolytischen Wirkung Von Schachtelhalmkraut (Herba Equiseti) [Hemolytic effect of horsetail (Herba equiseti)] *Scientia Pharmaceutica*, 1989; 57 (3):214-5.

Scott IM; Leduc RI; Burt AJ; Marles RJ; Arnason JT; Foster BC: The inhibition of human cytochrome P450 by ethanol extracts of North American botanicals. *Pharmaceutical Biology*, 2006; 44 (5):315-27.

Sparavigna A; Setaro M; Genet M; Frisenda L: Equisetum arvense in a new transungual technology improves nail structure and appearance; *Journal of Plastic Dermatology*, 2006; 2 (1):31-8. (Only abstract available)

Stajner D; Popovic BM; Canadanovic-Brunet J; Boza P: Free radical scavenging activity of three Equisetum species from Fruska gora mountain. *Fitoterapia*, 2006; 77 (7-8):601-4.

Suganda AG; Amoros M; Girre L; Fauconnier B: Inhibitory Effects of Some Crude and Semi-Purified Extracts of Plants Indigenous to France on the Replication of Human Herpes Virus 1 and Human Poliovirus 2 in Cell Culture. *Journal of Natural Products*, 1983; 46 (5):626-32.

The Committee of Experts on Cosmetic Products: Plants in cosmetics. Volume I: Ed. Council of Europe Publishing, 2001; 125-6.

Trouillas P; Calliste CA; Allais DP; Simon A; Marfak A; Delage C; Duroux JL: Antioxidant, anti-inflammatory and antiproliferative properties of sixteen water plant extracts used in the Limousin countryside as herbal teas: *Food Chemistry*, 2003; 80 (3):399-407.

Van Hellemont J: *Compendium de Phytotherapie*. Ed. APB, 1986; 144-6.

Veit M: Die Schachtelhalme (Equisetaceae). *Objekte Der Forschung und der Praxis: Equisetum spp.* *Deutsche Apotheker Zeitung*, 1987; 127 (41):2049-56.

Veit M: Probleme Bei der Bewertung pflanzlicher Diuretika. Als Beispiel Schachtelhalmkraut DAB 10 (Equiseti herba) [Problems of the evaluation of herbal diuretics. For example herb of Horsetail DAB 10 (Equiseti herba)] Zeitschrift für Phytotherapie, 1994; 15 (6):331-41.

Vollmer V; Hindemith H: Untersuchungen über die diuretische Wirkung von Ononis spinosa L. und Equisetum arvense L. an Kaninchen und Mäusen; Archiv für experimentelle Pathologie und Pharmakologie, 1937; 186:565-73.

Vollmer H; Hübner K: Untersuchung über die diuretische Wirkung der Fructus juniperi, Radix levistici, Radix ononidis, Folia betulae, Radix liquiritae und Herba equiseti an Ratten; Archiv für experimentelle Pathologie und Pharmakologie, 1937; 186:592-605.

Vollmer H: Bemerkungen zur Technik des Diureseversuches erläutert an Hand einiger Coffeinversuche-zugleich Bemerkungen über die diuretischen Wirkung von Hauhechel und Schachtelhalm. Archiv für experimentelle Pathologie und Pharmakologie 1939; 193:483-9.

Vollmer H: Untersuchungen über einige pflanzliche Diuretika; Fortschritte der Therapie, 1941; 19: 322.

Wachter R: Vergleichende Versuche über die diuretischen Wirkung einiger in der Volksmedizin gebräuchlicher Pflanzen, Dissertation, Gießen 1938

Whiting PW; Clouston A; Kerlin P: Black cohosh and other herbal remedies associated with acute hepatitis: Medical Journal of Australia, 2002; 177 (8):440-3.

Wichtl M; Anton R: Plantes thérapeutiques (2ème édition allemande – 2ème édition française) Ed. Tec & Doc – EM inter, 2003; 195-9.

Wichtl M.: Herbal drug and Phytopharmaceuticals (3rd English edition) Ed. MedPharm GmbH Scientific Publishers, 2004; 195-9.

Willuhn G: Phytopharmaka in der Dermatologie. [Phytomedicines in dermatology] Zeitschrift für Phytotherapie, 1995; 16 (6):325-42.

(2) References read but not included to the Assessment Report

Andrade-Cetto A; Heinrich M: Mexican plants with hypoglycaemic effect used in the treatment of diabetes. *Journal of Ethnopharmacology*, 2005; 99 (3):325-48.

Anonymous: Schachtelhalm (nicht nur) für die Niere. [Horsetail plant (not only) for the kidney]. *Schweizerische Rundschau für Medizin Praxis [Revue suisse de médecine Praxis]*, 2005; 94 (6):222-3.

Anonymous: Equisetum hyemale. In: List PH; Hörhammer L: *Hagers Handbuch der Pharmazeutischen Praxis*, Springer Verlag Berlin 1975, 4. Ed.; 793.

Anonymous: Equisetum L.. *Pharmazeutische Stoffliste* E 9/86, 1986; 136-7.

Berger F: Herba Equiseti majoris. In: Berger F (ed.): *Handbuch der Drogenkunde*. W Maudrich Verlag Wien 1954; 212-3.

Bühning U: Heilpflanzen bei Ulcus cruris [Medicinal plants for the treatment of leg ulcers]: *Zeitschrift für Phytotherapie*; 2005; 26 (4):185-7.

Carnat A; Petitjean-Freytet C; Muller D; Lamaison JL: Teneurs en principaux constituants de la prêle, Equisetum arvense L. [Main constituents of the sterile fronds of Equisetum arvense L.]. *Plantes Médicinales et Phytothérapie*, 1991; 25 (1):32-8.

Ganeva Y; Chanev C; Dentchev T: Triterpenoids and sterols from Equisetum arvense: *Dokladi na B"lgarskata Akademiya na Naukite*, 2001; 54 (2):53-6.

Gessner O; Orzechowski G (ed.): Equisetum. In: Gessner O; Orzechowski G (ed.): *Gift- und Arzneipflanzen von Mitteleuropa*, Carl Winter Verlag Heidelberg, 3rd Ed. ISBN 3533023729, 1974; 429-30.

Ghazala N; Iqbal SH; Bhutta AA: Equisetum: an excellent host for VAM infections. *Biologia Pakistan*, 1987; 33 (1):97-108.

Gräfe U; Gumbinger HG; Winterhoff H; Veit M: Urinary metabolites of phenolic compounds of Equisetum arvense. *2nd International Congress of Phytomedicine and 7th Congress of the Society of Phytotherapy in Cooperation with ESCOP 1996*; 82.

Gürocak S; Küpeli B: Consumption of Historical and Current Phytotherapeutic Agents for Urolithiasis: A Critical Review: *Journal of Urology*, 2006; 176 (2):450-5.

Hagenstroem UHT; Elze A: Bladder and Kidney Teas. *Phyto-galenical Aspects*. [Ger.]: *Deutsche Apothekerzeitung*, 1986; 126 (37):1917-21.

Ingram TJ: Compositions comprising equisetum and symphytum and methods for the treatment of symptoms associated with mycoplasma infection: *Official Gazette of the United States Patent and Trademark Office Patents*, 2002; 1260 (2): <http://www.uspto.gov/web/menu/patdata.html>

Koeppen BM; Stanton BA: *Renal Physiology* (2nd and 3rd edition). Mosby 2001.

Kolettis TM; Oikonomou G; Novas I; Sideris DA: Transient complete atrioventricular block associated with herb intake: *Europace*, 2005; 7 (3):225-6.

Koniger H: Method for combating summer eczema and malanders: *Official Gazette of the United States Patent and Trademark Office Patents*, 2003; 1272 (2) <http://www.uspto.gov/web/menu/patdata.html>

Martini MC, Seiler M: Actifs & additives en Cosmetologie. Ed. Lavoisier (TEC & DOC), 1992; 23-4.

Pourmorad F; Hosseinimehr SJ; Shahabimajd N: Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. African Journal of Biotechnology, 2006; 5 (11):1142-5.

Schilcher H: Pflanzliche Urologika [Plant products used in urology] Deutsche Apotheker Zeitung, 1984; 124 (47):2429-36

Senin P; Setnikar I; Rovati AL: Phytoestrogens and probiotic for women's health: Official Gazette of the United States Patent and Trademark Office Patents, 2006;

Steenkamp V: Phytomedicines for the prostate. Fitoterapia, 2003; 74 (6):545-52.

Veit M (1996) Zur Pharmakokinetik pflanzlicher Phenole. Pharmazeutische Zeitung 141(12): 52-3

Veit M; Cygan FC; Geiger H: Seasonal and intraspecific variations in flavonoids of Equisetum arvense. Planta Medica, 1991; 57 (Supplement issue 2) A 3.

Veit M; Bauer K; Geiger H; Cygan FC: Flavonoids of Equisetum hybrids in the subgenus Equisetum. Planta Medica, 1992; 58 (Supplement issue 1), A 697.

De Villar GPN; Loria V; Monereo S: Tratamientos «Alternativos» de la Obesidad: Mito y Realidad [«Alternative» drugs for obesity treatment: Myth and reality] Medicina Clinica, 2003; 121 (13):500-10

Weidner C; Veit M; Czygan FC: Accumulation Dynamics of Caffeic Acid Conjugates in Equisetum arvense. Planta Medica, 1991; 57 (Supplement issue 2), A37.

Zander R: Equisetum hyemale. In: Zander R: Handwörterbuch der Pflanzennamen. Eugen Ulmer Verlag Stuttgart, 1979; 276.