



EMA/COMP/407852/2011 Rev.2  
Committee for Orphan Medicinal Products

## Public summary of opinion on orphan designation

N-(cyanomethyl)-4-(2-{[4-(morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)benzamide, dihydrochloride salt for the treatment of post-essential thrombocythaemia myelofibrosis

First publication	16 August 2011
Rev.1: transfer of sponsorship	16 March 2012
Rev.2: transfer of sponsorship	5 June 2013
<b>Disclaimer</b> Please note that revisions to the Public Summary of Opinion are purely administrative updates. Therefore, the scientific content of the document reflects the outcome of the Committee for Orphan Medicinal Products (COMP) at the time of designation and is not updated after first publication.	

On 5 August 2011, orphan designation (EU/3/11/887) was granted by the European Commission to Cres Pharmaceuticals Limited, United Kingdom, for N-(cyanomethyl)-4-(2-{[4-(morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)benzamide, dihydrochloride salt for the treatment of post-essential thrombocythaemia myelofibrosis.

The sponsorship was transferred to YM BioSciences (UK) Limited, United Kingdom, in February 2012 and subsequently to Gilead Sciences International Ltd, United Kingdom, in May 2013.

### What is post-essential thrombocythaemia myelofibrosis?

Myelofibrosis is a disease in which the bone marrow (the spongy tissue inside the large bones) becomes dense and fibrous, and starts producing abnormal immature blood cells that replace the normal blood cells. It can develop following thrombocythaemia (overproduction of platelets, components that help the blood to clot). When the thrombocythaemia is not caused by any known condition, the disease is known as post-essential thrombocythaemia myelofibrosis.

In myelofibrosis, some immature blood cells migrate from the bone marrow to other organs, such as the spleen and liver, where they mature. This causes the organs to become enlarged. Patients with myelofibrosis can develop several symptoms, including bone pain, fever, tiredness, weakness, weight loss, infections and bleeding.



Post-essential thrombocythaemia myelofibrosis is a debilitating disease that is long lasting and life threatening because it can lead to severe anaemia (low red blood cell counts) and infections, and can result in leukaemia (cancer of the white blood cells).

### **What is the estimated number of patients affected by the condition?**

At the time of designation, post-essential thrombocythaemia myelofibrosis affected less than 0.15 in 10,000 people in the European Union (EU). This is equivalent to a total of fewer than 8,000 people\*, and is below the ceiling for orphan designation, which is 5 people in 10,000. This is based on the information provided by the sponsor and the knowledge of the Committee for Orphan Medicinal Products (COMP).

### **What treatments are available?**

At the time of designation, although hydroxycarbamide and busulfan were authorised in the EU for primary myelofibrosis (myelofibrosis of unknown cause), there were no treatments authorised specifically for post-essential thrombocythaemia myelofibrosis. Medicines were authorised to treat the symptoms, including erythropoietin (a hormone that stimulates the production of red blood cells) to treat anaemia, and surgery or radiation to remove or shrink the enlarged spleen. In some patients, allogeneic stem-cell transplantation was used to treat the disease. This is a complex procedure where the patient receives stem cells from a matched donor to help restore the bone marrow.

### **How is this medicine expected to work?**

This medicine is thought to work by blocking some enzymes known as Janus kinases (JAKs). These enzymes can be found in some receptors on the surface of cells and are involved in the reproduction and growth of blood cells. In myelofibrosis, JAKs are more active than normal. By blocking these enzymes, this medicine is expected to slow down the abnormal growth of blood cells, reducing the symptoms of the disease.

### **What is the stage of development of this medicine?**

The effects of N-(cyanomethyl)-4-(2-{[4-(morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)benzamide, dihydrochloride salt have been evaluated in experimental models.

At the time of submission of the application for orphan designation, clinical trials with the medicine in patients with myelofibrosis were ongoing.

At the time of submission, this medicine was not authorised anywhere in the EU for post-essential thrombocythaemia myelofibrosis. Orphan designation of the medicine had been granted in the United States of America for myelofibrosis.

In accordance with Regulation (EC) No 141/2000 of 16 December 1999, the COMP adopted a positive opinion on 5 May 2011 recommending the granting of this designation.

---

\*Disclaimer: For the purpose of the designation, the number of patients affected by the condition is estimated and assessed on the basis of data from the European Union (EU 27), Norway, Iceland and Liechtenstein. At the time of designation, this represented a population of 507,700,000 (Eurostat 2011).

Opinions on orphan medicinal product designations are based on the following three criteria:

- the seriousness of the condition;
- the existence of alternative methods of diagnosis, prevention or treatment;
- either the rarity of the condition (affecting not more than 5 in 10,000 people in the EU) or insufficient returns on investment.

Designated orphan medicinal products are products that are still under investigation and are considered for orphan designation on the basis of potential activity. An orphan designation is not a marketing authorisation. As a consequence, demonstration of quality, safety and efficacy is necessary before a product can be granted a marketing authorisation.

## **For more information**

Sponsor's contact details:

Gilead Sciences International Ltd.  
Flowers Building  
Granta Park  
Abington  
Cambridge CB21 6GT  
United Kingdom  
Telephone: + 44 1223 897 300  
Telefax: + 44 1223 897 284

For contact details of patients' organisations whose activities are targeted at rare diseases see:

- [Orphanet](#), a database containing information on rare diseases which includes a directory of patients' organisations registered in Europe.
- [European Organisation for Rare Diseases \(EURORDIS\)](#), a non-governmental alliance of patient organisations and individuals active in the field of rare diseases.

## Translations of the active ingredient and indication in all official EU languages<sup>1</sup>, Norwegian and Icelandic

Language	Active ingredient	Indication
English	N-(cyanomethyl)-4-(2-{ [4-(morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)benzamide, dihydrochloride salt	Treatment of post-essential thrombocythaemia myelofibrosis
Bulgarian	N-(цианометил)-4-(2-{ [4-(морфолин-4-ил)фенил]амино}пиримидин-4-ил)бензамид дихидрохлорид сол	Лечение на миелофиброза след есенциална тромбоцитемия
Czech	N-(kyanomethyl)-4-(2-{ [4-(morfolin-4-yl)fenyl]amino}pyrimidin-4-yl)benzamid dihydrochlorid sůl	Léčba post-esenciální trombocytémické myelofibrózy
Danish	N-(cyanomethyl)-4-(2-{ [4-(morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)benzamid-dihydrochlorid-salt	Behandling af post essentiel thrombocythæmi myelofibrose
Dutch	N-(cyanomethyl)-4-(2-{ [4-(morfoline-4-yl)fenyl]amino}pyrimidine-4-yl)benzamididihydrochloridezout	Behandeling van myelofibrosis volgend op essentiële trombocytemie
Estonian	N-(tsüanometüül)-4-(2-{ [4-(morfoliin-4-üül)fenüül]amiino}pürimidiin-4-üül)bensamiid dihidrokloriid sool	Postessentsiaalse trombotsüteemia müelofibroosi ravi
Finnish	N-(syanometyyli)-4-(2-{ [4-(morfoliini-4-yyli)fenyyli]amino}pyrimidiini-4-yyli)bensamididihydrokloridisuola	Essentiaalisen trombosytämian jälkeisen myelofibroosin hoito
French	N-(cyanométhyle)-4-(2-{ [4-(morpholine-4-yle)phényle]amino}pyrimidine-4-yle)benzamide dichlorhydrate	Traitement de la myélobiose consécutive à une thrombocytémie essentielle
German	N-(Cyanomethyl)-4-(2-{ [4-(Morpholin-4-yl)Phenyl]amino}pyrimidin-4-yl)Benzamid Dihydrochloridsalz	Behandlung einer Myelofibrose nach essentieller Thrombozythämie
Greek	N-(κυανομέθυλ)-4(2-{ [4-(μορφολινο-4-υλ)φαινυλ]αμινο}πυριμιδινο-4-υλ)βενζαμιδιο, άλας διυδροχλωρικό	Θεραπεία της μυελοϊνωσης από ιδιοπαθή θρομβοκυττάρωση
Hungarian	N-(cianometil)-4-(2-{ [4-(morfolin-4-yl)fenil]amino}pirimidin-4-yl)benzamid dihidroklorid só	Esszenciális thrombocytaemiát követő mielofibrózis kezelésére
Italian	N-(cianometil)-4-(2-{ [4-(morfolin-4-ile)fenil]ammino}pirimidin-4-ile)benzamide sale dicloridrato	Trattamento della mielofibrosi post-trombocitemia essenziale
Latvian	N-(ciānmetil)-4-(2-{ [4-(morfolīn-4-il)fenil]amino}pirimidin-4-il)benzamīda dihidrohlorīda sāls	Pēc-esenciālas trombocitēmijas mielofibrozes ārstēšana
Lithuanian	N-(cianometil)-4-(2-{ [4-(morfolin-4-il)fenil]amino}pirimidin-4-il)benzamidias dihidrochlorido druska	Mielofibrozes gydymas po esencialinės trombocitemijos

<sup>1</sup> At the time of designation

Language	Active ingredient	Indication
Maltese	Melħ dihydrochloride ta' N-(cyanomethyl)-4-(2-{ [4-(morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)benzamide	Kura tal-mjelofibrozi konsegwenti għal tromboċitemija essenzjali
Polish	N-(cyjanometylo)-4-(2-{ [4-(morfolin-4-ylo)fenylo]amino}pirymidyn-4-ylo)benzamid dichlorowodorek	Leczenie mielofibrozy wywołanej nadpłytkowością samoistną
Portuguese	N-(cianometil)-4-(2-{ [4-(morfolina-4-il)fenil]amino}pirimidina-4-il)cloridrato de benzamida	Tratamento da mielofibrose devida a trombocitémia essencial
Romanian	Diclorhidrat de N-(cianometil)-4-(2-{ [4-(morfolin-4-yl)fenil]amino}pirimidin-4-yl)benzamidă	Tratamentul mielofibrozei post-trombocitemie esențială
Slovak	N-(cyanometyl)-4-(2-{ [4-(morfolín-4-yl)fenyl]amino}pyrimidín-4-yl)benzamid dihydrochlorid	Liečba myelofibrózy po esenciálnej trombocytémii
Slovenian	N-(cianometil)-4-(2-{ [4-(morfolin-4-il)fenil]amino}pirimidin-4-il)benzamid dihidroklorid soli	Zdravljenje mielofibroze, nastale po esecialni trombocitemiji
Spanish	N-(cianometil)-4-(2-{ [4-(morfolina-4-il)fenil]amino}pirimidina-4-il)benzamida sal de dihidrocloruro	Tratamiento de la mielofibrosis secundaria a trombocitemia esencial
Swedish	N-(cyanometyl)-4-(2-{ [4-(morfolin-4-yl)fenyl]amino}pyrimidin-4-yl)benzamid dihydrokloridsalt	Behandling av post-essentiell trombocytemi myelofibros
Norwegian	N-(cyanometyl)-4-(2-{ [4-(morfolin-4-yl)fenyl]amino}pyrimidin-4-yl)benzamid dihydroklorid	Behandling av myelofibrose sekundært til essensiell trombocytemi
Icelandic	N-(sýanómetýl)-4-(2-{ [4-(morfoólin-4-ýl)fenýl]amínó}pýrimídín-4-ýl)benzamið díhýdróklóríð salt	Meðferð á myelófibrósu í kjölfar eðlislægs blóðflagnadreyra