



EUROPEAN MEDICINES AGENCY
SCIENCE MEDICINES HEALTH

5 September 2016
EMA/COMP/442574/2016
Committee for Orphan Medicinal Products

Public summary of opinion on orphan designation

Autologous Epstein-Barr virus specific T cells derived from peripheral blood mononuclear cells, expanded ex vivo for the treatment of extranodal NK/T-cell lymphoma, nasal type

On 14 July 2016, orphan designation (EU/3/16/1695) was granted by the European Commission to Cell Medica Ltd., United Kingdom, for autologous Epstein-Barr virus specific T cells derived from peripheral blood mononuclear cells, expanded ex vivo for the treatment of extranodal NK/T-cell lymphoma, nasal type.

What is extranodal NK/T-cell lymphoma, nasal type?

Extranodal NK/T-cell lymphoma, nasal type, is an aggressive cancer of a type of white blood cell called lymphocytes. Patients with this cancer have cancerous growths in the nose and surrounding areas, which can destroy bone, cartilage and soft tissue, disfiguring the face. The cancer is usually associated with infection with a type of virus called Epstein-Barr virus.

Extranodal NK/T-cell lymphoma, nasal type, is a debilitating and life-threatening disease that causes disfigurement and is associated with poor overall survival.

What is the estimated number of patients affected by the condition?

At the time of designation, extranodal NK/T-cell lymphoma, nasal type, affected approximately 0.08 in 10,000 people in the European Union (EU). This was equivalent to a total of around 4,000 people*, and is below the ceiling for orphan designation, which is 5 people in 10,000. This is based on the information provided by the sponsor and the knowledge of the Committee for Orphan Medicinal Products (COMP).

What treatments are available?

At the time of designation, various cancer medicines were authorised in the EU for the treatment of non-Hodgkins' lymphomas such as extranodal NK/T-cell lymphoma, nasal type.

*Disclaimer: For the purpose of the designation, the number of patients affected by the condition is estimated and assessed on the basis of data from the European Union (EU 28), Norway, Iceland and Liechtenstein. This represents a population of 513,700,000 (Eurostat 2016).



The sponsor has provided sufficient information to show that the medicine might be of significant benefit for patients with extranodal NK/T-cell lymphoma, nasal type because early clinical studies indicate that patients whose disease came back after previous treatment responded to this treatment. This assumption will need to be confirmed at the time of marketing authorisation, in order to maintain the orphan status.

How is this medicine expected to work?

The cancer cells in patients with extranodal NK/T cell lymphoma are nearly always infected with Epstein-Barr virus.

This medicine contains certain T cells obtained from the patient. T cells are cells of the immune system (the body's natural defences) that kill infected or abnormal cells. To make this medicine, the T cells are exposed to protein fragments from Epstein-Barr virus, so that they can recognise cells infected with the virus and attack them. When the T cells are injected back into the patient, they are expected to attach to infected cancer cells and kill them.

What is the stage of development of this medicine?

At the time of submission of the application for orphan designation, the evaluation of the effects of the medicine in experimental models was ongoing.

At the time of submission, clinical trials with the medicine in patients with extranodal NK/T cell lymphoma, nasal type were ongoing.

At the time of submission, the medicine was not authorised anywhere in the EU for extranodal NK/T cell lymphoma, nasal type or designated as an orphan medicinal product elsewhere for this condition.

In accordance with Regulation (EC) No 141/2000 of 16 December 1999, the COMP adopted a positive opinion on 16 June 2016 recommending the granting of this designation.

Opinions on orphan medicinal product designations are based on the following three criteria:

- the seriousness of the condition;
- the existence of alternative methods of diagnosis, prevention or treatment;
- either the rarity of the condition (affecting not more than 5 in 10,000 people in the EU) or insufficient returns on investment.

Designated orphan medicinal products are products that are still under investigation and are considered for orphan designation on the basis of potential activity. An orphan designation is not a marketing authorisation. As a consequence, demonstration of quality, safety and efficacy is necessary before a product can be granted a marketing authorisation.

For more information

Sponsor's contact details:

Contact details of the current sponsor for this orphan designation can be found on EMA website, on the medicine's [rare disease designations page](#).

For contact details of patients' organisations whose activities are targeted at rare diseases see:

- [Orphanet](#), a database containing information on rare diseases, which includes a directory of patients' organisations registered in Europe;
- [European Organisation for Rare Diseases \(EURORDIS\)](#), a non-governmental alliance of patient organisations and individuals active in the field of rare diseases.

Translations of the active ingredient and indication in all official EU languages¹, Norwegian and Icelandic

Language	Active ingredient	Indication
English	Autologous Epstein-Barr virus specific T-cells derived from peripheral blood mononuclear cells, expanded ex vivo	Treatment of extranodal NK/T-cell lymphoma, nasal type
Bulgarian	Автоложни Т-клетки, специфични за Epstein-Barr вирус, получени от мононуклеарни клетки от периферна кръв, култивирани ex vivo	Лечение на екстранодален NK/T-клетъчен лимфом, назален тип
Croatian	Autologne T-stanice specifične za Epstein-Barr virus derivirane iz mononuklearnih stanica periferne krvi, ekspandirane ex vivo	Liječenje ektranodalnog NK/T-staničnog limfoma, nazalni oblik
Czech	Autologní Epstein-Barrové virus specifické T buňky získané z periferních mononukleárů, expandované ex vivo	Léčba extranodálních NK/T nasálních lymfomů
Danish	Autolog Epstein-Barr virus specifikke T-celler, udvundet fra perifert blod mononukleare celler, ekspanderet ex vivo	Behandling af ektranodal T-NK-cellelymfom, nasal type
Dutch	Autologe Epstein-Barr virus specifieke T-cellen afgeleid uit perifeer bloed mononucleaire cellen, geëxpandeerd ex vivo	Behandeling van extranodaal NK/T-cel lymfoom, nasaal type
Estonian	Ex vivo paljundatud autoloogsed Epstein-Barri viiruse spetsiifilised T-rakud, mis on saadud perifeerse vere mononukleaarsetest rakkudest	Nina-tüüpi ektranodaalse NK/T-rakkümfoomi ravi
Finnish	Autologiset Epstein-Barr virukselle spesifiset t-solut, jotka ovat peräisin perifeerisen veren mononuklearisista, ex vivo monistetusta soluista	Ekstranodaalisen, nasaalityyppisen NK/T-solulymfooman hoito
French	Cellules T autologues spécifiques du virus d'Epstein-Barr, dérivées de cellules mononucléaires sanguines périphériques, expansées ex vivo	Traitement du lymphome T/NK extraganglionnaire, de type nasal
German	Autologe T-Zellen spezifisch gegen das Epstein-Barr Virus, gewonnen von mononukleäre Zellen des peripheren Blutes und expandiert ex vivo	Behandlung des extranodalen NK/T-Zell-Lymphom, nasaler Typ
Greek	Αυτόλογα Epstein-Barr ιού ειδικά Τ-κύτταρα που προέρχονται από μονοπύρρηνα κύτταρα περιφερικού αίματος τα οποία επεκτάθηκαν ex vivo	Θεραπεία του εξωλεμφαδενικού NK/T λεμφώματος, ρινικού τύπου
Hungarian	Ex vivo expandált, perifériális vér mononukleáris sejtekből származó autológ Epstein-Barr vírus specifikus T-sejtek	Nazális típusú extranodális NK/T-sejtes limfóma kezelése

¹ At the time of designation

Language	Active ingredient	Indication
Italian	Linfociti T autologhi specifici per il virus di Epstein-Barr, derivati da cellule mononucleari ematiche periferiche, espanse in vivo	Trattamento del linfoma T/NK extraganglionare, di tipo nasale
Latvian	Autologas pret Epšteina-Barra vīrusu specifiskas T šūnas, kas iegūtas no perifēro asiņu mononukleārajām šūnām un pavairotas ex vivo	Deguna tipa ekstranodālas NK/T-šūnu limfomas ārstēšana
Lithuanian	Autologinės, Epstein-Barr virusui specifinės T ląstelės, išskirtos iš periferinio kraujo mononuklearinių ląstelių, padaugintos ex vivo	Ekstranodalinės NK/T ląstelių limfomos (nazalinis tipas) gydymas
Maltese	Ċelluli T awtologuži speċifiċi għall-virus Epstein-Barr imnissla minn ċelluli mononukleari tad-demmi periferali mwassa' ex vivo	Kura tal-linfoma taċ-ċelluli NK/T extranodali, tat-tip nażali
Polish	Autologiczne komórki T specyficzne dla wirusa Epstein-Barr wyprowadzone z mononuklearnych komórek krwi obwodowej, ekspandowane ex vivo	Leczenie pozawęzłowego chłoniaka NK/T-komórkowego, typu nosowego
Portuguese	Células T autólogas específicas do vírus de Epstein-Barr derivadas de células mononucleares do sangue periférico, expandidas ex vivo	Tratamento de linfoma de células NK/T extranodal, de tipo nasal
Romanian	Celule T autologe specific pentru virusul Epstein-Barr derivate din celule mononucleare provenite din sângele periferic, expandate ex vivo	Tratamentul limfomului extranodal cu celule NK/T de tip nazal
Slovak	Autológne Epstein-Barr vírus špecifické T-bunky pochádzajúce z mononukleárných buniek periférnej krvi, expandované ex vivo	Liečba extranodálneho NK/T-bunkového lymfómu, nazálneho typu
Slovenian	Autologne, za virus Epstein-Barr specifične T-celice, ki izvirajo iz mononuklearnih celic periferne krvi, ekspandirane ex vivo	Zdravljenje ekstranodalnega NK/T-celičnega limfoma nazalnega tipa
Spanish	Celulas T Autologas expandido ex-vivo específicas para el virus Epstein Barr derivados de células mononucleares	Tratamiento del linfoma de célula NK/T extranodal tipo nasal
Swedish	Ex vivo expanderade autologa Epstein-Barrvirusspecifika T-celler från perifera mononukleära blodceller	Behandling av extranodalt NK/T-cellslymfom av nasal typ
Norwegian	Autologe Epstein-Barr virus spesifikke T-celler fra mononukleære celler i perifert blod, ekspandert ex vivo	Behandling av ekstranodalt NK/T-celle lymfom, nasal type
Icelandic	Samgena Epstein-Barr veiru sértækar T-frumur einangraðar úr einkjarna blóðfrumum og fjölgað ex vivo	Meðferð á NK/T-frumu eitlakrabbameini af nasal gerð