

## **ANNEX I**

### **SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Actos 15 mg tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 15 mg of pioglitazone (as hydrochloride).

Excipients:

Each tablet contains 92.87 mg of lactose monohydrate (see section 4.4).

For a full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Tablet.

The tablets are white to off-white, round, convex and marked '15' on one face and 'ACTOS' on the other face.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Pioglitazone is indicated as second or third line treatment of type 2 diabetes mellitus as described below:

as **monotherapy**

- in adult patients (particularly overweight patients) inadequately controlled by diet and exercise for whom metformin is inappropriate because of contraindications or intolerance

as **dual oral therapy** in combination with

- metformin, in adult patients (particularly overweight patients) with insufficient glycaemic control despite maximal tolerated dose of monotherapy with metformin
- a sulphonylurea, only in adult patients who show intolerance to metformin or for whom metformin is contraindicated, with insufficient glycaemic control despite maximal tolerated dose of monotherapy with a sulphonylurea.

as **triple oral therapy** in combination with

- metformin and a sulphonylurea, in adult patients (particularly overweight patients) with insufficient glycaemic control despite dual oral therapy.
- Pioglitazone is also indicated for combination with insulin in type 2 diabetes mellitus adult patients with insufficient glycaemic control on insulin for whom metformin is inappropriate because of contraindications or intolerance (see section 4.4).

After initiation of therapy with pioglitazone, patients should be reviewed after 3 to 6 months to assess adequacy of response to treatment (e.g. reduction in HbA1c). In patients who fail to show an adequate response, pioglitazone should be discontinued. In light of potential risks with prolonged therapy, prescribers should confirm at subsequent routine reviews that the benefit of pioglitazone is maintained (see section 4.4).

## **4.2 Posology and method of administration**

### Posology

Pioglitazone treatment may be initiated at 15 mg or 30 mg once daily. The dose may be increased in increments up to 45 mg once daily.

In combination with insulin, the current insulin dose can be continued upon initiation of pioglitazone therapy. If patients report hypoglycaemia, the dose of insulin should be decreased.

### Special population

#### *Elderly*

No dose adjustment is necessary for elderly patients (see section 5.2). Physicians should start treatment with the lowest available dose and increase the dose gradually, particularly when pioglitazone is used in combination with insulin (see section 4.4 Fluid retention and cardiac failure).

#### *Renal impairment*

No dose adjustment is necessary in patients with impaired renal function (creatinine clearance > 4 ml/min) (see section 5.2). No information is available from dialysed patients therefore pioglitazone should not be used in such patients.

#### *Hepatic impairment*

Pioglitazone should not be used in patients with hepatic impairment (see section 4.3 and 4.4).

### Paediatric population

The safety and efficacy of Actos in children and adolescents under 18 years of age have not been established. No data are available.

### Method of administration

Pioglitazone tablets are taken orally once daily with or without food. Tablets should be swallowed with a glass of water.

## **4.3 Contraindications**

Pioglitazone is contraindicated in patients with:

- hypersensitivity to the active substance or to any of the excipients
- cardiac failure or history of cardiac failure (NYHA stages I to IV)
- hepatic impairment
- diabetic ketoacidosis
- current bladder cancer or a history of bladder cancer
- uninvestigated macroscopic haematuria

## **4.4 Special warnings and precautions for use**

### *Fluid retention and cardiac failure*

Pioglitazone can cause fluid retention, which may exacerbate or precipitate heart failure. When treating patients who have at least one risk factor for development of congestive heart failure (e.g. prior myocardial infarction or symptomatic coronary artery disease or the elderly), physicians should start with the lowest available dose and increase the dose gradually. Patients should be observed for

signs and symptoms of heart failure, weight gain or oedema; particularly those with reduced cardiac reserve. There have been post-marketing cases of cardiac failure reported when pioglitazone was used in combination with insulin or in patients with a history of cardiac failure. Patients should be observed for signs and symptoms of heart failure, weight gain and oedema when pioglitazone is used in combination with insulin. Since insulin and pioglitazone are both associated with fluid retention, concomitant administration may increase the risk of oedema. Pioglitazone should be discontinued if any deterioration in cardiac status occurs.

A cardiovascular outcome study of pioglitazone has been performed in patients under 75 years with type 2 diabetes mellitus and pre-existing major macrovascular disease. Pioglitazone or placebo was added to existing antidiabetic and cardiovascular therapy for up to 3.5 years. This study showed an increase in reports of heart failure, however this did not lead to an increase in mortality in this study.

### *Elderly*

Combination use with insulin should be considered with caution in the elderly because of increased risk of serious heart failure.

In light of age-related risks (especially bladder cancer, fractures and heart failure), the balance of benefits and risks should be considered carefully both before and during treatment in the elderly.

### *Bladder Cancer*

Cases of bladder cancer were reported more frequently in a meta-analysis of controlled clinical trials with pioglitazone (19 cases from 12506 patients, 0.15%) than in control groups (7 cases from 10212 patients, 0.07%) HR=2.64 (95% CI 1.11-6.31, P=0.029). After excluding patients in whom exposure to study drug was less than one year at the time of diagnosis of bladder cancer, there were 7 cases (0.06%) on pioglitazone and 2 cases (0.02%) in control groups. Available epidemiological data also suggest a small increased risk of bladder cancer in diabetic patients treated with pioglitazone in particular in patients treated for the longest durations and with the highest cumulative doses. A possible risk after short term treatment cannot be excluded.

Risk factors for bladder cancer should be assessed before initiating pioglitazone treatment (risks include age, smoking history, exposure to some occupational or chemotherapy agents e.g. cyclophosphamide or prior radiation treatment in the pelvic region). Any macroscopic haematuria should be investigated before starting pioglitazone therapy.

Patients should be advised to promptly seek the attention of their physician if macroscopic haematuria or other symptoms such as dysuria or urinary urgency develop during treatment.

### *Monitoring of liver function*

There have been rare reports of hepatocellular dysfunction during post-marketing experience (see section 4.8). It is recommended, therefore, that patients treated with pioglitazone undergo periodic monitoring of liver enzymes. Liver enzymes should be checked prior to the initiation of therapy with pioglitazone in all patients. Therapy with pioglitazone should not be initiated in patients with increased baseline liver enzyme levels (ALT > 2.5 X upper limit of normal) or with any other evidence of liver disease.

Following initiation of therapy with pioglitazone, it is recommended that liver enzymes be monitored periodically based on clinical judgement. If ALT levels are increased to 3 X upper limit of normal during pioglitazone therapy, liver enzyme levels should be reassessed as soon as possible. If ALT levels remain > 3 X the upper limit of normal, therapy should be discontinued. If any patient develops symptoms suggesting hepatic dysfunction, which may include unexplained nausea, vomiting, abdominal pain, fatigue, anorexia and/or dark urine, liver enzymes should be checked. The decision whether to continue the patient on therapy with pioglitazone should be guided by clinical judgement pending laboratory evaluations. If jaundice is observed, the medicinal product should be discontinued.

### *Weight gain*

In clinical trials with pioglitazone there was evidence of dose related weight gain, which may be due to fat accumulation and in some cases associated with fluid retention. In some cases weight increase may be a symptom of cardiac failure, therefore weight should be closely monitored. Part of the treatment of diabetes is dietary control. Patients should be advised to adhere strictly to a calorie-controlled diet.

### *Haematology*

There was a small reduction in mean haemoglobin (4% relative reduction) and haematocrit (4.1% relative reduction) during therapy with pioglitazone, consistent with haemodilution. Similar changes were seen in metformin (haemoglobin 3-4% and haematocrit 3.6–4.1% relative reductions) and to a lesser extent sulphonylurea and insulin (haemoglobin 1–2% and haematocrit 1–3.2% relative reductions) treated patients in comparative controlled trials with pioglitazone.

### *Hypoglycaemia*

As a consequence of increased insulin sensitivity, patients receiving pioglitazone in dual or triple oral therapy with a sulphonylurea or in dual therapy with insulin may be at risk for dose-related hypoglycaemia, and a reduction in the dose of the sulphonylurea or insulin may be necessary.

### *Eye disorders*

Post-marketing reports of new-onset or worsening diabetic macular oedema with decreased visual acuity have been reported with thiazolidinediones, including pioglitazone. Many of these patients reported concurrent peripheral oedema. It is unclear whether or not there is a direct association between pioglitazone and macular oedema but prescribers should be alert to the possibility of macular oedema if patients report disturbances in visual acuity; an appropriate ophthalmological referral should be considered.

### *Others*

An increased incidence in bone fractures in women was seen in a pooled analysis of adverse reactions of bone fracture from randomised, controlled, double blind clinical trials in over 8100 pioglitazone and 7400 comparator treated patients, on treatment for up to 3.5 years.

Fractures were observed in 2.6% of women taking pioglitazone compared to 1.7% of women treated with a comparator. No increase in fracture rates was observed in men treated with pioglitazone (1.3%) versus comparator (1.5%).

The fracture incidence calculated was 1.9 fractures per 100 patient years in women treated with pioglitazone and 1.1 fractures per 100 patient years in women treated with a comparator. The observed excess risk of fractures for women in this dataset on pioglitazone is therefore 0.8 fractures per 100 patient years of use.

In the 3.5 year cardiovascular risk PROactive study, 44/870 (5.1%; 1.0 fractures per 100 patient years) of pioglitazone-treated female patients experienced fractures compared to 23/905 (2.5%; 0.5 fractures per 100 patient years) of female patients treated with comparator. No increase in fracture rates was observed in men treated with pioglitazone (1.7%) versus comparator (2.1%).

The risk of fractures should be considered in the long term care of women treated with pioglitazone.

As a consequence of enhancing insulin action, pioglitazone treatment in patients with polycystic ovarian syndrome may result in resumption of ovulation. These patients may be at risk of pregnancy.

Patients should be aware of the risk of pregnancy and if a patient wishes to become pregnant or if pregnancy occurs, the treatment should be discontinued (see section 4.6).

Pioglitazone should be used with caution during concomitant administration of cytochrome P450 2C8 inhibitors (e.g. gemfibrozil) or inducers (e.g. rifampicin). Glycaemic control should be monitored closely. Pioglitazone dose adjustment within the recommended posology or changes in diabetic treatment should be considered (see section 4.5).

Actos tablets contain lactose monohydrate and therefore should not be administered to patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Interaction studies have shown that pioglitazone has no relevant effect on either the pharmacokinetics or pharmacodynamics of digoxin, warfarin, phenprocoumon and metformin. Co-administration of pioglitazone with sulphonylureas does not appear to affect the pharmacokinetics of the sulphonylurea. Studies in man suggest no induction of the main inducible cytochrome P450, 1A, 2C8/9 and 3A4. *In vitro* studies have shown no inhibition of any subtype of cytochrome P450. Interactions with substances metabolised by these enzymes, e.g. oral contraceptives, cyclosporin, calcium channel blockers, and HMGCoA reductase inhibitors are not to be expected.

Co-administration of pioglitazone with gemfibrozil (an inhibitor of cytochrome P450 2C8) is reported to result in a 3-fold increase in AUC of pioglitazone. Since there is a potential for an increase in dose-related adverse events, a decrease in the dose of pioglitazone may be needed when gemfibrozil is concomitantly administered. Close monitoring of glycaemic control should be considered (see section 4.4). Co-administration of pioglitazone with rifampicin (an inducer of cytochrome P450 2C8) is reported to result in a 54% decrease in AUC of pioglitazone. The pioglitazone dose may need to be increased when rifampicin is concomitantly administered. Close monitoring of glycaemic control should be considered (see section 4.4).

#### **4.6 Fertility, pregnancy and lactation**

##### *Pregnancy*

There are no adequate human data to determine the safety of pioglitazone during pregnancy. Foetal growth restriction was apparent in animal studies with pioglitazone. This was attributable to the action of pioglitazone in diminishing the maternal hyperinsulinaemia and increased insulin resistance that occurs during pregnancy thereby reducing the availability of metabolic substrates for foetal growth. The relevance of such a mechanism in humans is unclear and pioglitazone should not be used in pregnancy.

##### *Breastfeeding*

Pioglitazone has been shown to be present in the milk of lactating rats. It is not known whether pioglitazone is secreted in human milk. Therefore, pioglitazone should not be administered to breast-feeding women.

##### *Fertility*

In animal fertility studies there was no effect on copulation, impregnation or fertility index.

#### **4.7 Effects on ability to drive and use machines**

Actos has no or negligible effect on the ability to drive and use machines. However patients who experience visual disturbance should be cautious when driving or using machines.

## 4.8 Undesirable effects

Adverse reactions reported in excess (> 0.5%) of placebo and as more than an isolated case in patients receiving pioglitazone in double-blind studies are listed below as MedDRA preferred term by system organ class and absolute frequency. Frequencies are defined as: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing incidence and seriousness.

| Adverse reaction                                                    | Frequency of adverse reactions of pioglitazone by treatment regimen |                |                    |                                  |              |
|---------------------------------------------------------------------|---------------------------------------------------------------------|----------------|--------------------|----------------------------------|--------------|
|                                                                     | Mono-therapy                                                        | Combination    |                    |                                  |              |
|                                                                     |                                                                     | with metformin | with sulphonylurea | with metformin and sulphonylurea | with insulin |
| <b>Infections and infestations</b>                                  |                                                                     |                |                    |                                  |              |
| upper respiratory tract infection                                   | common                                                              | common         | common             | common                           | common       |
| bronchitis                                                          |                                                                     |                |                    |                                  | common       |
| sinusitis                                                           | uncommon                                                            | uncommon       | uncommon           | uncommon                         | uncommon     |
| <b>Blood and lymphatic system disorders</b>                         |                                                                     |                |                    |                                  |              |
| anaemia                                                             |                                                                     | common         |                    |                                  |              |
| <b>Metabolism and nutrition disorders</b>                           |                                                                     |                |                    |                                  |              |
| hypo-glycaemia                                                      |                                                                     |                | uncommon           | very common                      | common       |
| appetite increased                                                  |                                                                     |                | uncommon           |                                  |              |
| <b>Nervous system disorders</b>                                     |                                                                     |                |                    |                                  |              |
| hypo-aesthesia                                                      | common                                                              | common         | common             | common                           | common       |
| headache                                                            |                                                                     | common         | uncommon           |                                  |              |
| dizziness                                                           |                                                                     |                | common             |                                  |              |
| insomnia                                                            | uncommon                                                            | uncommon       | uncommon           | uncommon                         | uncommon     |
| <b>Eye disorders</b>                                                |                                                                     |                |                    |                                  |              |
| visual disturbance <sup>1</sup>                                     | common                                                              | common         | uncommon           |                                  |              |
| macular oedema <sup>2</sup>                                         | not known                                                           | not known      | not known          | not known                        | not known    |
| <b>Ear and labyrinth disorders</b>                                  |                                                                     |                |                    |                                  |              |
| vertigo                                                             |                                                                     |                | uncommon           |                                  |              |
| <b>Cardiac disorders</b>                                            |                                                                     |                |                    |                                  |              |
| heart failure <sup>3</sup>                                          |                                                                     |                |                    |                                  | common       |
| <b>Neoplasms benign, malignant and unspecified (including cysts</b> |                                                                     |                |                    |                                  |              |

| Adverse reaction                                            | Frequency of adverse reactions of pioglitazone by treatment regimen |                |                    |                                  |              |
|-------------------------------------------------------------|---------------------------------------------------------------------|----------------|--------------------|----------------------------------|--------------|
|                                                             | Mono-therapy                                                        | Combination    |                    |                                  |              |
|                                                             |                                                                     | with metformin | with sulphonylurea | with metformin and sulphonylurea | with insulin |
| <b>and polyps)</b>                                          |                                                                     |                |                    |                                  |              |
| bladder cancer                                              | uncommon                                                            | uncommon       | uncommon           | uncommon                         | uncommon     |
| <b>Respiratory, thoracic and mediastinal disorders</b>      |                                                                     |                |                    |                                  |              |
| dyspnoea                                                    |                                                                     |                |                    |                                  | common       |
| <b>Gastrointestinal disorders</b>                           |                                                                     |                |                    |                                  |              |
| flatulence                                                  |                                                                     | uncommon       | common             |                                  |              |
| <b>Skin and subcutaneous tissue disorders</b>               |                                                                     |                |                    |                                  |              |
| sweating                                                    |                                                                     |                | uncommon           |                                  |              |
| <b>Musculoskeletal and connective tissue disorders</b>      |                                                                     |                |                    |                                  |              |
| fracture bone <sup>4</sup>                                  | common                                                              | common         | common             | common                           | common       |
| arthralgia                                                  |                                                                     | common         |                    | common                           | common       |
| back pain                                                   |                                                                     |                |                    |                                  | common       |
| <b>Renal and urinary disorders</b>                          |                                                                     |                |                    |                                  |              |
| haematuria                                                  |                                                                     | common         |                    |                                  |              |
| glycosuria                                                  |                                                                     |                | uncommon           |                                  |              |
| proteinuria                                                 |                                                                     |                | uncommon           |                                  |              |
| <b>Reproductive system and breast disorders</b>             |                                                                     |                |                    |                                  |              |
| erectile dysfunction                                        |                                                                     | common         |                    |                                  |              |
| <b>General disorders and administration site conditions</b> |                                                                     |                |                    |                                  |              |
| oedema                                                      |                                                                     |                |                    |                                  | very common  |
| fatigue                                                     |                                                                     |                | uncommon           |                                  |              |
| <b>Investigations</b>                                       |                                                                     |                |                    |                                  |              |
| weight increased <sup>5</sup>                               | common                                                              | common         | common             | common                           | common       |
| blood creatine phospho-kinase increased                     |                                                                     |                |                    | common                           |              |
| increased lactic dehydro-genase                             |                                                                     |                | uncommon           |                                  |              |
| Alanine aminotransferase                                    | not known                                                           | not known      | not known          | not known                        | not known    |

| Adverse reaction       | Frequency of adverse reactions of pioglitazone by treatment regimen |                |                    |                                  |              |
|------------------------|---------------------------------------------------------------------|----------------|--------------------|----------------------------------|--------------|
|                        | Mono-therapy                                                        | Combination    |                    |                                  |              |
|                        |                                                                     | with metformin | with sulphonylurea | with metformin and sulphonylurea | with insulin |
| increased <sup>6</sup> |                                                                     |                |                    |                                  |              |

<sup>1</sup> Visual disturbance has been reported mainly early in treatment and is related to changes in blood glucose due to temporary alteration in the turgidity and refractive index of the lens as seen with other hypoglycaemic treatments.

<sup>2</sup> Oedema was reported in 6–9% of patients treated with pioglitazone over one year in controlled clinical trials. The oedema rates for comparator groups (sulphonylurea, metformin) were 2–5%. The reports of oedema were generally mild to moderate and usually did not require discontinuation of treatment.

<sup>3</sup> In controlled clinical trials the incidence of reports of heart failure with pioglitazone treatment was the same as in placebo, metformin and sulphonylurea treatment groups, but was increased when used in combination therapy with insulin. In an outcome study of patients with pre-existing major macrovascular disease, the incidence of serious heart failure was 1.6% higher with pioglitazone than with placebo, when added to therapy that included insulin. However, this did not lead to an increase in mortality in this study. Heart failure has been reported rarely with marketing use of pioglitazone, but more frequently when pioglitazone was used in combination with insulin or in patients with a history of cardiac failure.

<sup>4</sup> A pooled analysis was conducted of adverse reactions of bone fractures from randomised, comparator controlled, double blind clinical trials in over 8100 patients in the pioglitazone-treated groups and 7400 in the comparator-treated groups of up to 3.5 years duration. A higher rate of fractures was observed in women taking pioglitazone (2.6%) versus comparator (1.7%). No increase in fracture rates was observed in men treated with pioglitazone (1.3%) versus comparator (1.5%). In the 3.5 year PROactive study, 44/870 (5.1%) of pioglitazone-treated female patients experienced fractures compared to 23/905 (2.5%) of female patients treated with comparator. No increase in fracture rates was observed in men treated with pioglitazone (1.7%) versus comparator (2.1%).

<sup>5</sup> In active comparator controlled trials mean weight increase with pioglitazone given as monotherapy was 2–3 kg over one year. This is similar to that seen in a sulphonylurea active comparator group. In combination trials pioglitazone added to metformin resulted in mean weight increase over one year of 1.5 kg and added to a sulphonylurea of 2.8 kg. In comparator groups addition of sulphonylurea to metformin resulted in a mean weight gain of 1.3 kg and addition of metformin to a sulphonylurea a mean weight loss of 1.0 kg.

<sup>6</sup> In clinical trials with pioglitazone the incidence of elevations of ALT greater than three times the upper limit of normal was equal to placebo but less than that seen in metformin or sulphonylurea comparator groups. Mean levels of liver enzymes decreased with treatment with pioglitazone. Rare cases of elevated liver enzymes and hepatocellular dysfunction have occurred in post-marketing experience. Although in very rare cases fatal outcome has been reported, causal relationship has not been established.

## 4.9 Overdose

In clinical studies, patients have taken pioglitazone at higher than the recommended highest dose of 45 mg daily. The maximum reported dose of 120 mg/day for four days, then 180 mg/day for seven days was not associated with any symptoms.

Hypoglycaemia may occur in combination with sulphonylureas or insulin. Symptomatic and general supportive measures should be taken in case of overdose.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Drugs used in diabetes, blood glucose lowering drugs, excl. insulins; ATC code: A10BG03.

Pioglitazone effects may be mediated by a reduction of insulin resistance. Pioglitazone appears to act via activation of specific nuclear receptors (peroxisome proliferator activated receptor gamma) leading to increased insulin sensitivity of liver, fat and skeletal muscle cells in animals. Treatment with pioglitazone has been shown to reduce hepatic glucose output and to increase peripheral glucose disposal in the case of insulin resistance.

Fasting and postprandial glycaemic control is improved in patients with type 2 diabetes mellitus. The improved glycaemic control is associated with a reduction in both fasting and postprandial plasma insulin concentrations. A clinical trial of pioglitazone vs. gliclazide as monotherapy was extended to two years in order to assess time to treatment failure (defined as appearance of  $HbA_{1c} \geq 8.0\%$  after the first six months of therapy). Kaplan-Meier analysis showed shorter time to treatment failure in patients treated with gliclazide, compared with pioglitazone. At two years, glycaemic control (defined as  $HbA_{1c} < 8.0\%$ ) was sustained in 69% of patients treated with pioglitazone, compared with 50% of patients on gliclazide. In a two-year study of combination therapy comparing pioglitazone with gliclazide when added to metformin, glycaemic control measured as mean change from baseline in  $HbA_{1c}$  was similar between treatment groups after one year. The rate of deterioration of  $HbA_{1c}$  during the second year was less with pioglitazone than with gliclazide.

In a placebo controlled trial, patients with inadequate glycaemic control despite a three month insulin optimisation period were randomised to pioglitazone or placebo for 12 months. Patients receiving pioglitazone had a mean reduction in  $HbA_{1c}$  of 0.45% compared with those continuing on insulin alone, and a reduction of insulin dose in the pioglitazone treated group.

HOMA analysis shows that pioglitazone improves beta cell function as well as increasing insulin sensitivity. Two-year clinical studies have shown maintenance of this effect.

In one year clinical trials, pioglitazone consistently gave a statistically significant reduction in the albumin/creatinine ratio compared to baseline.

The effect of pioglitazone (45 mg monotherapy vs. placebo) was studied in a small 18-week trial in type 2 diabetics. Pioglitazone was associated with significant weight gain. Visceral fat was significantly decreased, while there was an increase in extra-abdominal fat mass. Similar changes in body fat distribution on pioglitazone have been accompanied by an improvement in insulin sensitivity. In most clinical trials, reduced total plasma triglycerides and free fatty acids, and increased HDL-cholesterol levels were observed as compared to placebo, with small, but not clinically significant increases in LDL-cholesterol levels.

In clinical trials of up to two years duration, pioglitazone reduced total plasma triglycerides and free fatty acids, and increased HDL cholesterol levels, compared with placebo, metformin or gliclazide. Pioglitazone did not cause statistically significant increases in LDL cholesterol levels compared with placebo, whilst reductions were observed with metformin and gliclazide. In a 20-week study, as well as reducing fasting triglycerides, pioglitazone reduced post prandial hypertriglyceridaemia through an effect on both absorbed and hepatically synthesised triglycerides. These effects were independent of pioglitazone's effects on glycaemia and were statistically significant different to glibenclamide.

In PROactive, a cardiovascular outcome study, 5238 patients with type 2 diabetes mellitus and pre-existing major macrovascular disease were randomised to pioglitazone or placebo in addition to existing antidiabetic and cardiovascular therapy, for up to 3.5 years. The study population had an average age of 62 years; the average duration of diabetes was 9.5 years. Approximately one third of patients were receiving insulin in combination with metformin and/or a sulphonylurea. To be eligible patients had to have had one or more of the following: myocardial infarction, stroke, percutaneous cardiac intervention or coronary artery bypass graft, acute coronary syndrome, coronary artery disease, or peripheral arterial obstructive disease. Almost half of the patients had a previous myocardial infarction and approximately 20% had had a stroke. Approximately half of the study population had at least two of the cardiovascular history entry criteria. Almost all subjects (95%) were receiving cardiovascular medicinal products (beta blockers, ACE inhibitors, angiotensin II antagonists, calcium channel blockers, nitrates, diuretics, aspirin, statins, fibrates).

Although the study failed regarding its primary endpoint, which was a composite of all-cause mortality, non-fatal myocardial infarction, stroke, acute coronary syndrome, major leg amputation, coronary revascularisation and leg revascularisation, the results suggest that there are no long-term cardiovascular concerns regarding use of pioglitazone. However, the incidences of oedema, weight gain and heart failure were increased. No increase in mortality from heart failure was observed.

#### *Paediatric population*

The European Medicines Agency has waived the obligation to submit the results of studies with Actos in all subsets of the paediatric population in Type 2 Diabetes Mellitus. See section 4.2 for information on paediatric use.

## **5.2 Pharmacokinetic properties**

### *Absorption*

Following oral administration, pioglitazone is rapidly absorbed, and peak plasma concentrations of unchanged pioglitazone are usually achieved 2 hours after administration. Proportional increases of the plasma concentration were observed for doses from 2–60 mg. Steady state is achieved after 4–7 days of dosing. Repeated dosing does not result in accumulation of the compound or metabolites. Absorption is not influenced by food intake. Absolute bioavailability is greater than 80%.

### *Distribution*

The estimated volume of distribution in humans is 0.25 l/kg.

Pioglitazone and all active metabolites are extensively bound to plasma protein (> 99%).

### *Biotransformation*

Pioglitazone undergoes extensive hepatic metabolism by hydroxylation of aliphatic methylene groups. This is predominantly via cytochrome P450 2C8 although other isoforms may be involved to a lesser degree. Three of the six identified metabolites are active (M-II, M-III, and M-IV). When activity, concentrations and protein binding are taken into account, pioglitazone and metabolite M-III contribute equally to efficacy. On this basis M-IV contribution to efficacy is approximately three-fold that of pioglitazone, whilst the relative efficacy of M-II is minimal.

*In vitro* studies have shown no evidence that pioglitazone inhibits any subtype of cytochrome P450. There is no induction of the main inducible P450 isoenzymes 1A, 2C8/9, and 3A4 in man.

Interaction studies have shown that pioglitazone has no relevant effect on either the pharmacokinetics or pharmacodynamics of digoxin, warfarin, phenprocoumon and metformin. Concomitant administration of pioglitazone with gemfibrozil (an inhibitor of cytochrome P450 2C8) or with

rifampicin (an inducer of cytochrome P450 2C8) is reported to increase or decrease, respectively, the plasma concentration of pioglitazone (see section 4.5).

### *Elimination*

Following oral administration of radiolabelled pioglitazone to man, recovered label was mainly in faeces (55%) and a lesser amount in urine (45%). In animals, only a small amount of unchanged pioglitazone can be detected in either urine or faeces. The mean plasma elimination half-life of unchanged pioglitazone in man is 5 to 6 hours and for its total active metabolites 16 to 23 hours.

### *Elderly*

Steady state pharmacokinetics are similar in patients age 65 and over and young subjects.

### *Patients with renal impairment*

In patients with renal impairment, plasma concentrations of pioglitazone and its metabolites are lower than those seen in subjects with normal renal function, but oral clearance of parent substance is similar. Thus free (unbound) pioglitazone concentration is unchanged.

### *Patients with hepatic impairment*

Total plasma concentration of pioglitazone is unchanged, but with an increased volume of distribution. Intrinsic clearance is therefore reduced, coupled with a higher unbound fraction of pioglitazone.

## **5.3 Preclinical safety data**

In toxicology studies, plasma volume expansion with haemodilution, anaemia, and reversible eccentric cardiac hypertrophy was consistently apparent after repeated dosing of mice, rats, dogs, and monkeys. In addition, increased fatty deposition and infiltration were observed. These findings were observed across species at plasma concentrations  $\leq 4$  times the clinical exposure. Foetal growth restriction was apparent in animal studies with pioglitazone. This was attributable to the action of pioglitazone in diminishing the maternal hyperinsulinaemia and increased insulin resistance that occurs during pregnancy thereby reducing the availability of metabolic substrates for foetal growth.

Pioglitazone was devoid of genotoxic potential in a comprehensive battery of *in vivo* and *in vitro* genotoxicity assays. An increased incidence of hyperplasia (males and females) and tumours (males) of the urinary bladder epithelium was apparent in rats treated with pioglitazone for up to 2 years.

The formation and presence of urinary calculi with subsequent irritation and hyperplasia was postulated as the mechanistic basis for the observed tumourigenic response in the male rat. A 24-month mechanistic study in male rats demonstrated that administration of pioglitazone resulted in an increased incidence of hyperplastic changes in the bladder. Dietary acidification significantly decreased but did not abolish the incidence of tumours. The presence of microcrystals exacerbated the hyperplastic response but was not considered to be the primary cause of hyperplastic changes. The relevance to humans of the tumourigenic findings in the male rat cannot be excluded.

There was no tumorigenic response in mice of either sex. Hyperplasia of the urinary bladder was not seen in dogs or monkeys treated with pioglitazone for up to 12 months.

In an animal model of familial adenomatous polyposis (FAP), treatment with two other thiazolidinediones increased tumour multiplicity in the colon. The relevance of this finding is unknown.

Environmental Risk Assessment: no environmental impact is anticipated from the clinical use of pioglitazone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Carmellose calcium  
Hyprolose  
Lactose monohydrate  
Magnesium stearate

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years.

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

Aluminium/aluminium blisters, packs of 14, 28, 30, 50, 56, 84, 90, 98, 112 and 196 tablets.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

No special requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Takeda Global Research and Development Centre (Europe) Ltd  
61 Aldwych  
London  
WC2B 4AE  
United Kingdom

## **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/150/001  
EU/1/00/150/002  
EU/1/00/150/003  
EU/1/00/150/007  
EU/1/00/150/009  
EU/1/00/150/016  
EU/1/00/150/017  
EU/1/00/150/018  
EU/1/00/150/025  
EU/1/00/150/026

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 13/10/2000

Date of last renewal: 31/08/2010

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>

## 1. NAME OF THE MEDICINAL PRODUCT

Actos 30 mg tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 30 mg of pioglitazone (as hydrochloride).

Excipients:

Each tablet contains 76.34 mg of lactose monohydrate (see section 4.4).

For a full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Tablet.

The tablets are white to off-white, round, flat and marked '30' on one face and 'ACTOS' on the other face.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Pioglitazone is indicated as second or third line treatment of type 2 diabetes mellitus as described below:

as **monotherapy**

- in adult patients (particularly overweight patients) inadequately controlled by diet and exercise for whom metformin is inappropriate because of contraindications or intolerance

as **dual oral therapy** in combination with

- metformin, in adult patients (particularly overweight patients) with insufficient glycaemic control despite maximal tolerated dose of monotherapy with metformin
- a sulphonylurea, only in adult patients who show intolerance to metformin or for whom metformin is contraindicated, with insufficient glycaemic control despite maximal tolerated dose of monotherapy with a sulphonylurea.

as **triple oral therapy** in combination with

- metformin and a sulphonylurea, in adult patients (particularly overweight patients) with insufficient glycaemic control despite dual oral therapy.
- Pioglitazone is also indicated for combination with insulin in type 2 diabetes mellitus adult patients with insufficient glycaemic control on insulin for whom metformin is inappropriate because of contraindications or intolerance (see section 4.4).

After initiation of therapy with pioglitazone, patients should be reviewed after 3 to 6 months to assess adequacy of response to treatment (e.g. reduction in HbA1c). In patients who fail to show an adequate response, pioglitazone should be discontinued. In light of potential risks with prolonged therapy, prescribers should confirm at subsequent routine reviews that the benefit of pioglitazone is maintained (see section 4.4).

## **4.2 Posology and method of administration**

### Posology

Pioglitazone treatment may be initiated at 15 mg or 30 mg once daily. The dose may be increased in increments up to 45 mg once daily.

In combination with insulin, the current insulin dose can be continued upon initiation of pioglitazone therapy. If patients report hypoglycaemia, the dose of insulin should be decreased.

### Special population

#### *Elderly*

No dose adjustment is necessary for elderly patients (see section 5.2). Physicians should start treatment with the lowest available dose and increase the dose gradually, particularly when pioglitazone is used in combination with insulin (see section 4.4 Fluid retention and cardiac failure).

#### *Renal impairment*

No dose adjustment is necessary in patients with impaired renal function (creatinine clearance > 4 ml/min) (see section 5.2). No information is available from dialysed patients therefore pioglitazone should not be used in such patients.

#### *Hepatic impairment*

Pioglitazone should not be used in patients with hepatic impairment (see section 4.3 and 4.4).

### Paediatric population

The safety and efficacy of Actos in children and adolescents under 18 years of age have not been established. No data are available.

### Method of administration

Pioglitazone tablets are taken orally once daily with or without food. Tablets should be swallowed with a glass of water.

## **4.3 Contraindications**

Pioglitazone is contraindicated in patients with:

- hypersensitivity to the active substance or to any of the excipients
- cardiac failure or history of cardiac failure (NYHA stages I to IV)
- hepatic impairment
- diabetic ketoacidosis
- current bladder cancer or a history of bladder cancer
- uninvestigated macroscopic haematuria

## **4.4 Special warnings and precautions for use**

### *Fluid retention and cardiac failure*

Pioglitazone can cause fluid retention, which may exacerbate or precipitate heart failure. When treating patients who have at least one risk factor for development of congestive heart failure (e.g. prior myocardial infarction or symptomatic coronary artery disease or the elderly), physicians should start with the lowest available dose and increase the dose gradually. Patients should be observed for signs and symptoms of heart failure, weight gain or oedema; particularly those with reduced cardiac

reserve. There have been post-marketing cases of cardiac failure reported when pioglitazone was used in combination with insulin or in patients with a history of cardiac failure. Patients should be observed for signs and symptoms of heart failure, weight gain and oedema when pioglitazone is used in combination with insulin. Since insulin and pioglitazone are both associated with fluid retention, concomitant administration may increase the risk of oedema. Pioglitazone should be discontinued if any deterioration in cardiac status occurs.

A cardiovascular outcome study of pioglitazone has been performed in patients under 75 years with type 2 diabetes mellitus and pre-existing major macrovascular disease. Pioglitazone or placebo was added to existing antidiabetic and cardiovascular therapy for up to 3.5 years. This study showed an increase in reports of heart failure, however this did not lead to an increase in mortality in this study.

### *Elderly*

Combination use with insulin should be considered with caution in the elderly because of increased risk of serious heart failure.

In light of age-related risks (especially bladder cancer, fractures and heart failure), the balance of benefits and risks should be considered carefully both before and during treatment in the elderly.

### *Bladder cancer*

Cases of bladder cancer were reported more frequently in a meta-analysis of controlled clinical trials with pioglitazone (19 cases from 12506 patients, 0.15%) than in control groups (7 cases from 10212 patients, 0.07%) HR=2.64 (95% CI 1.11-6.31, P=0.029). After excluding patients in whom exposure to study drug was less than one year at the time of diagnosis of bladder cancer, there were 7 cases (0.06%) on pioglitazone and 2 cases (0.02%) in control groups. Available epidemiological data also suggest a small increased risk of bladder cancer in diabetic patients treated with pioglitazone in particular in patients treated for the longest durations and with the highest cumulative doses. A possible risk after short term treatment cannot be excluded.

Risk factors for bladder cancer should be assessed before initiating pioglitazone treatment (risks include age, smoking history, exposure to some occupational or chemotherapy agents e.g. cyclophosphamide or prior radiation treatment in the pelvic region). Any macroscopic haematuria should be investigated before starting pioglitazone therapy.

Patients should be advised to promptly seek the attention of their physician if macroscopic haematuria or other symptoms such as dysuria or urinary urgency develop during treatment.

### *Monitoring of liver function*

There have been rare reports of hepatocellular dysfunction during post-marketing experience (see section 4.8). It is recommended, therefore, that patients treated with pioglitazone undergo periodic monitoring of liver enzymes. Liver enzymes should be checked prior to the initiation of therapy with pioglitazone in all patients. Therapy with pioglitazone should not be initiated in patients with increased baseline liver enzyme levels (ALT > 2.5 X upper limit of normal) or with any other evidence of liver disease.

Following initiation of therapy with pioglitazone, it is recommended that liver enzymes be monitored periodically based on clinical judgement. If ALT levels are increased to 3 X upper limit of normal during pioglitazone therapy, liver enzyme levels should be reassessed as soon as possible. If ALT levels remain > 3 X the upper limit of normal, therapy should be discontinued. If any patient develops symptoms suggesting hepatic dysfunction, which may include unexplained nausea, vomiting, abdominal pain, fatigue, anorexia and/or dark urine, liver enzymes should be checked. The decision whether to continue the patient on therapy with pioglitazone should be guided by clinical judgement pending laboratory evaluations. If jaundice is observed, the medicinal product should be discontinued.

### *Weight gain*

In clinical trials with pioglitazone there was evidence of dose related weight gain, which may be due to fat accumulation and in some cases associated with fluid retention. In some cases weight increase may be a symptom of cardiac failure, therefore weight should be closely monitored. Part of the treatment of diabetes is dietary control. Patients should be advised to adhere strictly to a calorie-controlled diet.

### *Haematology*

There was a small reduction in mean haemoglobin (4% relative reduction) and haematocrit (4.1% relative reduction) during therapy with pioglitazone, consistent with haemodilution. Similar changes were seen in metformin (haemoglobin 3-4% and haematocrit 3.6-4.1% relative reductions) and to a lesser extent sulphonylurea and insulin (haemoglobin 1-2% and haematocrit 1-3.2% relative reductions) treated patients in comparative controlled trials with pioglitazone.

### *Hypoglycaemia*

As a consequence of increased insulin sensitivity, patients receiving pioglitazone in dual or triple oral therapy with a sulphonylurea or in dual therapy with insulin may be at risk for dose-related hypoglycaemia, and a reduction in the dose of the sulphonylurea or insulin may be necessary.

### *Eye disorders*

Post-marketing reports of new-onset or worsening diabetic macular oedema with decreased visual acuity have been reported with thiazolidinediones, including pioglitazone. Many of these patients reported concurrent peripheral oedema. It is unclear whether or not there is a direct association between pioglitazone and macular oedema but prescribers should be alert to the possibility of macular oedema if patients report disturbances in visual acuity; an appropriate ophthalmological referral should be considered.

### *Others*

An increased incidence in bone fractures in women was seen in a pooled analysis of adverse reactions of bone fracture from randomised, controlled, double blind clinical trials in over 8100 pioglitazone and 7400 comparator treated patients, on treatment for up to 3.5 years.

Fractures were observed in 2.6% of women taking pioglitazone compared to 1.7% of women treated with a comparator. No increase in fracture rates was observed in men treated with pioglitazone (1.3%) versus comparator (1.5%).

The fracture incidence calculated was 1.9 fractures per 100 patient years in women treated with pioglitazone and 1.1 fractures per 100 patient years in women treated with a comparator. The observed excess risk of fractures for women in this dataset on pioglitazone is therefore 0.8 fractures per 100 patient years of use.

In the 3.5 year cardiovascular risk PROactive study, 44/870 (5.1%; 1.0 fractures per 100 patient years) of pioglitazone-treated female patients experienced fractures compared to 23/905 (2.5%; 0.5 fractures per 100 patient years) of female patients treated with comparator. No increase in fracture rates was observed in men treated with pioglitazone (1.7%) versus comparator (2.1%).

The risk of fractures should be considered in the long term care of women treated with pioglitazone.

As a consequence of enhancing insulin action, pioglitazone treatment in patients with polycystic ovarian syndrome may result in resumption of ovulation. These patients may be at risk of pregnancy. Patients should be aware of the risk of pregnancy and if a patient wishes to become pregnant or if pregnancy occurs, the treatment should be discontinued (see section 4.6).

Pioglitazone should be used with caution during concomitant administration of cytochrome P450 2C8 inhibitors (e.g. gemfibrozil) or inducers (e.g. rifampicin). Glycaemic control should be monitored closely. Pioglitazone dose adjustment within the recommended posology or changes in diabetic treatment should be considered (see section 4.5).

Actos tablets contain lactose monohydrate and therefore should not be administered to patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Interaction studies have shown that pioglitazone has no relevant effect on either the pharmacokinetics or pharmacodynamics of digoxin, warfarin, phenprocoumon and metformin. Co-administration of pioglitazone with sulphonylureas does not appear to affect the pharmacokinetics of the sulphonylurea. Studies in man suggest no induction of the main inducible cytochrome P450, 1A, 2C8/9 and 3A4. *In vitro* studies have shown no inhibition of any subtype of cytochrome P450. Interactions with substances metabolised by these enzymes, e.g. oral contraceptives, cyclosporin, calcium channel blockers, and HMGCoA reductase inhibitors are not to be expected.

Co-administration of pioglitazone with gemfibrozil (an inhibitor of cytochrome P450 2C8) is reported to result in a 3-fold increase in AUC of pioglitazone. Since there is a potential for an increase in dose-related adverse events, a decrease in the dose of pioglitazone may be needed when gemfibrozil is concomitantly administered. Close monitoring of glycaemic control should be considered (see section 4.4). Co-administration of pioglitazone with rifampicin (an inducer of cytochrome P450 2C8) is reported to result in a 54% decrease in AUC of pioglitazone. The pioglitazone dose may need to be increased when rifampicin is concomitantly administered. Close monitoring of glycaemic control should be considered (see section 4.4).

#### **4.6 Fertility, pregnancy and lactation**

##### *Pregnancy*

There are no adequate human data to determine the safety of pioglitazone during pregnancy. Foetal growth restriction was apparent in animal studies with pioglitazone. This was attributable to the action of pioglitazone in diminishing the maternal hyperinsulinaemia and increased insulin resistance that occurs during pregnancy thereby reducing the availability of metabolic substrates for foetal growth. The relevance of such a mechanism in humans is unclear and pioglitazone should not be used in pregnancy.

##### *Breastfeeding*

Pioglitazone has been shown to be present in the milk of lactating rats. It is not known whether pioglitazone is secreted in human milk. Therefore, pioglitazone should not be administered to breast-feeding women.

##### *Fertility*

In animal fertility studies there was no effect on copulation, impregnation or fertility index.

#### **4.7 Effects on ability to drive and use machines**

Actos has no or negligible effect on the ability to drive and use machines. However patients who experience visual disturbance should be cautious when driving or using machines.

## 4.8 Undesirable effects

Adverse reactions reported in excess ( $\geq 0.5\%$ ) of placebo and as more than an isolated case in patients receiving pioglitazone in double-blind studies are listed below as MedDRA preferred term by system organ class and absolute frequency. Frequencies are defined as: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing incidence and seriousness.

| Adverse reaction                                   | Frequency of adverse reactions of pioglitazone by treatment regimen |                |                    |                                  |              |
|----------------------------------------------------|---------------------------------------------------------------------|----------------|--------------------|----------------------------------|--------------|
|                                                    | Mono-therapy                                                        | Combination    |                    |                                  |              |
|                                                    |                                                                     | with metformin | with sulphonylurea | with metformin and sulphonylurea | with insulin |
| <b>Infections and infestations</b>                 |                                                                     |                |                    |                                  |              |
| upper respiratory tract infection                  | common                                                              | common         | common             | common                           | common       |
| bronchitis                                         |                                                                     |                |                    |                                  | common       |
| sinusitis                                          | uncommon                                                            | uncommon       | uncommon           | uncommon                         | uncommon     |
| <b>Blood and lymphatic system disorders</b>        |                                                                     |                |                    |                                  |              |
| anaemia                                            |                                                                     | common         |                    |                                  |              |
| <b>Metabolism and nutrition disorders</b>          |                                                                     |                |                    |                                  |              |
| hypo-glycaemia                                     |                                                                     |                | uncommon           | very common                      | common       |
| appetite increased                                 |                                                                     |                | uncommon           |                                  |              |
| <b>Nervous system disorders</b>                    |                                                                     |                |                    |                                  |              |
| hypo-aesthesia                                     | common                                                              | common         | common             | common                           | common       |
| headache                                           |                                                                     | common         | uncommon           |                                  |              |
| dizziness                                          |                                                                     |                | common             |                                  |              |
| insomnia                                           | uncommon                                                            | uncommon       | uncommon           | uncommon                         | uncommon     |
| <b>Eye disorders</b>                               |                                                                     |                |                    |                                  |              |
| visual disturbance <sup>1</sup>                    | common                                                              | common         | uncommon           |                                  |              |
| macular oedema <sup>2</sup>                        | not known                                                           | not known      | not known          | not known                        | not known    |
| <b>Ear and labyrinth disorders</b>                 |                                                                     |                |                    |                                  |              |
| vertigo                                            |                                                                     |                | uncommon           |                                  |              |
| <b>Cardiac disorders</b>                           |                                                                     |                |                    |                                  |              |
| heart failure <sup>3</sup>                         |                                                                     |                |                    |                                  | common       |
| <b>Neoplasms benign, malignant and unspecified</b> |                                                                     |                |                    |                                  |              |

| Adverse reaction                                            | Frequency of adverse reactions of pioglitazone by treatment regimen |                |                    |                                  |              |
|-------------------------------------------------------------|---------------------------------------------------------------------|----------------|--------------------|----------------------------------|--------------|
|                                                             | Mono-therapy                                                        | Combination    |                    |                                  |              |
|                                                             |                                                                     | with metformin | with sulphonylurea | with metformin and sulphonylurea | with insulin |
| <b>(including cysts and polyps)</b>                         |                                                                     |                |                    |                                  |              |
| bladder cancer                                              | uncommon                                                            | uncommon       | uncommon           | uncommon                         | uncommon     |
| <b>Respiratory, thoracic and mediastinal disorders</b>      |                                                                     |                |                    |                                  |              |
| dyspnoea                                                    |                                                                     |                |                    |                                  | common       |
| <b>Gastrointestinal disorders</b>                           |                                                                     |                |                    |                                  |              |
| flatulence                                                  |                                                                     | uncommon       | common             |                                  |              |
| <b>Skin and subcutaneous tissue disorders</b>               |                                                                     |                |                    |                                  |              |
| sweating                                                    |                                                                     |                | uncommon           |                                  |              |
| <b>Musculoskeletal and connective tissue disorders</b>      |                                                                     |                |                    |                                  |              |
| fracture bone <sup>4</sup>                                  | common                                                              | common         | common             | common                           | common       |
| arthralgia                                                  |                                                                     | common         |                    | common                           | common       |
| back pain                                                   |                                                                     |                |                    |                                  | common       |
| <b>Renal and urinary disorders</b>                          |                                                                     |                |                    |                                  |              |
| haematuria                                                  |                                                                     | common         |                    |                                  |              |
| glycosuria                                                  |                                                                     |                | uncommon           |                                  |              |
| proteinuria                                                 |                                                                     |                | uncommon           |                                  |              |
| <b>Reproductive system and breast disorders</b>             |                                                                     |                |                    |                                  |              |
| erectile dysfunction                                        |                                                                     | common         |                    |                                  |              |
| <b>General disorders and administration site conditions</b> |                                                                     |                |                    |                                  |              |
| oedema                                                      |                                                                     |                |                    |                                  | very common  |
| fatigue                                                     |                                                                     |                | uncommon           |                                  |              |
| <b>Investigations</b>                                       |                                                                     |                |                    |                                  |              |
| weight increased <sup>5</sup>                               | common                                                              | common         | common             | common                           | common       |
| blood creatine phospho-kinase increased                     |                                                                     |                |                    | common                           |              |
| increased lactic dehydro-genase                             |                                                                     |                | uncommon           |                                  |              |

| Adverse reaction                                | Frequency of adverse reactions of pioglitazone by treatment regimen |                |                    |                                  |              |
|-------------------------------------------------|---------------------------------------------------------------------|----------------|--------------------|----------------------------------|--------------|
|                                                 | Mono-therapy                                                        | Combination    |                    |                                  |              |
|                                                 |                                                                     | with metformin | with sulphonylurea | with metformin and sulphonylurea | with insulin |
| Alanine aminotransferase increased <sup>6</sup> | not known                                                           | not known      | not known          | not known                        | not known    |

<sup>1</sup> Visual disturbance has been reported mainly early in treatment and is related to changes in blood glucose due to temporary alteration in the turgidity and refractive index of the lens as seen with other hypoglycaemic treatments.

<sup>2</sup> Oedema was reported in 6–9% of patients treated with pioglitazone over one year in controlled clinical trials. The oedema rates for comparator groups (sulphonylurea, metformin) were 2–5%. The reports of oedema were generally mild to moderate and usually did not require discontinuation of treatment.

<sup>3</sup> In controlled clinical trials the incidence of reports of heart failure with pioglitazone treatment was the same as in placebo, metformin and sulphonylurea treatment groups, but was increased when used in combination therapy with insulin. In an outcome study of patients with pre-existing major macrovascular disease, the incidence of serious heart failure was 1.6% higher with pioglitazone than with placebo, when added to therapy that included insulin. However, this did not lead to an increase in mortality in this study. Heart failure has been reported rarely with marketing use of pioglitazone, but more frequently when pioglitazone was used in combination with insulin or in patients with a history of cardiac failure.

<sup>4</sup> A pooled analysis was conducted of adverse reactions of bone fractures from randomised, comparator controlled, double blind clinical trials in over 8100 patients in the pioglitazone-treated groups and 7400 in the comparator-treated groups of up to 3.5 years duration. A higher rate of fractures was observed in women taking pioglitazone (2.6%) versus comparator (1.7%). No increase in fracture rates was observed in men treated with pioglitazone (1.3%) versus comparator (1.5%). In the 3.5 year PROactive study, 44/870 (5.1%) of pioglitazone-treated female patients experienced fractures compared to 23/905 (2.5%) of female patients treated with comparator. No increase in fracture rates was observed in men treated with pioglitazone (1.7%) versus comparator (2.1%).

<sup>5</sup> In active comparator controlled trials mean weight increase with pioglitazone given as monotherapy was 2–3 kg over one year. This is similar to that seen in a sulphonylurea active comparator group. In combination trials pioglitazone added to metformin resulted in mean weight increase over one year of 1.5 kg and added to a sulphonylurea of 2.8 kg. In comparator groups addition of sulphonylurea to metformin resulted in a mean weight gain of 1.3 kg and addition of metformin to a sulphonylurea a mean weight loss of 1.0 kg.

<sup>6</sup> In clinical trials with pioglitazone the incidence of elevations of ALT greater than three times the upper limit of normal was equal to placebo but less than that seen in metformin or sulphonylurea comparator groups. Mean levels of liver enzymes decreased with treatment with pioglitazone. Rare cases of elevated liver enzymes and hepatocellular dysfunction have occurred in post-marketing experience. Although in very rare cases fatal outcome has been reported, causal relationship has not been established.

## 4.9 Overdose

In clinical studies, patients have taken pioglitazone at higher than the recommended highest dose of 45 mg daily. The maximum reported dose of 120 mg/day for four days, then 180 mg/day for seven days was not associated with any symptoms.

Hypoglycaemia may occur in combination with sulphonylureas or insulin. Symptomatic and general supportive measures should be taken in case of overdose.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Drugs used in diabetes, blood glucose lowering drugs, excl. insulins;  
ATC code: A10BG03.

Pioglitazone effects may be mediated by a reduction of insulin resistance. Pioglitazone appears to act via activation of specific nuclear receptors (peroxisome proliferator activated receptor gamma) leading to increased insulin sensitivity of liver, fat and skeletal muscle cells in animals. Treatment with pioglitazone has been shown to reduce hepatic glucose output and to increase peripheral glucose disposal in the case of insulin resistance.

Fasting and postprandial glycaemic control is improved in patients with type 2 diabetes mellitus. The improved glycaemic control is associated with a reduction in both fasting and postprandial plasma insulin concentrations. A clinical trial of pioglitazone vs. gliclazide as monotherapy was extended to two years in order to assess time to treatment failure (defined as appearance of  $HbA_{1c} \geq 8.0\%$  after the first six months of therapy). Kaplan-Meier analysis showed shorter time to treatment failure in patients treated with gliclazide, compared with pioglitazone. At two years, glycaemic control (defined as  $HbA_{1c} < 8.0\%$ ) was sustained in 69% of patients treated with pioglitazone, compared with 50% of patients on gliclazide. In a two-year study of combination therapy comparing pioglitazone with gliclazide when added to metformin, glycaemic control measured as mean change from baseline in  $HbA_{1c}$  was similar between treatment groups after one year. The rate of deterioration of  $HbA_{1c}$  during the second year was less with pioglitazone than with gliclazide.

In a placebo controlled trial, patients with inadequate glycaemic control despite a three month insulin optimisation period were randomised to pioglitazone or placebo for 12 months. Patients receiving pioglitazone had a mean reduction in  $HbA_{1c}$  of 0.45% compared with those continuing on insulin alone, and a reduction of insulin dose in the pioglitazone treated group.

HOMA analysis shows that pioglitazone improves beta cell function as well as increasing insulin sensitivity. Two-year clinical studies have shown maintenance of this effect.

In one year clinical trials, pioglitazone consistently gave a statistically significant reduction in the albumin/creatinine ratio compared to baseline.

The effect of pioglitazone (45 mg monotherapy vs. placebo) was studied in a small 18-week trial in type 2 diabetics. Pioglitazone was associated with significant weight gain. Visceral fat was significantly decreased, while there was an increase in extra-abdominal fat mass. Similar changes in body fat distribution on pioglitazone have been accompanied by an improvement in insulin sensitivity. In most clinical trials, reduced total plasma triglycerides and free fatty acids, and increased HDL-cholesterol levels were observed as compared to placebo, with small, but not clinically significant increases in LDL-cholesterol levels.

In clinical trials of up to two years duration, pioglitazone reduced total plasma triglycerides and free fatty acids, and increased HDL cholesterol levels, compared with placebo, metformin or gliclazide. Pioglitazone did not cause statistically significant increases in LDL cholesterol levels compared with

placebo, whilst reductions were observed with metformin and gliclazide. In a 20-week study, as well as reducing fasting triglycerides, pioglitazone reduced post prandial hypertriglyceridaemia through an effect on both absorbed and hepatically synthesised triglycerides. These effects were independent of pioglitazone's effects on glycaemia and were statistically significant different to glibenclamide.

In PROactive, a cardiovascular outcome study, 5238 patients with type 2 diabetes mellitus and pre-existing major macrovascular disease were randomised to pioglitazone or placebo in addition to existing antidiabetic and cardiovascular therapy, for up to 3.5 years. The study population had an average age of 62 years; the average duration of diabetes was 9.5 years. Approximately one third of patients were receiving insulin in combination with metformin and/or a sulphonylurea. To be eligible patients had to have had one or more of the following: myocardial infarction, stroke, percutaneous cardiac intervention or coronary artery bypass graft, acute coronary syndrome, coronary artery disease, or peripheral arterial obstructive disease. Almost half of the patients had a previous myocardial infarction and approximately 20% had had a stroke. Approximately half of the study population had at least two of the cardiovascular history entry criteria. Almost all subjects (95%) were receiving cardiovascular medicinal products (beta blockers, ACE inhibitors, angiotensin II antagonists, calcium channel blockers, nitrates, diuretics, aspirin, statins, fibrates).

Although the study failed regarding its primary endpoint, which was a composite of all-cause mortality, non-fatal myocardial infarction, stroke, acute coronary syndrome, major leg amputation, coronary revascularisation and leg revascularisation, the results suggest that there are no long-term cardiovascular concerns regarding use of pioglitazone. However, the incidences of oedema, weight gain and heart failure were increased. No increase in mortality from heart failure was observed.

#### *Paediatric population*

The European Medicines Agency has waived the obligation to submit the results of studies with Actos in all subsets of the paediatric population in Type 2 Diabetes Mellitus. See section 4.2 for information on paediatric use.

## **5.2 Pharmacokinetic properties**

### *Absorption*

Following oral administration, pioglitazone is rapidly absorbed, and peak plasma concentrations of unchanged pioglitazone are usually achieved 2 hours after administration. Proportional increases of the plasma concentration were observed for doses from 2–60 mg. Steady state is achieved after 4–7 days of dosing. Repeated dosing does not result in accumulation of the compound or metabolites. Absorption is not influenced by food intake. Absolute bioavailability is greater than 80%.

### *Distribution*

The estimated volume of distribution in humans is 0.25 l/kg.

Pioglitazone and all active metabolites are extensively bound to plasma protein (> 99%).

### *Biotransformation*

Pioglitazone undergoes extensive hepatic metabolism by hydroxylation of aliphatic methylene groups. This is predominantly via cytochrome P450 2C8 although other isoforms may be involved to a lesser degree. Three of the six identified metabolites are active (M-II, M-III, and M-IV). When activity, concentrations and protein binding are taken into account, pioglitazone and metabolite M-III contribute equally to efficacy. On this basis M-IV contribution to efficacy is approximately three-fold that of pioglitazone, whilst the relative efficacy of M-II is minimal.

*In vitro* studies have shown no evidence that pioglitazone inhibits any subtype of cytochrome P450. There is no induction of the main inducible P450 isoenzymes 1A, 2C8/9, and 3A4 in man.

Interaction studies have shown that pioglitazone has no relevant effect on either the pharmacokinetics or pharmacodynamics of digoxin, warfarin, phenprocoumon and metformin. Concomitant administration of pioglitazone with gemfibrozil (an inhibitor of cytochrome P450 2C8) or with rifampicin (an inducer of cytochrome P450 2C8) is reported to increase or decrease, respectively, the plasma concentration of pioglitazone (see section 4.5).

#### *Elimination:*

Following oral administration of radiolabelled pioglitazone to man, recovered label was mainly in faeces (55%) and a lesser amount in urine (45%). In animals, only a small amount of unchanged pioglitazone can be detected in either urine or faeces. The mean plasma elimination half-life of unchanged pioglitazone in man is 5 to 6 hours and for its total active metabolites 16 to 23 hours.

#### *Elderly*

Steady state pharmacokinetics are similar in patients age 65 and over and young subjects.

#### *Patients with renal impairment*

In patients with renal impairment, plasma concentrations of pioglitazone and its metabolites are lower than those seen in subjects with normal renal function, but oral clearance of parent substance is similar. Thus free (unbound) pioglitazone concentration is unchanged.

#### *Patients with hepatic impairment*

Total plasma concentration of pioglitazone is unchanged, but with an increased volume of distribution. Intrinsic clearance is therefore reduced, coupled with a higher unbound fraction of pioglitazone.

### **5.3 Preclinical safety data**

In toxicology studies, plasma volume expansion with haemodilution, anaemia, and reversible eccentric cardiac hypertrophy was consistently apparent after repeated dosing of mice, rats, dogs, and monkeys. In addition, increased fatty deposition and infiltration were observed. These findings were observed across species at plasma concentrations  $\leq 4$  times the clinical exposure. Foetal growth restriction was apparent in animal studies with pioglitazone. This was attributable to the action of pioglitazone in diminishing the maternal hyperinsulinaemia and increased insulin resistance that occurs during pregnancy thereby reducing the availability of metabolic substrates for foetal growth.

Pioglitazone was devoid of genotoxic potential in a comprehensive battery of *in vivo* and *in vitro* genotoxicity assays. An increased incidence of hyperplasia (males and females) and tumours (males) of the urinary bladder epithelium was apparent in rats treated with pioglitazone for up to 2 years.

The formation and presence of urinary calculi with subsequent irritation and hyperplasia was postulated as the mechanistic basis for the observed tumourigenic response in the male rat. A 24-month mechanistic study in male rats demonstrated that administration of pioglitazone resulted in an increased incidence of hyperplastic changes in the bladder. Dietary acidification significantly decreased but did not abolish the incidence of tumours. The presence of microcrystals exacerbated the hyperplastic response but was not considered to be the primary cause of hyperplastic changes. The relevance to humans of the tumourigenic findings in the male rat cannot be excluded.

There was no tumorigenic response in mice of either sex. Hyperplasia of the urinary bladder was not seen in dogs or monkeys treated with pioglitazone for up to 12 months.

In an animal model of familial adenomatous polyposis (FAP), treatment with two other thiazolidinediones increased tumour multiplicity in the colon. The relevance of this finding is unknown.

Environmental Risk Assessment: no environmental impact is anticipated from the clinical use of pioglitazone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Carmellose calcium  
Hyprolose  
Lactose monohydrate  
Magnesium stearate

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years.

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

Aluminium/aluminium blisters, packs of 14, 28, 30, 50, 56, 84, 90, 98, 112 and 196 tablets.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

No special requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Takeda Global Research and Development Centre (Europe) Ltd  
61 Aldwych  
London  
WC2B 4AE  
United Kingdom

## **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/150/004  
EU/1/00/150/005  
EU/1/00/150/006  
EU/1/00/150/008  
EU/1/00/150/010  
EU/1/00/150/019  
EU/1/00/150/020  
EU/1/00/150/021  
EU/1/00/150/027

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 13/10/2000

Date of last renewal: 31/08/2010

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>

## 1. NAME OF THE MEDICINAL PRODUCT

Actos 45 mg tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 45 mg of pioglitazone (as hydrochloride).

Excipients:

Each tablet contains 114.51 mg of lactose monohydrate (see section 4.4).

For a full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Tablet.

The tablets are white to off-white, round, flat and marked '45' on one face and 'ACTOS' on the other face.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Pioglitazone is indicated as second or third line treatment of type 2 diabetes mellitus as described below:

as **monotherapy**

- in adult patients (particularly overweight patients) inadequately controlled by diet and exercise for whom metformin is inappropriate because of contraindications or intolerance

as **dual oral therapy** in combination with

- metformin, in adult patients (particularly overweight patients) with insufficient glycaemic control despite maximal tolerated dose of monotherapy with metformin
- a sulphonylurea, only in adult patients who show intolerance to metformin or for whom metformin is contraindicated, with insufficient glycaemic control despite maximal tolerated dose of monotherapy with a sulphonylurea.

as **triple oral therapy** in combination with

- metformin and a sulphonylurea, in adult patients (particularly overweight patients) with insufficient glycaemic control despite dual oral therapy.
- Pioglitazone is also indicated for combination with insulin in type 2 diabetes mellitus adult patients with insufficient glycaemic control on insulin for whom metformin is inappropriate because of contraindications or intolerance (see section 4.4).

After initiation of therapy with pioglitazone, patients should be reviewed after 3 to 6 months to assess adequacy of response to treatment (e.g. reduction in HbA1c). In patients who fail to show an adequate response, pioglitazone should be discontinued. In light of potential risks with prolonged therapy, prescribers should confirm at subsequent routine reviews that the benefit of pioglitazone is maintained (see section 4.4).

## **4.2 Posology and method of administration**

### Posology

Pioglitazone treatment may be initiated at 15 mg or 30 mg once daily. The dose may be increased in increments up to 45 mg once daily.

In combination with insulin, the current insulin dose can be continued upon initiation of pioglitazone therapy. If patients report hypoglycaemia, the dose of insulin should be decreased.

### Special population

#### *Elderly*

No dose adjustment is necessary for elderly patients (see section 5.2). Physicians should start treatment with the lowest available dose and increase the dose gradually, particularly when pioglitazone is used in combination with insulin (see section 4.4 Fluid retention and cardiac failure).

#### *Renal impairment*

No dose adjustment is necessary in patients with impaired renal function (creatinine clearance > 4 ml/min) (see section 5.2). No information is available from dialysed patients therefore pioglitazone should not be used in such patients.

#### *Hepatic impairment*

Pioglitazone should not be used in patients with hepatic impairment (see section 4.3 and 4.4).

### Paediatric population

The safety and efficacy of Actos in children and adolescents under 18 years of age have not been established. No data are available.

### Method of administration

Pioglitazone tablets are taken orally once daily with or without food. Tablets should be swallowed with a glass of water.

## **4.3 Contraindications**

Pioglitazone is contraindicated in patients with:

- hypersensitivity to the active substance or to any of the excipients
- cardiac failure or history of cardiac failure (NYHA stages I to IV)
- hepatic impairment
- diabetic ketoacidosis
- current bladder cancer or a history of bladder cancer
- uninvestigated macroscopic haematuria

## **4.4 Special warnings and precautions for use**

### *Fluid retention and cardiac failure*

Pioglitazone can cause fluid retention, which may exacerbate or precipitate heart failure. When treating patients who have at least one risk factor for development of congestive heart failure (e.g. prior myocardial infarction or symptomatic coronary artery disease or the elderly), physicians should start with the lowest available dose and increase the dose gradually. Patients should be observed for

signs and symptoms of heart failure, weight gain or oedema; particularly those with reduced cardiac reserve. There have been post-marketing cases of cardiac failure reported when pioglitazone was used in combination with insulin or in patients with a history of cardiac failure. Patients should be observed for signs and symptoms of heart failure, weight gain and oedema when pioglitazone is used in combination with insulin. Since insulin and pioglitazone are both associated with fluid retention, concomitant administration may increase the risk of oedema. Pioglitazone should be discontinued if any deterioration in cardiac status occurs.

A cardiovascular outcome study of pioglitazone has been performed in patients under 75 years with type 2 diabetes mellitus and pre-existing major macrovascular disease. Pioglitazone or placebo was added to existing antidiabetic and cardiovascular therapy for up to 3.5 years. This study showed an increase in reports of heart failure, however this did not lead to an increase in mortality in this study..

### *Elderly*

Combination use with insulin should be considered with caution in the elderly because of increased risk of serious heart failure.

In light of age- related risks (especially bladder cancer, fractures and heart failure), the balance of benefits and risks should be considered carefully both before and during treatment in the elderly.

### *Bladder cancer*

Cases of bladder cancer were reported more frequently in a meta-analysis of controlled clinical trials with pioglitazone (19 cases from 12506 patients, 0.15%) than in control groups (7 cases from 10212 patients, 0.07%) HR=2.64 (95% CI 1.11-6.31, P=0.029). After excluding patients in whom exposure to study drug was less than one year at the time of diagnosis of bladder cancer, there were 7 cases (0.06%) on pioglitazone and 2 cases (0.02%) in control groups. Available epidemiological data also suggest a small increased risk of bladder cancer in diabetic patients treated with pioglitazone in particular in patients treated for the longest durations and with the highest cumulative doses. A possible risk after short term treatment cannot be excluded.

Risk factors for bladder cancer should be assessed before initiating pioglitazone treatment (risks include age, smoking history, exposure to some occupational or chemotherapy agents e.g. cyclophosphamide or prior radiation treatment in the pelvic region). Any macroscopic haematuria should be investigated before starting pioglitazone therapy.

Patients should be advised to promptly seek the attention of their physician if macroscopic haematuria or other symptoms such as dysuria or urinary urgency develop during treatment.

### *Monitoring of liver function*

There have been rare reports of hepatocellular dysfunction during post-marketing experience (see section 4.8). It is recommended, therefore, that patients treated with pioglitazone undergo periodic monitoring of liver enzymes. Liver enzymes should be checked prior to the initiation of therapy with pioglitazone in all patients. Therapy with pioglitazone should not be initiated in patients with increased baseline liver enzyme levels (ALT > 2.5 X upper limit of normal) or with any other evidence of liver disease.

Following initiation of therapy with pioglitazone, it is recommended that liver enzymes be monitored periodically based on clinical judgement. If ALT levels are increased to 3 X upper limit of normal during pioglitazone therapy, liver enzyme levels should be reassessed as soon as possible. If ALT levels remain > 3 X the upper limit of normal, therapy should be discontinued. If any patient develops symptoms suggesting hepatic dysfunction, which may include unexplained nausea, vomiting, abdominal pain, fatigue, anorexia and/or dark urine, liver enzymes should be checked. The decision whether to continue the patient on therapy with pioglitazone should be guided by clinical judgement pending laboratory evaluations. If jaundice is observed, the medicinal product should be discontinued.

### *Weight gain*

In clinical trials with pioglitazone there was evidence of dose related weight gain, which may be due to fat accumulation and in some cases associated with fluid retention. In some cases weight increase may be a symptom of cardiac failure, therefore weight should be closely monitored. Part of the treatment of diabetes is dietary control. Patients should be advised to adhere strictly to a calorie-controlled diet.

### *Haematology*

There was a small reduction in mean haemoglobin (4% relative reduction) and haematocrit (4.1% relative reduction) during therapy with pioglitazone, consistent with haemodilution. Similar changes were seen in metformin (haemoglobin 3-4% and haematocrit 3.6–4.1% relative reductions) and to a lesser extent sulphonylurea and insulin (haemoglobin 1–2% and haematocrit 1–3.2% relative reductions) treated patients in comparative controlled trials with pioglitazone.

### *Hypoglycaemia*

As a consequence of increased insulin sensitivity, patients receiving pioglitazone in dual or triple oral therapy with a sulphonylurea or in dual therapy with insulin may be at risk for dose-related hypoglycaemia, and a reduction in the dose of the sulphonylurea or insulin may be necessary.

### *Eye disorders*

Post-marketing reports of new-onset or worsening diabetic macular oedema with decreased visual acuity have been reported with thiazolidinediones, including pioglitazone. Many of these patients reported concurrent peripheral oedema. It is unclear whether or not there is a direct association between pioglitazone and macular oedema but prescribers should be alert to the possibility of macular oedema if patients report disturbances in visual acuity; an appropriate ophthalmological referral should be considered.

### *Others*

An increased incidence in bone fractures in women was seen in a pooled analysis of adverse reactions of bone fracture from randomised, controlled, double blind clinical trials in over 8100 pioglitazone and 7400 comparator treated patients, on treatment for up to 3.5 years.

Fractures were observed in 2.6% of women taking pioglitazone compared to 1.7% of women treated with a comparator. No increase in fracture rates was observed in men treated with pioglitazone (1.3%) versus comparator (1.5%).

The fracture incidence calculated was 1.9 fractures per 100 patient years in women treated with pioglitazone and 1.1 fractures per 100 patient years in women treated with a comparator. The observed excess risk of fractures for women in this dataset on pioglitazone is therefore 0.8 fractures per 100 patient years of use.

In the 3.5 year cardiovascular risk PROactive study, 44/870 (5.1%; 1.0 fractures per 100 patient years) of pioglitazone-treated female patients experienced fractures compared to 23/905 (2.5%; 0.5 fractures per 100 patient years) of female patients treated with comparator. No increase in fracture rates was observed in men treated with pioglitazone (1.7%) versus comparator (2.1%).

The risk of fractures should be considered in the long term care of women treated with pioglitazone.

As a consequence of enhancing insulin action, pioglitazone treatment in patients with polycystic ovarian syndrome may result in resumption of ovulation. These patients may be at risk of pregnancy.

Patients should be aware of the risk of pregnancy and if a patient wishes to become pregnant or if pregnancy occurs, the treatment should be discontinued (see section 4.6).

Pioglitazone should be used with caution during concomitant administration of cytochrome P450 2C8 inhibitors (e.g. gemfibrozil) or inducers (e.g. rifampicin). Glycaemic control should be monitored closely. Pioglitazone dose adjustment within the recommended posology or changes in diabetic treatment should be considered (see section 4.5).

Actos tablets contain lactose monohydrate and therefore should not be administered to patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Interaction studies have shown that pioglitazone has no relevant effect on either the pharmacokinetics or pharmacodynamics of digoxin, warfarin, phenprocoumon and metformin. Co-administration of pioglitazone with sulphonylureas does not appear to affect the pharmacokinetics of the sulphonylurea. Studies in man suggest no induction of the main inducible cytochrome P450, 1A, 2C8/9 and 3A4. *In vitro* studies have shown no inhibition of any subtype of cytochrome P450. Interactions with substances metabolised by these enzymes, e.g. oral contraceptives, cyclosporin, calcium channel blockers, and HMGCoA reductase inhibitors are not to be expected.

Co-administration of pioglitazone with gemfibrozil (an inhibitor of cytochrome P450 2C8) is reported to result in a 3-fold increase in AUC of pioglitazone. Since there is a potential for an increase in dose-related adverse events, a decrease in the dose of pioglitazone may be needed when gemfibrozil is concomitantly administered. Close monitoring of glycaemic control should be considered (see section 4.4). Co-administration of pioglitazone with rifampicin (an inducer of cytochrome P450 2C8) is reported to result in a 54% decrease in AUC of pioglitazone. The pioglitazone dose may need to be increased when rifampicin is concomitantly administered. Close monitoring of glycaemic control should be considered (see section 4.4).

#### **4.6 Fertility, pregnancy and lactation**

##### *Pregnancy*

There are no adequate human data to determine the safety of pioglitazone during pregnancy. Foetal growth restriction was apparent in animal studies with pioglitazone. This was attributable to the action of pioglitazone in diminishing the maternal hyperinsulinaemia and increased insulin resistance that occurs during pregnancy thereby reducing the availability of metabolic substrates for foetal growth. The relevance of such a mechanism in humans is unclear and pioglitazone should not be used in pregnancy.

##### *Breastfeeding*

Pioglitazone has been shown to be present in the milk of lactating rats. It is not known whether pioglitazone is secreted in human milk. Therefore, pioglitazone should not be administered to breast-feeding women.

##### *Fertility*

In animal fertility studies there was no effect on copulation, impregnation or fertility index

#### **4.7 Effects on ability to drive and use machines**

Actos has no or negligible effect on the ability to drive and use machines. However patients who experience visual disturbance should be cautious when driving or using machines.

## 4.8 Undesirable effects

Adverse reactions reported in excess ( $\geq 0.5\%$ ) of placebo and as more than an isolated case in patients receiving pioglitazone in double-blind studies are listed below as MedDRA preferred term by system organ class and absolute frequency. Frequencies are defined as: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing incidence and seriousness.

| Adverse reaction                            | Frequency of adverse reactions of pioglitazone by treatment regimen |                |                    |                                  |              |
|---------------------------------------------|---------------------------------------------------------------------|----------------|--------------------|----------------------------------|--------------|
|                                             | Mono-therapy                                                        | Combination    |                    |                                  |              |
|                                             |                                                                     | with metformin | with sulphonylurea | with metformin and sulphonylurea | with insulin |
| <b>Infections and infestations</b>          |                                                                     |                |                    |                                  |              |
| upper respiratory tract infection           | common                                                              | common         | common             | common                           | common       |
| bronchitis                                  |                                                                     |                |                    |                                  | common       |
| sinusitis                                   | uncommon                                                            | uncommon       | uncommon           | uncommon                         | uncommon     |
| <b>Blood and lymphatic system disorders</b> |                                                                     |                |                    |                                  |              |
| anaemia                                     |                                                                     | common         |                    |                                  |              |
| <b>Metabolism and nutrition disorders</b>   |                                                                     |                |                    |                                  |              |
| hypo-glycaemia                              |                                                                     |                | uncommon           | very common                      | common       |
| appetite increased                          |                                                                     |                | uncommon           |                                  |              |
| <b>Nervous system disorders</b>             |                                                                     |                |                    |                                  |              |
| hypo-aesthesia                              | common                                                              | common         | common             | common                           | common       |
| headache                                    |                                                                     | common         | uncommon           |                                  |              |
| dizziness                                   |                                                                     |                | common             |                                  |              |
| insomnia                                    | uncommon                                                            | uncommon       | uncommon           | uncommon                         | uncommon     |
| <b>Eye disorders</b>                        |                                                                     |                |                    |                                  |              |
| visual disturbance <sup>1</sup>             | common                                                              | common         | uncommon           |                                  |              |
| macular oedema <sup>2</sup>                 | not known                                                           | not known      | not known          | not known                        | not known    |
| <b>Ear and labyrinth disorders</b>          |                                                                     |                |                    |                                  |              |
| vertigo                                     |                                                                     |                | uncommon           |                                  |              |
| <b>Cardiac disorders</b>                    |                                                                     |                |                    |                                  |              |
| heart failure <sup>3</sup>                  |                                                                     |                |                    |                                  | common       |
| <b>Neoplasms benign,</b>                    |                                                                     |                |                    |                                  |              |

| Adverse reaction                                              | Frequency of adverse reactions of pioglitazone by treatment regimen |                |                    |                                  |              |
|---------------------------------------------------------------|---------------------------------------------------------------------|----------------|--------------------|----------------------------------|--------------|
|                                                               | Mono-therapy                                                        | Combination    |                    |                                  |              |
|                                                               |                                                                     | with metformin | with sulphonylurea | with metformin and sulphonylurea | with insulin |
| <b>malignant and unspecified (including cysts and polyps)</b> |                                                                     |                |                    |                                  |              |
| bladder cancer                                                | uncommon                                                            | uncommon       | uncommon           | uncommon                         | uncommon     |
| <b>Respiratory, thoracic and mediastinal disorders</b>        |                                                                     |                |                    |                                  |              |
| dyspnoea                                                      |                                                                     |                |                    |                                  | common       |
| <b>Gastrointestinal disorders</b>                             |                                                                     |                |                    |                                  |              |
| flatulence                                                    |                                                                     | uncommon       | common             |                                  |              |
| <b>Skin and subcutaneous tissue disorders</b>                 |                                                                     |                |                    |                                  |              |
| sweating                                                      |                                                                     |                | uncommon           |                                  |              |
| <b>Musculoskeletal and connective tissue disorders</b>        |                                                                     |                |                    |                                  |              |
| fracture bone <sup>4</sup>                                    | common                                                              | common         | common             | common                           | common       |
| arthralgia                                                    |                                                                     | common         |                    | common                           | common       |
| back pain                                                     |                                                                     |                |                    |                                  | common       |
| <b>Renal and urinary disorders</b>                            |                                                                     |                |                    |                                  |              |
| haematuria                                                    |                                                                     | common         |                    |                                  |              |
| glycosuria                                                    |                                                                     |                | uncommon           |                                  |              |
| proteinuria                                                   |                                                                     |                | uncommon           |                                  |              |
| <b>Reproductive system and breast disorders</b>               |                                                                     |                |                    |                                  |              |
| erectile dysfunction                                          |                                                                     | common         |                    |                                  |              |
| <b>General disorders and administration site conditions</b>   |                                                                     |                |                    |                                  |              |
| oedema                                                        |                                                                     |                |                    |                                  | very common  |
| fatigue                                                       |                                                                     |                | uncommon           |                                  |              |
| <b>Investigations</b>                                         |                                                                     |                |                    |                                  |              |
| weight increased <sup>5</sup>                                 | common                                                              | common         | common             | common                           | common       |
| blood creatine phospho-kinase                                 |                                                                     |                |                    | common                           |              |

| Adverse reaction                                | Frequency of adverse reactions of pioglitazone by treatment regimen |                |                    |                                  |              |
|-------------------------------------------------|---------------------------------------------------------------------|----------------|--------------------|----------------------------------|--------------|
|                                                 | Mono-therapy                                                        | Combination    |                    |                                  |              |
|                                                 |                                                                     | with metformin | with sulphonylurea | with metformin and sulphonylurea | with insulin |
| increased                                       |                                                                     |                |                    |                                  |              |
| increased lactic dehydro-genase                 |                                                                     |                | uncommon           |                                  |              |
| Alanine aminotransferase increased <sup>6</sup> | not known                                                           | not known      | not known          | not known                        | not known    |

<sup>1</sup> Visual disturbance has been reported mainly early in treatment and is related to changes in blood glucose due to temporary alteration in the turgidity and refractive index of the lens as seen with other hypoglycaemic treatments.

<sup>2</sup> Oedema was reported in 6–9% of patients treated with pioglitazone over one year in controlled clinical trials. The oedema rates for comparator groups (sulphonylurea, metformin) were 2–5%. The reports of oedema were generally mild to moderate and usually did not require discontinuation of treatment.

<sup>3</sup> In controlled clinical trials the incidence of reports of heart failure with pioglitazone treatment was the same as in placebo, metformin and sulphonylurea treatment groups, but was increased when used in combination therapy with insulin. In an outcome study of patients with pre-existing major macrovascular disease, the incidence of serious heart failure was 1.6% higher with pioglitazone than with placebo, when added to therapy that included insulin. However, this did not lead to an increase in mortality in this study. Heart failure has been reported rarely with marketing use of pioglitazone, but more frequently when pioglitazone was used in combination with insulin or in patients with a history of cardiac failure.

<sup>4</sup> A pooled analysis was conducted of adverse reactions of bone fractures from randomised, comparator controlled, double blind clinical trials in over 8100 patients in the pioglitazone-treated groups and 7400 in the comparator-treated groups of up to 3.5 years duration. A higher rate of fractures was observed in women taking pioglitazone (2.6%) versus comparator (1.7%). No increase in fracture rates was observed in men treated with pioglitazone (1.3%) versus comparator (1.5%). In the 3.5 year PROactive study, 44/870 (5.1%) of pioglitazone-treated female patients experienced fractures compared to 23/905 (2.5%) of female patients treated with comparator. No increase in fracture rates was observed in men treated with pioglitazone (1.7%) versus comparator (2.1%).

<sup>5</sup> In active comparator controlled trials mean weight increase with pioglitazone given as monotherapy was 2–3 kg over one year. This is similar to that seen in a sulphonylurea active comparator group. In combination trials pioglitazone added to metformin resulted in mean weight increase over one year of 1.5 kg and added to a sulphonylurea of 2.8 kg. In comparator groups addition of sulphonylurea to metformin resulted in a mean weight gain of 1.3 kg and addition of metformin to a sulphonylurea a mean weight loss of 1.0 kg.

<sup>6</sup> In clinical trials with pioglitazone the incidence of elevations of ALT greater than three times the upper limit of normal was equal to placebo but less than that seen in metformin or sulphonylurea comparator groups. Mean levels of liver enzymes decreased with treatment with pioglitazone. Rare cases of elevated liver enzymes and hepatocellular dysfunction have occurred in post-marketing experience. Although in very rare cases fatal outcome has been reported, causal relationship has not been established.

## 4.9 Overdose

In clinical studies, patients have taken pioglitazone at higher than the recommended highest dose of 45 mg daily. The maximum reported dose of 120 mg/day for four days, then 180 mg/day for seven days was not associated with any symptoms.

Hypoglycaemia may occur in combination with sulphonylureas or insulin. Symptomatic and general supportive measures should be taken in case of overdose.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Drugs used in diabetes, blood glucose lowering drugs, excl. insulins;  
ATC code: A10BG03.

Pioglitazone effects may be mediated by a reduction of insulin resistance. Pioglitazone appears to act via activation of specific nuclear receptors (peroxisome proliferator activated receptor gamma) leading to increased insulin sensitivity of liver, fat and skeletal muscle cells in animals. Treatment with pioglitazone has been shown to reduce hepatic glucose output and to increase peripheral glucose disposal in the case of insulin resistance.

Fasting and postprandial glycaemic control is improved in patients with type 2 diabetes mellitus. The improved glycaemic control is associated with a reduction in both fasting and postprandial plasma insulin concentrations. A clinical trial of pioglitazone vs. gliclazide as monotherapy was extended to two years in order to assess time to treatment failure (defined as appearance of  $HbA_{1c} \geq 8.0\%$  after the first six months of therapy). Kaplan-Meier analysis showed shorter time to treatment failure in patients treated with gliclazide, compared with pioglitazone. At two years, glycaemic control (defined as  $HbA_{1c} < 8.0\%$ ) was sustained in 69% of patients treated with pioglitazone, compared with 50% of patients on gliclazide. In a two-year study of combination therapy comparing pioglitazone with gliclazide when added to metformin, glycaemic control measured as mean change from baseline in  $HbA_{1c}$  was similar between treatment groups after one year. The rate of deterioration of  $HbA_{1c}$  during the second year was less with pioglitazone than with gliclazide.

In a placebo controlled trial, patients with inadequate glycaemic control despite a three month insulin optimisation period were randomised to pioglitazone or placebo for 12 months. Patients receiving pioglitazone had a mean reduction in  $HbA_{1c}$  of 0.45% compared with those continuing on insulin alone, and a reduction of insulin dose in the pioglitazone treated group.

HOMA analysis shows that pioglitazone improves beta cell function as well as increasing insulin sensitivity. Two-year clinical studies have shown maintenance of this effect.

In one year clinical trials, pioglitazone consistently gave a statistically significant reduction in the albumin/creatinine ratio compared to baseline.

The effect of pioglitazone (45 mg monotherapy vs. placebo) was studied in a small 18-week trial in type 2 diabetics. Pioglitazone was associated with significant weight gain. Visceral fat was significantly decreased, while there was an increase in extra-abdominal fat mass. Similar changes in body fat distribution on pioglitazone have been accompanied by an improvement in insulin sensitivity. In most clinical trials, reduced total plasma triglycerides and free fatty acids, and increased HDL-cholesterol levels were observed as compared to placebo, with small, but not clinically significant increases in LDL-cholesterol levels.

In clinical trials of up to two years duration, pioglitazone reduced total plasma triglycerides and free fatty acids, and increased HDL cholesterol levels, compared with placebo, metformin or gliclazide.

Pioglitazone did not cause statistically significant increases in LDL cholesterol levels compared with placebo, whilst reductions were observed with metformin and gliclazide. In a 20-week study, as well as reducing fasting triglycerides, pioglitazone reduced post prandial hypertriglyceridaemia through an effect on both absorbed and hepatically synthesised triglycerides. These effects were independent of pioglitazone's effects on glycaemia and were statistically significant different to glibenclamide.

In PROactive, a cardiovascular outcome study, 5238 patients with type 2 diabetes mellitus and pre-existing major macrovascular disease were randomised to pioglitazone or placebo in addition to existing antidiabetic and cardiovascular therapy, for up to 3.5 years. The study population had an average age of 62 years; the average duration of diabetes was 9.5 years. Approximately one third of patients were receiving insulin in combination with metformin and/or a sulphonylurea. To be eligible patients had to have had one or more of the following: myocardial infarction, stroke, percutaneous cardiac intervention or coronary artery bypass graft, acute coronary syndrome, coronary artery disease, or peripheral arterial obstructive disease. Almost half of the patients had a previous myocardial infarction and approximately 20% had had a stroke. Approximately half of the study population had at least two of the cardiovascular history entry criteria. Almost all subjects (95%) were receiving cardiovascular medicinal products (beta blockers, ACE inhibitors, angiotensin II antagonists, calcium channel blockers, nitrates, diuretics, aspirin, statins, fibrates).

Although the study failed regarding its primary endpoint, which was a composite of all-cause mortality, non-fatal myocardial infarction, stroke, acute coronary syndrome, major leg amputation, coronary revascularisation and leg revascularisation, the results suggest that there are no long-term cardiovascular concerns regarding use of pioglitazone. However, the incidences of oedema, weight gain and heart failure were increased. No increase in mortality from heart failure was observed.

#### *Paediatric population*

The European Medicines Agency has waived the obligation to submit the results of studies with Actos in all subsets of the paediatric population in Type 2 Diabetes Mellitus. See section 4.2 for information on paediatric use.

## **5.2 Pharmacokinetic properties**

### *Absorption*

Following oral administration, pioglitazone is rapidly absorbed, and peak plasma concentrations of unchanged pioglitazone are usually achieved 2 hours after administration. Proportional increases of the plasma concentration were observed for doses from 2–60 mg. Steady state is achieved after 4–7 days of dosing. Repeated dosing does not result in accumulation of the compound or metabolites. Absorption is not influenced by food intake. Absolute bioavailability is greater than 80%.

### *Distribution*

The estimated volume of distribution in humans is 0.25 l/kg.

Pioglitazone and all active metabolites are extensively bound to plasma protein (> 99%).

### *Biotransformation*

Pioglitazone undergoes extensive hepatic metabolism by hydroxylation of aliphatic methylene groups. This is predominantly via cytochrome P450 2C8 although other isoforms may be involved to a lesser degree. Three of the six identified metabolites are active (M-II, M-III, and M-IV). When activity, concentrations and protein binding are taken into account, pioglitazone and metabolite M-III contribute equally to efficacy. On this basis M-IV contribution to efficacy is approximately three-fold that of pioglitazone, whilst the relative efficacy of M-II is minimal.

*In vitro* studies have shown no evidence that pioglitazone inhibits any subtype of cytochrome P450. There is no induction of the main inducible P450 isoenzymes 1A, 2C8/9, and 3A4 in man.

Interaction studies have shown that pioglitazone has no relevant effect on either the pharmacokinetics or pharmacodynamics of digoxin, warfarin, phenprocoumon and metformin. Concomitant administration of pioglitazone with gemfibrozil (an inhibitor of cytochrome P450 2C8) or with rifampicin (an inducer of cytochrome P450 2C8) is reported to increase or decrease, respectively, the plasma concentration of pioglitazone (see section 4.5).

#### *Elimination*

Following oral administration of radiolabelled pioglitazone to man, recovered label was mainly in faeces (55%) and a lesser amount in urine (45%). In animals, only a small amount of unchanged pioglitazone can be detected in either urine or faeces. The mean plasma elimination half-life of unchanged pioglitazone in man is 5 to 6 hours and for its total active metabolites 16 to 23 hours.

#### *Elderly*

Steady state pharmacokinetics are similar in patients age 65 and over and young subjects.

#### *Patients with renal impairment*

In patients with renal impairment, plasma concentrations of pioglitazone and its metabolites are lower than those seen in subjects with normal renal function, but oral clearance of parent substance is similar. Thus free (unbound) pioglitazone concentration is unchanged.

#### *Patients with hepatic impairment*

Total plasma concentration of pioglitazone is unchanged, but with an increased volume of distribution. Intrinsic clearance is therefore reduced, coupled with a higher unbound fraction of pioglitazone.

### **5.3 Preclinical safety data**

In toxicology studies, plasma volume expansion with haemodilution, anaemia, and reversible eccentric cardiac hypertrophy was consistently apparent after repeated dosing of mice, rats, dogs, and monkeys. In addition, increased fatty deposition and infiltration were observed. These findings were observed across species at plasma concentrations  $\leq 4$  times the clinical exposure. Foetal growth restriction was apparent in animal studies with pioglitazone. This was attributable to the action of pioglitazone in diminishing the maternal hyperinsulinaemia and increased insulin resistance that occurs during pregnancy thereby reducing the availability of metabolic substrates for foetal growth.

Pioglitazone was devoid of genotoxic potential in a comprehensive battery of *in vivo* and *in vitro* genotoxicity assays. An increased incidence of hyperplasia (males and females) and tumours (males) of the urinary bladder epithelium was apparent in rats treated with pioglitazone for up to 2 years.

The formation and presence of urinary calculi with subsequent irritation and hyperplasia was postulated as the mechanistic basis for the observed tumourigenic response in the male rat. A 24-month mechanistic study in male rats demonstrated that administration of pioglitazone resulted in an increased incidence of hyperplastic changes in the bladder. Dietary acidification significantly decreased but did not abolish the incidence of tumours. The presence of microcrystals exacerbated the hyperplastic response but was not considered to be the primary cause of hyperplastic changes. The relevance to humans of the tumourigenic findings in the male rat cannot be excluded.

There was no tumorigenic response in mice of either sex. Hyperplasia of the urinary bladder was not seen in dogs or monkeys treated with pioglitazone for up to 12 months.

In an animal model of familial adenomatous polyposis (FAP), treatment with two other thiazolidinediones increased tumour multiplicity in the colon. The relevance of this finding is unknown.

Environmental Risk Assessment: no environmental impact is anticipated from the clinical use of pioglitazone.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Carmellose calcium  
Hyprolose  
Lactose monohydrate  
Magnesium stearate

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years.

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

Aluminium/aluminium blisters, packs of 14, 28, 30, 50, 56, 84, 90, 98, 112 and 196 tablets.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

No special requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Takeda Global Research and Development Centre (Europe) Ltd  
61 Aldwych  
London  
WC2B 4AE  
United Kingdom

## **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/150/011  
EU/1/00/150/012  
EU/1/00/150/013  
EU/1/00/150/014  
EU/1/00/150/015  
EU/1/00/150/022

EU/1/00/150/023  
EU/1/00/150/024  
EU/1/00/150/029  
EU/1/00/150/030

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 13/10/2000  
Date of last renewal: 31/08/2010

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>

## **ANNEX II**

- A. MANUFACTURING AUTHORISATION HOLDERS  
RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND  
USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE  
MARKETING AUTHORISATION**

## **A. MANUFACTURING AUTHORISATION HOLDERS RESPONSIBLE FOR BATCH RELEASE**

### Name and address of the manufacturers responsible for batch release

Takeda Ireland Limited.  
Bray Business Park, Kilruddery, County Wicklow.  
Ireland.

Lilly S.A.  
Avda. de la Industria 30  
28108 Alcobendas (Madrid)  
Spain

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to medical prescription.

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

### Pharmacovigilance system

The MAH must ensure that the system of pharmacovigilance presented in Module 1.8.1. of the Marketing Authorisation is in place and functioning before and whilst the product is on the market.

### Risk Management Plan

The MAH shall submit within 1 month of the Commission decision an updated Risk Management Plan which shall include a risk minimisation plan which includes additional risk minimisation measures to address the identified risks of bladder cancer and heart failure, as detailed below.

The MAH shall perform the pharmacovigilance activities detailed in the Pharmacovigilance Plan, as agreed in the Risk Management Plan presented in Module 1.8.2 of the Marketing Authorisation and any subsequent updates of the RMP agreed by the Committee for Medicinal Products for Human Use (CHMP).

As per the CHMP Guideline on Risk Management Systems for medicinal products for human use, the updated RMP should be submitted at the same time as the next Periodic Safety Update Report (PSUR).

In addition, an updated RMP should be submitted:

- When new information is received that may impact on the current Safety Specification, Pharmacovigilance Plan or risk minimisation activities
- Within 60 days of an important (pharmacovigilance or risk minimisation) milestone being reached
- At the request of the European Medicines Agency

## PSURs

The PSUR cycle for the medicinal product should follow a half-yearly cycle until otherwise agreed by the CHMP.

### Conditions or restrictions with regard to the safe and effective use of the medicinal product

The MAH shall provide an educational pack targeting all physicians who are expected to prescribe/use Pioglitazone. Prior to distribution of the prescriber guide in each Member State, the MAH must agree the content and format of the educational material, together with a communication plan, with the national competent authority.

- This educational pack is aimed at strengthening awareness of important identified risks of bladder cancer and heart failure and the overall recommendations intended to optimise the benefit-risk margin at the patient level.
- The physician educational pack should contain: The Summary of Product Characteristics, package leaflet, and a Prescriber Guide.

The Prescriber Guide should highlight the following:

- Patient selection criteria including that Pioglitazone should not be used as first line therapy and emphasising the need for regular review of treatment benefit.
- The risk of bladder cancer and relevant risk minimisation advice.
- The risk of heart failure and relevant risk minimisation advice.
- Caution in use in the elderly in light of age related risks (in particular bladder cancer, fractures and heart failure).

**ANNEX III**

**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****CARTON****1. NAME OF THE MEDICINAL PRODUCT**

Actos 15 mg tablets

Pioglitazone

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each tablet contains 15 mg pioglitazone (as hydrochloride).

**3. LIST OF EXCIPIENTS**

Contains lactose monohydrate. See leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

14 tablets  
28 tablets  
30 tablets  
50 tablets  
56 tablets  
84 tablets  
90 tablets  
98 tablets  
112 tablets  
196 tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS****10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Takeda Global Research and Development Centre (Europe) Ltd  
61 Aldwych  
London  
WC2B 4AE  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/150/007 14 tablets  
EU/1/00/150/001 28 tablets  
EU/1/00/150/016 30 tablets  
EU/1/00/150/002 50 tablets  
EU/1/00/150/009 56 tablets  
EU/1/00/150/017 84 tablets  
EU/1/00/150/018 90 tablets  
EU/1/00/150/003 98 tablets  
EU/1/00/150/025 112 tablets  
EU/1/00/150/026 196 tablets

**13. BATCH NUMBER**

Batch

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Actos 15 mg

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****CARTON****1. NAME OF THE MEDICINAL PRODUCT**

Actos 30 mg tablets

Pioglitazone

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each tablet contains 30 mg pioglitazone (as hydrochloride).

**3. LIST OF EXCIPIENTS**

Contains lactose monohydrate. See leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

14 tablets  
28 tablets  
30 tablets  
50 tablets  
56 tablets  
84 tablets  
90 tablets  
98 tablets  
112 tablets  
196 tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS****10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Takeda Global Research and Development Centre (Europe) Ltd  
61 Aldwych  
London  
WC2B 4AE  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/150/008 14 tablets  
EU/1/00/150/004 28 tablets  
EU/1/00/150/019 30 tablets  
EU/1/00/150/005 50 tablets  
EU/1/00/150/010 56 tablets  
EU/1/00/150/020 84 tablets  
EU/1/00/150/021 90 tablets  
EU/1/00/150/006 98 tablets  
EU/1/00/150/027 112 tablets  
EU/1/00/150/028 196 tablets

**13. BATCH NUMBER**

Batch

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Actos 30 mg

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****CARTON****1. NAME OF THE MEDICINAL PRODUCT**

Actos 45 mg tablets

Pioglitazone

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each tablet contains 45 mg pioglitazone (as hydrochloride).

**3. LIST OF EXCIPIENTS**

Contains lactose monohydrate. See leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

14 tablets  
28 tablets  
30 tablets  
50 tablets  
56 tablets  
84 tablets  
90 tablets  
98 tablets  
112 tablets  
196 tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE REACH AND SIGHT OF CHILDREN**

Keep out of the reach and sight of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS****10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Takeda Global Research and Development Centre (Europe) Ltd  
61 Aldwych  
London  
WC2B 4AE  
United Kingdom

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/00/150/011 14 tablets  
EU/1/00/150/012 28 tablets  
EU/1/00/150/022 30 tablets  
EU/1/00/150/013 50 tablets  
EU/1/00/150/014 56 tablets  
EU/1/00/150/023 84 tablets  
EU/1/00/150/024 90 tablets  
EU/1/00/150/015 98 tablets  
EU/1/00/150/029 112 tablets  
EU/1/00/150/030 196 tablets

**13. BATCH NUMBER**

Batch

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Actos 45 mg

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER**

**1. NAME OF THE MEDICINAL PRODUCT**

Actos 15 mg tablets

Pioglitazone

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Takeda (logo)

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Batch

**5. OTHER (FOR CALENDARISED PACKS)**

MON  
TUE  
WED  
THU  
FRI  
SAT  
SUN

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS****BLISTER****1. NAME OF THE MEDICINAL PRODUCT**

Actos 30 mg tablets

Pioglitazone

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Takeda (logo)

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Batch

**5. OTHER (FOR CALENDARISED PACKS)**

MON  
TUE  
WED  
THU  
FRI  
SAT  
SUN

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS****BLISTER****1. NAME OF THE MEDICINAL PRODUCT**

Actos 45 mg tablets

Pioglitazone

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Takeda (logo)

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Batch

**5. OTHER (FOR CALENDARISED PACKS)**

MON  
TUE  
WED  
THU  
FRI  
SAT  
SUN

## **B. PACKAGE LEAFLET**

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### Actos 15 mg tablets

Pioglitazone

**Read all of this leaflet carefully before you start taking this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What Actos is and what it is used for
2. Before you take Actos
3. How to take Actos
4. Possible side effects
5. How to store Actos
6. Further information

## 1. WHAT ACTOS IS AND WHAT IT IS USED FOR

Actos contains pioglitazone. It is an anti-diabetic medicine used to treat type 2 (non-insulin dependent) diabetes mellitus, when metformin is not suitable or has failed to work adequately. This is the diabetes that usually develops in adulthood.

Actos helps control the level of sugar in your blood when you have type 2 diabetes by helping your body make better use of the insulin it produces. Your doctor will check whether Actos is working 3 to 6 months after you start taking it.

Actos may be used on its own in patients who are unable to take metformin, and where treatment with diet and exercise has failed to control blood sugar or may be added to other therapies (such as metformin, sulphonylurea or insulin) which have failed to provide sufficient control in blood sugar.

## 2. BEFORE YOU TAKE ACTOS

**Do not take Actos**

- if you are hypersensitive (allergic) to pioglitazone or any of the other ingredients of Actos.
- if you have heart failure or have had heart failure in the past.
- if you have liver disease.
- if you have had diabetic ketoacidosis (a complication of diabetes causing rapid weight loss, nausea or vomiting).
- if you have or have ever had bladder cancer.
- if you have blood in your urine that your doctor has not checked.

**Take special care with Actos**

Tell your doctor before you start to take this medicine

- if you retain water (fluid retention) or have heart failures problems in particular if you are over 75 years old.
- if you have a special type of diabetic eye disease called macular oedema (swelling of the back of the eye).
- if you have cysts on your ovaries (polycystic ovary syndrome). There may be an increased possibility of becoming pregnant because you may ovulate again when you take Actos. If this

- applies to you, use appropriate contraception to avoid the possibility of an unplanned pregnancy.
- if you have a problem with your liver or heart. Before you start taking Actos you will have a blood sample taken to check your liver function. This check may be repeated at intervals. Some patients with long-standing type 2 diabetes mellitus and heart disease or previous stroke who were treated with Actos and insulin experienced the development of heart failure. Inform your doctor as soon as possible if you experience signs of heart failure such as unusual shortness of breath or rapid increase in weight or localised swelling (oedema).

If you take Actos with other medicines for diabetes, it is more likely that your blood sugar could fall below the normal level (hypoglycaemia).

You may also experience a reduction in blood count (anaemia).

#### **Broken bones**

A higher number of bone fractures was seen in women (but not in men) taking pioglitazone. Your doctor will take this into account when treating your diabetes.

#### **Children**

Use in children under 18 years is not recommended.

#### **Taking other medicines**

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

You can usually continue to take other medicines whilst you are being treated with Actos. However, certain medicines are especially likely to affect the amount of sugar in your blood:

- gemfibrozil (used to lower cholesterol)
- rifampicin (used to treat tuberculosis and other infections)

Tell your doctor or pharmacist if you are taking any of these. Your blood sugar will be checked, and your dose of Actos may need to be changed.

#### **Taking Actos with food and drink**

You may take your tablets with or without food. You should swallow the tablets with a glass of water.

#### **Pregnancy and breastfeeding**

Tell your doctor if

- you are, you think you might be or are planning to become pregnant.
- you are breast-feeding or if you are planning to breast-feed your baby.

Your doctor will advise you to discontinue this medicine.

#### **Driving and using machines**

Pioglitazone will not affect your ability to drive or use machines but take care if you experience abnormal vision.

#### **Important information about some of the ingredients of Actos**

This medicine contains lactose monohydrate. If you have been told by your doctor that you have intolerance to some sugars, contact your doctor before taking Actos.

### **3. HOW TO TAKE ACTOS**

One tablet of 15 mg of pioglitazone should be taken once daily. If necessary your doctor may tell you to take a different dose.

If you have the impression that the effect of Actos is too weak, talk to your doctor.

When Actos is taken in combination with other medicines used to treat diabetes (such as insulin, chlorpropamide, glibenclamide, gliclazide, tolbutamide) your doctor will tell you whether you need to take a smaller dose of your medicines.

Your doctor will ask you to have blood tests periodically during treatment with Actos. This is to check that your liver is working normally.

If you are following a diabetic diet, you should continue with this while you are taking Actos.

Your weight should be checked at regular intervals; if your weight increases, inform your doctor.

**If you take more Actos than you should**

If you accidentally take too many tablets, or if someone else or a child takes your medicine, talk to a doctor or pharmacist immediately. Your blood sugar could fall below the normal level and can be increased by taking sugar. It is recommended that you carry some sugar lumps, sweets, biscuits or sugary fruit juice.

**If you forget to take Actos**

Take Actos daily as prescribed. However if you miss a dose, just carry on with the next dose as normal. Do not take a double dose to make up for a forgotten tablet.

**If you stop taking Actos**

Actos should be used every day to work properly. If you stop using Actos, your blood sugar may go up. Talk to your doctor before stopping this treatment.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### **4. POSSIBLE SIDE EFFECTS**

Like all medicines, Actos can cause side effects, although not everybody gets them.

In particular, patients have experienced the following serious side effects:

Heart failure has been experienced commonly (1 to 10 users in 100) in patients taking Actos in combination with insulin. Symptoms are unusual shortness of breath or rapid increase in weight or localised swelling (oedema). If you experience any of these, especially if you are over the age of 65, seek medical advice straight away.

Bladder cancer has been experienced uncommonly (1 to 10 users in 1000) in patients taking Actos. Signs and symptoms include blood in your urine, pain when urinating or a sudden need to urinate. If you experience any of these, talk to your doctor as soon as possible.

Localised swelling (oedema) has also been experienced very commonly in patients taking Actos in combination with insulin. If you experience this side effect, talk to your doctor as soon as possible.

Broken bones have been reported commonly (1 to 10 users in 100) in women patients taking Actos. If you experience this side effect, talk to your doctor as soon as possible.

Blurred vision due to swelling (or fluid) at the back of the eye (frequency not known) has also been reported in patients taking Actos. If you experience this symptom for the first time, talk to your doctor as soon as possible. Also, if you already have blurred vision and the symptom gets worse, talk to your doctor as soon as possible.

The other side effects that have been experienced by some patients taking Actos are:

common (affects 1 to 10 users in 100)

- respiratory infection
- abnormal vision
- weight gain
- numbness

uncommon (affects 1 to 10 users in 1,000)

- inflammation of the sinuses (sinusitis)
- difficulty sleeping (insomnia)

not known (frequency cannot be estimated from the available data)

- increase in liver enzymes

The other side effects that have been experienced by some patients when Actos is taken with other antidiabetic medicines are:

very common (affects more than 1 user in 10)

- decreased blood sugar (hypoglycaemia)

common (affects 1 to 10 users in 100)

- headache
- dizziness
- joint pain
- impotence
- back pain
- shortness of breath
- small reduction in red blood cell count
- flatulence

uncommon (affects 1 to 10 users in 1,000)

- sugar in urine, proteins in urine
- increase in enzymes
- spinning sensation (vertigo)
- sweating
- tiredness
- increased appetite

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE ACTOS**

Keep out of the reach and sight of children.

Do not use Actos after the expiry date which is stated on the carton and the blister pack after the word "EXP". The expiry date refers to the last day of that month.

This medicine does not require any special storage precautions.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## 6. FURTHER INFORMATION

### What Actos contains

- The active substance is pioglitazone. Each tablet contains 15 mg of pioglitazone (as hydrochloride).
- The other ingredients are lactose monohydrate, hypolose, carmellose calcium and magnesium stearate.

### What Actos looks like and contents of the pack

Actos tablets are white to off white, round, convex tablets marked '15' on one face and 'ACTOS' on the other face. The tablets are supplied in blister packs of 14, 28, 30, 50, 56, 84, 90, 98, 112 or 196 tablets. Not all the pack sizes may be marketed.

### Marketing Authorisation Holder and Manufacturer

Marketing authorisation holder

Takeda Global Research and Development Centre (Europe) Ltd, 61 Aldwych, London, WC2B 4AE, United Kingdom.

Manufacturer

Takeda Ireland Limited, Bray Business Park, Kilruddery, County Wicklow, Ireland.

Lilly S.A, Avda. de la Industria 30, 28108 Alcobendas (Madrid), Spain

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### **België/Belgique/Belgien**

Takeda Pharmaceuticals Benelux SPRL  
België/Belgique/Belgien  
Tél/Tel: + 44 (0)20 3116 8953

#### **Luxembourg/Luxemburg**

Takeda Pharmaceuticals Benelux SPRL  
Belgique/Belgien  
Tél/Tel: + 44 (0)20 3116 8954

#### **България**

Takeda Global R & D Centre (Europe),  
Обединеното кралство  
Тел.: +44 (0)20 3116 8000

#### **Magyarország**

Lilly Hungária Kft.  
Tel: + 36 1 328 5100

#### **Česká republika**

ELI LILLY ČR, s.r.o.  
Tel: + 420 234 664 111

#### **Malta**

Takeda Italia Farmaceutici SpA  
Tel: + 39 06 5026 01

#### **Danmark**

Takeda Pharmaceuticals Nordics AB  
Sverige  
Tlf: + 46 (0)8 58 61 33 80

#### **Nederland**

Eli Lilly Nederland B.V.  
Tel: + 31 (0) 30 60 25 800

#### **Deutschland**

Takeda Pharma GmbH  
Tel. + 49 (0) 241 941-0

#### **Norge**

Takeda Pharmaceuticals Nordics AB  
Sverige  
Tlf: + 46 (0)8 58 61 33 80

**Eesti**

Eli Lilly Holdings Limited Eesti filiaal  
Tel: +3726441100

**Ελλάδα**

ΦΑΡΜΑΣΕΡΒ-ΛΙΛΛΥ Α.Ε.Β.Ε.  
Τηλ: +30 210 629 4600

**España**

Lilly S.A.  
Tel: + 34 (91) 663 50 00

**France**

Laboratoires Takeda  
Tél: +33 (0)1 46 25 16 16

**Ireland**

Takeda UK Ltd  
Tel: + 44 (0)1628 537 900

**Ísland**

Icepharma hf.  
Sími: +354 540 8000

**Italia**

Takeda Italia Farmaceutici SpA  
Tel: + 39 06 5026 01

**Κύπρος**

Takeda Global R & D Centre (Europe),  
Ηνωμένο Βασίλειο  
Τηλ: +44 (0)20 3116 8000

**Latvija**

Eli Lilly Holdings Limited pārstāvniecība  
Latvijā  
Tel: +371 7364000

**Lietuva**

Eli Lilly Holdings Limited atstovybė  
Tel. +370 (5) 2649600

**Österreich**

Takeda Pharma Ges.m.b.H.  
Tel: + 43 (1) 524 40 64

**Polska**

Takeda Global R & D Centre (Europe), Wielka  
Brytania  
Tel: +44 (0)20 3116 8000

**Portugal**

Lilly Portugal - Produtos Farmacêuticos, Lda  
Tel: + 351 21 412 6600

**România**

Eli Lilly România S.R.L.  
Tel: +40 21 4023000

**Slovenija**

Eli Lilly farmacevtska družba, d.o.o.  
Tel: +386 (0)1 580 00 10

**Slovenská republika**

Eli Lilly Slovakia, s.r.o.  
Tel: + 421 220 663 111

**Suomi/Finland**

Oy Eli Lilly Finland Ab  
Puh/Tel: + 358 (0)9 8545250

**Sverige**

Takeda Pharmaceuticals Nordics AB  
Sverige  
Tel: + 46 (0)8 58 61 33 80

**United Kingdom**

Takeda UK Ltd  
Tel: + 44 (0)1628 537 900

**This leaflet was last approved in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.

## **PACKAGE LEAFLET: INFORMATION FOR THE USER**

### **Actos 30 mg tablets**

Pioglitazone

**Read all of this leaflet carefully before you start taking this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**In this leaflet:**

1. What Actos is and what it is used for
2. Before you take Actos
3. How to take Actos
4. Possible side effects
5. How to store Actos
6. Further information

## **1. WHAT ACTOS IS AND WHAT IT IS USED FOR**

Actos contains pioglitazone. It is an anti-diabetic medicine used to treat type 2 (non-insulin dependent) diabetes mellitus, when metformin is not suitable or has failed to work adequately. This is the diabetes that usually develops in adulthood.

Actos helps control the level of sugar in your blood when you have type 2 diabetes by helping your body make better use of the insulin it produces. Your doctor will check whether Actos is working 3 to 6 months after you start taking it.

Actos may be used on its own in patients who are unable to take metformin, and where treatment with diet and exercise has failed to control blood sugar or may be added to other therapies (such as metformin, sulphonylurea or insulin) which have failed to provide sufficient control of blood sugar.

## **2. BEFORE YOU TAKE ACTOS**

**Do not take Actos**

- if you are hypersensitive (allergic) to pioglitazone or any of the other ingredients Actos.
- if you have heart failure or have had heart failure in the past.
- if you have liver disease.
- if you have had diabetic ketoacidosis (a complication of diabetes causing rapid weight loss, nausea or vomiting).
- if you have or have ever had bladder cancer.
- if you have blood in your urine that your doctor has not checked.

**Take special care with Actos**

Tell your doctor before you start to take this medicine

- if you retain water (fluid retention) or have heart failures problems in particular if you are over 75 years old.
- if you have a special type of diabetic eye disease called macular oedema (swelling of the back of the eye).
- if you have cysts on your ovaries (polycystic ovary syndrome). There may be an increased possibility of becoming pregnant because you may ovulate again when you take Actos. If this

- applies to you, use appropriate contraception to avoid the possibility of an unplanned pregnancy.
- if you have a problem with your liver or heart. Before you start taking Actos you will have a blood sample taken to check your liver function. This check may be repeated at intervals. Some patients with long-standing type 2 diabetes mellitus and heart disease or previous stroke who were treated with Actos and insulin experienced the development of heart failure. Inform your doctor as soon as possible if you experience signs of heart failure such as unusual shortness of breath or rapid increase in weight or localised swelling (oedema).

If you take Actos with other medicines for diabetes, it is more likely that your blood sugar could fall below the normal level (hypoglycaemia).

You may also experience a reduction in blood count (anaemia).

#### **Broken bones**

A higher number of bone fractures was seen in women (but not in men) taking pioglitazone. Your doctor will take this into account when treating your diabetes.

#### **Children**

Use in children under 18 years is not recommended.

#### **Taking other medicines**

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

You can usually continue to take other medicines whilst you are being treated with Actos. However, certain medicines are especially likely to affect the amount of sugar in your blood:

- gemfibrozil (used to lower cholesterol)
- rifampicin (used to treat tuberculosis and other infections)

Tell your doctor or pharmacist if you are taking any of these. Your blood sugar will be checked, and your dose of Actos may need to be changed.

#### **Taking Actos with food and drink**

You may take your tablets with or without food. You should swallow the tablets with a glass of water.

#### **Pregnancy and breastfeeding**

Tell your doctor if

- you are, you think you might be or are planning to become pregnant.
- you are breast-feeding or if you are planning to breast-feed your baby.

Your doctor will advise you to discontinue this medicine.

#### **Driving and using machines**

Pioglitazone will not affect your ability to drive or use machines but take care if you experience abnormal vision.

#### **Important information about some of the ingredients of Actos**

This medicine contains lactose monohydrate. If you have been told by your doctor that you have intolerance to some sugars, contact your doctor before taking Actos.

### **3. HOW TO TAKE ACTOS**

One tablet of 30 mg of pioglitazone should be taken once daily. If necessary your doctor may tell you to take a different dose.

If you have the impression that the effect of Actos is too weak, talk to your doctor.

When Actos is taken in combination with other medicines used to treat diabetes (such as insulin, chlorpropamide, glibenclamide, gliclazide, tolbutamide) your doctor will tell you whether you need to take a smaller dose of your medicines.

Your doctor will ask you to have blood tests periodically during treatment with Actos. This is to check that your liver is working normally.

If you are following a diabetic diet, you should continue with this while you are taking Actos.

Your weight should be checked at regular intervals; if your weight increases, inform your doctor.

**If you take more Actos than you should**

If you accidentally take too many tablets, or if someone else or a child takes your medicine, talk to a doctor or pharmacist immediately. Your blood sugar could fall below the normal level and can be increased by taking sugar. It is recommended that you carry some sugar lumps, sweets, biscuits or sugary fruit juice.

**If you forget to take Actos**

Take Actos daily as prescribed. However if you miss a dose, just carry on with the next dose as normal. Do not take a double dose to make up a forgotten tablet.

**If you stop taking Actos**

Actos should be used every day to work properly. If you stop using Actos, your blood sugar may go up. Talk to your doctor before stopping this treatment.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### **4. POSSIBLE SIDE EFFECTS**

Like all medicines, Actos can cause side effects, although not everybody gets them.

In particular, patients have experienced the following serious side effects:

Heart failure has been experienced commonly (1 to 10 users in 100) in patients taking Actos in combination with insulin. Symptoms are unusual shortness of breath or rapid increase in weight or localised swelling (oedema). If you experience any of these, especially if you are over the age of 65, seek medical advice straight away.

Bladder cancer has been experienced uncommonly (1 to 10 users in 1000) in patients taking Actos. Signs and symptoms include blood in your urine, pain when urinating or a sudden need to urinate. If you experience any of these, talk to your doctor as soon as possible.

Localised swelling (oedema) has also been experienced very commonly in patients taking Actos in combination with insulin. If you experience this side effect, talk to your doctor as soon as possible.

Broken bones have been reported commonly (1 to 10 users in 100) in women patients taking Actos. If you experience this side effect, talk to your doctor as soon as possible.

Blurred vision due to swelling (or fluid) at the back of the eye (frequency not known) has also been reported in patients taking Actos. If you experience this symptom for the first time, talk to your doctor as soon as possible. Also, if you already have blurred vision and the symptom gets worse, talk to your doctor as soon as possible.

The other side effects that have been experienced by some patients taking Actos are:

common (affects 1 to 10 users in 100)

- respiratory infection
- abnormal vision
- weight gain
- numbness

uncommon (affects 1 to 10 users in 1,000)

- inflammation of the sinuses (sinusitis)
- difficulty sleeping (insomnia)

not known (frequency cannot be estimated from the available data)

- increase in liver enzymes

The other side effects that have been experienced by some patients when Actos is taken with other antidiabetic medicines are:

very common (affects more than 1 user in 10)

- decreased blood sugar (hypoglycaemia)

common (affects 1 to 10 users in 100)

- headache
- dizziness
- joint pain
- impotence
- back pain
- shortness of breath
- small reduction in red blood cell count
- flatulence

uncommon (affects 1 to 10 users in 1,000)

- sugar in urine, proteins in urine
- increase in enzymes
- spinning sensation (vertigo)
- sweating
- tiredness
- increased appetite

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE ACTOS**

Keep out of the reach and sight of children.

Do not use Actos after the expiry date which is stated on the carton and the blister pack after the word "EXP". The expiry date refers to the last day of that month.

This medicine does not require any special storage precautions.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## 6. FURTHER INFORMATION

### What Actos contains

- The active substance in Actos is pioglitazone. Each tablet contains 30 mg of pioglitazone (as hydrochloride).
- The other ingredients are lactose monohydrate, hypolose, carmellose calcium and magnesium stearate.

### What Actos looks like and contents of the pack

Actos tablets are white to off white, round, flat tablets marked '30' on one face and 'ACTOS' on the other face. The tablets are supplied in blister packs of 14, 28, 30, 50, 56, 84, 90, 98, 112 or 196 tablets. Not all the pack sizes may be marketed.

### Marketing Authorisation Holder and Manufacturer

Marketing authorisation holder

Takeda Global Research and Development Centre (Europe) Ltd, 61 Aldwych, London, WC2B 4AE, United Kingdom.

Manufacturer

Takeda Ireland Limited, Bray Business Park, Kilruddery, County Wicklow, Ireland.

Lilly S.A, Avda. de la Industria 30, 28108 Alcobendas (Madrid), Spain

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### **België/Belgique/Belgien**

Takeda Pharmaceuticals Benelux SPRL  
België/Belgique/Belgien  
Tél/Tel: + 44 (0)20 3116 8953

#### **Luxembourg/Luxemburg**

Takeda Pharmaceuticals Benelux SPRL  
Belgique/Belgien  
Tél/Tel: + 44 (0)20 3116 8954

#### **България**

Takeda Global R & D Centre (Europe),  
Обединеното кралство  
Тел.: +44 (0)20 3116 8000

#### **Magyarország**

Lilly Hungária Kft.  
Tel: + 36 1 328 5100

#### **Česká republika**

ELI LILLY ČR, s.r.o.  
Tel: + 420 234 664 111

#### **Malta**

Takeda Italia Farmaceutici SpA  
Tel: + 39 06 5026 01

#### **Danmark**

Takeda Pharmaceuticals Nordics AB  
Sverige  
Tlf: + 46 (0)8 58 61 33 80

#### **Nederland**

Eli Lilly Nederland B.V.  
Tel: + 31 (0) 30 60 25 800

#### **Deutschland**

Takeda Pharma GmbH  
Tel. + 49 (0) 241 941-0

#### **Norge**

Takeda Pharmaceuticals Nordics AB  
Sverige  
Tlf: + 46 (0)8 58 61 33 80

**Eesti**

Eli Lilly Holdings Limited Eesti filiaal  
Tel: +3726441100

**Ελλάδα**

ΦΑΡΜΑΣΕΡΒ-ΛΙΛΛΥ Α.Ε.Β.Ε.  
Τηλ: +30 210 629 4600

**España**

Lilly S.A.  
Tel: + 34 (91) 663 50 00

**France**

Laboratoires Takeda  
Tél: +33 (0)1 46 25 16 16

**Ireland**

Takeda UK Ltd  
Tel: + 44 (0)1628 537 900

**Ísland**

Icepharma hf.  
Sími: +354 540 8000

**Italia**

Takeda Italia Farmaceutici SpA  
Tel: + 39 06 5026 01

**Κύπρος**

Takeda Global R & D Centre (Europe),  
Ηνωμένο Βασίλειο  
Τηλ: +44 (0)20 3116 8000

**Latvija**

Eli Lilly Holdings Limited pārstāvniecība  
Latvijā  
Tel: +371 7364000

**Lietuva**

Eli Lilly Holdings Limited atstovybė  
Tel. +370 (5) 2649600

**Österreich**

Takeda Pharma Ges.m.b.H.  
Tel: + 43 (1) 524 40 64

**Polska**

Takeda Global R & D Centre (Europe), Wielka  
Brytania  
Tel: +44 (0)20 3116 8000

**Portugal**

Lilly Portugal - Produtos Farmacêuticos, Lda  
Tel: + 351 21 412 6600

**România**

Eli Lilly România S.R.L.  
Tel: +40 21 4023000

**Slovenija**

Eli Lilly farmacevtska družba, d.o.o.  
Tel: +386 (0)1 580 00 10

**Slovenská republika**

Eli Lilly Slovakia, s.r.o.  
Tel: + 421 220 663 111

**Suomi/Finland**

Oy Eli Lilly Finland Ab  
Puh/Tel: + 358 (0)9 8545250

**Sverige**

Takeda Pharmaceuticals Nordics AB  
Sverige  
Tel: + 46 (0)8 58 61 33 80

**United Kingdom**

Takeda UK Ltd  
Tel: + 44 (0)1628 537 900

**This leaflet was last approved in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### Actos 45 mg tablets

Pioglitazone

**Read all of this leaflet carefully before you start taking this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

#### **In this leaflet:**

1. What Actos is and what it is used for
2. Before you take Actos
3. How to take Actos
4. Possible side effects
5. How to store Actos
6. Further information

## **1. WHAT ACTOS IS AND WHAT IT IS USED FOR**

Actos contains pioglitazone. It is an anti-diabetic medicine used to treat type 2 (non-insulin dependent) diabetes mellitus, when metformin is not suitable or has failed to work adequately. This is the diabetes that usually develops in adulthood.

Actos helps control the level of sugar in your blood when you have type 2 diabetes by helping your body make better use of the insulin it produces. Your doctor will check whether Actos is working 3 to 6 months after you start taking it.

Actos may be used on its own in patients who are unable to take metformin, and where treatment with diet and exercise has failed to control blood sugar or may be added to other therapies (such as metformin, sulphonylurea or insulin) which have failed to provide sufficient control of blood sugar.

## **2. BEFORE YOU TAKE ACTOS**

### **Do not take Actos**

- if you are hypersensitive (allergic) to pioglitazone or any of the other ingredients of Actos.
- if you have heart failure or have had heart failure in the past.
- if you have liver disease.
- if you have had diabetic ketoacidosis (a complication of diabetes causing rapid weight loss, nausea or vomiting).
- if you have or have ever had bladder cancer.
- if you have blood in your urine that your doctor has not checked.

### **Take special care with Actos**

Tell your doctor before you start to take this medicine

- if you retain water (fluid retention) or have heart failures problems in particular if you are over 75 years old.
- if you have a special type of diabetic eye disease called macular oedema (swelling of the back of the eye).
- if you have cysts on your ovaries (polycystic ovary syndrome). There may be an increased possibility of becoming pregnant because you may ovulate again when you take Actos. If this

- applies to you, use appropriate contraception to avoid the possibility of an unplanned pregnancy.
- if you have a problem with your liver or heart. Before you start taking Actos you will have a blood sample taken to check your liver function. This check may be repeated at intervals. Some patients with long-standing type 2 diabetes mellitus and heart disease or previous stroke who were treated with Actos and insulin experienced the development of heart failure. Inform your doctor as soon as possible if you experience signs of heart failure such as unusual shortness of breath or rapid increase in weight or localised swelling (oedema).

If you take Actos with other medicines for diabetes, it is more likely that your blood sugar could fall below the normal level (hypoglycaemia).

You may also experience a reduction in blood count (anaemia).

#### **Broken bones**

A higher number of bone fractures was seen in women (but not in men) taking pioglitazone. Your doctor will take this into account when treating your diabetes.

#### **Children**

Use in children under 18 years is not recommended.

#### **Taking other medicines**

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

You can usually continue to take other medicines whilst you are being treated with Actos. However, certain medicines are especially likely to affect the amount of sugar in your blood:

- gemfibrozil (used to lower cholesterol)
- rifampicin (used to treat tuberculosis and other infections)

Tell your doctor or pharmacist if you are taking any of these. Your blood sugar will be checked, and your dose of Actos may need to be changed.

#### **Taking Actos with food and drink**

You may take your tablets with or without food. You should swallow the tablets with a glass of water.

#### **Pregnancy and breastfeeding**

Tell your doctor if

- you are, you think you might be or are planning to become pregnant.
- you are breast-feeding or if you are planning to breast-feed your baby.

Your doctor will advise you to discontinue this medicine.

#### **Driving and using machines**

Pioglitazone will not affect your ability to drive or use machines but take care if you experience abnormal vision.

#### **Important information about some of the ingredients of Actos**

This medicine contains lactose monohydrate. If you have been told by your doctor that you have intolerance to some sugars, contact your doctor before taking Actos.

### **3. HOW TO TAKE ACTOS**

One tablet of 45 mg of pioglitazone should be taken once daily. If necessary your doctor may tell you to take a different dose.

If you have the impression that the effect of Actos is too weak, talk to your doctor.

When Actos is taken in combination with other medicines used to treat diabetes (such as insulin, chlorpropamide, glibenclamide, gliclazide, tolbutamide) your doctor will tell you whether you need to take a smaller dose of your medicines.

Your doctor will ask you to have blood tests periodically during treatment with Actos. This is to check that your liver is working normally.

If you are following a diabetic diet, you should continue with this while you are taking Actos.

Your weight should be checked at regular intervals; if your weight increases, inform your doctor.

**If you take more Actos than you should**

If you accidentally take too many tablets, or if someone else or a child takes your medicine, talk to a doctor or pharmacist immediately. Your blood sugar could fall below the normal level and can be increased by taking sugar. It is recommended that you carry some sugar lumps, sweets, biscuits or sugary fruit juice.

**If you forget to take Actos**

Take Actos daily as prescribed. However if you miss a dose, just carry on with the next dose as normal. Do not take a double dose to make up for a forgotten tablet.

**If you stop taking Actos**

Actos should be used every day to work properly. If you stop using Actos, your blood sugar may go up. Talk to your doctor before stopping this treatment.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### **4. POSSIBLE SIDE EFFECTS**

Like all medicines, Actos can cause side effects, although not everybody gets them.

In particular, patients have experienced the following serious side effects:

Heart failure has been experienced commonly (1 to 10 users in 100) in patients taking Actos in combination with insulin. Symptoms are unusual shortness of breath or rapid increase in weight or localised swelling (oedema). If you experience any of these, especially if you are over the age of 65, seek medical advice straight away.

Bladder cancer has been experienced uncommonly (1 to 10 users in 1000) in patients taking Actos. Signs and symptoms include blood in your urine, pain when urinating or a sudden need to urinate. If you experience any of these, talk to your doctor as soon as possible.

Localised swelling (oedema) has also been experienced very commonly in patients taking Actos in combination with insulin. If you experience this side effect, talk to your doctor as soon as possible.

Broken bones have been reported commonly (1 to 10 users in 100) in women patients taking Actos. If you experience this side effect, talk to your doctor as soon as possible.

Blurred vision due to swelling (or fluid) at the back of the eye (frequency not known) has also been reported in patients taking Actos. If you experience this symptom for the first time, talk to your doctor as soon as possible. Also, if you already have blurred vision and the symptom gets worse, talk to your doctor as soon as possible.

The other side effects that have been experienced by some patients taking Actos are:

common (affects 1 to 10 users in 100)

- respiratory infection
- abnormal vision
- weight gain
- numbness

uncommon (affects 1 to 10 users in 1,000)

- inflammation of the sinuses (sinusitis)
- difficulty sleeping (insomnia)

not known (frequency cannot be estimated from the available data)

- increase in liver enzymes

The other side effects that have been experienced by some patients when Actos is taken with other antidiabetic medicines are:

very common (affects more than 1 user in 10)

- decreased blood sugar (hypoglycaemia)

common (affects 1 to 10 users in 100)

- headache
- dizziness
- joint pain
- impotence
- back pain
- shortness of breath
- small reduction in red blood cell count
- flatulence

uncommon (affects 1 to 10 users in 1,000)

- sugar in urine, proteins in urine
- increase in enzymes
- spinning sensation (vertigo)
- sweating
- tiredness
- increased appetite

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## **5. HOW TO STORE ACTOS**

Keep out of the reach and sight of children.

Do not use Actos after the expiry date which is stated on the carton and the blister pack after the word "EXP". The expiry date refers to the last day of that month.

This medicine does not require any special storage precautions.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## **6. FURTHER INFORMATION**

**What Actos contains**

- The active substance in Actos is pioglitazone. Each tablet contains 45 mg of pioglitazone (as hydrochloride).
- The other ingredients are lactose monohydrate, hypollose, carmellose calcium and magnesium stearate.

**What Actos looks like and contents of the pack**

Actos tablets are white to off white, round, flat tablets marked '45' on one face and 'ACTOS' on the other face. The tablets are supplied in blister packs of 14, 28, 30, 50, 56, 84, 90, 98, 112 or 196 tablets. Not all pack sizes may be marketed.

**Marketing Authorisation Holder and Manufacturer**

Marketing authorisation holder

Takeda Global Research and Development Centre (Europe) Ltd, 61 Aldwych, London, WC2B 4AE, United Kingdom.

Manufacturer

Takeda Ireland Limited, Bray Business Park, Kilruddery, County Wicklow, Ireland.

Lilly S.A, Avda. de la Industria 30, 28108 Alcobendas (Madrid), Spain

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Takeda Pharmaceuticals Benelux SPRL

België/Belgique/Belgien

Tél/Tel: + 44 (0)20 3116 8953

**Luxembourg/Luxemburg**

Takeda Pharmaceuticals Benelux SPRL

Belgique/Belgien

Tél/Tel: + 44 (0)20 3116 8954

**България**

Takeda Global R & D Centre (Europe),

Обединеното кралство

Тел.: +44 (0)20 3116 8000

**Magyarország**

Lilly Hungária Kft.

Tel: + 36 1 328 5100

**Česká republika**

ELI LILLY ČR, s.r.o.

Tel: + 420 234 664 111

**Malta**

Takeda Italia Farmaceutici SpA

Tel: + 39 06 5026 01

**Danmark**

Takeda Pharmaceuticals Nordics AB

Sverige

Tlf: + 46 (0)8 58 61 33 80

**Nederland**

Eli Lilly Nederland B.V.

Tel: + 31 (0) 30 60 25 800

**Deutschland**

Takeda Pharma GmbH

Tel. + 49 (0) 241 941-0

**Norge**

Takeda Pharmaceuticals Nordics AB

Sverige

Tlf: + 46 (0)8 58 61 33 80

**Eesti**

Eli Lilly Holdings Limited Eesti filiaal  
Tel: +3726441100

**Ελλάδα**

ΦΑΡΜΑΣΕΡΒ-ΛΙΛΛΥ Α.Ε.Β.Ε.  
Τηλ: +30 210 629 4600

**España**

Lilly S.A.  
Tel: + 34 (91) 663 50 00

**France**

Laboratoires Takeda  
Tél: +33 (0)1 46 25 16 16

**Ireland**

Takeda UK Ltd  
Tel: + 44 (0)1628 537 900

**Ísland**

Icepharma hf.  
Sími: +354 540 8000

**Italia**

Takeda Italia Farmaceutici SpA  
Tel: + 39 06 5026 01

**Κύπρος**

Takeda Global R & D Centre (Europe),  
Ηνωμένο Βασίλειο  
Τηλ: +44 (0)20 3116 8000

**Latvija**

Eli Lilly Holdings Limited pārstāvniecība  
Latvijā  
Tel: +371 7364000

**Lietuva**

Eli Lilly Holdings Limited atstovybė  
Tel. +370 (5) 2649600

**Österreich**

Takeda Pharma Ges.m.b.H.  
Tel: + 43 (1) 524 40 64

**Polska**

Takeda Global R & D Centre (Europe), Wielka  
Brytania  
Tel: +44 (0)20 3116 8000

**Portugal**

Lilly Portugal - Produtos Farmacêuticos, Lda  
Tel: + 351 21 412 6600

**România**

Eli Lilly România S.R.L.  
Tel: +40 21 4023000

**Slovenija**

Eli Lilly farmacevtska družba, d.o.o  
Tel: +386 (0)1 580 00 10

**Slovenská republika**

Eli Lilly Slovakia, s.r.o.  
Tel: + 421 220 663 111

**Suomi/Finland**

Oy Eli Lilly Finland Ab  
Puh/Tel: + 358 (0)9 8545250

**Sverige**

Takeda Pharmaceuticals Nordics AB  
Sverige  
Tel: + 46 (0)8 58 61 33 80

**United Kingdom**

Takeda UK Ltd  
Tel: + 44 (0)1628 537 900

**This leaflet was last approved in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.