



EUROPEAN MEDICINES AGENCY
SCIENCE MEDICINES HEALTH

EMADOC-1700519818-2155553

European Medicines Agency decision

CW/0001/2025

of 2 June 2025

on a class waiver on condition(s) in accordance with Regulation (EC) No 1901/2006 of the European Parliament and of the Council

Only the English text is authentic.



European Medicines Agency decision

CW/0001/2025

of 2 June 2025

on a class waiver on condition(s) in accordance with Regulation (EC) No 1901/2006 of the European Parliament and of the Council

The European Medicines Agency,

Having regard to the Treaty on the Functioning of the European Union,

Having regard to Regulation (EC) No 1901/2006 of the European Parliament and of the Council of 12 December 2006 on medicinal products for paediatric use and amending Regulation (EEC) No. 1768/92, Directive 2001/20/EC, Directive 2001/83/EC and Regulation (EC) No 726/2004¹,

Having regard to Regulation (EC) No 726/2004 of the European Parliament and of the Council of 31 March 2004 laying down Union procedures for the authorisation and supervision of medicinal products for human use and establishing a European Medicines Agency²,

Having regard to the decision of the European Medicines Agency P/1/2007 on a class waiver on conditions and the subsequent decisions on a class waiver on conditions: decision P/17/2008 of 21 April 2008, decision P/47/2008 of 14 July 2008, decision P/65/2009 of 13 April 2009, decision P/146/2009 of 26 October 2009, decision CW/0001/2015 of 23 July 2015,

Having regard to the opinion of the Paediatric Committee of the European Medicines Agency, issued of its own motion on 23 May 2025 in accordance with Article 12 of Regulation (EC) No 1901/2006,

Having regard to Article 25 of Regulation (EC) No 1901/2006,

Whereas:

- (1) The Paediatric Committee of the European Medicines Agency has given an opinion of its own motion on the granting of a waiver on condition(s).
- (2) It is therefore appropriate to adopt a decision granting a waiver on condition(s).

¹ OJ L 378, 27.12.2006, p.1, as amended.

² OJ L 136, 30.4.2004, p. 1, as amended.

Has adopted this decision:

Article 1

A waiver on condition(s), the details of which are set out in the opinion of the Paediatric Committee of the European Medicines Agency annexed hereto, together with its appendices, is hereby granted.

Article 2

This decision is applicable to all natural and legal persons intending to market a product falling within the scope of Article 1 of this decision.

Article 3

The decision CW/0001/2015 of the European Medicines Agency adopted on 23 July 2015 is hereby superseded and replaced by this decision.

EMADOC-1700519818-2008870

Amsterdam, 23 May 2025

Opinion of the Paediatric Committee on a class waiver on condition(s)

Waived conditions

See Annex I

Basis for opinion

On 3 December 2007, the European Medicines Agency adopted decision P/1/2007 on a class waiver on conditions. Subsequently, the following decisions on a class waiver on conditions were adopted: decision P/17/2008 of 21 April 2008, decision P/47/2008 of 14 July 2008, decision P/65/2009 of 13 April 2009, decision P/146/2009 of 26 October 2009, and decision CW/0001/2015 of 23 July 2015.

According to Article 12 of Regulation (EC) No 1901/2006, the Paediatric Committee may, of its own motion, adopt an opinion on granting of a class waiver.

Opinion

1. The Paediatric Committee, having considered the conditions listed in Annex I, recommends:
 - pursuant to Article 12 of Regulation (EC) No 1901/2006, to grant a waiver for medicinal products intended for the diagnosis of IgE-mediated allergy in all subsets of the paediatric population, on the grounds that the specific medicinal product does not represent a significant therapeutic benefit over existing treatments for paediatric patients in accordance with Article 11(1)(c) of said Regulation.

The Paediatric Committee member of Norway agrees with the above-mentioned recommendation of the Paediatric Committee.

2. The conditions waived are set out in the Annex I.
3. The grounds for the granting of a waiver for the waived condition are set out in the Annex II.

This opinion is forwarded to the Executive Director of the European Medicines Agency.

Annex I

List of conditions waived

List of conditions waived

Conditions

1. the classes of androgen receptor modulator, of oestrogen receptor modulator, of growth and sex hormone as well as their releasing or inhibiting factors, and of sex hormone-metabolism modulator medicinal products for treatment of breast malignant neoplasms, prostate malignant neoplasms and neuroendocrine malignant neoplasms;
2. the class of first-generation taxoid medicinal products for treatment of breast malignant neoplasms, gynaecological epithelial malignant neoplasms, prostate malignant neoplasms, intestinal malignant neoplasms, pancreatic malignant neoplasms, head and neck epithelial malignant neoplasms as well as lung malignant neoplasms;
3. the class of Ecteinascidin medicinal products for treatment of gynaecological epithelial malignant neoplasms;
4. the class of Her- / epidermal growth factor-receptor antibody medicinal products for treatment of breast malignant neoplasms, intestinal malignant neoplasms and head and neck epithelial malignant neoplasms;
5. the class of thymidylate synthase inhibitor medicinal products for the treatment of intestinal malignant neoplasms and lung malignant neoplasms;

on the ground that these classes of medicinal products are likely to be ineffective in all of the paediatric population in the concerned condition, in accordance with Article 11.1 (a) of said Regulation;

6. the class of Colchicum alkaloids (colchicine and its derivatives) medicinal products for the treatment of primary gout;
7. the class of androgen antagonist medicinal products for treatment of benign prostatic hyperplasia;
8. the class of pyrimidine- and pyrimidine analogue-containing medicinal products for treatment of breast malignant neoplasms, intestinal malignant neoplasms, lung malignant neoplasms, pancreatic malignant neoplasms, head and neck epithelial malignant neoplasms, skin malignant neoplasms and actinic keratosis;
9. the class of first- and second-generation platinum-containing medicinal products for treatment of urinary tract malignant neoplasms, head and neck epithelial malignant neoplasms and lung malignant neoplasms;
10. the class of alkylating-methylating medicinal products for treatment of skin malignant neoplasms;
11. the class of ribonucleotide reductase-beta-2 inhibitor medicinal products for treatment of myeloproliferative neoplasms;
12. the class of primarily alkylating medicinal products for treatment of myeloproliferative neoplasms and mature B, T and NK cell neoplasms;
13. the class of photosensitising medicinal products for treatment of head and neck epithelial malignant neoplasms;

14. the class of retinoic X receptor-activating medicinal products for treatment of mature B, T and NK cell neoplasms;
15. the class of immunomodulatory cytokine medicinal products for treatment of neuroendocrine malignant neoplasms, skin malignant neoplasms, myeloproliferative neoplasms and mature B, T and NK cell neoplasms;

16. all classes of medicinal products for diagnosis of IgE-mediated allergy;

on the ground that the classes of medicinal products do not represent a significant therapeutic benefit over existing treatments for paediatric patients in the concerned condition, in accordance with Article 11.1 (c) of said Regulation;

17. the class of peroxisome proliferator-activated receptor (PPAR)-gamma modulators, including dual and multiple PPAR modulator (e.g., thiazolidinediones, glitazars, triple modulators) medicinal products for treatment of type II diabetes mellitus;

on the ground that these classes of medicinal products are likely to be unsafe in all of the paediatric population in the concerned condition, in accordance with Article 11.1 (a) of said Regulation;

18. all classes of medicinal products for treatment of primary and secondary osteoarthritis;
19. all classes of medicinal products for treatment of organic amnesic syndrome (excluding amnesic syndrome caused by alcohol and other psychoactive substances);
20. all classes of medicinal products for treatment of age-related macular degeneration and diabetic macular oedema;
21. all classes of medicinal products for treatment of climacteric symptoms associated with decreased oestrogen levels, as occurring at menopause;
22. all classes of medicinal products for treatment of Alzheimer's disease;
23. all classes of medicinal products for treatment of erectile dysfunction;
24. all classes of medicinal products for treatment of chronic obstructive pulmonary disease (COPD) (excluding chronic lung diseases associated with long-term airflow limitation, such as asthma, bronchopulmonary dysplasia, primary cilia dyskinesia, obstructive lung disease related to graft-versus-host disease after [bone-marrow] transplantation);
25. all classes of medicinal products for treatment of vulvar intraepithelial neoplasia;

on the ground that the disease or condition for which the specific medicinal product is intended occurs only in adult populations, in accordance with Article 11.1 (b) of said Regulation.

Annex II

Grounds for the granting of a waiver

Grounds for the granting of a waiver

Scope

Clinical studies with medicinal products, intended to diagnose IgE-mediated allergy, are not expected to bring a significant therapeutic benefit to or to fulfil a therapeutic need of the paediatric population. This is because sensitivity and specificity of diagnostic allergens can be extrapolated from adults to children.

This will avoid unnecessary trials in the paediatric population.

A medicinal product also developed for another condition not listed in Annex I will not be waived from the requirements of the Regulation (EC) No 1901/2006.

The conditions listed in Annex I should not prevent an applicant from considering the development of a medicinal product in different or related conditions/indications for use in the paediatric population.

The PDCO may adopt further opinions recommending the granting of waivers in other conditions.

In addition, the PDCO may adopt an opinion advocating the review of granted waivers as per Article 14(2) of Regulation (EC) No 1901/2006.

Voluntary submission of a paediatric investigation plan for a waived condition is still possible.

Grounds

The references provided were identified through systematic search and may not be exhaustive or necessarily available for the EU population.

The PDCO acknowledges that there may be anecdotal cases of occurrence of waived conditions in the paediatric population.

Medicinal products that are likely to be ineffective in the paediatric population

| Class of medicinal product | References | Group of conditions where the medicinal product is authorised for treatment |
|--|---|--|
| Androgen receptor modulator, oestrogen receptor modulator, growth and sex hormone as well as their releasing or inhibiting factors, sex hormone-metabolism modulator medicinal products ¹ | (Bourdeaut et al. 2009) (Janem et al. 2010) (Skapek et al. 2013) (Michalski et al. 2010) | <ul style="list-style-type: none">• Breast malignant neoplasms• Prostate malignant neoplasms• Neuroendocrine malignant neoplasms |

¹ Examples: Tamoxifen, Toremifene, Fulvestrant, Degarelix, Enzalutamide, Abiraterone, Somatostatin, Octreotide

| | | |
|--|--|---|
| First-generation taxoid medicinal products ² | (Horton et al. 2008) (Norwegian Medicines Agency 2010) (Sanofi 2012) (Jacobs et al. 2010) | <ul style="list-style-type: none"> • Breast malignant neoplasms • Gynaecological epithelial malignant neoplasms • Prostate malignant neoplasms • Intestinal malignant neoplasms • Head and neck epithelial malignant neoplasms • Lung malignant neoplasms • Pancreatic malignant neoplasms |
| Her-/Epidermal growth factor-receptor antibody medicinal products ³ | (Trippett et al. 2009) (Ebb et al. 2012) (European Medicines Agency 2009) | <ul style="list-style-type: none"> • Breast malignant neoplasms • Intestinal malignant neoplasms • Head and neck epithelial malignant neoplasms |
| Ecteinascidin medicinal products | (Baruchel et al. 2012) (European Medicines Agency 2014) | <ul style="list-style-type: none"> • Gynaecological epithelial malignant neoplasms |
| Thymidylate synthase inhibitor medicinal product ⁴ | (Warwick et al. 2013) (Horton et al. 2005) (Malempati et al. 2007) | <ul style="list-style-type: none"> • Intestinal malignant neoplasms • Lung malignant neoplasms |

² Examples: Paclitaxel, Docetaxel, Ixabepilone

³ Examples: Trastuzumab, Pertuzumab

⁴ Examples: Pemetrexed, Raltitrexed

Medicinal products that lack significant therapeutic benefit over existing treatments for paediatric population

| Class of medicinal product | Reference(s) | Group of conditions where the medicinal product is used for the treatment of |
|--|---|---|
| Pyrimidine- and pyrimidine analogue-containing medicinal products ⁵ | (Kilburn et al. 2013) (Hoffmann-La Roche 2014) (Trobaugh-Lotrario & Katzenstein 2012) (Buehrlen et al. 2012) | <ul style="list-style-type: none"> • Breast malignant neoplasms • Intestinal malignant neoplasms • Head and neck epithelial malignant neoplasms • Lung malignant neoplasms • Pancreatic malignant neoplasms • Skin malignant neoplasms • Actinic keratosis |
| First- and second-generation platinum-containing medicinal products ⁶ | (Pizzo & Poplack 2011, p.301) | <ul style="list-style-type: none"> • Urinary tract malignant neoplasms • Intestinal malignant neoplasms • Head and neck epithelial malignant neoplasms • Lung malignant neoplasms |
| Immunomodulatory cytokine medicinal products ⁷ | (Buehrlen et al. 2012) | <ul style="list-style-type: none"> • Neuroendocrine malignant neoplasms • Skin malignant neoplasms • Myeloproliferative neoplasms • Mature B, T and NK cell neoplasms • Neuroblastoma |
| Photosensitising medicinal products | (Basset-Seguin et al. 2014) | <ul style="list-style-type: none"> • Head and neck epithelial malignant neoplasms |
| Colchicum alkaloids (colchicine and its derivatives) | (Medicines & Healthcare products Regulatory Agency (MHRA) 2012) (Kallinich et al. 2007) | <ul style="list-style-type: none"> • Primary gout • Auto-inflammatory syndromes |

⁵ Examples: 5-Fluorouracil, Capecitabine

⁶ Examples: Cisplatin, carboplatin, oxaliplatin

⁷ Examples: Interleukin 2, Interferon alpha

| | | |
|--|--|---|
| Alkylating-methylating medicinal products ⁸ | (Pizzo & Poplack 2011, pp.304, 649, 290) | <ul style="list-style-type: none"> • Skin malignant neoplasms • Brain tumours • Neuroblastoma • Sarcoma • Hodgkin lymphoma |
| Androgen antagonists ⁹ | (Speiser et al. 2010) | <ul style="list-style-type: none"> • Benign prostatic hyperplasia • Congenital adrenal hyperplasia • Hyperandrogenism |
| Ribonucleotide reductase beta-2 inhibitor medicinal products ¹⁰ | (Sharma et al. 2014, sec. O-079) (Pizzo & Poplack 2011, p 621) | <ul style="list-style-type: none"> • Myeloproliferative neoplasms • Leukaemia |
| Primarily alkylating medicinal products | (Pizzo & Poplack 2011, pp 290 and 299) (Michel et al. 2011) | <ul style="list-style-type: none"> • Myeloproliferative neoplasms • Mature B, T and NK cell neoplasms • Hodgkin lymphoma |
| Retinoic X receptor-activating medicinal products ¹¹ | (Mehta et al. 2012) | <ul style="list-style-type: none"> • Mature B, T and NK cell neoplasms |
| Diagnosis of IgE-mediated allergy medicinal products | (Bousquet et al. 2012) (Ansotegui et al. 2020) (Heinzerling et al. 2013) (Riggioni et al. 2023) | <ul style="list-style-type: none"> • Diagnosis of IgE-mediated allergy |

⁸ Examples: Dacarbazine (DTIC), Temozolomide

⁹ Example: Finasteride

¹⁰ Example: Hydroxycarbamide (hydroxyurea)

¹¹ Examples: Bendamustine, Carmustine (BCNU); Temoporfin; Bexarotene

Previously granted, revoked and revised waivers

| Condition in previous class condition waiver | Groups of condition or condition referred to in revised class waiver | Class(es) of medicinal product referred to in revised class waiver | Scientific assessment conclusion |
|--|---|--|---|
| Oropharyngeal, laryngeal or nasal epithelial carcinoma (excluding nasopharyngeal carcinoma or lymphoepithelioma) | Head and neck epithelial malignant neoplasms | First-generation taxoid medicinal products Her- / epidermal growth factor-receptor antibody medicinal products Pyrimidine- and pyrimidine analogue-containing medicinal products Photosensitising medicinal products First- and second generation platinum-containing medicinal products | Likely ineffective Likely ineffective Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits |
| Lung carcinoma (small cell and non-small cell carcinoma) | Lung malignant neoplasms | First-generation taxoid medicinal products Thymidylate synthase inhibitor medicinal products First- and second-generation platinum-containing medicinal products | Likely ineffective Likely ineffective Do not represent significant therapeutic benefits |
| Basal cell carcinoma | Skin malignant neoplasm | Pyrimidine- and pyrimidine analogue-containing medicines Alkylating-methylating medicines Immunomodulatory cytokine medicinal products | Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits |

| | | | |
|---|---|--|--|
| Breast carcinoma | Breast malignant neoplasms | <p>Androgen receptor modulator, of oestrogen receptor modulator, of growth and sex hormone as well as their releasing or inhibiting factors, and of sex hormone-metabolism modulator medicinal products</p> <p>First-generation taxoid medicinal products</p> <p>Her- / epidermal growth factor-receptor antibody medicinal products</p> <p>Pyrimidine- and pyrimidine analogue-containing medicines</p> | <p>Likely ineffective</p> <p>Likely ineffective</p> <p>Likely ineffective</p> <p>Do not represent significant therapeutic benefits</p> |
| Ovarian carcinoma (excluding rhabdomyosarcoma and germ cell tumours) | Gynaecological epithelial malignant neoplasms | <p>First-generation taxoid medicinal products</p> <p>Ecteinascidin medicinal products</p> | <p>Likely ineffective</p> <p>Likely ineffective</p> |
| Fallopian tube carcinoma (excluding rhabdomyosarcoma and germ cell tumours) | Gynaecological epithelial malignant neoplasms | <p>First-generation taxoid medicines</p> <p>Ecteinascidin medicinal products</p> | <p>Likely ineffective</p> <p>Likely ineffective</p> |
| Endometrial carcinoma | Gynaecological epithelial malignant neoplasms | <p>First-generation taxoid medicinal products</p> <p>Ecteinascidin medicinal products</p> | <p>Likely ineffective</p> <p>Likely ineffective</p> |
| Peritoneal carcinoma (excluding blastomas and sarcomas) | Gynaecological epithelial malignant neoplasms | <p>First-generation taxoid medicinal products</p> <p>Ecteinascidin medicinal products</p> | <p>Likely ineffective</p> <p>Likely ineffective</p> |
| Prostate carcinoma (excluding rhabdomyosarcoma) | Prostate malignant neoplasms | <p>Androgen receptor modulator, of oestrogen receptor modulator, of growth and sex hormone as well as their releasing or inhibiting factors, and of sex hormone-</p> | <p>Likely ineffective</p> |

| | | | |
|---|--|--|---|
| | | metabolism modulator medicinal products First-generation taxoid medicinal products | Likely ineffective |
| Hairy cell leukaemia | Mature B, T and NK cell neoplasms | Primarily alkylating medicinal products Retinoic X receptor-activating medicinal products Immunomodulatory cytokine medicinal products | Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits |
| Multiple myeloma | Mature B, T and NK cell neoplasms | Primarily alkylating medicinal products Retinoic X receptor-activating medicines Immunomodulatory cytokine medicinal products | Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits |
| Alzheimer's disease | Alzheimer's disease | | Confirmed |
| Vascular dementia and vascular cognitive disorder/impairment | Coronary atherosclerosis, peripheral atherosclerosis, vascular dementia and vascular cognitive disorder / impairment | | Revocation |
| Organic amnesic syndrome (excluding amnesic syndrome caused by alcohol and other psychoactive substances) | Organic amnesic syndrome (excluding amnesic syndrome caused by alcohol and other psychoactive substances) | | Confirmed |
| Amyotrophic lateral sclerosis | Amyotrophic lateral sclerosis | | Revocation |
| Parkinson's disease (non-juvenile) | Parkinson disease (non-juvenile) | | Revocation |

| | | | |
|---|--|---|---|
| Age-related macular degeneration | Age-related macular degeneration and diabetic macular oedema | | Confirmed |
| Climacteric symptoms associated with decreased oestrogen levels, as occurring at menopause | Climacteric symptoms associated with decreased oestrogen levels, as occurring at menopause | | Confirmed |
| Chronic Obstructive Pulmonary Disease (COPD) (excluding chronic lung diseases associated with long-term airflow limitation, such as asthma, bronchopulmonary dysplasia, primary cilia dyskinesia, obstructive lung disease related to graft-versus-host disease after (bone-marrow) transplantation). | Chronic obstructive pulmonary disease (COPD) (excluding chronic lung diseases associated with long-term airflow limitation, such as asthma, bronchopulmonary dysplasia, primary cilia dyskinesia, obstructive lung disease related to graft-versus-host disease after [bone-marrow] transplantation) | | Confirmed |
| Liver and intrahepatic bile duct carcinoma (excluding hepatoblastoma) | Liver and intrahepatic bile duct carcinoma | | Revocation |
| Adenocarcinoma of the pancreas | Pancreatic malignant neoplasms | First-generation taxoid medicinal products Pyrimidine- and pyrimidine analogue-containing medicines | Likely ineffective Do not represent significant therapeutic benefits |
| Gastroenteropancreatic neuroendocrine tumours (excluding neuroblastoma, neuroganglioblastoma, pheochromocytoma) | Neuroendocrine malignant neoplasms | Androgen receptor modulators, of oestrogen receptor modulators, of growth and sex hormones as well as their releasing or inhibiting factors, and of sex hormone-metabolism modulators Immunomodulatory cytokine medicinal products | Likely ineffective Do not represent significant therapeutic benefits |

| | | | |
|--|------------------------------------|--|--|
| Gastric carcinoids | Neuroendocrine malignant neoplasms | Androgen receptor modulator, of oestrogen receptor modulator, of growth and sex hormone as well as their releasing or inhibiting factors, and of sex hormone-metabolism modulator medicinal products Immunomodulatory cytokine medicinal products | Likely ineffective Do not represent significant therapeutic benefits |
| Adenocarcinoma of the colon and rectum | Intestinal malignant neoplasms | First-generation taxoid medicinal products Her- / epidermal growth factor-receptor antibody medicinal products Thymidylate synthase inhibitor medicinal products First- and second generation platinum-containing medicinal products Pyrimidine- and pyrimidine analogue-containing medicinal products | Likely ineffective Likely ineffective Likely ineffective Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits |
| Ureter and bladder carcinoma | Ureter and bladder carcinoma | First- and second generation platinum-containing medicinal products | Do not represent significant therapeutic benefits |
| Kidney and renal pelvis carcinoma (excluding nephroblastoma, nephroblastomatosis, clear cell sarcoma, mesoblastic nephroma, renal medullary carcinoma and rhabdoid tumour of the kidney) | Kidney and renal pelvis carcinoma | | Revocation |

| | | | |
|---|---|---|--|
| Vaginal and vulvar carcinoma (excluding rhabdomyosarcoma and soft tissue sarcoma) | Gynaecological malignant neoplasms | First-generation taxoid medicinal products Ecteinascidin medicinal products | Likely ineffective Likely ineffective |
| Melanoma (from 0 to less than 12 years) | Skin malignant neoplasm | Pyrimidine- and pyrimidine analogue-containing medicinal products Alkylating-methylating medicinal products Immunomodulatory cytokine medicinal products | Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits |
| Gastric adenocarcinoma | Intestinal malignant neoplasms | First-generation taxoid medicinal products Her- / epidermal growth factor-receptor antibody medicinal products Thymidylate synthase inhibitor medicinal products First- and second generation platinum-containing Pyrimidine- and pyrimidine analogue-containing medicinal products | Likely ineffective Likely ineffective Likely ineffective Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits |
| Chronic lymphocytic leukaemia | Mature B, T and NK cell neoplasms | Primarily alkylating medicinal products Retinoic X receptor-activating medicinal products Do not represent significant therapeutic benefits | Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits |
| Cervix and corpus uteri carcinoma | Gynaecological epithelial malignant neoplasms | First-generation taxoid medicinal products | Likely ineffective |

| | | | |
|---|--|--|---|
| | | Ecteinascidin medicinal products | Likely ineffective |
| Follicular lymphoma | Mature B, T and NK cell neoplasms | Primarily alkylating medicinal products Retinoic X receptor-activating medicinal products Immunomodulatory cytokine medicinal products | Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits Do not represent significant therapeutic benefits |
| Primary and secondary osteoarthritis | Primary and secondary osteoarthritis | | Confirmed |
| Coronary atherosclerosis | Coronary atherosclerosis, peripheral atherosclerosis, vascular dementia and vascular cognitive disorder / impairment | | Revocation |
| Peripheral atherosclerosis | Coronary atherosclerosis, peripheral atherosclerosis, vascular dementia and vascular cognitive disorder / impairment | | Revocation |
| Huntington chorea | Huntington chorea | | Revocation |
| Benign prostatic hyperplasia | Benign prostatic hyperplasia | Androgen antagonist medicinal products | Do not represent significant therapeutic benefits |
| Erectile dysfunction | Erectile dysfunction | | Confirmed |
| Primary gout (excluding Lesch-Nyhan syndrome and other secondary forms of gout) | Primary gout | Colchicum alkaloid (colchicine and its derivatives) medicinal products | Do not represent significant therapeutic benefits |
| Primary myelofibrosis | Myeloproliferative neoplasms | Ribonucleotide reductase beta-2 inhibitor medicinal products | Do not represent significant therapeutic benefits |

| | | | |
|----------------------------------|--|---|---|
| | | Primarily alkylating medicinal products | Do not represent significant therapeutic benefits |
| | | Immunomodulatory cytokine medicinal products | Do not represent significant therapeutic benefits |
| Diabetic macular oedema | Age-related macular degeneration and diabetic macular oedema | | Confirmed |
| Mesothelioma | Lung malignant neoplasms | First-generation taxoid medicinal products Thymidylate synthase inhibitor medicinal products | Likely ineffective Likely ineffective |
| Actinic keratosis | Actinic keratosis | Pyrimidine- and pyrimidine analogue-containing medicines | Do not represent significant therapeutic benefits |
| Vulvar intraepithelial neoplasia | Vulvar intraepithelial neoplasia | | Confirmed |

References

Adamson PC, 2013. Unintended consequences of regulatory initiatives in childhood cancer drug development. *JAMA Pediatrics*, 167(10), pp.886–887. Available at: <http://dx.doi.org/10.1001/jamapediatrics.2013.2488> [Accessed August 19, 2013].

Ansotegui I. J. et al., 2020. IgE allergy diagnostics and other relevant tests in allergy, a World Allergy Organization position paper. *World Allergy Organ J.*, 13(2):100080. Available at: <https://doi.org/10.1016/j.waojou.2019.100080>. [Accessed May 20, 2025].

Barosi, G., Lupo, L. & Rosti, V., 2012. Management of myeloproliferative neoplasms: from academic guidelines to clinical practice. *Current Hematologic Malignancy Reports*, 7(1), pp.50–56.

Baruchel, S. et al., 2012. A phase 2 trial of trabectedin in children with recurrent rhabdomyosarcoma, Ewing sarcoma and non-rhabdomyosarcoma soft tissue sarcomas: a report from the Children's Oncology Group. *European Journal of Cancer (Oxford, England: 1990)*, 48(4), pp.579–585.

Basset-Seguin, N. et al., 2014. Consensus recommendations for the treatment of basal cell carcinomas in Gorlin syndrome with topical methylnolaevulinate-photodynamic therapy. *Journal of the European Academy of Dermatology and Venereology: JEADV*, 28(5), pp.626–632.

Berg, S.L. et al., 2011. Safety, pharmacokinetics, and immunomodulatory effects of lenalidomide in children and adolescents with relapsed/refractory solid tumors or myelodysplastic syndrome: a Children's Oncology Group Phase I Consortium report. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*, 29(3), pp.316–323.

- Bertini, E., 2014. Orphanet: Juvenile amyotrophic lateral sclerosis. Available at: http://www.orpha.net/consor/cgi-bin/OC_Exp.php?lng=en&Expert=300605 [Accessed November 21, 2014].
- Bousquet J. et al., 2012. Practical guide to skin prick tests in allergy to aeroallergens. Position paper EAACI. Allergy, 67(1), pp. 18-24. Available at: <https://onlinelibrary.wiley.com/doi/epdf/10.1111/j.1398-9995.2011.02728.x> [Accessed May 20, 2025].
- Bourdeaut, F. et al., 2009. VIP hypersecretion as primary or secondary syndrome in neuroblastoma: A retrospective study by the Société Française des Cancers de l'Enfant (SFCE). Pediatric Blood & Cancer, 52(5), pp.585–590. Available at: <http://onlinelibrary.wiley.com/doi/10.1002/pbc.21912/abstract> [Accessed November 25, 2014].
- Buehrlen, M. et al., 2012. Multimodal treatment, including interferon beta, of nasopharyngeal carcinoma in children and young adults: preliminary results from the prospective, multicenter study NPC-2003-GPOH/DCOG. Cancer, 118(19), pp.4892–4900.
- Cote, A.T. et al., 2013. Childhood obesity and cardiovascular dysfunction. Journal of the American College of Cardiology, 62(15), pp.1309–1319.
- Ebb, D. et al., 2012. Phase II Trial of Trastuzumab in Combination With Cytotoxic Chemotherapy for Treatment of Metastatic Osteosarcoma With Human Epidermal Growth Factor Receptor 2 Overexpression: A Report From the Children's Oncology Group. Journal of Clinical Oncology, 30(20), pp.2545–2551. Available at: <http://jco.ascopubs.org/content/30/20/2545> [Accessed June 4, 2013].
- European Medicines Agency, 2009. European Medicines Agency - Human medicines - Withdrawn applications - Theraloc. Available at: http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/000931/wapp/Initial_authorisation/human_wapp_000043.jsp&mid=WC0b01ac058001d128&source=homeMedSearch&category=human [Accessed November 25, 2014].
- European Medicines Agency, 2014. Summary of product characteristics - European public assessment report - Yondelis. Available at: http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/000773/human_med_001165.jsp&mid=WC0b01ac058001d124 [Accessed April 19, 2015].
- Finsterer, J. & Burgunder, J.-M., 2014. Recent progress in the genetics of motor neuron disease. European Journal of Medical Genetics, 57(2-3), pp.103–112.
- Geller, J.I. et al., 2015. Characterization of adolescent and pediatric renal cell carcinoma: A report from the Children's Oncology Group study AREN03B2. Cancer, 121(14), pp.2457–2464. Available at: <http://onlinelibrary.wiley.com/doi/10.1002/cncr.29368/abstract> [Accessed July 16, 2015].
- Hargrave, D., Bartels, U. & Bouffet, E., 2006. Diffuse brainstem glioma in children: critical review of clinical trials. The Lancet Oncology, 7(3), pp.241–248. Available at: [http://www.thelancet.com/journals/lanonc/article/PIIS1470-2045\(06\)70615-5/fulltext](http://www.thelancet.com/journals/lanonc/article/PIIS1470-2045(06)70615-5/fulltext) [Accessed February 17, 2010].
- Heinzerling L. et al., 2013. The skin prick test - European standards. Clin Transl Allergy. 2013 3(1), p.3. Available at: <https://onlinelibrary.wiley.com/doi/epdf/10.1186/2045-7022-3-3> [Accessed May 20, 2025].
- Hoffmann-La Roche, 2014. A Study of Capecitabine (Xeloda) and Concomitant Radiation Therapy in Children With Newly Diagnosed Brainstem Gliomas - Full Text View - ClinicalTrials.gov. Available at: <http://clinicaltrials.gov/ct2/show/study/NCT01118377> [Accessed November 25, 2014].

Horton, T.M. et al., 2008. A Phase 1 and pharmacokinetic clinical trial of paclitaxel for the treatment of refractory leukemia in children: a Children's Oncology Group study. *Pediatric Blood & Cancer*, 50(4), pp.788–792.

Horton, T.M. et al., 2005. Phase I trial and pharmacokinetic study of raltitrexed in children with recurrent or refractory leukemia: a pediatric oncology group study. *Clinical Cancer Research: An Official Journal of the American Association for Cancer Research*, 11(5), pp.1884–1889.

Institute of Cancer Research, 2014. EU rules are denying children latest cancer drugs. Available at: http://www.icr.ac.uk/press/press_archive/press_releases_2014/24538.shtml [Accessed February 20, 2014].

International Agency for Research on Cancer, 2012. GLOBOCAN 2012 age-specific table. Available at: http://globocan.iarc.fr/old/age-specific_table_n.asp?selection=63990&title=European+Union+%28EU-28%29&sex=0&type=0&stat=1&>window=1&sort=0&submit=%C2%A0Execute%C2%A0 [Accessed November 25, 2014].

Jacobs, S. et al., 2010. Phase II trial of ixabepilone administered daily for five days in children and young adults with refractory solid tumors: a report from the children's oncology group. *Clinical Cancer Research: An Official Journal of the American Association for Cancer Research*, 16(2), pp.750–754.

Janem, W. et al., 2010. Malignant insulinoma in a child. *Pediatric Blood & Cancer*, 55(7), pp.1423–1426. Available at: <http://onlinelibrary.wiley.com/doi/10.1002/pbc.22784/abstract> [Accessed November 25, 2014].

Jansen, M.H.A. et al., 2012. Diffuse intrinsic pontine gliomas: a systematic update on clinical trials and biology. *Cancer Treatment Reviews*, 38(1), pp.27–35. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21764221> [Accessed February 8, 2012].

Kallinich, T. et al., 2007. Colchicine Use in Children and Adolescents With Familial Mediterranean Fever: Literature Review and Consensus Statement. *Pediatrics*, 119(2), pp.e474–e483. Available at: <http://pediatrics.aappublications.org/content/119/2/e474> [Accessed November 26, 2014].

Kilburn, L.B. et al., 2013. Phase I trial of capecitabine rapidly disintegrating tablets and concomitant radiation therapy in children with newly diagnosed brainstem gliomas and high-grade gliomas. *Neuro-Oncology*, 15(6), pp.759–766.

Lansbury, L. et al., 2013. Interventions for non-metastatic squamous cell carcinoma of the skin: systematic review and pooled analysis of observational studies. *BMJ*, 347, p.f6153. Available at: <http://www.bmj.com/content/347/bmj.f6153> [Accessed April 22, 2015].

Malempati, S. et al., 2007. Phase I trial and pharmacokinetic study of pemetrexed in children with refractory solid tumors: the Children's Oncology Group. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology*, 25(12), pp.1505–1511.

Masters, G.A. et al., 2015. Clinical Cancer Advances 2015: Annual Report on Progress Against Cancer From the American Society of Clinical Oncology. *Journal of Clinical Oncology*, 33(7), pp.786–809. Available at: <http://jco.ascopubs.org/content/33/7/786> [Accessed February 28, 2015].

MEB The Netherlands & MPA Sweden, 2006. Paraplatin INN carboplatin - Paediatric Public Assessment Report EU Work Sharing Procedure, Available at: http://www.hma.eu/fileadmin/dateien/Human_Medicines/CMD_h_/Paediatric_Regulation/Assessment_Reports/Previous_worksharing_project/Paraplatin_carboplatin_PaedPAR.pdf [Accessed July 5, 2015].

Medicines & Healthcare products Regulatory Agency (MHRA), 2012. Public Assessment Report for paediatric studies submitted in accordance with Article 45 of Regulation (EC) No1901/2006, as

amended COLCHICINE UK/W/015/pdWS/001. Available at:
http://www.hma.eu/fileadmin/dateien/Human_Medicines/CMD_h_/Paediatric_Regulation/Assessment_Reports/Article_45_work-sharing/Colchicine_2012_02_45_PdAR.pdf [Accessed April 22, 2015].

Mehta, N. et al., 2012. Bexarotene is active against subcutaneous panniculitis-like T-cell lymphoma in adult and pediatric populations. *Clinical Lymphoma, Myeloma & Leukemia*, 12(1), pp.20–25.

Menda, Y. et al., 2010. Phase I Trial of 90Y-DOTA0-Tyr3-Octreotide Therapy in Children and Young Adults with Refractory Solid Tumors That Express Somatostatin Receptors. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine*, 51(10), pp.1524–1531. Available at:
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3753801/> [Accessed April 18, 2015].

Michalski, A. et al., 2010. The addition of high-dose tamoxifen to standard radiotherapy does not improve the survival of patients with diffuse intrinsic pontine glioma. *Journal of Neuro-Oncology*, 100(1), pp.81–88.

Michel, G. et al., 2011. Weight-based strategy of dose administration in children using intravenous busulfan: Clinical and pharmacokinetic results. *Pediatric Blood & Cancer*. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/21254374> [Accessed January 25, 2011].

Muggia, F. & Kudlowitz, D., 2014. Novel taxanes. *Anti-Cancer Drugs*, 25(5), pp.593–598.

Norwegian Medicines Agency, 2010. Taxol Paclitaxel - Rapporteur's Public Assessment Report for paediatric studies submitted in accordance with Article 45 of Regulation (EC) No. 1901/2006. Available at:
http://www.hma.eu/fileadmin/dateien/Human_Medicines/CMD_h_/Paediatric_Regulation/Assessment_Reports/Article_45_work-sharing/Paclitaxel_2010_04_45PaedPdAR.pdf [Accessed November 25, 2014].

Orban, P. et al., 2007. Chapter 15 Juvenile amyotrophic lateral sclerosis. *Handbook of Clinical Neurology*, 82, pp.301–312.

Pastore, G. et al., 2006. Malignant renal tumours incidence and survival in European children (1978-1997): report from the Automated Childhood Cancer Information System project. *European Journal of Cancer (Oxford, England: 1990)*, 42(13), pp.2103–2114.

Pizzo, P. & Poplack, D., 2011. *Principles and Practice of Pediatric Oncology* 6th ed., Philadelphia: Lippincott Williams & Wilkins.

Riggioni C. et al., 2023. Systematic review and meta-analyses on the accuracy of diagnostic tests for IgE-mediated food allergy. *Allergy*. 2024;79(2), pp. 324-352. Available at:
<https://onlinelibrary.wiley.com/doi/epdf/10.1111/all.15939> [Accessed May 20, 2025].

Rodrigues, A.N. et al., 2013. Cardiovascular risk factor investigation: a pediatric issue. *International Journal of General Medicine*, 6, pp.57–66.

Rodriguez-Galindo, C. et al., 2013. Children's Oncology Group's 2013 blueprint for research: Rare tumors. *Pediatric Blood & Cancer*, 60(6), pp.1016–1021. Available at:
<http://onlinelibrary.wiley.com/doi/10.1002/pbc.24428/abstract> [Accessed April 19, 2015].

Rodriguez-Galindo, C. et al., 2005. Preradiation chemotherapy with methotrexate, cisplatin, 5-fluorouracil, and leucovorin for pediatric nasopharyngeal carcinoma. *Cancer*, 103(4), pp.850–7.

Roos, R., 2011. Juvenile Huntington disease. Available at: http://www.orpha.net/consor/cgi-bin/Disease_Search.php?Ing=EN&data_id=19562 [Accessed October 24, 2014].

- Sanofi, 2012. Docetaxel in Combination With Cisplatin-5-fluorouracil for the Induction Treatment of Nasopharyngeal Carcinoma in Children and Adolescents - Full Text View - ClinicalTrials.gov. Available at: <http://clinicaltrials.gov/ct2/show/NCT00565448> [Accessed November 25, 2014].
- Sharma, M. et al., 2014. To compare role of hydroxyurea and hyperhydration versus hyperhydration alone to decrease total leukocyte count in children of acute leukemia with hyperleukocytosis: a randomized control trial [abstract O-079]. In *Pediatric Blood & Cancer*. pp. S105–S433. Available at: <http://onlinelibrary.wiley.com/doi/10.1002/pbc.25314/abstract> [Accessed November 25, 2014].
- Siegel, D.A. et al., 2014. Cancer Incidence Rates and Trends Among Children and Adolescents in the United States, 2001–2009. *Pediatrics*, 134(4), pp.e945–e955. Available at: <http://pediatrics.aappublications.org/content/134/4/e945> [Accessed October 23, 2014].
- Skapek, S.X. et al., 2013. Safety and efficacy of high-dose tamoxifen and sulindac for desmoid tumor in children: results of a Children’s Oncology Group (COG) phase II study. *Pediatric Blood & Cancer*, 60(7), pp.1108–1112.
- Speiser, P.W. et al., 2010. Congenital Adrenal Hyperplasia Due to Steroid 21-Hydroxylase Deficiency: An Endocrine Society Clinical Practice Guideline. *The Journal of Clinical Endocrinology and Metabolism*, 95(9), pp.4133–4160. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2936060/> [Accessed April 22, 2015].
- Spreafico, F., 2015. Paediatrics: Towards evidence-based management of paediatric RCC. *Nature Reviews Urology*, advance online publication. Available at: <http://www.nature.com/nrurol/journal/vaop/ncurrent/full/nrurol.2015.158.html#access> [Accessed July 16, 2015].
- Swerdlow, S. et al., 2008. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues, Fourth Edition, Available at: <http://apps.who.int/bookorders/anglais/detart1.jsp?codlan=1&codcol=70&codcch=4002>.
- Thomsen, T.R. & Rodnitzky, R.L., 2010. Juvenile parkinsonism: epidemiology, diagnosis and treatment. *CNS drugs*, 24(6), pp.467–477.
- Trippett, T.M. et al., 2009. Phase I and Pharmacokinetic Study of Cetuximab and Irinotecan in Children With Refractory Solid Tumors: A Study of the Pediatric Oncology Experimental Therapeutic Investigators’ Consortium. *Journal of Clinical Oncology*, 27(30), pp.5102–5108. Available at: <http://jco.ascopubs.org/content/27/30/5102> [Accessed May 19, 2014].
- Trobaugh-Lotrario, A.D. & Katzenstein, H.M., 2012. Chemotherapeutic approaches for newly diagnosed hepatoblastoma: past, present, and future strategies. *Pediatric Blood & Cancer*, 59(5), pp.809–812.
- Vassal, G., 2009. Will children with cancer benefit from the new European Paediatric Medicines Regulation? *European Journal of Cancer* (Oxford, England: 1990), 45(9), pp.1535–1546. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19419857> [Accessed May 11, 2009].
- Vassal, G., Geoerger, B. & Morland, B., 2013. Is the European Pediatric Medicine Regulation Working for Children and Adolescents with Cancer? *Clinical Cancer Research*, 19(6), pp.1315–1325. Available at: <http://clincancerres.aacrjournals.org/cgi/doi/10.1158/1078-0432.CCR-12-2551> [Accessed November 4, 2013].
- Veal, G.J. et al., 2007. Adaptive dosing and platinum-DNA adduct formation in children receiving high-dose carboplatin for the treatment of solid tumours. *British Journal of Cancer*, 96(5), pp.725–731.

Warwick, A.B. et al., 2013. Phase 2 Trial of Pemetrexed in Children and Adolescents with Refractory Solid Tumors: a Children's Oncology Group Study. *Pediatric blood & cancer*, 60(2), pp.237–241. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3463652/> [Accessed April 22, 2015].

Whelan, J.S. et al., 2015. EURAMOS-1, an international randomised study for osteosarcoma: results from pre-randomisation treatment. *Annals of Oncology*, 26(2), pp.407–414. Available at: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4304379/> [Accessed March 24, 2015].