

12 June 2025  
 EMA/354670/2023  
 Scientific Evidence Generation

## Register of deadlines to put a medicinal product on the market

In accordance with Article 33 of the REGULATION (EC) No 1901/2006 OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 12 December 2006

### Disclaimer

This Register may not be exhaustive for non-centrally authorised products.

Product invented name	Paediatric procedure number	Active substance(s)	Authorised paediatric indication(s)	Initial Marketing Authorisation date	Variation date for paediatric indication	Deadline to place product on the market	Product with paediatric indication placed on the market*
Cancidas	EMA-000010-PIP01-07-M01	casprofungin	Treatment of invasive aspergillosis in adult or paediatric patients who are refractory to or intolerant of amphotericin B, lipid formulations of amphotericin B and / or itraconazole. Refractoriness is defined as progression of infection or failure to improve after a minimum of 7 days of prior therapeutic doses of effective antifungal therapy. Empirical therapy for presumed fungal infections (such as Candida or Aspergillus) in febrile, neutropaenic adult or paediatric patients.	24/10/2001	26/11/2008	26/11/2010	Yes
PegIntron ViraferonPeg	EMA-000071-PIP01-07	peginterferon alfa-2b	PegIntron/ViraferonPeg is indicated in a combination regimen with ribavirin for the treatment of children 3 years of age and older and adolescents, who have chronic hepatitis C, not previously treated, without liver decompensation, and who are positive for HCV-RNA. When deciding not to defer treatment until adulthood, it is important to consider that the combination therapy induced a growth inhibition. The reversibility of growth inhibition is uncertain. The decision to treat should be made on a case by case basis (see section 4.4)."	25/05/2000	11/11/2009	11/11/2011	Yes.
PegIntron ViraferonPeg	EMA-C-000384-PIP01-08	Peginterferon alfa-2b	PegIntron/ViraferonPeg is indicated in a combination regimen with ribavirin for the treatment of children 3 years of age and older and adolescents, who have chronic hepatitis C, not previously treated, without liver decompensation, and who are positive for HCV-RNA. When deciding not to defer treatment until adulthood, it is important to consider that the combination therapy induced a growth inhibition. The reversibility of growth inhibition is uncertain. The decision to treat should be made on a case by case basis (see section 4.4)."	25/05/2000	11/11/2009	11/11/2011	Yes.
Rebetol	EMA-000070-PIP01-07	ribavirin	Rebetol is indicated, in a combination regimen with peginterferon alfa-2b or interferon alfa-2b, for the treatment of children 3 years of age and older and adolescents, who have chronic hepatitis C, not previously treated, without liver decompensation, and who are positive for HCV-RNA. 3 When deciding to not to defer treatment until adulthood, it is important to consider that the combination therapy induced a growth inhibition. The reversibility of growth inhibition is uncertain. The decision to treat should be made on a case by case basis (see section 4.4).	07/05/1999	11/11/2009	11/11/2011	Yes
Orencia	EMA-000118-PIP01-07-M01	abatacept	Orencia in combination with methotrexate is indicated for the treatment of moderate to severe active polyarticular juvenile idiopathic arthritis (JIA) in paediatric patients 6 years of age and older who have had an insufficient response to other DMARDs including at least one TNF inhibitor. Orencia can be given as monotherapy in case of intolerance to methotrexate or when treatment with methotrexate is inappropriate.	21/05/2007	20/01/2010	20/01/2012	Yes
Cozaar	EMA-000008-PIP01-07	losartan potassium	Treatment of essential hypertension in adults and children and adolescents 6-18 years of age.	23/02/2010	23/02/2010	23/02/2012	Yes
Xalatan	EMA-000011-PIP01-07-M03	latanoprost	Reduction of elevated intraocular pressure in paediatric patients with elevated intraocular pressure and paediatric glaucoma.	29/05/1997	15/10/2010	15/10/2012	Yes

Pediacel	EMA-000278-PIP01-08-M01	Purified diphtheria toxoid, Purified tetanus toxoid, Five component acellular pertussis [Purified Pertussis Toxoid (PT), Purified Filamentous Haemagglutinin (FHA), Purified Fimbriae Types 2 and 3 (FIM) and Purified Pertactin (PRN)], Inactivated poliomyelitis vaccine (Vero) – Type 1 (Mahoney), Type 2 (MEF-1) and Type 3 (Saukett), Purified polyribosylribitol phosphate capsular polysaccharide of Haemophilus influenzae type b covalently bound to Tetanus protein (PRP-T)	Pediacel is indicated for primary and booster vaccination against diphtheria, tetanus, pertussis, poliomyelitis and invasive Haemophilus influenzae type b disease in infants and children from the age of 6 weeks up to the fourth birthday. Pediacel should be used in accordance with applicable official recommendations.	03/12/2010	03/12/2010	03/12/2012	Yes.
Nexium and associated names	EMA-000331-PIP01-08-M01	esomeprazole sodium / esomeprazole magnesium trihydrate	Children and adolescents with duodenal ulcers caused by H. Pylori infection.	09/12/2000	15/04/2011	15/04/2013	Yes
Sortis and associated names, As well as: Lipitor, Tahor, Xarator, Zarator, Liprimar, Totalip, Torvast, Cardyl	EMA-000073-PIP01-07	atorvastatin calcium (trihydrate)	Atorvastatin is indicated as an adjunct to diet for reduction of elevated total cholesterol, apolipoprotein B, and triglycerides in patients with primary hypercholesterolaemia including familial hypercholesterolaemia (heterozygous variant) or combined (mixed) hyperlipidaemia (Corresponding to Types IIa and IIb of the Fredrickson classification) when response to diet and other nonpharmacological measures is inadequate. Atorvastatin is also indicated to reduce total-C and LDL-C in patients with homozygous familial hypercholesterolaemia as an adjunct to other lipid-lowering treatments (e.g. LDL apheresis) or if such treatments are unavailable.	03/08/2010	05/05/2011	05/05/2013	Yes
Diovan	EMA-000005-PIP01-07-M01	valsartan	Treatment of hypertension in children and adolescents 6 - 18 years of age.	12/05/2010	11/05/2011	11/05/2013	Yes
Viramune	EMA-000391-PIP01-08-M01	nevirapine	Tablets and oral suspension Viramune is indicated in combination with other antiretroviral medicinal products for the treatment of HIV-1-infected adults, adolescents, and children of any age (see section 4.4). Prolonged-release tablets Viramune is indicated in combination with other antiretroviral medicinal products for the treatment of HIV-1-infected adolescents and children three years and above and able to swallow tablets (see section 4.2 and 4.4).	05/02/1998	05/09/2011	05/09/2013	Yes
Gardasil	EMA-000375-PIP01-08-M02	human papillomavirus type 6 L1 protein / human papillomavirus type 11 L1 protein / human papillomavirus type 16 L1 protein / human Papillomavirus type 18 L1 protein	Prevention of premalignant genital lesions (cervical, vulvar, vaginal, anal, perineal, perianal, penile), cervical, anal, perineal, and perianal cancer, and external genital warts (condyloma acuminata) causally related to human papillomavirus (HPV) types 6, 11, 16 and 18 in males.	20/09/2006	16/11/2011	16/11/2013	Yes
Remicade	EMA-000549-PIP01-09-M01	infliximab	Paediatric Crohn's disease Remicade is indicated for treatment of severe, active Crohn's disease, in children and adolescents aged six to 17 years, who have not responded to conventional therapy including a corticosteroid, an immunomodulator and primary nutrition therapy; or who are intolerant to or have contraindications for such therapies. Remicade has been studied only in combination with conventional immunosuppressive therapy. Paediatric ulcerative colitis Remicade is indicated for treatment of severely active ulcerative colitis, in paediatric patients aged six to 17 years, who have had an inadequate response to conventional therapy including corticosteroids and 6-MP or AZA, or who are intolerant to or have medical contraindications for such therapies.	13/08/1999	21/02/2012	21/02/2014	Yes
RotaTeq	EMA-000967-PIP01-10-M01	rotavirus type P1A[8]/rotavirus type G3/rotavirus type G1/rotavirus type G4/rotavirus type G2	RotaTeq is indicated for the active immunisation of infants from the age of six weeks to 32 weeks for prevention of gastroenteritis due to rotavirus infection. RotaTeq is to be used on the basis of official recommendations.	27/06/2006	21/02/2012	21/02/2014	Yes

Enbrel	EMA-000299-PIP01-08-M03	etanercept	Juvenile idiopathic arthritis Treatment of polyarthritis (rheumatoid factor positive or negative) and extended oligoarthritis in children and adolescents from the age of 2 years who have had an inadequate response to, or who have proved intolerant of, methotrexate. Treatment of psoriatic arthritis in adolescents from the age of 12 years who have had an inadequate response to, or who have proved intolerant of, methotrexate. Treatment of enthesitis-related arthritis in adolescents from the age of 12 years who have had an inadequate response to, or who have proved intolerant of, conventional therapy. Enbrel has not been studied in children aged less than 2 years. Paediatric plaque psoriasis Treatment of chronic severe plaque psoriasis in children and adolescents from the age of 6 years who are inadequately controlled by, or are intolerant to, other systemic therapies or phototherapies.	03/02/2000	31/07/2012	31/07/2014	Yes
Glivec	EMA-000463-PIP01-08-M03	imatinib mesilate	Treatment of paediatric patients with newly diagnosed Philadelphia chromosome positive acute lymphoblastic leukaemia (Ph+ALL) integrated with chemotherapy.	07/11/2001	27/06/2013	27/06/2015	Yes
Ezetrol, Ezetimibe MSD SP and associated names, Viemm and associated names, Zient and associated names	EMA-000007-PIP01-07-M02	ezetimibe	Treatment of primary hypercholesterolaemia Ezetimibe, co-administered with an HMG-CoA reductase inhibitor (statin), is indicated as adjunctive therapy to diet for use in patients with primary (heterozygous familial and non-familial) hypercholesterolaemia who are not appropriately controlled with a statin alone. Ezetimibe monotherapy is indicated as adjunctive therapy to diet for use in patients with primary (heterozygous familial and non-familial) hypercholesterolaemia in whom a statin is considered inappropriate or is not tolerated. Treatment of homozygous familial hypercholesterolaemia (HoFH) Authorised indications: Ezetimibe, co-administered with a statin, is indicated as adjunctive therapy to diet for use in patients with HoFH. Patients may also receive adjunctive treatments (e.g. LDL apheresis). Treatment of homozygous sitosterolaemia (phytosterolaemia) Authorised indication: Ezetimibe is indicated as adjunctive therapy to diet for use in patients with homozygous familial sitosterolaemia. <b>Children and adolescents ≥ 10 years</b> (pubertal status: boys Tanner Stage II and above and girls who are at least one year post-menarche): No dosage adjustment is required (see section 5.2). The clinical experience in paediatric and adolescent patients (aged 10-17 years old) is, however, limited. When Ezetrol is administered with simvastatina statin, the dosage instructions for simvathe statin, in adolescents children should be consulted. <b>Children &gt;6 and &lt;10 years:</b> There is limited data on safety and efficacy in this age group. Ezetrol is not recommended for use in children below age 10 due to insufficient data on safety and efficacy (see section s 5.1 and 5.2). <b>Children &lt;6 years:</b> There is no available data on use of Ezetrol in this age group.	17/10/2002	30/07/2013	30/07/2015	Yes
Prezista	EMA-000038-PIP01-07-M03	darunavir (as ethanolate)	Prezista, co-administered with low dose ritonavir is indicated in combination with other antiretroviral medicinal products for the treatment of human immunodeficiency virus (HIV-1) infection in adult and paediatric patients from the age of 3 years and at least 15 kg body weight (see section 4.2). In deciding to initiate treatment with Prezista co-administered with low dose ritonavir, careful consideration should be given to the treatment history of the individual patient and the patterns of mutations associated with different agents. Genotypic or phenotypic testing (when available) and treatment history should guide the use of Prezista.	12/02/2007	19/09/2013	19/09/2015	Yes
Invega	EMA-000014-PIP01-07-M06	paliperidone palmitate/paliperidone	Invega is indicated for the treatment of schizophrenia in adults and in adolescents 15 years and older.	25/06/2007	23/05/2014	23/05/2016	Yes
Baraclude	EMA-000339-PIP02-09-M03	entecavir monohydrate	Baraclude is also indicated for the treatment of chronic HBV infection in nucleoside naive paediatric patients from 2 to < 18 years of age with compensated liver disease who have evidence of active viral replication and persistently elevated serum ALT levels, or histological evidence of moderate to severe inflammation and/or fibrosis.	26/06/2006	22/08/2014	22/08/2016	Yes

Travatan	EMA-001271-PIP01-12-M01	travoprost	Decrease of elevated intraocular pressure in paediatric patients aged 2 months to < 18 years with ocular hypertension or paediatric glaucoma (see section 5.1).	27/11/2001	19/12/2014	19/12/2016	Yes
Emend	EMA-000144-PIP01-07-M05	aprepitant	Capsules: Prevention nausea and vomiting associated with highly and moderately emetogenic cancer chemotherapy in adults and adolescents from the age of 12. EMEND 125 mg/80 mg is given as part of combination therapy Powder for oral suspension: Prevention of nausea and vomiting associated with highly and moderately emetogenic cancer chemotherapy in children, toddlers and infants from the age of 6 months to less than 12 years. EMEND powder for oral suspension is given as part of combination therapy.	11/11/2003	17/12/2015	17/12/2017	No
Revolade	EMA-000170-PIP01-07-M03	eltrombopag (olamine)	Revolade is indicated for chronic immune (idiopathic) thrombocytopenic purpura (ITP) patients aged 1 year and above who are refractory to other treatments (e.g. corticosteroids, immunoglobulins) (see sections 4.2 and 5.1)	11/03/2010	04/04/2016	04/04/2018	No
HyQvia	EMA-000872-PIP01-10-M03	human normal immunoglobulin	Replacement therapy in adults, children and adolescents (0-18 years) ( $\geq 18$ years) in: • Primary immunodeficiency syndromes with impaired antibody production (see section 4.4). • Hypogammaglobulinaemia and recurrent bacterial infections in patients with chronic lymphocytic leukaemia (CLL), in whom prophylactic antibiotics have failed or are contra-indicated. • Hypogammaglobulinaemia and recurrent bacterial infections in multiple myeloma (MM) patients. • Hypogammaglobulinaemia in patients pre- and post-allogeneic hematopoietic stem cell transplantation (HSCT).	16/05/2013	01/06/2016	01/06/2018	Yes
Ryzodeg	EMA-000479-PIP01-08-M03	insulin degludec/insulin aspart	Treatment of diabetes mellitus in adults, adolescents and children from the age of 2 years	21/01/2013	22/07/2016	22/07/2018	Yes
Caprelsa	EMA-000052-PIP01-07-M03	vandetanib	Caprelsa is indicated for the treatment of aggressive and symptomatic medullary thyroid cancer (MTC) in patients with unresectable locally advanced or metastatic disease. Caprelsa is indicated in adults, children and adolescents aged 5 years and older. For patients in whom re-arranged-during-transfection (RET) mutation is not known or is negative, a possible lower benefit should be taken into account before individual treatment decision.	17/02/2012	16/12/2016	16/12/2018	Yes
Celsentri	EMA-000020-PIP01-07-M05	maraviroc	Celsentri, in combination with other antiretroviral medicinal products, is indicated for treatment-experienced adults, adolescents and children of 2 years of age and older and weighing at least 10 kg infected with only CCR5-tropic HIV-1 detectable	18/09/2007	06/07/2017	06/07/2019	No
Kaletra	EMA-001005-PIP01-10-M01	lopinavir/ritonavir	Kaletra is indicated in combination with other antiretroviral medicinal products for the treatment of human immunodeficiency virus (HIV-1) infected adults, adolescents and children from 14 days and older. The choice of Kaletra to treat protease inhibitor experienced HIV-1 infected patients should be based on individual viral resistance testing and treatment history of patients (see sections 4.4 and 5.1).	19/03/2001	26/07/2017	26/07/2019	No
Mimpara	EMA-000078-PIP01-07-M08	cinacalcet (as hydrochloride)	Treatment of secondary hyperparathyroidism (HPT) in children aged 3 years and older with end-stage renal disease (ESRD) on maintenance dialysis therapy in whom secondary HPT is not adequately controlled with standard of care therapy.	22/10/2004	28/08/2017	28/08/2019	No

Humira	EMA-000366-PIP05-12-M02	adalimumab	<p>Juvenile idiopathic arthritis</p> <p>Polyarticular juvenile idiopathic arthritis: Humira in combination with methotrexate is indicated for the treatment of active polyarticular juvenile idiopathic arthritis, in patients from the age of 2 years who have had an inadequate response to one or more disease-modifying anti-rheumatic drugs (DMARDs). Humira can be given as monotherapy in case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate (for the efficacy in monotherapy see section 5.1). Humira has not been studied in patients aged less than 2 years.</p> <p>Enthesitis-related arthritis: Humira is indicated for the treatment of active enthesitis-related arthritis in patients, 6 years of age and older, who have had an inadequate response to, or who are intolerant of, conventional therapy (see section 5.1).</p> <p>Paediatric plaque psoriasis</p> <p>Humira is indicated for the treatment of severe chronic plaque psoriasis in children and adolescents from 4 years of age who have had an inadequate response to or are inappropriate candidates for topical therapy and phototherapies.</p> <p>Paediatric Crohn's disease</p> <p>Humira is indicated for the treatment of moderately to severely active Crohn's disease in paediatric patients (from 6 years of age) who have had an inadequate response to conventional therapy including primary nutrition therapy and a corticosteroid and/or an immunomodulator, or who are intolerant to or have contraindications for such therapies.</p> <p>Paediatric Uveitis</p> <p>Humira is indicated for the treatment of paediatric chronic non-infectious anterior uveitis in patients from 2 years of age who have had an inadequate response to or are intolerant to conventional therapy, or in whom conventional therapy is inappropriate.</p>	08/09/2003	05/09/2017	05/09/2019	No
Firazyr	EMA-000408-PIP01-08-M05	icatibant	Firazyr is indicated for symptomatic treatment of acute attacks of hereditary angioedema (HAE) in adults, adolescents and children aged 2 years and older, with C1-esterase-inhibitor deficiency.	11/07/2008	19/10/2017	19/10/2019	Yes
Stribild	EMA-00970-PIP01-10-M01	elvitegravir / cobicistat / emtricitabine / tenofovir disoproxil	Stribild is indicated for the treatment of HIV-1 infection in adolescents aged 12 to < 18 years weighing $\geq$ 35 kg who are infected with HIV-1 without known mutations associated with resistance to any of the three antiretroviral agents in Stribild and who have experienced toxicities which preclude the use of other regimens that do not contain tenofovir disoproxil.	24/05/2013	19/10/2017	19/10/2019	Yes
Exjade	EMA-001103-PIP01-10-M03	<input type="checkbox"/> deferasirox	<p>EXJADE is indicated for the treatment of chronic iron overload due to frequent blood transfusions (<math>\geq</math> 7 ml/kg/month of packed red blood cells) in patients with beta thalassaemia major aged 6 years and older.</p> <p>EXJADE is also indicated for the treatment of chronic iron overload due to blood transfusions when deferoxamine therapy is contraindicated or inadequate in the following patient groups:</p>	28/08/2006	11/11/2017	11/11/2019	Pending
Tasigna	EMA-000290-PIP01-08-M04	nilotinib	<p>Tasigna is indicated for the treatment of:</p> <ul style="list-style-type: none"> <li>• adult and paediatric patients with newly diagnosed Philadelphia chromosome positive chronic myelogenous leukaemia (CML) in the chronic phase,</li> <li>• paediatric patients with chronic phase Philadelphia chromosome positive CML with resistance or intolerance to prior therapy including imatinib.</li> </ul>	19/11/2007	15/11/2017	15/11/2019	No
Yervoy	EMA-000117-PIP02-10-M07	ipilimumab	Yervoy as monotherapy is indicated for the treatment of advanced (unresectable or metastatic) melanoma in adults, and adolescents 12 years of age and older (see section 4.4).	13/07/2011	18/01/2018	18/01/2020	Yes
Truvada	EMA-001091-PIP02-15	tenofovir disoproxil fumarate / emtricitabine	<p>Treatment of HIV-1 infection:</p> <p>Truvada is indicated in antiretroviral combination therapy for the treatment of HIV-1 infected adults (see section 5.1).</p> <p>Truvada is also indicated for the treatment of HIV-1 infected adolescents, with NRTI resistance or toxicities precluding the use of first line agents (see sections 4.2, 4.4 and 5.1).</p> <p>Pre-exposure prophylaxis (PrEP):</p> <p>Truvada is indicated in combination with safer sex practices for pre-exposure prophylaxis to reduce the risk of sexually acquired HIV-1 infection in adults and adolescents at high risk (see sections 4.2, 4.4 and 5.1).</p>	20/02/2005	05/02/2018	05/02/2020	Yes

Kineret	EMA-001212-PIP01-11	anakinra	<p>Periodic fever syndromes</p> <p>Kineret is indicated for the treatment of the following autoinflammatory periodic fever syndromes in adults, adolescents, children and infants aged 8 months and older with a body weight of 10 kg or above:</p> <p>1. Cryopyrin-Associated Periodic Syndromes (CAPS)</p> <p>Kineret is indicated for the treatment of CAPS, including:</p> <ul style="list-style-type: none"> <li>• Neonatal-Onset Multisystem Inflammatory Disease (NOMID) / Chronic Infantile Neurological, Cutaneous, Articular Syndrome (CINCA)</li> <li>• Muckle-Wells Syndrome (MWS)</li> <li>• Familial Cold Autoinflammatory Syndrome (FCAS)</li> </ul> <p>2. Familial Mediterranean Fever (FMF)</p> <p>Kineret is indicated for the treatment of Familial Mediterranean Fever (FMF). Kineret should be given in combination with colchicine, if appropriate.</p> <p>Still's Disease</p> <p>Kineret is indicated in adults, adolescents, children and infants aged 8 months and older with a body weight of 10 kg or above for the treatment of Still's disease, including Systemic Juvenile Idiopathic Arthritis (SJIA) and Adult-Onset Still's Disease (AOSD), with active systemic features of moderate to high disease activity, or in patients with continued disease activity after treatment with non-steroidal anti-inflammatory drugs (NSAIDs) or glucocorticoids.</p> <p>Kineret can be given as monotherapy or in combination with other anti-inflammatory drugs and disease-modifying antirheumatic drugs (DMARDs).</p>	08/03/2002	06/04/2018	06/04/2020	Yes
Ivemend	EMA-000406-PIP01-08-M04	fosaprepitant dimeglumine	Prevention of nausea and vomiting associated with highly and moderately emetogenic cancer chemotherapy in adults and paediatric patients aged 6 months and older.	11/01/2008	30/04/2018	30/04/2020	Yes
Inovelon	EMA-000709-PIP01-09-M05	rufinamide	Inovelon is indicated as adjunctive therapy in the treatment of seizures associated with Lennox-Gastaut syndrome in patients 1 years of age and older.	16/01/2007	03/08/2018	03/08/2020	No
Isentress	EMA-000279-PIP01-08-M05	raltegravir	Isentress is indicated in combination with other anti-retroviral medicinal products for the treatment of human immunodeficiency virus (HIV-1) infection / ISENTRESS 600 mg film-coated tablets is indicated in combination with other anti-retroviral medicinal products for the treatment of human immunodeficiency virus (HIV-1) infection in adults, and paediatric patients weighing at least 40 kg.	19/12/2007	23/03/2018	23/03/2020	No
RoActemra	EMA-000309-PIP01-08-M07	tocilizumab	<p>RoActemra is indicated for the treatment of active systemic juvenile idiopathic arthritis (sJIA) in patients 2 years of age and older, who have responded inadequately to previous therapy with NSAIDs and systemic corticosteroids. RoActemra can be given as monotherapy (in case of intolerance to MTX or where treatment with MTX is inappropriate) or in combination with MTX.</p> <p>RoActemra in combination with methotrexate (MTX) is indicated for the treatment of juvenile idiopathic polyarthritis (pJIA; rheumatoid factor positive or negative and extended oligoarthritis) in patients 2 years of age and older, who have responded inadequately to previous therapy with MTX.</p> <p>RoActemra can be given as monotherapy in case of intolerance to MTX or where continued treatment with MTX is inappropriate.</p> <p>RoActemra is indicated for the treatment and prophylaxis of chimeric antigen receptor (CAR) T cell-induced severe or life-threatening cytokine release syndrome (CRS) in adults and paediatric patients 2 years of age and older.</p>	15/01/2009	12/04/2018	12/04/2020	Yes
Sprycel	EMA-000567-PIP01-09-M05	dasatinib (as monohydrate)	<p>Sprycel is indicated for the treatment of paediatric patients with:</p> <ul style="list-style-type: none"> <li>• newly diagnosed Ph+ CML in chronic phase (Ph+ CML-CP) or Ph+ CML-CP resistant or intolerant to prior therapy including imatinib.</li> <li>• newly diagnosed Ph+ ALL in combination with chemotherapy.</li> </ul>	20/11/2006	02/07/2018	02/07/2020	No
Coagadex	EMA-000971-PIP01-10-M03	human coagulation factor X	Coagadex is indicated for the treatment and prophylaxis of bleeding episodes and for perioperative management in patients with hereditary factor X deficiency. Coagadex is indicated in all age groups.	16/03/2016	27/08/2018	27/08/2020	Yes
Nucala	EMA-000069-PIP02-10-M08	mepolizumab	Nucala is indicated as an add-on treatment for severe refractory eosinophilic asthma in adults, adolescents and children aged 6 years and older.	01/12/2015	27/08/2018	27/08/2020	Yes

Gilenya	EMA-000087-PIP01-07-M05	fingolimod (hydrochloride)	Gilenya is indicated as single disease modifying therapy in highly active relapsing remitting multiple sclerosis for the following groups of adult patients and paediatric patients aged 10 years and older: <ul style="list-style-type: none"> <li>Patients with highly active disease despite a full and adequate course of treatment with at least one disease modifying therapy (for exceptions and information about washout periods see sections 4.4 and 5.1).</li> </ul> or <ul style="list-style-type: none"> <li>Patients with rapidly evolving severe relapsing remitting multiple sclerosis defined by 2 or more disabling relapses in one year, and with 1 or more Gadolinium enhancing lesions on brain MRI or a significant increase in T2 lesion load as compared to a previous recent MRI.</li> </ul>	17/03/2011	22/11/2018	22/11/2020	No
Ravicti	EMA-000297-PIP02-12-M02	glycerol phenylbutyrate	Ravicti is indicated for use as adjunctive therapy for chronic management of patients with urea cycle disorders (UCDs) including deficiencies of carbamoyl phosphate synthetase I (CPS), ornithine carbamoyltransferase (OTC), argininosuccinate synthetase (ASS), argininosuccinate lyase (ASL), arginase I (ARG) and ornithine translocase deficiency hyperornithinaemia-hyperammonaemia homocitrullinuria syndrome (HHH) who cannot be managed by dietary protein restriction and/or amino acid supplementation alone. Ravicti must be used with dietary protein restriction and, in some cases, dietary supplements (e.g., essential amino acids, arginine, citrulline, protein-free calorie supplements).	26/11/2015	18/12/2018	18/12/2020	Yes
Simponi	EMA-000265-PIP01-08-M04	golimumab	Simponi in combination with MTX is indicated for the treatment of polyarticular juvenile idiopathic arthritis in children 2 years of age, who have responded inadequately to previous therapy with MTX.	01/10/2009	18/02/2019	18/02/2021	No
Orencia	EMA-000118-PIP02-10-M03	abatacept	Orencia in combination with methotrexate is indicated for the treatment of moderate to severe active polyarticular juvenile idiopathic arthritis (pJIA) in paediatric patients 2 years of age and older who have had an inadequate response to previous DMARD therapy. Orencia can be given as monotherapy in case of intolerance to methotrexate or when treatment with methotrexate is inappropriate.	21/05/2007	08/04/2019	08/04/2021	No
Mozobil	EMA-000174-PIP01-07-M03	plerixafor	Mozobil is indicated in combination with G-CSF to enhance mobilisation of haematopoietic stem cells to the peripheral blood for collection and subsequent autologous transplantation in children with lymphoma or solid malignant tumours, either: <ul style="list-style-type: none"> <li>pre-emptively, when circulating stem cell count on the predicted day of collection after adequate mobilization with G-CSF (with or without chemotherapy) is expected to be insufficient with regards to desired hematopoietic stem cells yield, or</li> <li>who previously failed to collect sufficient haematopoietic stem cells (see section 4.2).</li> </ul>	30/07/2009	13/05/2019	13/05/2021	Yes
Zinforo	EMA-C-000769-PIP01-09-M08	ceftaroline fosamil	Zinforo is indicated for the treatment of the following infections in neonates, infants, children, adolescents and adults(see sections 4.4 and 5.1): <ul style="list-style-type: none"> <li>Complicated skin and soft tissue infections (cSSTI)</li> <li>Community-acquired pneumonia (CAP)</li> </ul> Consideration should be given to official guidance on the appropriate use of antibacterial agents.	22/08/2012	25/07/2019	25/07/2021	Yes
Victoza	EMA-000128-PIP01-07-M08	liraglutide	Victoza is indicated for the treatment of adults, adolescents and children aged 10 years and above with insufficiently controlled type 2 diabetes mellitus as an adjunct to diet and exercise: <ul style="list-style-type: none"> <li>as monotherapy when metformin is considered inappropriate due to intolerance or contraindications</li> <li>in addition to other medicinal products for the treatment of diabetes.</li> </ul> For study results with respect to combinations, effects on glycaemic control and cardiovascular events, and the populations studied, see sections 4.4, 4.5 and 5.1.	30/06/2009	09/08/2019	09/08/2021	yes
Lucentis	EMA-000527-PIP04-13-M01	ranibizumab	Lucentis is indicated in preterm infants for the treatment of retinopathy of prematurity (ROP) with zone I (stage 1+, 2+, 3 or 3+), zone II (stage 3+) or AP-ROP (aggressive posterior ROP) disease.	22/01/2007	03/09/2019	03/09/2021	Yes
Benlysta	EMA-000520-PIP01-08	belimumab	Benlysta is indicated as add-on therapy in patients aged 5 years and older with active, autoantibody positive systemic lupus erythematosus (SLE) with a high degree of disease activity (e.g., positive anti dsDNA and low complement) despite standard therapy.	13/07/2011	21/10/2019	21/10/2021	Yes

Stelara	EMA-000311-PIP01-08-M04	ustekinumab	Stelara is indicated for the treatment of moderate to severe plaque psoriasis in children and adolescent patients from the age of 6 years and older, who are inadequately controlled by, or are intolerant to, other systemic therapies or phototherapies.	15/01/2009	20/01/2020	20/01/2022	Yes
Dificlir	EMA-000636-PIP01-09-M07	fidaxomicin	Dificlir film-coated tablets is indicated for the treatment of Clostridioides difficile infections (CDI) also known as C. difficile-associated diarrhoea (CDAD) in adult and paediatric patients with a body weight of at least 12.5 kg (see section 4.2 and 5.1). DIFICLIR granules for oral suspension is indicated for the treatment of Clostridioides difficile infections (CDI) also known as C. difficile-associated diarrhoea (CDAD) in adults	05/12/2011	13/02/2020	13/02/2022	No
MabThera	EMA-000308-PIP01-08-M04	rituximab	MabThera in combination with chemotherapy is indicated for the treatment of paediatric patients (aged ≥ 6 months to < 18 years old) with previously untreated advanced stage CD20 positive diffuse large B-cell lymphoma (DLBCL), Burkitt lymphoma (BL)/Burkitt leukaemia (mature B-cell acute leukaemia) (BAL) or Burkitt-like lymphoma (BLL).	02/06/1998 21/03/2014 (new formulation)	03/03/2020	03/03/2022	Yes
MabThera	EMA-000308-PIP02-11-M01	rituximab	MabThera in combination with chemotherapy is indicated for the treatment of paediatric patients (aged ≥ 6 months to < 18 years old) with previously untreated advanced stage CD20 positive diffuse large B-cell lymphoma (DLBCL), Burkitt lymphoma (BL)/Burkitt leukaemia (mature B-cell acute leukaemia) (BAL) or Burkitt-like lymphoma (BLL). MabThera, in combination with glucocorticoids, is indicated for the induction of remission in paediatric patients (aged ≥ 2 to < 18 years old) with severe, active granulomatosis with polyangiitis (Wegener's) (GPA) and microscopic polyangiitis (MPA).	02/06/1998 21/03/2014 (new formulation)	03/03/2020	03/03/2022	Yes
Intelence	EMA-000222-PIP01-08-M09	etravirine	Intelence, in combination with a boosted protease inhibitor and other antiretroviral medicinal products, is indicated for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in antiretroviral treatment-experienced adult patients and in antiretroviral treatment-experienced paediatric patients from 2 years of age (see sections 4.4, 4.5 and 5.1).	28/08/2008	28/04/2020	28/04/2022	No
Ruconest	EMA-000367-PIP01-08-M08	conestat alfa	Ruconest is indicated for treatment of acute angioedema attacks in adults, adolescents, and children (aged 2 years and above) with hereditary angioedema (HAE) due to C1 esterase inhibitor deficiency.	28/10/2010	28/04/2020	28/04/2022	Yes
Ecalta	EMA-000469-PIP01-08-M07	anidulafungin	Treatment of invasive candidiasis in adults and paediatric patients aged 1 month to < 18 years.	20/09/2007	03/06/2020	03/06/2022	Yes
Cablivi	EMA-001157-PIP01-11-M02	caplacizumab	Cablivi is indicated for the treatment of adults and adolescents of 12 years of age and older weighing at least 40 kg experiencing an episode of acquired thrombotic thrombocytopenic purpura (aTTP), in conjunction with plasma exchange and immunosuppression.	30/08/2018	09/06/2020	09/06/2022	Yes
Sovaldi	EMA-001276-PIP01-12-M02	sofosbuvir	Sovaldi is indicated in combination with other medicinal products for the treatment of chronic hepatitis C (CHC) in adults and paediatric patients aged 3 years and above (see sections 4.2, 4.4 and 5.1). For hepatitis C virus (HCV) genotype specific activity, see sections 4.4 and 5.1.	16/01/2014	25/06/2020	25/06/2022	No
Harvoni	EMA-001411-PIP01-12-M04	sofosbuvir / ledipasvir	Harvoni is indicated for the treatment of chronic hepatitis C (CHC) in adult and paediatric patients aged 3 years and above (see sections 4.2, 4.4 and 5.1). For hepatitis C virus (HCV) genotype-specific activity see sections 4.4 and 5.1.	17/11/2014	03/07/2020	03/07/2022	No
Latuda	EMA-001230-PIP01-11-M05	lurasidone hydrochloride	Latuda is indicated for the treatment of schizophrenia in adults and adolescent aged 13 years and over.	21/03/2014	25/08/2020	25/08/2022	Yes
Velphoro	EMA-001061-PIP01-10-M03	sucroferric oxyhydroxide	Velphoro is indicated for the control of serum phosphorus levels in paediatric patients 2 years of age and older with CKD stages 4-5 (defined by a glomerular filtration rate < 30 mL/min/1.73 m <sup>2</sup> ) or with CKD on dialysis.	26/08/2014	16/11/2020	16/11/2022	No

Humira	EMA-000366-PIP02-09-M06	adalimumab	<p>Juvenile idiopathic arthritis</p> <ul style="list-style-type: none"> <li>• Polyarticular juvenile idiopathic arthritis: Humira in combination with methotrexate is indicated for the treatment of active polyarticular juvenile idiopathic arthritis, in patients from the age of 2 years who have had an inadequate response to one or more disease-modifying anti-rheumatic drugs (DMARDs). Humira can be given as monotherapy in case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate (for the efficacy in monotherapy see section 5.1). Humira has not been studied in patients aged less than 2 years.</li> <li>• Enthesitis-related arthritis: Humira is indicated for the treatment of active enthesitis-related arthritis in patients, 6 years of age and older, who have had an inadequate response to, or who are intolerant of, conventional therapy (see section 5.1).</li> </ul> <p>Paediatric plaque psoriasis: Humira is indicated for the treatment of severe chronic plaque psoriasis in children and adolescents from 4 years of age who have had an inadequate response to or are inappropriate candidates for topical therapy and phototherapies.</p> <p>Paediatric Crohn's disease: Humira is indicated for the treatment of moderately to severely active Crohn's disease in paediatric patients (from 6 years of age) who have had an inadequate response to conventional therapy including primary nutrition therapy and a corticosteroid and/or an immunomodulator, or who are intolerant to or have contraindications for such therapies.</p> <p>Paediatric Uveitis: Humira is indicated for the treatment of paediatric chronic non-infectious anterior uveitis in patients from 2 years of age who have had an inadequate response to or are intolerant to conventional therapy, or in whom conventional therapy is inappropriate.</p> <p>Paediatric ulcerative colitis: Humira is indicated for the treatment of moderately to severely active ulcerative colitis in paediatric patients (from 6 years of age) who have had an inadequate response to conventional therapy including corticosteroids and/or 6-mercaptopurine (6-MP) or azathioprine (AZA), or who are intolerant to or have medical contraindications for such therapies.</p>	08/09/2003	20/11/2020	20/11/2022	Yes
Pradaxa	EMA-000081-PIP01-07-M11	dabigatran etexilate mesilate	<p>Treatment of VTE and prevention of recurrent VTE in paediatric patients from birth to less than 18 years of age.</p> <p>For age appropriate dose forms, see section 4.2.</p>	17/03/2008	11/01/2021	11/01/2023	Yes
Xarelto	EMA-000430-PIP01-08-M11	rivaroxaban	Xarelto, co-administered with acetylsalicylic acid (ASA), is indicated for the prevention of atherothrombotic events in adult patients with coronary artery disease (CAD) or symptomatic peripheral artery disease (PAD) at high risk of ischaemic events.	30/09/2008	21/01/2021	21/01/2023	No
Maviret	EMA-001832-PIP01-15-M02	glecaprevir / pibrentasvir	Maviret is indicated for the treatment of chronic hepatitis C virus (HCV) infection in adults and children aged 3 years and older.	26/07/2017	21/06/2021	21/06/2023	No
Spherox <sup>1</sup>	EMA-001264-PIP01-12-M02	spheroids of human autologous matrix-associated chondrocytes	Repair of symptomatic articular cartilage defects of the femoral condyle and the patella of the knee (International Cartilage Regeneration & Joint Preservation Society [ICRS] grade III or IV) with defect sizes up to 10 cm <sup>2</sup> in adults and adolescents with closed epiphyseal growth plate in the affected joint.	10/07/2017	28/06/2021	28/06/2023	Yes
Deltyba	EMA-001113-PIP01-10-M06	delamanid	Deltyba is indicated for use as part of an appropriate combination regimen for pulmonary multidrug resistant tuberculosis (MDR-TB) in adults, adolescents, children and infants with a body weight of at least 10 kg when an effective treatment regimen cannot otherwise be composed for reasons of resistance or tolerability.	27/04/2014	16/09/2021	16/09/2023	No
Vosevi	EMA-001822-PIP01-15-M01	sofosbuvir / voxilaprevir / velpatasvir	Vosevi is indicated for the treatment of chronic hepatitis C virus (HCV) infection in patients aged 12 years and older and weighing at least 30 kg.	26/07/2017	16/09/2021	16/09/2023	No
Cosentyx	EMA-000380-PIP01-08-M04	secukinumab	<ul style="list-style-type: none"> <li>- Paediatric plaque psoriasis: Cosentyx is indicated for the treatment of moderate to severe plaque psoriasis in children and adolescents from the age of 6 years who are candidates for systemic therapy.</li> <li>- Juvenile idiopathic arthritis (JIA): Enthesitis-related arthritis (ERA)</li> </ul>	14/01/2015	16/07/2021	16/07/2023	Pending

Zepatier	EMA-001604-PIP01-13-M03	elbasvir / grazoprevir	ZEPATIER is indicated for the treatment of chronic hepatitis C (CHC) in adult and paediatric patients 12 years of age and older who weigh at least 30 kg	22/07/2016	22/10/2021	22/10/2023	Yes
Nucala	EMA-000069-PIP04-13-M02	mepolizumab	- Severe eosinophilic asthma: Nucala is indicated as an add-on treatment for severe refractory eosinophilic asthma in adults, adolescents and children aged 6 years and older. - Eosinophilic granulomatosis with polyangiitis (EGPA): Nucala is indicated as an add-on treatment for patients aged 6 years and older with relapsing-remitting or refractory eosinophilic granulomatosis with polyangiitis (EGPA)	01/12/2015	12/11/2021	12/11/2023	Yes
Forxiga	EMA-000694-PIP01-09-M08	dapagliflozin propanediol monohydrate	Type 2 diabetes mellitus: Forxiga is indicated in adults and children aged 10 years and above for the treatment of insufficiently controlled type 2 diabetes mellitus as an adjunct to diet and exercise - as monotherapy when metformin is considered inappropriate due to intolerance. - in addition to other medicinal products for the treatment of type 2 diabetes.	11/11/2012	15/11/2021	15/11/2023	Yes
Epclusa	EMA-001646-PIP01-14-M02	Sofosbuvir / velpatasvir	Epclusa is indicated for the treatment of chronic hepatitis C virus (HCV) infection in patients 3 years of age and older	06/07/2016	07/01/2022	07/01/2024	Yes
Clensia	EMA-001356-PIP02-12-M04	Macrogol 4000/Sodium sulphate (as sodium sulfate anhydrous)/Simeticone/Sodium citrate/Citric acid (as citric acid anhydrous)/Sodium chloride/Potassium chloride dimethyl fumarate	Bowel cleansing prior to any clinical procedures requiring a clean bowel e.g. bowel endoscopy or radiology. Clensia is indicated for use in adults, adolescents and children aged 6 years and older.	09/09/2016	11/04/2024	11/04/2026	Yes
Tecfidera	EMA-000832-PIP01-10-M05	dimethyl fumarate	Tecfidera is indicated for the treatment of adult and paediatric patients aged 13 years and older with relapsing remitting multiple sclerosis (RRMS).	30/01/2014	13/05/2022	13/05/2024	Yes
Adjupanrix	EMA-000160-PIP01-07-M05	split influenza virus, inactivated, containing antigen: A/VietNam/1194/2004 (H5N1) like strain used (NIBRG-14)	Prophylaxis of influenza in an officially declared pandemic situation.	10/10/2009	24/05/2022	24/05/2024	Yes
Bydureon	EMA-000689-PIP01-09-M11	exenatide	Bydureon is indicated in adults, adolescents and children aged 10 years and above with type 2 diabetes mellitus to improve glycaemic control in combination with other glucose-lowering medicinal products including basal insulin, when the therapy in use, together with diet and exercise, does not provide adequate glycaemic control.	17/06/2011	30/05/2022	30/05/2024	Yes
Cosentyx	EMA-000380-PIP02-09-M04	secukinumab	- Paediatric plaque psoriasis: Cosentyx is indicated for the treatment of moderate to severe plaque psoriasis in children and adolescents from the age of 6 years who are candidates for systemic therapy. - Juvenile idiopathic arthritis (JIA): Enthesitis-related arthritis (ERA)	14/01/2015	20/06/2022	20/06/2024	Pending
Elonva	EMA-000306-PIP01-08-M04	corifollitropin alfa	Elonva is indicated for the treatment of adolescent males (14 years and older) with hypogonadotropic hypogonadism, in combination with human Chorionic Gonadotropin (hCG).	25/01/2010	21/06/2022	21/06/2024	Yes
Lonquex	EMA-001019-PIP01-10-M05	lipegfilgrastim	Lonquex is indicated in adults and in children 2 years of age and older for reduction in the duration of neutropenia and the incidence of febrile neutropenia in patients treated with cytotoxic chemotherapy for malignancy (with the exception of chronic myeloid leukaemia and myelodysplastic syndromes).	25/07/2013	22/07/2022	22/07/2024	Pending
Zerbaxa	EMA-001142-PIP01-11-M04	ceftolozane sulfate / tazobactam sodium	Zerbaxa is indicated for the treatment of the following infections in adult and paediatric patients: - Complicated intra-abdominal infections; - Acute pyelonephritis; - Complicated urinary tract infections.	18/09/2015	25/07/2022	25/07/2024	Yes

Genvoya	EMA-001460-PIP01-13-M05	elvitegravir / cobicistat / emtricitabine / tenofovir alafenamide	Genvoya is indicated for the treatment of human immunodeficiency virus-1 (HIV-1) infection without any known mutations associated with resistance to the integrase inhibitor class, emtricitabine or tenofovir in adults and paediatric patients aged from 2 years and with body weight at least 14 kg.	19/11/2015	03/10/2022	03/10/2024	Pending
Vaxneuvance	EMA-002215-PIP01-17-M03	pneumococcal polysaccharide conjugate vaccine (adsorbed)	Vaxneuvance is indicated for active immunisation for the prevention of invasive disease, pneumonia and acute otitis media caused by Streptococcus pneumoniae in infants, children and adolescents from 6 weeks to less than 18 years of age.	13/12/2021	21/10/2022	21/10/2024	Pending
Eylea	EMA-000236-PIP05-18	aflibercept	EYLEA is indicated in preterm infants for the treatment of <ul style="list-style-type: none"> <li>retinopathy of prematurity (ROP) with zone I (stage 1+, 2+, 3 or 3+), zone II (stage 2+ or 3+) or AP-ROP (aggressive posterior ROP) disease.</li> </ul>	21/11/2012	09/12/2022	09/12/2024	Pending
Adcirca	EMA-000452-PIP02-10-M06	tadalafil	Treatment of paediatric patients aged 2 years and above with pulmonary arterial hypertension (PAH) classified as WHO functional class II and III.	01/10/2008	24/02/2023	24/02/2025	Pending
Wakix	EMA-001176-PIP01-11-M06	pitolisant	Wakix is indicated in adults, adolescents and children from the age of 6 years for the treatment of narcolepsy with or without cataplexy.	31/03/2016	24/02/2023	24/02/2025	Yes
Trulicity	EMA-000783-PIP01-09-M06	dulaglutide	Trulicity is indicated for the treatment of patients 10 years and above with insufficiently controlled type 2 diabetes mellitus as an adjunct to diet and exercise <ul style="list-style-type: none"> <li>as monotherapy when metformin is considered inappropriate due to intolerance or contraindications</li> <li>in addition to other medicinal products for the treatment of diabetes.</li> </ul>	21/11/2014	06/03/2023	06/03/2025	Yes
Treosulfan	EMA-000883-PIP01-10-M05	Treosulfan	Treosulfan in combination with fludarabine is indicated as part of conditioning treatment prior to allogeneic haematopoietic stem cell transplantation (alloHSCT) in adult patients and in paediatric patients older than one month with malignant and non-malignant diseases.	20/06/2019	06/03/2023	06/03/2025	Yes
Dupilixent	EMA-001501-PIP01-13-M07	dupilumab	- Atopic dermatitis Adults and adolescents: Dupixent is indicated for the treatment of moderate-to-severe atopic dermatitis in adults and adolescents 12 years and older who are candidates for systemic therapy. Children 6 months to 11 years of age: Dupixent is indicated for the treatment of severe atopic dermatitis in children 6 months to 11 years old.	26/09/2017	15/03/2023	15/03/2025	Pending
Adempas	EMA-000718-PIP01-09-M06	Riociguat	Adempas is indicated for the treatment of PAH in paediatric patients aged less than 18 years of age and body weight $\geq$ 50 kg with WHO Functional Class (FC) II to III in combination with endothelin receptor antagonists.	27/03/2014	31/05/2023	31/05/2025	Pending
Opdivo	EMA-001407-PIP01-12-M03	nivolumab	Melanoma: OPDIVO as monotherapy or in combination with ipilimumab is indicated for the treatment of advanced (unresectable or metastatic) melanoma in adults and adolescents 12 years of age and older. Relative to nivolumab monotherapy, an increase in progression-free survival (PFS) and overall survival (OS) for the combination of nivolumab with ipilimumab is established only in patients with low tumour PD-L1.	19/06/2015	31/05/2023	31/05/2025	Yes
SOLIRIS	EMA-000876-PIP05-15-M05	eculizumab	Soliris is indicated in adults and children for the treatment of: <ul style="list-style-type: none"> <li>Paroxysmal nocturnal haemoglobinuria (PNH).</li> </ul> Evidence of clinical benefit is demonstrated in patients with haemolysis with clinical symptom(s) indicative of high disease activity, regardless of transfusion history. <ul style="list-style-type: none"> <li>Atypical haemolytic uremic syndrome (aHUS)</li> </ul>	20/06/2007	24/07/2023	24/07/2025	Yes
Mircera	EMA-000172-PIP01-07-M03	Methoxy polyethylene glycol-epoetin beta	Treatment of symptomatic anaemia associated with chronic kidney disease (CKD) in paediatric patients from 3 months to less than 18 years of age who are converting from another erythropoiesis stimulating agent (ESA) after their haemoglobin level was stabilised with the previous ESA	20/07/2007	26/07/2023	26/07/2025	Pending

Refixia	EMA-000731-PIP01-09-M03	Nonacog beta pegol (glycopegylated recombinant coagulation factor IX)	Treatment and prophylaxis of bleeding in patients with haemophilia B (congenital factor IX deficiency). Refixia can be used for all age groups.	02/06/2017	04/08/2023	04/08/2025	Yes
Olumiant	EMA-001220-PIP03-16-M02	baricitinib	Atopic dermatitis Baricitinib is indicated for the treatment of moderate to severe atopic dermatitis in adult and paediatric patients 2 years of age and older who are candidates for systemic therapy.	13/02/2017	18/10/2023	18/10/2025	Pending
TAKHZYRO	EMA-001864-PIP01-15-M07	Ianadelumab	TAKHZYRO is indicated for routine prevention of recurrent attacks of hereditary angioedema (HAE) in patients aged 2 years and older.	22/11/2018	15/11/2023	15/11/2025	Pending
Praluent	EMA-001169-PIP01-11-M05	alirocumab	Praluent is indicated in adults with primary hypercholesterolaemia (heterozygous familial and non-familial) or mixed dyslipidaemia, and in paediatric patients 8 years of age and older with heterozygous familial hypercholesterolaemia (HeFH) as an adjunct to diet: - in combination with a statin or statin with other lipid lowering therapies in patients unable to reach LDL-C goals with the maximum tolerated dose of a statin or, - alone or in combination with other lipid-lowering therapies in patients who are statin-	23/09/2015	15/11/2023	15/11/2025	Yes
Jardiance	EMA-000828-PIP01-09-M09	Empagliflozin	Type 2 diabetes mellitus Jardiance is indicated in adults and children aged 10 years and above for the treatment of insufficiently controlled type 2 diabetes mellitus as an adjunct to diet and exercise - as monotherapy when metformin is considered inappropriate due to intolerance - in addition to other medicinal products for the treatment of diabetes	22/05/2014	07/12/2023	07/12/2025	Yes
VeraSeal	EMA-001598-PIP01-13-M03	Human Fibrinogen/Human Thrombin	Supportive treatment in patients where standard surgical techniques are insufficient: - for improvement of haemostasis. - as suture support: in vascular surgery. VeraSeal is indicated in all age groups.	10/11/2017	25/01/2024	25/01/2026	Yes
Zinplava	EMA-001645-PIP01-14-M04	bezlotoxumab	ZINPLAVA is indicated for the prevention of recurrence of Clostridioides difficile infection (CDI) in adult and paediatric patients 1 year of age and older at high risk for recurrence of CDI.	18/01/2017	26/01/2024	26/01/2026	Pending
Kalydeco	EMA-000335-PIP01-08-M15	Ivacaftor	Kalydeco tablets are indicated: • As monotherapy for the treatment of adults, adolescents, and children aged 6 years and older and weighing 25 kg or more with cystic fibrosis (CF) who have an R117H CFTR mutation or one of the following gating (class III) mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene: G551D, G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N or S549R (see sections 4.4 and 5.1)	23/07/2012	25/04/2024	25/04/2026	Pending
Triumeq	EMA-001219-PIP01-11-M06	Dolutegravir/Abacavir/lamivudine	Triumeq is indicated for the treatment of Human Immunodeficiency Virus type 1 (HIV-1) infected adults, adolescents and children weighing at least 25 kg Triumeq is indicated for the treatment of Human Immunodeficiency Virus type 1 (HIV-1) infected children of at least 3 months of age and weighing at least 6 kg to less than 25 kg	31/08/2014	27/05/2024	27/05/2026	Pending

\*: As declared by Marketing Authorisation Holders; the product needs to be on the market in all Member States where the adult product was already commercialised and within the deadline as per Article 33 of the Paediatric Regulation (No 1906/2006).  
From 2023 AR, responses have been updated to indicate 'Yes' - The product has been placed in the market in all applicable Member States by the established deadline.;  
'No' - The product has not been placed on the market in all applicable Member States, or the deadline was not met; 'Pending' - The deadline for placing the product on the market has not yet been reached.

<sup>1</sup> : Name change due to Marketing Authorisation; former name: (co.don) chondrosphere