

EMA/104834/2016

European Medicines Agency decision

P/0071/2016

of 18 March 2016

on the acceptance of a modification of an agreed paediatric investigation plan for naltrexone (hydrochloride) / bupropion (hydrochloride) (Mysimba), (EMA-001373-PIP01-12-M01) in accordance with Regulation (EC) No 1901/2006 of the European Parliament and of the Council

Disclaimer

This Decision does not constitute entitlement to the rewards and incentives referred to in Title V of Regulation (EC) No 1901/2006.

Only the English text is authentic.

European Medicines Agency decision

P/0071/2016

of 18 March 2016

on the acceptance of a modification of an agreed paediatric investigation plan for naltrexone (hydrochloride) / bupropion (hydrochloride) (Mysimba), (EMA-001373-PIP01-12-M01) in accordance with Regulation (EC) No 1901/2006 of the European Parliament and of the Council

The European Medicines Agency,

Having regard to the Treaty on the Functioning of the European Union,

Having regard to Regulation (EC) No 1901/2006 of the European Parliament and of the Council of 12 December 2006 on medicinal products for paediatric use and amending Regulation (EEC) No. 1768/92, Directive 2001/20/EC, Directive 2001/83/EC and Regulation (EC) No 726/2004¹,

Having regard to Regulation (EC) No 726/2004 of the European Parliament and of the Council of 31 March 2004 laying down Community procedures for the authorisation and supervision of medicinal products for human and veterinary use and establishing a European Medicines Agency²,

Having regard to the European Medicines Agency's decision P/0188/2013 issued on 8 August 2013,

Having regard to the application submitted by Orexigen Therapeutics Ireland Limited on 9 November 2015 under Article 22 of Regulation (EC) No 1901/2006 proposing changes to the agreed paediatric investigation plan with a deferral and a waiver,

Having regard to the opinion of the Paediatric Committee of the European Medicines Agency, issued on 29 January 2016, in accordance with Article 22 of Regulation (EC) No 1901/2006,

Having regard to Article 25 of Regulation (EC) No 1901/2006,

Whereas:

- (1) The Paediatric Committee of the European Medicines Agency has given an opinion on the acceptance of changes to the agreed paediatric investigation plan.
- (2) It is therefore appropriate to adopt a decision on the acceptance of changes to the agreed paediatric investigation plan.

¹ OJ L 378, 27.12.2006, p.1.

² OJ L 136, 30.4.2004, p. 1.

Has adopted this decision:

Article 1

Changes to the agreed paediatric investigation plan for naltrexone (hydrochloride) / bupropion (hydrochloride) (Mysimba), prolonged-release tablet, oral use, are hereby accepted in the scope set out in the opinion of the Paediatric Committee of the European Medicines Agency annexed hereto, together with its appendices.

Article 2

This decision is addressed to Orexigen Therapeutics Ireland Limited, 2nd Floor Palmerston House, Fenian Street, 2 - Dublin, Ireland.

Done at London, 18 March 2016

For the European Medicines Agency
Zaïde Frias
Head of Division
Human Medicines Research and Development Support
(Signature on file)



EUROPEAN MEDICINES AGENCY
SCIENCE MEDICINES HEALTH

EMA/PDCO/743321/2015

London, 29 January 2016

Opinion of the Paediatric Committee on the acceptance of a modification of an agreed Paediatric Investigation Plan

EMA-001373-PIP01-12-M01

Scope of the application

Active substance(s):

Naltrexone (hydrochloride) / bupropion (hydrochloride)

Invented name:

Mysimba

Condition(s):

Treatment of obesity

Authorised indication(s):

See Annex II

Pharmaceutical form(s):

Prolonged-release tablet

Route(s) of administration:

Oral use

Name / corporate name of the PIP applicant:

Orexigen Therapeutics Ireland Limited

Information about the authorised medicinal product:

See Annex II



Basis for opinion

Pursuant to Article 22 of Regulation (EC) No 1901/2006 as amended, Orexigen Therapeutics Ireland Limited submitted to the European Medicines Agency on 9 November 2015 an application for modification of the agreed paediatric investigation plan with a deferral and a waiver as set out in the European Medicines Agency's decision P/0188/2013 issued on 8 August 2013.

The application for modification proposed changes to the agreed paediatric investigation plan.

The procedure started on 30 November 2015.

Scope of the modification

Some measures and timelines of the Paediatric Investigation Plan have been modified.

Opinion

1. The Paediatric Committee, having assessed the application in accordance with Article 22 of Regulation (EC) No 1901/2006 as amended, recommends as set out in the appended summary report:
 - to agree to changes to the paediatric investigation plan in the scope set out in the Annex I of this opinion.

The Norwegian Paediatric Committee member agrees with the above-mentioned recommendation of the Paediatric Committee.

2. The measures and timelines of the paediatric investigation plan and the subset(s) of the paediatric population and condition(s) covered by the waiver are set out in the Annex I.

This opinion is forwarded to the applicant and the Executive Director of the European Medicines Agency, together with its annexes and appendix.

Annex I

The subset(s) of the paediatric population and condition(s) covered by the waiver and the measures and timelines of the agreed paediatric investigation plan (PIP)

1. Waiver

1.1. Condition

Treatment of obesity

The waiver applies to:

- the paediatric population from birth to less than 2 years of age;
- for prolonged-release tablet, for oral use;
- on the grounds that the condition for which the specific medicinal product is intended does not occur in the specified paediatric subsets.

And to:

- the paediatric population from 2 to less than 6 years of age;
- for prolonged-release tablet, for oral use;
- on the grounds that the specific medicinal product does not represent a significant therapeutic benefit over existing treatments.

2. Paediatric Investigation Plan

2.1. Condition

Treatment of obesity

2.1.1. Indication targeted by the PIP

Treatment of obesity

2.1.2. Subsets of the paediatric population concerned by the paediatric development

From 6 to less than 18 years of age

2.1.3. Pharmaceutical form

Prolonged-release tablet

2.1.4. Measures

Area	Number of measures	Description
Quality	1	Measure 1 Development of an age-appropriate oral preparation solid form, for oral use in children from 6 to less than 12 years of age
Non-clinical	1	Measure 2 Mice juvenile toxicity study to assess toxicokinetics, CNS parameters, learning, memory, behaviour, and sexual maturation

		Measure 3 Removed in procedure EMEA-001373-PIP01-12-M01
Clinical	4	Measure 4 Double-blind, randomised, multicentre, placebo-controlled, multiple dose study to assess pharmacokinetics, pharmacodynamics and tolerability of naltrexone and bupropion as fixed-dose combination, in post-pubertal obese adolescents from 12 to less than 18 years of age Measure 5 Double-blind, randomised, multi-centre, placebo-controlled study to assess safety and efficacy of naltrexone and bupropion as fixed-dose combination, in post-pubertal obese adolescents from 12 to less than 18 years of age Measure 6 Double-blind, randomised, multicentre, placebo-controlled, multiple dose study to assess pharmacokinetics, pharmacodynamics and tolerability of naltrexone and bupropion as fixed-dose combination, in pre-pubertal obese children from 6 to less than 12 years of age Measure 7 Double-blind, randomised, multi-centre, placebo-controlled study to assess safety and efficacy of naltrexone and bupropion as fixed-dose combination, in pre-pubertal obese children from 6 to less than 12 years of age

3. Follow-up, completion and deferral of PIP

Concerns on potential long term safety issues in relation to paediatric use:	Yes
Date of completion of the paediatric investigation plan:	By January 2024
Deferral for one or more measures contained in the paediatric investigation plan:	Yes

Annex II

Information about the authorised medicinal product

Condition(s) and authorised indication(s):

1. Treatment of obesity

Authorised indication:

- management of weight in adult patients (≥ 18 years) with an initial Body Mass Index (BMI) of
 - ≥ 30 kg/m² (obese), or
 - ≥ 27 kg/m² to < 30 kg/m² (overweight) in the presence of one or more weight-related co-morbidities (e.g., type 2 diabetes, dyslipidaemia, or controlled hypertension)

Treatment with Mysimba should be discontinued after 16 weeks if patients have not lost at least 5% of their initial body weight.

Authorised pharmaceutical form(s):

Prolonged-release tablet

Blue, biconvex, round tablet of 11.9 mm diameter debossed with "NB-890" on one side

Authorised route(s) of administration:

Oral use