



EUROPEAN MEDICINES AGENCY  
SCIENCE MEDICINES HEALTH

## VANFLYTA

Procedural steps taken and scientific information after the authorisation

Application number	Scope	Opinion/ Notification <sup>1</sup> issued on	Commission Decision Issued <sup>2</sup> / amended on	Product Information affected <sup>3</sup>	Summary
IB/0005	B.II.f.1.b.1 - Stability of FP - Extension of the shelf life of the finished product - As packaged for sale (supported by real time data)	16/12/2024		SmPC and PL	
IB/0003	C.I.z - Changes (Safety/Efficacy) of Human and Veterinary Medicinal Products - Other variation	01/08/2024		SmPC	

<sup>1</sup> Notifications are issued for type I variations and Article 61(3) notifications (unless part of a group including a type II variation or extension application or a worksharing application). Opinions are issued for all other procedures.

<sup>2</sup> A Commission decision (CD) is issued for procedures that affect the terms of the marketing authorisation (e.g. summary of product characteristics, annex II, labelling, package leaflet). The CD is issued within two months of the opinion for variations falling under the scope of Article 23.1a(a) of Regulation (EU) No. 712/2012, or within one year for other procedures.

<sup>3</sup> SmPC (Summary of Product Characteristics), Annex II, Labelling, PL (Package Leaflet).



PSUSA/176/202312	Periodic Safety Update EU Single assessment - quizartinib	11/07/2024	n/a		PRAC Recommendation - maintenance
II/0002	<p>To update section 4.5 and 5.2 of the SmPC in order to add information on interaction with Breast cancer resistant protein (BCRP) substrates based on results from study GE-2161 – Inhibitory Effects of Quizartinib on the Transport Activity of BCRP (REC). In addition, the MAH is taking this opportunity to introduce editorial changes to the PI.</p> <p>C.I.4 - Change(s) in the SPC, Labelling or PL due to new quality, preclinical, clinical or pharmacovigilance data</p>	23/05/2024		SmPC	<p>SmPC new text:</p> <p>In vitro data indicate that quizartinib is an inhibitor of Breast cancer resistance protein (BCRP). The clinical relevance is currently not known. Caution should be used when quizartinib is co-administered with medicinal products that are substrates of BCRP.</p> <p>Quizartinib inhibits BCRP with an estimated in vitro IC50 of 0.813 µM. As no clinical data is available, it cannot be excluded that quizartinib could inhibit this transporter at the recommended doses.</p>