

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

## **1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 75 mg film-coated tablets  
Darunavir Mylan 150 mg film-coated tablets  
Darunavir Mylan 300 mg film-coated tablets  
Darunavir Mylan 600 mg film-coated tablets

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

### Darunavir Mylan 75 mg film-coated tablets

Each film-coated tablet contains 75 mg of darunavir.

### Darunavir Mylan 150 mg film-coated tablets

Each film-coated tablet contains 150 mg of darunavir.

### Darunavir Mylan 300 mg film-coated tablets

Each film-coated tablet contains 300 mg of darunavir.

### Darunavir Mylan 600 mg film-coated tablets

Each film-coated tablet contains 600 mg of darunavir.

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

### Darunavir Mylan 75 mg film-coated tablets

Film-coated tablet.

White to off-white, oval shaped, biconvex film-coated tablets approximately 9.7 mm by 4.6 mm, debossed with 'M' on one side and 'DV1' on the other side.

### Darunavir Mylan 150 mg film-coated tablets

Film-coated tablet.

White to off-white, capsule shaped, biconvex film-coated tablets approximately 12.75 mm by 6.3 mm debossed with 'M' on one side and 'DV2' on the other side.

### Darunavir Mylan 300 mg film-coated tablets

Film-coated tablet.

White to off-white, oval shaped, biconvex film coated tablets approximately 16.5 mm by 8.2 mm, debossed with 'M' on one side of the tablet and 'DV3' on other side.

### Darunavir Mylan 600 mg film-coated tablets

Film-coated tablet.

White to off-white, oval shaped, biconvex film-coated tablets approximately 21.2 mm by 10.6 mm, debossed with 'M' on one side and 'DV5' on the other side.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Darunavir, co-administered with low dose ritonavir is indicated in combination with other antiretroviral medicinal products for the treatment of patients with human immunodeficiency virus (HIV-1) infection (see section 4.2).

Darunavir Mylan 75 mg, 150 mg, 300 mg and 600 mg tablets may be used to provide suitable dose regimens (see section 4.2):

- For the treatment of HIV-1 infection in antiretroviral treatment (ART)-experienced adult patients, including those that have been highly pre-treated.
- For the treatment of HIV-1 infection in paediatric patients from the age of 3 years and at least 15 kg body weight.

In deciding to initiate treatment with darunavir co-administered with low dose ritonavir, careful consideration should be given to the treatment history of the individual patient and the patterns of mutations associated with different agents. Genotypic or phenotypic testing (when available) and treatment history should guide the use of darunavir (see sections 4.2, 4.4 and 5.1).

### 4.2 Posology and method of administration

Therapy should be initiated by a healthcare provider experienced in the management of HIV infection. After therapy with darunavir has been initiated, patients should be advised not to alter the dosage, dose form or discontinue therapy without discussing with their healthcare provider.

#### Posology

Darunavir Mylan must always be given orally with low dose ritonavir as a pharmacokinetic enhancer and in combination with other antiretroviral medicinal products. The Summary of Product Characteristics of ritonavir must, therefore, be consulted prior to initiation of therapy with darunavir.

#### *ART-experienced adult patients*

The recommended dose regimen is 600 mg twice daily taken with ritonavir 100 mg twice daily taken with food. Darunavir Mylan 75 mg, 150 mg, 300 mg and 600 mg tablets can be used to construct the twice daily 600 mg regimen.

The use of 75 mg and 150 mg tablets to achieve the recommended dose is appropriate when there is a difficulty in swallowing the 300 mg or 600 mg tablets. Before prescribing darunavir tablets, young children should be assessed for the ability to swallow intact tablets. For young children unable to swallow tablets, more suitable formulations containing darunavir should be checked for their availability.

#### *ART-naïve adult patients*

For dosage recommendations in ART-naïve patients see the Summary of Product Characteristics for Darunavir Mylan 400 mg and 800 mg tablets.

#### *ART-naïve paediatric patients (3 to 17 years of age and weighing at least 15 kg)*

The weight-based dose of darunavir and ritonavir in paediatric patients is provided in the table below.

<b>Recommended dose for treatment-naïve paediatric patients (3 to 17 years) with darunavir tablets and ritonavir<sup>a</sup></b>	
<b>Body weight (kg)</b>	<b>Dose (once daily with food)</b>
≥ 15 kg to < 30 kg	600 mg darunavir/100 mg ritonavir once daily
≥ 30 kg to < 40 kg	675 mg darunavir/100 mg ritonavir once daily
≥ 40 kg	800 mg darunavir/100 mg ritonavir once daily

<sup>a</sup> ritonavir oral solution: 80 mg/ml

*ART-experienced paediatric patients (3 to 17 years of age and weighing at least 15 kg)*

Darunavir twice daily taken with ritonavir taken with food is usually recommended.

A once daily dose regimen of darunavir taken with ritonavir taken with food may be used in patients with prior exposure to antiretroviral medicinal products but without darunavir resistance associated mutations (DRV-RAMs)\* and who have plasma HIV-1 RNA < 100,000 copies/ml and CD4+ cell count ≥ 100 cells x 10<sup>6</sup>/L.

\* DRV-RAMs: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

The weight-based dose of darunavir and ritonavir in paediatric patients is provided in the table below. The recommended dose of darunavir with low dose ritonavir should not exceed the recommended adult dose (600/100 mg twice daily or 800/100 mg once daily).

<b>Recommended dose for treatment-experienced paediatric patients (3 to 17 years) with darunavir tablets and ritonavir<sup>a</sup></b>		
<b>Body weight (kg)</b>	<b>Dose (once daily with food)</b>	<b>Dose (twice daily with food)</b>
≥ 15 kg–< 30 kg	600 mg darunavir/100 mg ritonavir once daily	375 mg darunavir/50 mg ritonavir twice daily
≥ 30 kg–< 40 kg	675 mg darunavir/100 mg ritonavir once daily	450 mg darunavir/60 mg ritonavir twice daily
≥ 40 kg	800 mg darunavir/100 mg ritonavir once daily	600 mg darunavir/100 mg ritonavir twice daily

<sup>a</sup> ritonavir oral solution: 80 mg/ml

For ART-experienced paediatric patients HIV genotypic testing is recommended. However, when HIV genotypic testing is not feasible, the darunavir/ritonavir once daily dosing regimen is recommended in HIV protease inhibitor-naïve paediatric patients and the twice daily dosing regimen is recommended in HIV protease inhibitor-experienced patients.

#### *Advice on missed doses*

In case a dose of darunavir and/or ritonavir is missed within 6 hours of the time it is usually taken, patients should be instructed to take the prescribed dose of darunavir and ritonavir with food as soon as possible. If this is noticed later than 6 hours after the time it is usually taken, the missed dose should not be taken and the patient should resume the usual dosing schedule.

This guidance is based on the 15 hour half-life of darunavir in the presence of ritonavir and the recommended dosing interval of approximately 12 hours.

If a patient vomits within 4 hours of taking the medicine, another dose of Darunavir Mylan with ritonavir should be taken with food as soon as possible. If a patient vomits more than 4 hours after taking the medicine, the patient does not need to take another dose of Darunavir Mylan with ritonavir until the next regularly scheduled time.

#### Special populations

##### *Elderly*

Limited information is available in this population, and therefore, darunavir should be used with caution in this age group (see sections 4.4 and 5.2).

### *Hepatic impairment*

Darunavir is metabolised by the hepatic system. No dose adjustment is recommended in patients with mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment, however, darunavir should be used with caution in these patients. No pharmacokinetic data are available in patients with severe hepatic impairment. Severe hepatic impairment could result in an increase of darunavir exposure and a worsening of its safety profile. Therefore, darunavir must not be used in patients with severe hepatic impairment (Child-Pugh Class C) (see sections 4.3, 4.4 and 5.2).

### *Renal impairment*

No dose adjustment is required in patients with renal impairment (see sections 4.4 and 5.2).

### *Paediatric population*

Darunavir/ritonavir should not be used in children with a body weight of less than 15 kg as the dose for this population has not been established in a sufficient number of patients (see section 5.1).

Darunavir/ritonavir should not be used in children below 3 years of age because of safety concerns (see sections 4.4 and 5.3).

The weight-based dose regimen for darunavir and ritonavir is provided in the tables above.

### *Pregnancy and postpartum*

No dose adjustment is required for darunavir/ritonavir during pregnancy and postpartum.

Darunavir/ritonavir should be used during pregnancy only if the potential benefit justifies the potential risk (see sections 4.4, 4.6 and 5.2).

### Method of administration

Patients should be instructed to take Darunavir Mylan with low dose ritonavir within 30 minutes after completion of a meal. The type of food does not affect the exposure to darunavir (see sections 4.4, 4.5 and 5.2).

## **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Patients with severe (Child-Pugh Class C) hepatic impairment.

Combination of rifampicin with darunavir with concomitant low dose ritonavir (see section 4.5).

Co-administration with the combination product lopinavir/ritonavir (see section 4.5).

Co-administration with herbal preparations containing St John's wort (*Hypericum perforatum*) (see section 4.5).

Co-administration of darunavir with low dose ritonavir, with active substances that are highly dependent on CYP3A for clearance and for which elevated plasma concentrations are associated with serious and/or life-threatening events. These active substances include e.g.:

- alfuzosin
- amiodarone, bepridil, dronedarone, ivabradine, quinidine, ranolazine
- astemizole, terfenadine
- colchicine when used in patients with renal and/or hepatic impairment (see section 4.5)
- ergot derivatives (e.g. dihydroergotamine, ergometrine, ergotamine, methylergonovine)
- elbasvir/grazoprevir
- cisapride
- dapoxetine
- domperidone
- naloxegol
- lurasidone, pimozide, quetiapine, sertindole (see section 4.5)

- triazolam, midazolam administered orally (for caution on parenterally administered midazolam, see section 4.5)
- sildenafil - when used for the treatment of pulmonary arterial hypertension, avanafil
- simvastatin, lovastatin and lomitapide (see section 4.5)
- ticagrelor (see section 4.5).

#### **4.4 Special warnings and precautions for use**

Regular assessment of virological response is advised. In the setting of lack or loss of virological response, resistance testing should be performed.

Darunavir must always be given orally with low dose ritonavir as a pharmacokinetic enhancer and in combination with other antiretroviral medicinal products (see section 5.2). The Summary of Product Characteristics of ritonavir as appropriate, must therefore be consulted prior to initiation of therapy with darunavir.

Increasing the dose of ritonavir from that recommended in section 4.2 did not significantly affect darunavir concentrations. It is not recommended to alter the dose of ritonavir.

Darunavir binds predominantly to  $\alpha_1$ -acid glycoprotein. This protein binding is concentration-dependent indicative for saturation of binding. Therefore, protein displacement of medicinal products highly bound to  $\alpha_1$ -acid glycoprotein cannot be ruled out (see section 4.5).

##### ART-experienced patients - once daily dosing

Darunavir used in combination with cobicistat or low dose ritonavir once daily in ART-experienced patients should not be used in patients with one or more darunavir resistance associated mutations (DRV-RAMs) or HIV-1 RNA  $\geq 100,000$  copies/ml or CD4+ cell count  $< 100$  cells  $\times 10^6/L$  (see section 4.2). Combinations with optimised background regimen (OBRs) other than  $\geq 2$  NRTIs have not been studied in this population. Limited data are available in patients with HIV-1 clades other than B (see section 5.1).

##### Paediatric population

Darunavir is not recommended for use in paediatric patients below 3 years of age or less than 15 kg body weight (see sections 4.2 and 5.3).

##### Pregnancy

Darunavir/ritonavir should be used during pregnancy only if the potential benefit justifies the potential risk. Caution should be used in pregnant women with concomitant medications which may further decrease darunavir exposure (see sections 4.5 and 5.2).

##### Elderly

As limited information is available on the use of darunavir in patients aged 65 and over, caution should be exercised in the administration of darunavir in elderly patients, reflecting the greater frequency of decreased hepatic function and of concomitant disease or other therapy (see sections 4.2 and 5.2).

##### Severe skin reactions

During the darunavir/ritonavir clinical development program (N=3,063), severe skin reactions, which may be accompanied with fever and/or elevations of transaminases, have been reported in 0.4% of patients. DRESS (Drug Rash with Eosinophilia and Systemic Symptoms) and Stevens-Johnson syndrome has been rarely ( $< 0.1\%$ ) reported, and during post-marketing experience toxic epidermal necrolysis and acute generalised exanthematous pustulosis have been reported. Darunavir should be

discontinued immediately if signs or symptoms of severe skin reactions develop. These can include, but are not limited to, severe rash or rash accompanied by fever, general malaise, fatigue, muscle or joint aches, blisters, oral lesions, conjunctivitis, hepatitis and/or eosinophilia.

Rash occurred more commonly in treatment-experienced patients receiving regimens containing darunavir/ritonavir + raltegravir compared to patients receiving darunavir/ritonavir without raltegravir or raltegravir without darunavir (see section 4.8).

Darunavir contains a sulphonamide moiety. Darunavir Mylan should be used with caution in patients with a known sulphonamide allergy.

### Hepatotoxicity

Drug-induced hepatitis (e.g. acute hepatitis, cytolytic hepatitis) has been reported with darunavir. During the darunavir/ritonavir clinical development program (N=3,063), hepatitis was reported in 0.5% of patients receiving combination antiretroviral therapy with darunavir/ritonavir. Patients with pre-existing liver dysfunction, including chronic active hepatitis B or C, have an increased risk for liver function abnormalities including severe and potentially fatal hepatic adverse reactions. In case of concomitant antiviral therapy for hepatitis B or C, please refer to the relevant product information for these medicinal products.

Appropriate laboratory testing should be conducted prior to initiating therapy with darunavir/ritonavir and patients should be monitored during treatment. Increased AST/ALT monitoring should be considered in patients with underlying chronic hepatitis, cirrhosis, or in patients who have pre-treatment elevations of transaminases, especially during the first several months of darunavir/ritonavir treatment.

If there is evidence of new or worsening liver dysfunction (including clinically significant elevation of liver enzymes and/or symptoms such as fatigue, anorexia, nausea, jaundice, dark urine, liver tenderness, hepatomegaly) in patients using darunavir/ritonavir, interruption or discontinuation of treatment should be considered promptly.

### Patients with coexisting conditions

#### *Hepatic impairment*

The safety and efficacy of darunavir have not been established in patients with severe underlying liver disorders and darunavir is therefore contraindicated in patients with severe hepatic impairment. Due to an increase in the unbound darunavir plasma concentrations, darunavir should be used with caution in patients with mild or moderate hepatic impairment (see sections 4.2, 4.3 and 5.2).

#### *Renal impairment*

No special precautions or dose adjustments for darunavir/ritonavir are required in patients with renal impairment. As darunavir and ritonavir are highly bound to plasma proteins, it is unlikely that they will be significantly removed by haemodialysis or peritoneal dialysis. Therefore, no special precautions or dose adjustments are required in these patients (see sections 4.2 and 5.2).

#### *Haemophiliac patients*

There have been reports of increased bleeding, including spontaneous skin haematomas and haemarthrosis in patients with haemophilia type A and B treated with PIs. In some patients additional factor VIII was given. In more than half of the reported cases, treatment with PIs was continued or reintroduced if treatment had been discontinued. A causal relationship has been suggested, although the mechanism of action has not been elucidated. Haemophiliac patients should, therefore, be made aware of the possibility of increased bleeding.

#### *Weight and metabolic parameters*

An increase in weight and in levels of blood lipids and glucose may occur during antiretroviral therapy. Such changes may in part be linked to disease control and life style. For lipids, there is in

some cases evidence for a treatment effect, while for weight gain there is no strong evidence relating this to any particular treatment. For monitoring of blood lipids and glucose reference is made to established HIV treatment guidelines. Lipid disorders should be managed as clinically appropriate.

### Osteonecrosis

Although the aetiology is considered to be multifactorial (including corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index), cases of osteonecrosis have been reported particularly in patients with advanced HIV disease and/or long-term exposure to combination antiretroviral therapy (CART). Patients should be advised to seek medical advice if they experience joint aches and pain, joint stiffness or difficulty in movement.

### Immune reconstitution inflammatory syndrome

In HIV infected patients with severe immune deficiency at the time of initiation of combination antiretroviral therapy (CART), an inflammatory reaction to asymptomatic or residual opportunistic pathogens may arise and cause serious clinical conditions, or aggravation of symptoms. Typically, such reactions have been observed within the first weeks or months of initiation of CART. Relevant examples are cytomegalovirus retinitis, generalised and/or focal mycobacterial infections and pneumonia caused by *Pneumocystis jirovecii* (formerly known as *Pneumocystis carinii*). Any inflammatory symptoms should be evaluated and treatment instituted when necessary. In addition, reactivation of herpes simplex and herpes zoster has been observed in clinical studies with darunavir co-administered with low dose ritonavir.

Autoimmune disorders (such as Graves' disease and autoimmune hepatitis) have also been reported to occur in the setting of immune reactivation; however, the reported time to onset is more variable and these events can occur many months after initiation of treatment (see section 4.8).

### Interactions with medicinal products

Several of the interaction studies have been performed with darunavir at lower than recommended doses. The effects on co-administered medicinal products may thus be underestimated and clinical monitoring of safety may be indicated. For full information on interactions with other medicinal products see section 4.5.

Efavirenz in combination with boosted darunavir once daily may result in sub-optimal darunavir  $C_{min}$ . If efavirenz is to be used in combination with darunavir, the darunavir/ritonavir 600/100 mg twice daily regimen should be used (see section 4.5).

Life-threatening and fatal drug interactions have been reported in patients treated with colchicine and strong inhibitors of CYP3A and P-glycoprotein (P-gp; see sections 4.3 and 4.5).

### Darunavir Mylan contains sodium

Darunavir Mylan 75 mg, 150 mg, 300 mg and 600 mg film-coated tablets contain less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

## **4.5 Interaction with other medicinal products and other forms of interaction**

Interaction studies have only been performed in adults.

### Medicinal products that may be affected by darunavir boosted with ritonavir

Darunavir and ritonavir are inhibitors of CYP3A, CYP2D6 and P-gp. Co-administration of darunavir/ritonavir with medicinal products primarily metabolised by CYP3A and/or CYP2D6 or transported by P-gp may result in increased systemic exposure to such medicinal products, which could increase or prolong their therapeutic effect and adverse reactions.



Co-administration of darunavir/ritonavir with drugs that have active metabolite(s) formed by CYP3A may result in reduced plasma concentrations of these active metabolite(s), potentially leading to loss of their therapeutic effect (see the Interaction table below).

Darunavir co-administered with low dose ritonavir must not be combined with medicinal products that are highly dependent on CYP3A for clearance and for which increased systemic exposure is associated with serious and/or life-threatening events (narrow therapeutic index) (see section 4.3).

The overall pharmacokinetic enhancement effect by ritonavir was an approximate 14-fold increase in the systemic exposure of darunavir when a single dose of 600 mg darunavir was given orally in combination with ritonavir at 100 mg twice daily. Therefore, darunavir must only be used in combination with low dose ritonavir as a pharmacokinetic enhancer (see sections 4.4 and 5.2).

A clinical study utilising a cocktail of medicinal products that are metabolised by cytochromes CYP2C9, CYP2C19 and CYP2D6 demonstrated an increase in CYP2C9 and CYP2C19 activity and inhibition of CYP2D6 activity in the presence of darunavir/ritonavir, which may be attributed to the presence of low dose ritonavir. Co-administration of darunavir and ritonavir with medicinal products which are primarily metabolised by CYP2D6 (such as flecainide, propafenone, metoprolol) may result in increased plasma concentrations of these medicinal products, which could increase or prolong their therapeutic effect and adverse reactions. Co-administration of darunavir and ritonavir with medicinal products primarily metabolised by CYP2C9 (such as warfarin) and CYP2C19 (such as methadone) may result in decreased systemic exposure to such medicinal products, which could decrease or shorten their therapeutic effect.

Although the effect on CYP2C8 has only been studied *in vitro*, co-administration of darunavir and ritonavir and medicinal products primarily metabolised by CYP2C8 (such as paclitaxel, rosiglitazone, repaglinide) may result in decreased systemic exposure to such medicinal products, which could decrease or shorten their therapeutic effect.

Ritonavir inhibits the transporters P-glycoprotein, OATP1B1 and OATP1B3, and co-administration with substrates of these transporters can result in increased plasma concentrations of these compounds (e.g. dabigatran etexilate, digoxin, statins and bosentan; see the Interaction table below).

#### *Medicinal products that affect darunavir/ritonavir exposure*

Darunavir and ritonavir are metabolised by CYP3A. Medicinal products that induce CYP3A activity would be expected to increase the clearance of darunavir and ritonavir, resulting in lowered plasma concentrations of darunavir and ritonavir (e.g. rifampicin, St John's wort, lopinavir).

Co-administration of darunavir and ritonavir and other medicinal products that inhibit CYP3A may decrease the clearance of darunavir and ritonavir and may result in increased plasma concentrations of darunavir and ritonavir (e.g. indinavir, azole antifungals like clotrimazole). These interactions are described in the interaction table below.

#### Interaction table

Interactions between darunavir/ritonavir and antiretroviral and non-antiretroviral medicinal products are listed in the table below. The direction of the arrow for each pharmacokinetic parameter is based on the 90% confidence interval of the geometric mean ratio being within ( $\leftrightarrow$ ), below ( $\downarrow$ ) or above ( $\uparrow$ ) the 80-125% range (not determined as "ND").

Several of the interaction studies (indicated by # in the table below) have been performed at lower than recommended doses of darunavir or with a different dosing regimen (see section 4.2 Posology). The effects on co-administered medicinal products may thus be underestimated and clinical monitoring of safety may be indicated.

The below list of examples of drug-drug interactions is not comprehensive and therefore the label of each drug that is co-administered with darunavir should be consulted for information related to the

route of metabolism, interaction pathways, potential risks, and specific actions to be taken with regards to co-administration.

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
<b>HIV ANTIRETROVIRALS</b>		
<b><i>Integrase strand transfer inhibitors</i></b>		
Dolutegravir	dolutegravir AUC ↓ 22% dolutegravir C <sub>24h</sub> ↓ 38% dolutegravir C <sub>max</sub> ↓ 11% darunavir ↔* * Using cross-study comparisons to historical pharmacokinetic data	Darunavir co-administered with low dose ritonavir and dolutegravir can be used without dose adjustment.
Raltegravir	Some clinical studies suggest raltegravir may cause a modest decrease in darunavir plasma concentrations.	At present the effect of raltegravir on darunavir plasma concentrations does not appear to be clinically relevant. Darunavir co-administered with low dose ritonavir and raltegravir can be used without dose adjustments.
<b><i>Nucleo(s)ide reverse transcriptase inhibitors (NRTIs)</i></b>		
Didanosine 400 mg once daily	didanosine AUC ↓ 9% didanosine C <sub>min</sub> ND didanosine C <sub>max</sub> ↓ 16% darunavir AUC ↔ darunavir C <sub>min</sub> ↔ darunavir C <sub>max</sub> ↔	Darunavir co-administered with low dose ritonavir and didanosine can be used without dose adjustments. Didanosine is to be administered on an empty stomach, thus it should be administered 1 hour before or 2 hours after darunavir/ritonavir given with food.
Tenofovir disoproxil 245 mg once daily <sup>‡</sup>	tenofovir AUC ↑ 22% tenofovir C <sub>min</sub> ↑ 37% tenofovir C <sub>max</sub> ↑ 24% #darunavir AUC ↑ 21% #darunavir C <sub>min</sub> ↑ 24% #darunavir C <sub>max</sub> ↑ 16% (↑ tenofovir from effect on MDR-1 transport in the renal tubules)	Monitoring of renal function may be indicated when darunavir co-administered with low dose ritonavir is given in combination with tenofovir disoproxil, particularly in patients with underlying systemic or renal disease, or in patients taking nephrotoxic agents.
Emtricitabine/tenofovir alafenamide	Tenofovir alafenamide ↔ Tenofovir ↑	The recommended dose of emtricitabine/tenofovir alafenamide is 200/10 mg once daily when used with darunavir with low dose ritonavir.
Abacavir Emtricitabine Lamivudine Stavudine Zidovudine	Not studied. Based on the different elimination pathways of the other NRTIs zidovudine, emtricitabine, stavudine, lamivudine, that are primarily renally excreted, and abacavir for which metabolism is not mediated by CYP450, no interactions are expected for these medicinal compounds and darunavir co-administered with low dose ritonavir.	Darunavir co-administered with low dose ritonavir can be used with these NRTIs without dose adjustment.

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
<b>Non-nucleo(s)ide reverse transcriptase inhibitors (NNRTIs)</b>		
Efavirenz 600 mg once daily	efavirenz AUC ↑ 21% efavirenz C <sub>min</sub> ↑ 17% efavirenz C <sub>max</sub> ↑ 15% #darunavir AUC ↓ 13% #darunavir C <sub>min</sub> ↓ 31% #darunavir C <sub>max</sub> ↓ 15% (↑ efavirenz from CYP3A inhibition) (↓ darunavir from CYP3A induction)	Clinical monitoring for central nervous system toxicity associated with increased exposure to efavirenz may be indicated when darunavir co-administered with low dose ritonavir is given in combination with efavirenz.  Efavirenz in combination with darunavir/ritonavir 800/100 mg once daily may result in sub-optimal darunavir C <sub>min</sub> . If efavirenz is to be used in combination with darunavir/ritonavir, the darunavir/ritonavir 600/100 mg twice daily regimen should be used (see section 4.4).
Etravirine 100 mg twice daily	etravirine AUC ↓ 37% etravirine C <sub>min</sub> ↓ 49% etravirine C <sub>max</sub> ↓ 32% darunavir AUC ↑ 15% darunavir C <sub>min</sub> ↔ darunavir C <sub>max</sub> ↔	Darunavir co-administered with low dose ritonavir and etravirine 200 mg twice daily can be used without dose adjustments.
Nevirapine 200 mg twice daily	nevirapine AUC ↑ 27% nevirapine C <sub>min</sub> ↑ 47% nevirapine C <sub>max</sub> ↑ 18% #darunavir: concentrations were consistent with historical data (↑ nevirapine from CYP3A inhibition)	Darunavir co-administered with low dose ritonavir and nevirapine can be used without dose adjustments.
Rilpivirine 150 mg once daily	rilpivirine AUC ↑ 130% rilpivirine C <sub>min</sub> ↑ 178% rilpivirine C <sub>max</sub> ↑ 79% darunavir AUC ↔ darunavir C <sub>min</sub> ↓ 11% darunavir C <sub>max</sub> ↔	Darunavir co-administered with low dose ritonavir and rilpivirine can be used without dose adjustments.
<b>HIV Protease inhibitors (PIs) - without additional co-administration of low dose ritonavir<sup>†</sup></b>		
Atazanavir 300 mg once daily	atazanavir AUC ↔ atazanavir C <sub>min</sub> ↑ 52% atazanavir C <sub>max</sub> ↓ 11% #darunavir AUC ↔ #darunavir C <sub>min</sub> ↔ #darunavir C <sub>max</sub> ↔  Atazanavir: comparison of atazanavir/ritonavir 300/100 mg once daily vs. atazanavir 300 mg once daily in combination with darunavir/ritonavir 400/100 mg twice daily. Darunavir: comparison of darunavir/ritonavir 400/100 mg twice daily vs. darunavir/ritonavir 400/100 mg twice daily in combination with atazanavir 300 mg once daily.	Darunavir co-administered with low dose ritonavir and atazanavir can be used without dose adjustments.

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
Indinavir 800 mg twice daily	indinavir AUC ↑ 23% indinavir C <sub>min</sub> ↑ 125% indinavir C <sub>max</sub> ↔ #darunavir AUC ↑ 24% #darunavir C <sub>min</sub> ↑ 44% #darunavir C <sub>max</sub> ↑ 11%  Indinavir: comparison of indinavir/ritonavir 800/100 mg twice daily vs. indinavir/darunavir/ritonavir 800/400/100 mg twice daily. Darunavir: comparison of darunavir/ritonavir 400/100 mg twice daily vs. darunavir/ritonavir 400/100 mg in combination with indinavir 800 mg twice daily.	When used in combination with darunavir co-administered with low dose ritonavir, dose adjustment of indinavir from 800 mg twice daily to 600 mg twice daily may be warranted in case of intolerance.
Saquinavir 1,000 mg twice daily	#darunavir AUC ↓ 26% #darunavir C <sub>min</sub> ↓ 42% #darunavir C <sub>max</sub> ↓ 17% saquinavir AUC ↓ 6% saquinavir C <sub>min</sub> ↓ 18% saquinavir C <sub>max</sub> ↓ 6%  Saquinavir: comparison of saquinavir/ritonavir 1,000/100 mg twice daily vs. saquinavir/darunavir/ritonavir 1,000/400/100 mg twice daily. Darunavir: comparison of darunavir/ritonavir 400/100 mg twice daily vs. darunavir/ritonavir 400/100 mg in combination with saquinavir 1,000 mg twice daily.	It is not recommended to combine darunavir co-administered with low dose ritonavir with saquinavir.
<b>HIV Protease inhibitors (PIs) - with co-administration of low dose ritonavir<sup>†</sup></b>		
Lopinavir/ritonavir 400/100 mg twice daily  Lopinavir/ritonavir 533/133.3 mg twice daily	lopinavir AUC ↑ 9% lopinavir C <sub>min</sub> ↑ 23% lopinavir C <sub>max</sub> ↓ 2% darunavir AUC ↓ 38% <sup>‡</sup> darunavir C <sub>min</sub> ↓ 51% <sup>‡</sup> darunavir C <sub>max</sub> ↓ 21% <sup>‡</sup> lopinavir AUC ↔ lopinavir C <sub>min</sub> ↑ 13% lopinavir C <sub>max</sub> ↑ 11% darunavir AUC ↓ 41% darunavir C <sub>min</sub> ↓ 55% darunavir C <sub>max</sub> ↓ 21% <sup>‡</sup> based upon non dose normalised values	Due to a decrease in the exposure (AUC) of darunavir by 40%, appropriate doses of the combination have not been established. Hence, concomitant use of darunavir co-administered with low dose ritonavir and the combination product lopinavir/ritonavir is contraindicated (see section 4.3).
<b>CCR5 ANTAGONIST</b>		
Maraviroc 150 mg twice daily	maraviroc AUC ↑ 305% maraviroc C <sub>min</sub> ND maraviroc C <sub>max</sub> ↑ 129% darunavir, ritonavir concentrations were consistent with historical data	The maraviroc dose should be 150 mg twice daily when co-administered with darunavir with low dose ritonavir.
<b>α1-ADRENORECEPTOR ANTAGONIST</b>		
Alfuzosin	Based on theoretical considerations darunavir is expected to increase alfuzosin plasma concentrations. (CYP3A inhibition)	Co-administration of darunavir with low dose ritonavir and alfuzosin is contraindicated (see section 4.3).

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
<b>ANAESTHETIC</b>		
Alfentanil	Not studied. The metabolism of alfentanil is mediated via CYP3A, and may as such be inhibited by darunavir co-administered with low dose ritonavir.	The concomitant use with darunavir and low dose ritonavir may require to lower the dose of alfentanil and requires monitoring for risks of prolonged or delayed respiratory depression.
<b>ANTIANGINA/ANTIARRHYTHMIC</b>		
Disopyramide Flecainide Lidocaine (systemic) Mexiletine Propafenone  Amiodarone Bepiridil Dronedarone Ivabradine, Quinidine Ranolazine	Not studied. Darunavir is expected to increase these antiarrhythmic plasma concentrations. (CYP3A and/or CYP2D6 inhibition)	Caution is warranted and therapeutic concentration monitoring, if available, is recommended for these antiarrhythmics when co-administered with darunavir with low dose ritonavir.  Darunavir co-administered with low dose ritonavir and amiodarone, bepridil, dronedarone, ivabradine, quinidine, or ranolazine is contraindicated (see section 4.3).
Digoxin 0.4 mg single dose	digoxin AUC ↑ 61% digoxin C <sub>min</sub> ND digoxin C <sub>max</sub> ↑ 29% (↑ digoxin from probable inhibition of P-gp)	Given that digoxin has a narrow therapeutic index, it is recommended that the lowest possible dose of digoxin should initially be prescribed in case digoxin is given to patients on darunavir/ritonavir therapy. The digoxin dose should be carefully titrated to obtain the desired clinical effect while assessing the overall clinical state of the subject.
<b>ANTIBIOTIC</b>		
Clarithromycin 500 mg twice daily	clarithromycin AUC ↑ 57% clarithromycin C <sub>min</sub> ↑ 174% clarithromycin C <sub>max</sub> ↑ 26% #darunavir AUC ↓ 13% #darunavir C <sub>min</sub> ↑ 1% #darunavir C <sub>max</sub> ↓ 17% 14-OH-clarithromycin concentrations were not detectable when combined with darunavir/ritonavir. (↑ clarithromycin from CYP3A inhibition and possible P-gp inhibition)	Caution should be exercised when clarithromycin is combined with darunavir co-administered with low dose ritonavir.  For patients with renal impairment the Summary of Product Characteristics for clarithromycin should be consulted for the recommended dose.
<b>ANTICOAGULANT/PLATELET AGGREGATION INHIBITOR</b>		
Apixaban Rivaroxaban	Not studied. Co-administration of boosted darunavir with these anticoagulants may increase concentrations of the anticoagulant. (CYP3A and/or P-gp inhibition).	The use of boosted darunavir with a direct anticoagulant (DOAC) that is metabolized by CYP3A4 and transported by P-gp is not recommended as this may lead to an increased bleeding risk.

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
Dabigatran etexilate Edoxaban	dabigatran etexilate (150 mg): darunavir/ritonavir 800/100 mg single dose: dabigatran AUC ↑ 72% dabigatran C <sub>max</sub> ↑ 64%	Darunavir/ritonavir: Clinical monitoring and/or dose reduction of the DOAC should be considered when a DOAC transported by P-gp but not metabolised by CYP3A4, including dabigatran etexilate and edoxaban, is co-administered with Darunavir/rtv.
Ticagrelor	darunavir/ritonavir 800/100 mg once daily: dabigatran AUC ↑ 18% dabigatran C <sub>max</sub> ↑ 22% Based on theoretical considerations, co-administration of boosted darunavir with ticagrelor may increase concentrations of ticagrelor (CYP3A and/or P-glycoprotein inhibition).	Concomitant administration of boosted darunavir with ticagrelor is contraindicated (see section 4.3).
Clopidogrel	Not studied. Co-administration of clopidogrel with boosted darunavir is expected to decrease clopidogrel active metabolite plasma concentration, which may reduce the antiplatelet activity of clopidogrel.	Co-administration of clopidogrel with boosted darunavir is not recommended.  Use of other antiplatelets not affected by CYP inhibition or induction (e.g. prasugrel) is recommended.
Warfarin	Not studied. Warfarin concentrations may be affected when co-administered with darunavir with low dose ritonavir.	It is recommended that the international normalised ratio (INR) be monitored when warfarin is combined with darunavir co-administered with low dose ritonavir.
<b>ANTICONVULSANTS</b>		
Phenobarbital Phenytoin	Not studied. Phenobarbital and phenytoin are expected to decrease plasma concentrations of darunavir and its pharmacoenhancer. (induction of CYP450 enzymes)	Darunavir co-administered with low dose ritonavir should not be used in combination with these medicines.
Carbamazepine 200 mg twice daily	carbamazepine AUC ↑ 45% carbamazepine C <sub>min</sub> ↑ 54% carbamazepine C <sub>max</sub> ↑ 43% darunavir AUC ↔ darunavir C <sub>min</sub> ↓ 15% darunavir C <sub>max</sub> ↔	No dose adjustment for darunavir/ritonavir is recommended. If there is a need to combine darunavir/ritonavir and carbamazepine, patients should be monitored for potential carbamazepine-related adverse events. Carbamazepine concentrations should be monitored and its dose should be titrated for adequate response. Based upon the findings, the carbamazepine dose may need to be reduced by 25% to 50% in the presence of darunavir/ritonavir.
Clonazepam	Not studied. Co-administration of boosted darunavir with clonazepam may increase concentrations of clonazepam. (CYP3A inhibition)	Clinical monitoring is recommended when co-administering boosted darunavir with clonazepam.

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
<b>ANTIDEPRESSANTS</b>		
<p>Paroxetine 20 mg once daily</p> <p>Sertraline 50 mg once daily</p> <p>Amitriptyline Desipramine Imipramine Nortriptyline Trazodone</p>	<p>paroxetine AUC ↓ 39% paroxetine C<sub>min</sub> ↓ 37% paroxetine C<sub>max</sub> ↓ 36% #darunavir AUC ↔ #darunavir C<sub>min</sub> ↔ #darunavir C<sub>max</sub> ↔ sertraline AUC ↓ 49% sertraline C<sub>min</sub> ↓ 49% sertraline C<sub>max</sub> ↓ 44% #darunavir AUC ↔ #darunavir C<sub>min</sub> ↓ 6% #darunavir C<sub>max</sub> ↔</p> <p>Concomitant use of darunavir co-administered with low dose ritonavir and these antidepressants may increase concentrations of the antidepressant. (CYP2D6 and/or CYP3A inhibition)</p>	<p>If antidepressants are co-administered with darunavir with low dose ritonavir, the recommended approach is a dose titration of the antidepressant based on a clinical assessment of antidepressant response. In addition, patients on a stable dose of these antidepressants who start treatment with darunavir with low dose ritonavir should be monitored for antidepressant response.</p> <p>Clinical monitoring is recommended when co-administering darunavir with low dose ritonavir with these antidepressants and a dose adjustment of the antidepressant may be needed.</p>
<b>ANTIEMETICS</b>		
Domperidone	Not studied.	Co-administration of domperidone with boosted darunavir is contraindicated.
<b>ANTIFUNGALS</b>		
Voriconazole	Not studied. Ritonavir may decrease plasma concentrations of voriconazole. (induction of CYP450 enzymes)	Voriconazole should not be combined with darunavir co-administered with low dose ritonavir unless an assessment of the benefit/risk ratio justifies the use of voriconazole.
<p>Fluconazole Isavuconazole Itraconazole Posaconazole</p> <p>Clotrimazole</p>	<p>Not studied. Darunavir may increase antifungal plasma concentrations and posaconazole, isavuconazole, itraconazole, or fluconazole may increase darunavir concentrations. (CYP3A and/or P-gp inhibition)</p> <p>Not studied. Concomitant systemic use of clotrimazole and darunavir co-administered with low dose ritonavir may increase plasma concentrations of darunavir and/or clotrimazole. Darunavir AUC<sub>24h</sub> ↑ 33% (based on population pharmacokinetic model)</p>	Caution is warranted and clinical monitoring is recommended. When co-administration is required the daily dose of itraconazole should not exceed 200 mg.

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
<b>ANTIGOUT MEDICINES</b>		
Colchicine	Not studied. Concomitant use of colchicine and darunavir co-administered with low dose ritonavir may increase the exposure to colchicine. (CYP3A and/ or P-gp inhibition)	A reduction in colchicine dosage or an interruption of colchicine treatment is recommended in patients with normal renal or hepatic function if treatment with darunavir co-administered with low dose ritonavir is required. For patients with renal or hepatic impairment colchicine with darunavir co-administered with low dose ritonavir is contraindicated (see sections 4.3 and 4.4).
<b>ANTIMALARIALS</b>		
Artemether/Lumefantrine 80/480 mg, 6 doses at 0, 8, 24, 36, 48, and 60 hours	artemether AUC ↓ 16% artemether C <sub>min</sub> ↔ artemether C <sub>max</sub> ↓ 18% dihydroartemisinin AUC ↓ 18% dihydroartemisinin C <sub>min</sub> ↔ dihydroartemisinin C <sub>max</sub> ↓ 18% lumefantrine AUC ↑ 175% lumefantrine C <sub>min</sub> ↑ 126% lumefantrine C <sub>max</sub> ↑ 65% darunavir AUC ↔ darunavir C <sub>min</sub> ↓ 13% darunavir C <sub>max</sub> ↔	The combination of darunavir and artemether/lumefantrine can be used without dose adjustments; however, due to the increase in lumefantrine exposure, the combination should be used with caution.
<b>ANTIMYCOBACTERIALS</b>		
Rifampicin Rifapentine	Not studied. Rifapentine and rifampicin are strong CYP3A inducers and have been shown to cause profound decreases in concentrations of other protease inhibitors, which can result in virological failure and resistance development (CYP450 enzyme induction). During attempts to overcome the decreased exposure by increasing the dose of other protease inhibitors with low dose ritonavir, a high frequency of liver reactions was seen with rifampicin.	The combination of rifapentine and darunavir with concomitant low dose ritonavir is not recommended.  The combination of rifampicin and darunavir with concomitant low dose ritonavir is contraindicated (see section 4.3).



INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
Rifabutin 150 mg once every other day	<p>rifabutin AUC** ↑ 55%</p> <p>rifabutin C<sub>min</sub>** ↑ ND</p> <p>rifabutin C<sub>max</sub>** ↔</p> <p>darunavir AUC ↑ 53%</p> <p>darunavir C<sub>min</sub> ↑ 68%</p> <p>darunavir C<sub>max</sub> ↑ 39%</p> <p>** sum of active moieties of rifabutin (parent drug + 25-<i>O</i>-desacetyl metabolite)</p> <p>The interaction trial showed a comparable daily systemic exposure for rifabutin between treatment at 300 mg once daily alone and 150 mg once every other day in combination with darunavir/ritonavir (600/100 mg twice daily) with an about 10-fold increase in the daily exposure to the active metabolite 25-<i>O</i>-desacetyl-rifabutin. Furthermore, AUC of the sum of active moieties of rifabutin (parent drug + 25-<i>O</i>-desacetyl metabolite) was increased 1.6-fold, while C<sub>max</sub> remained comparable.</p> <p>Data on comparison with a 150 mg once daily reference dose is lacking.</p> <p>(Rifabutin is an inducer and substrate of CYP3A.) An increase of systemic exposure to darunavir was observed when darunavir co-administered with 100 mg ritonavir was co-administered with rifabutin (150 mg once every other day).</p>	<p>A dosage reduction of rifabutin by 75% of the usual dose of 300 mg/day (i.e. rifabutin 150 mg once every other day) and increased monitoring for rifabutin related adverse events is warranted in patients receiving the combination with darunavir co-administered with ritonavir. In case of safety issues, a further increase of the dosing interval for rifabutin and/or monitoring of rifabutin levels should be considered.</p> <p>Consideration should be given to official guidance on the appropriate treatment of tuberculosis in HIV infected patients.</p> <p>Based upon the safety profile of darunavir/ritonavir, the increase in darunavir exposure in the presence of rifabutin does not warrant a dose adjustment for darunavir/ritonavir.</p> <p>Based on pharmacokinetic modeling, this dosage reduction of 75% is also applicable if patients receive rifabutin at doses other than 300 mg/day.</p>
<b>ANTINEOPLASTICS</b>		
Dasatinib Nilotinib Vinblastine Vincristine	Not studied. Darunavir is expected to increase these antineoplastic plasma concentrations. (CYP3A inhibition)	Concentrations of these medicinal products may be increased when co-administered with darunavir with low dose ritonavir resulting in the potential for increased adverse events usually associated with these agents. Caution should be exercised when combining one of these antineoplastic agents with darunavir with low dose ritonavir.
Everolimus Irinotecan		Concomitant use of everolimus or Irinotecan and darunavir co-administered with low dose ritonavir is not recommended.
<b>ANTIPSYCHOTICS/NEUROLEPTICS</b>		
Quetiapine	Not studied. Darunavir is expected to increase these antipsychotic plasma concentrations. (CYP3A inhibition)	Concomitant administration of darunavir with low dose ritonavir and quetiapine is contraindicated as it may increase quetiapine-related toxicity. Increased concentrations of quetiapine may lead to coma (see section 4.3).

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
Perphenazine Risperidone Thioridazine	Not studied. Darunavir is expected to increase these antipsychotic plasma concentrations. (CYP3A, CYP2D6 and/or P-gp inhibition)	A dose decrease may be needed for these drugs when co-administered with darunavir co-administered with low dose ritonavir.
Lurasidone Pimozide Sertindole		Concomitant administration of darunavir with low dose ritonavir and lurasidone, pimozide or sertindole is contraindicated (see section 4.3).
<b>β-BLOCKERS</b>		
Carvedilol Metoprolol Timolol	Not studied. Darunavir is expected to increase these β-blocker plasma concentrations. (CYP2D6 inhibition)	Clinical monitoring is recommended when co-administering darunavir with β-blockers. A lower dose of the β-blocker should be considered.
<b>CALCIUM CHANNEL BLOCKERS</b>		
Amlodipine Diltiazem Felodipine Nicardipine Nifedipine Verapamil	Not studied. Darunavir co-administered with low dose ritonavir can be expected to increase the plasma concentrations of calcium channel blockers. (CYP3A and/or CYP2D6 inhibition)	Clinical monitoring of therapeutic and adverse effects is recommended when these medicines are concomitantly administered with darunavir with low dose ritonavir.
<b>CORTICOSTEROIDS</b>		
Corticosteroids primarily metabolised by CYP3A (including betamethasone, budesonide, fluticasone, mometasone, prednisone, triamcinolone)	<p>Fluticasone: in a clinical study where ritonavir 100 mg capsules twice daily were co-administered with 50 µg intranasal fluticasone propionate (4 times daily) for 7 days in healthy subjects, fluticasone propionate plasma concentrations increased significantly, whereas the intrinsic cortisol levels decreased by approximately 86% (90% CI 82-89%). Greater effects may be expected when fluticasone is inhaled. Systemic corticosteroid effects including Cushing's syndrome and adrenal suppression have been reported in patients receiving ritonavir and inhaled or intranasally administered fluticasone. The effects of high fluticasone systemic exposure on ritonavir plasma levels are unknown.</p> <p>Other corticosteroids: interaction not studied. Plasma concentrations of these medicinal products may be increased when co-administered with darunavir with low dose ritonavir, resulting in reduced serum cortisol concentrations.</p>	<p>Concomitant use of darunavir with low dose ritonavir and corticosteroids (all routes of administration) that are metabolised by CYP3A may increase the risk of development of systemic corticosteroid effects, including Cushing's syndrome and adrenal suppression.</p> <p>Co-administration with CYP3A-metabolised corticosteroids is not recommended unless the potential benefit to the patient outweighs the risk, in which case patients should be monitored for systemic corticosteroid effects.</p> <p>Alternative corticosteroids which are less dependent on CYP3A metabolism e.g. beclomethasone should be considered, particularly for long term use.</p>
Dexamethasone (systemic)	Not studied. Dexamethasone may decrease plasma concentrations of darunavir. (CYP3A induction)	Systemic dexamethasone should be used with caution when combined with darunavir co-administered with low dose ritonavir.

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
<b>ENDOTHELIN RECEPTOR ANTAGONISTS</b>		
Bosentan	Not studied. Concomitant use of bosentan and darunavir co-administered with low dose ritonavir may increase plasma concentrations of bosentan. Bosentan is expected to decrease plasma concentrations of darunavir and/or its pharmacoenhancer. (CYP3A induction)	When administered concomitantly with darunavir and low dose ritonavir, the patient's tolerability of bosentan should be monitored.
<b>HEPATITIS C VIRUS (HCV) DIRECT-ACTING ANTIVIRALS</b>		
<b><i>NS3-4A protease inhibitors</i></b>		
Elbasvir/grazoprevir	Darunavir with low dose ritonavir may increase the exposure to grazoprevir. (CYP3A and OATP1B inhibition).	Concomitant use of darunavir with low dose ritonavir and elbasvir/grazoprevir is contraindicated (see section 4.3).
Glecaprevir/pibrentasvir	Based on theoretical considerations boosted darunavir may increase the exposure to glecaprevir and pibrentasvir. (P-gp, BCRP and/or OATP1B1/3 inhibition)	It is not recommended to co-administer boosted darunavir with glecaprevir/pibrentasvir.
<b>HERBAL PRODUCTS</b>		
St John's wort ( <i>Hypericum perforatum</i> )	Not studied. St John's wort is expected to decrease the plasma concentrations of darunavir and ritonavir. (CYP450 induction)	Darunavir co-administered with low dose ritonavir must not be used concomitantly with products containing St John's wort ( <i>Hypericum perforatum</i> ) (see section 4.3). If a patient is already taking St John's wort, stop St John's wort and if possible check viral levels. Darunavir exposure (and also ritonavir exposure) may increase on stopping St John's wort. The inducing effect may persist for at least 2 weeks after cessation of treatment with St John's wort.
<b>HMG CO-A REDUCTASE INHIBITORS</b>		
Lovastatin Simvastatin	Not studied. Lovastatin and simvastatin are expected to have markedly increased plasma concentrations when co-administered with darunavir co-administered with low dose ritonavir. (CYP3A inhibition)	Increased plasma concentrations of lovastatin or simvastatin may cause myopathy, including rhabdomyolysis. Concomitant use of darunavir co-administered with low dose ritonavir with lovastatin and simvastatin is therefore contraindicated (see section 4.3).
Atorvastatin 10 mg once daily	atorvastatin AUC ↑ 3-4 fold atorvastatin C <sub>min</sub> ↑ ≈5.5-10 fold atorvastatin C <sub>max</sub> ↑ ≈2 fold #darunavir/ritonavir	When administration of atorvastatin and darunavir co-administered with low dose ritonavir is desired, it is recommended to start with an atorvastatin dose of 10 mg once daily. A gradual dose increase of atorvastatin may be tailored to the clinical response.

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
Pravastatin 40 mg single dose	pravastatin AUC ↑ 81% <sup>¶</sup> pravastatin C <sub>min</sub> ND pravastatin C <sub>max</sub> ↑ 63% <sup>¶</sup> an up to five-fold increase was seen in a limited subset of subjects	When administration of pravastatin and darunavir co-administered with low dose ritonavir is required, it is recommended to start with the lowest possible dose of pravastatin and titrate up to the desired clinical effect while monitoring for safety.
Rosuvastatin 10 mg once daily	rosuvastatin AUC ↑ 48% <sup>  </sup> rosuvastatin C <sub>max</sub> ↑ 144% <sup>  </sup> <sup>  </sup> based on published data with darunavir/ritonavir	When administration of rosuvastatin and darunavir co-administered with low dose ritonavir is required, it is recommended to start with the lowest possible dose of rosuvastatin and titrate up to the desired clinical effect while monitoring for safety.
<b>OTHER LIPID MODIFYING AGENTS</b>		
Lomitapide	Based on theoretical considerations boosted darunavir is expected to increase the exposure of lomitapide when co-administered. (CYP3A inhibition)	Co-administration is contraindicated (see section 4.3).
<b>H2-RECEPTOR ANTAGONISTS</b>		
Ranitidine 150 mg twice daily	<sup>#</sup> darunavir AUC ↔ <sup>#</sup> darunavir C <sub>min</sub> ↔ <sup>#</sup> darunavir C <sub>max</sub> ↔	Darunavir co-administered with low dose ritonavir can be co-administered with H2-receptor antagonists without dose adjustments.
<b>IMMUNOSUPPRESSANTS</b>		
Ciclosporin Sirolimus Tacrolimus  Everolimus	Not studied. Exposure to these immunosuppressants will be increased when co-administered with darunavir co-administered with low dose ritonavir. (CYP3A inhibition)	Therapeutic drug monitoring of the immunosuppressive agent must be done when co-administration occurs.  Concomitant use of everolimus and darunavir co-administered with low dose ritonavir is not recommended.
<b>INHALED BETA AGONISTS</b>		
Salmeterol	Not studied. Concomitant use of salmeterol and darunavir co-administered with low dose ritonavir may increase plasma concentrations of salmeterol.	Concomitant use of salmeterol and darunavir co-administered with low dose ritonavir is not recommended. The combination may result in increased risk of cardiovascular adverse event with salmeterol, including QT prolongation, palpitations and sinus tachycardia.
<b>NARCOTIC ANALGESICS / TREATMENT OF OPIOID DEPENDENCE</b>		
Methadone individual dose ranging from 55 mg to 150 mg once daily	R(-) methadone AUC ↓ 16% R(-) methadone C <sub>min</sub> ↓ 15% R(-) methadone C <sub>max</sub> ↓ 24%	No adjustment of methadone dosage is required when initiating co-administration with darunavir/ritonavir. However, increased methadone dose may be necessary when concomitantly administered for a longer period of time due to induction of metabolism by ritonavir. Therefore, clinical monitoring is recommended, as maintenance therapy may need to be adjusted in some patients.

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
Buprenorphine/naloxone 8/2 mg–16/4 mg once daily	buprenorphine AUC ↓ 11% buprenorphine C <sub>min</sub> ↔ buprenorphine C <sub>max</sub> ↓ 8% norbuprenorphine AUC ↑ 46% norbuprenorphine C <sub>min</sub> ↑ 71% norbuprenorphine C <sub>max</sub> ↑ 36% naloxone AUC ↔ naloxone C <sub>min</sub> ND naloxone C <sub>max</sub> ↔	The clinical relevance of the increase in norbuprenorphine pharmacokinetic parameters has not been established. Dose adjustment for buprenorphine may not be necessary when co-administered with darunavir/ritonavir but a careful clinical monitoring for signs of opiate toxicity is recommended.
Fentanyl Oxycodone Tramadol	Based on theoretical considerations boosted darunavir may increase plasma concentrations of these analgesics. (CYP2D6 and/or CYP3A inhibition)	Clinical monitoring is recommended when co-administering boosted darunavir with these analgesics.
<b>OESTROGEN-BASED CONTRACEPTIVES</b>		
Drospirenone Ethinylestradiol (3 mg/0.02 mg once daily)  Ethinylestradiol Norethindrone 35 µg/1 mg once daily	Not studied with darunavir/ritonavir.  ethinylestradiol AUC ↓ 44% <sup>β</sup> ethinylestradiol C <sub>min</sub> ↓ 62% <sup>β</sup> ethinylestradiol C <sub>max</sub> ↓ 32% <sup>β</sup> norethindrone AUC ↓ 14% <sup>β</sup> norethindrone C <sub>min</sub> ↓ 30% <sup>β</sup> norethindrone C <sub>max</sub> ↔ <sup>β</sup> <sup>β</sup> with darunavir/ritonavir	When darunavir is coadministered with a drospirenone-containing product, clinical monitoring is recommended due to the potential for hyperkalaemia.  Alternative or additional contraceptive measures are recommended when oestrogen-based contraceptives are co-administered with darunavir and low dose ritonavir.  Patients using oestrogens as hormone replacement therapy should be clinically monitored for signs of oestrogen deficiency.
<b>OPIOID ANTAGONIST</b>		
Naloxegol	Not studied.	Co-administration of boosted darunavir and naloxegol is contraindicated.
<b>PHOSPHODIESTERASE, TYPE 5 (PDE-5) INHIBITORS</b>		
For the treatment of erectile dysfunction Avanafil Sildenafil Tadalafil Vardenafil	In an interaction study <sup>#</sup> , a comparable systemic exposure to sildenafil was observed for a single intake of 100 mg sildenafil alone and a single intake of 25 mg sildenafil co-administered with darunavir and low dose ritonavir.	The combination of avanafil and darunavir with low dose ritonavir is contraindicated (see section 4.3). Concomitant use of other PDE-5 inhibitors for the treatment of erectile dysfunction with darunavir co-administered with low dose ritonavir should be done with caution. If concomitant use of darunavir co-administered with low dose ritonavir with sildenafil, vardenafil or tadalafil is indicated, sildenafil at a single dose not exceeding 25 mg in 48 hours, vardenafil at a single dose not exceeding 2.5 mg in 72 hours or tadalafil at a single dose not exceeding 10 mg in 72 hours is recommended.

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
For the treatment of pulmonary arterial hypertension Sildenafil Tadalafil	Not studied. Concomitant use of sildenafil or tadalafil for the treatment of pulmonary arterial hypertension and darunavir co-administered with low dose ritonavir may increase plasma concentrations of sildenafil or tadalafil. (CYP3A inhibition)	A safe and effective dose of sildenafil for the treatment of pulmonary arterial hypertension co-administered with darunavir and low dose ritonavir has not been established. There is an increased potential for sildenafil- associated adverse events (including visual disturbances, hypotension, prolonged erection and syncope). Therefore, co- administration of darunavir with low dose ritonavir and sildenafil when used for the treatment of pulmonary arterial hypertension is contraindicated (see section 4.3). Co-administration of tadalafil for the treatment of pulmonary arterial hypertension with darunavir and low dose ritonavir is not recommended.
<b>PROTON PUMP INHIBITORS</b>		
Omeprazole 20 mg once daily	#darunavir AUC ↔ #darunavir C <sub>min</sub> ↔ #darunavir C <sub>max</sub> ↔	Darunavir co-administered with low dose ritonavir can be co-administered with proton pump inhibitors without dose adjustments.
<b>SEDATIVES/HYPNOTICS</b>		
Buspirone Clorazepate Diazepam Estazolam Flurazepam Midazolam (parenteral) Zolpidem          Midazolam (oral) Triazolam	Not studied. Sedative/hypnotics are extensively metabolised by CYP3A. Co-administration with darunavir/ritonavir may cause a large increase in the concentration of these medicines.  If parenteral midazolam is co-administered with darunavir co-administered with low dose ritonavir it may cause a large increase in the concentration of this benzodiazepine. Data from concomitant use of parenteral midazolam with other protease inhibitors suggest a possible 3-4 fold increase in midazolam plasma levels.	Clinical monitoring is recommended when co-administering darunavir with these sedatives/hypnotics and a lower dose of the sedatives/hypnotics should be considered.  If parenteral midazolam is co-administered with darunavir with low dose ritonavir, it should be done in an intensive care unit (ICU) or similar setting, which ensures close clinical monitoring and appropriate medical management in case of respiratory depression and/or prolonged sedation. Dose adjustment for midazolam should be considered, especially if more than a single dose of midazolam is administered.  Darunavir co-administered with low dose ritonavir is contraindicated with triazolam or oral midazolam (see section 4.3).
<b>TREATMENT FOR PREMATURE EJACULATION</b>		
Dapoxetine	Not studied.	Co-administration of boosted darunavir with dapoxetine is contraindicated.

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
<b>UROLOGICAL DRUGS</b>		
Fesoterodine Solifenacin	Not studied.	Use with caution. Monitor for fesoterodine or solifenacin adverse reactions, dose reduction of fesoterodine or solifenacin may be necessary.

# Studies have been performed at lower than recommended doses of darunavir or with a different dosing regimen (see section 4.2 Posology).

† The efficacy and safety of the use of darunavir with 100 mg ritonavir and any other HIV PI (e.g. (fos)amprenavir and tipranavir) has not been established in HIV patients. According to current treatment guidelines, dual therapy with protease inhibitors is generally not recommended.

‡ Study was conducted with tenofovir disoproxil fumarate 300 mg once daily.

## 4.6 Fertility, pregnancy and lactation

### Pregnancy

As a general rule, when deciding to use antiretroviral agents for the treatment of HIV infection in pregnant women and consequently for reducing the risk of HIV vertical transmission to the newborn, the animal data as well as the clinical experience in pregnant women should be taken into account.

There are no adequate and well controlled studies on pregnancy outcome with darunavir in pregnant women. Studies in animals do not indicate direct harmful effects with respect to pregnancy, embryonal/foetal development, parturition or postnatal development (see section 5.3).

Darunavir co-administered with low dose ritonavir should be used during pregnancy only if the potential benefit justifies the potential risk.

### Breast-feeding

It is not known whether darunavir is excreted in human milk. Studies in rats have demonstrated that darunavir is excreted in milk and at high levels (1,000 mg/kg/day) resulted in toxicity of the offspring.

Because of the potential for adverse reactions in breast-fed infants, women should be instructed not to breast-feed if they are receiving darunavir.

In order to avoid transmission of HIV to the infant it is recommended that women living with HIV do not breast-feed.

### Fertility

No human data on the effect of darunavir on fertility are available. There was no effect on mating or fertility with darunavir treatment in rats (see section 5.3).

## 4.7 Effects on ability to drive and use machines

Darunavir in combination with ritonavir has no or negligible influence on the ability to drive and use machines. However, dizziness has been reported in some patients during treatment with regimens containing darunavir co-administered with low dose ritonavir and should be borne in mind when considering a patient's ability to drive or operate machinery (see section 4.8).

## 4.8 Undesirable effects

### Summary of the safety profile

During the clinical development program (N=2,613 treatment-experienced subjects who initiated therapy with darunavir/ ritonavir 600/100 mg twice daily), 51.3% of subjects experienced at least one adverse reaction. The total mean treatment duration for subjects was 95.3 weeks. The most frequent adverse reactions reported in clinical trials and as spontaneous reports are diarrhoea, nausea, rash, headache and vomiting. The most frequent serious reactions are acute renal failure, myocardial infarction, immune reconstitution inflammatory syndrome, thrombocytopenia, osteonecrosis, diarrhoea, hepatitis and pyrexia.

In the 96 week analysis, the safety profile of darunavir/ ritonavir 800/100 mg once daily in treatment-naïve subjects was similar to that seen with darunavir/ ritonavir 600/100 mg twice daily in treatment-experienced subjects except for nausea which was observed more frequently in treatment-naïve subjects. This was driven by mild intensity nausea. No new safety findings were identified in the 192 week analysis of the treatment-naïve subjects in which the mean treatment duration of darunavir/ ritonavir 800/100 mg once daily was 162.5 weeks.

### Tabulated list of adverse reactions

Adverse reactions are listed by system organ class (SOC) and frequency category. Within each frequency category, adverse reactions are presented in order of decreasing seriousness. Frequency categories are defined as follows: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ) and not known (frequency cannot be estimated from the available data).

#### *Adverse reactions observed with darunavir/ritonavir in clinical trials and post-marketing*

MedDRA system organ class Frequency category	Adverse reaction
<i>Infections and infestations</i>	
Uncommon	herpes simplex
<i>Blood and lymphatic system disorders</i>	
Uncommon	thrombocytopenia, neutropenia, anaemia, leucopenia
Rare	increased eosinophil count
<i>Immune system disorders</i>	
Uncommon	immune reconstitution inflammatory syndrome, (drug) hypersensitivity
<i>Endocrine disorders</i>	
Uncommon	hypothyroidism, increased blood thyroid stimulating hormone
<i>Metabolism and nutrition disorders</i>	
Common	diabetes mellitus, hypertriglyceridaemia, hypercholesterolaemia, hyperlipidaemia
Uncommon	gout, anorexia, decreased appetite, decreased weight, increased weight, hyperglycaemia, insulin resistance, decreased high density lipoprotein, increased appetite, polydipsia, increased blood lactate dehydrogenase



MedDRA system organ class Frequency category	Adverse reaction
<i>Psychiatric disorders</i>	
Common	insomnia
Uncommon	depression, disorientation, anxiety, sleep disorder, abnormal dreams, nightmare, decreased libido
Rare	confusional state, altered mood, restlessness
<i>Nervous system disorders</i>	
Common	headache, peripheral neuropathy, dizziness
Uncommon	lethargy, paraesthesia, hypoaesthesia, dysgeusia, disturbance in attention, memory impairment, somnolence
Rare	syncope, convulsion, ageusia, sleep phase rhythm disturbance
<i>Eye disorders</i>	
Uncommon	conjunctival hyperaemia, dry eye
Rare	visual disturbance
<i>Ear and labyrinth disorders</i>	
Uncommon	vertigo
<i>Cardiac disorders</i>	
Uncommon	myocardial infarction, angina pectoris, prolonged electrocardiogram QT, tachycardia
Rare	acute myocardial infarction, sinus bradycardia, palpitations
<i>Vascular disorders</i>	
Uncommon	hypertension, flushing
<i>Respiratory, thoracic and mediastinal disorders</i>	
Uncommon	dyspnoea, cough, epistaxis, throat irritation
Rare	rhinorrhoea
<i>Gastrointestinal disorders</i>	
Very common	diarrhoea
Common	vomiting, nausea, abdominal pain, increased blood amylase, dyspepsia, abdominal distension, flatulence
Uncommon	pancreatitis, gastritis, gastrooesophageal reflux disease, aphthous stomatitis, retching, dry mouth, abdominal discomfort, constipation, increased lipase, eructation, oral dysaesthesia
Rare	stomatitis, haematemesis, cheilitis, dry lip, coated tongue

MedDRA system organ class Frequency category	Adverse reaction
<i>Hepatobiliary disorders</i>	
Common	increased alanine aminotransferase
Uncommon	hepatitis, cytolytic hepatitis, hepatic steatosis, hepatomegaly, increased transaminase, increased aspartate aminotransferase, increased blood bilirubin, increased blood alkaline phosphatase, increased gamma-glutamyltransferase
<i>Skin and subcutaneous tissue disorders</i>	
Common	rash (including macular, maculopapular, papular, erythematous and pruritic rash), pruritus
Uncommon	angioedema, generalised rash, allergic dermatitis, urticaria, eczema, erythema, hyperhidrosis, night sweats, alopecia, acne, dry skin, nail pigmentation
Rare	DRESS, Stevens-Johnson syndrome, erythema multiforme, dermatitis, seborrhoeic dermatitis, skin lesion, xeroderma
Not known	toxic epidermal necrolysis, acute generalised exanthematous pustulosis
<i>Musculoskeletal and connective tissue disorders</i>	
Uncommon	myalgia, osteonecrosis, muscle spasms, muscular weakness, arthralgia, pain in extremity, osteoporosis, increased blood creatine phosphokinase
Rare	musculoskeletal stiffness, arthritis, joint stiffness
<i>Renal and urinary disorders</i>	
Uncommon	acute renal failure, renal failure, nephrolithiasis, increased blood creatinine, proteinuria, bilirubinuria, dysuria, nocturia, pollakiuria
Rare	decreased creatinine renal clearance
Rare	crystal nephropathy <sup>§</sup>
<i>Reproductive system and breast disorders</i>	
Uncommon	erectile dysfunction, gynaecomastia
<i>General disorders and administration site conditions</i>	
Common	asthenia, fatigue
Uncommon	pyrexia, chest pain, peripheral oedema, malaise, feeling hot, irritability, pain
Rare	chills, abnormal feeling, xerosis

<sup>§</sup> adverse reaction identified in the post-marketing setting. Per the guideline on Summary of Product Characteristics (Revision 2, September 2009), the frequency of this adverse reaction in the post-marketing setting was determined using the "Rule of 3".

## Description of selected adverse reactions

### *Rash*

In clinical trials, rash was mostly mild to moderate, often occurring within the first four weeks of treatment and resolving with continued dosing. In cases of severe skin reaction see the warning in section 4.4.

During the clinical development program of raltegravir in treatment-experienced patients, rash, irrespective of causality, was more commonly observed with regimens containing darunavir/ritonavir + raltegravir compared to those containing darunavir/ritonavir without raltegravir or raltegravir without darunavir/ritonavir. Rash considered by the investigator to be drug-related occurred at similar rates. The exposure-adjusted rates of rash (all causality) were 10.9, 4.2, and 3.8 per 100 patient-years (PYR), respectively; and for drug-related rash were 2.4, 1.1, and 2.3 per 100 PYR, respectively. The rashes observed in clinical studies were mild to moderate in severity and did not result in discontinuation of therapy (see section 4.4).

### *Metabolic parameters*

Weight and levels of blood lipids and glucose may increase during antiretroviral therapy (see section 4.4).

### *Musculoskeletal abnormalities*

Increased CPK, myalgia, myositis and rarely, rhabdomyolysis have been reported with the use of protease inhibitors, particularly in combination with NRTIs.

Cases of osteonecrosis have been reported, particularly in patients with generally acknowledged risk factors, advanced HIV disease or long-term exposure to combination antiretroviral therapy (CART). The frequency of this is unknown (see section 4.4).

### *Immune reconstitution inflammatory syndrome*

In HIV infected patients with severe immune deficiency at the time of initiation of combination antiretroviral therapy (CART), an inflammatory reaction to asymptomatic or residual opportunistic infections may arise. Autoimmune disorders (such as Graves' disease and autoimmune hepatitis) have also been reported; however, the reported time to onset is more variable and these events can occur many months after initiation of treatment (see section 4.4).

### *Bleeding in haemophiliac patients*

There have been reports of increased spontaneous bleeding in haemophiliac patients receiving antiretroviral protease inhibitors (see section 4.4).

## Paediatric population

The safety assessment in paediatric patients is based on the 48-week analysis of safety data from three Phase II trials. The following patient populations were evaluated (see section 5.1):

- 80 ART-experienced HIV-1 infected paediatric patients aged from 6 to 17 years and weighing at least 20 kg who received darunavir tablets with low dose ritonavir twice daily in combination with other antiretroviral agents.
- 21 ART-experienced HIV-1 infected paediatric patients aged from 3 to < 6 years and weighing 10 kg to < 20 kg (16 participants from 15 kg to < 20 kg) who received darunavir oral suspension with low dose ritonavir twice daily in combination with other antiretroviral agents.
- 12 ART-naïve HIV-1 infected paediatric patients aged from 12 to 17 years and weighing at least 40 kg who received darunavir tablets with low dose ritonavir once daily in combination with other antiretroviral agents (see section 5.1).

Overall, the safety profile in these paediatric patients was similar to that observed in the adult population.

## Other special populations

### *Patients co-infected with hepatitis B and/or hepatitis C virus*

Among 1,968 treatment-experienced patients receiving darunavir co-administered with ritonavir 600/100 mg twice daily, 236 patients were co-infected with hepatitis B or C. Co-infected patients were more likely to have baseline and treatment emergent hepatic transaminase elevations than those without chronic viral hepatitis (see section 4.4).

## Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

## **4.9 Overdose**

Human experience of acute overdose with darunavir co-administered with low dose ritonavir is limited. Single doses up to 3,200 mg of darunavir as oral solution alone and up to 1,600 mg of the tablet formulation of darunavir in combination with ritonavir have been administered to healthy volunteers without untoward symptomatic effects.

There is no specific antidote for overdose with darunavir. Treatment of overdose with darunavir consists of general supportive measures including monitoring of vital signs and observation of the clinical status of the patient. Since darunavir is highly protein bound, dialysis is unlikely to be beneficial in significant removal of the active substance.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Antivirals for systemic use, protease inhibitors, ATC code: J05AE10.

#### Mechanism of action

Darunavir is an inhibitor of the dimerisation and of the catalytic activity of the HIV-1 protease (KD of  $4.5 \times 10^{-12}$  M). It selectively inhibits the cleavage of HIV encoded Gag-Pol polyproteins in virus infected cells, thereby preventing the formation of mature infectious virus particles.

#### Antiviral activity *in vitro*

Darunavir exhibits activity against laboratory strains and clinical isolates of HIV-1 and laboratory strains of HIV-2 in acutely infected T-cell lines, human peripheral blood mononuclear cells and human monocytes/macrophages with median EC<sub>50</sub> values ranging from 1.2 to 8.5 nM (0.7 to 5.0 ng/ml). Darunavir demonstrates antiviral activity *in vitro* against a broad panel of HIV-1 group M (A, B, C, D, E, F, G) and group O primary isolates with EC<sub>50</sub> values ranging from < 0.1 to 4.3 nM.

These EC<sub>50</sub> values are well below the 50% cellular toxicity concentration range of 87 µM to > 100 µM.

#### Resistance

*In vitro* selection of darunavir-resistant virus from wild type HIV-1 was lengthy (> 3 years). The selected viruses were unable to grow in the presence of darunavir concentrations above 400 nM. Viruses selected in these conditions and showing decreased susceptibility to darunavir (range: 23-50-fold) harboured 2 to 4 amino acid substitutions in the protease gene. The decreased susceptibility to

darunavir of the emerging viruses in the selection experiment could not be explained by the emergence of these protease mutations.

The clinical trial data from ART-experienced patients (*TITAN* trial and the pooled analysis of the *POWER* 1, 2 and 3 and *DUET* 1 and 2 trials) showed that virologic response to darunavir co-administered with low dose ritonavir was decreased when 3 or more darunavir RAMs (V11I, V32I, L33F, I47V, I50V, I54L or M, T74P, L76V, I84V and L89V) were present at baseline or when these mutations developed during treatment.

Increasing baseline darunavir fold change in  $EC_{50}$  (FC) was associated with decreasing virologic response. A lower and upper clinical cut-off of 10 and 40 were identified. Isolates with baseline FC  $\leq 10$  are susceptible; isolates with FC  $> 10$  to 40 have decreased susceptibility; isolates with FC  $> 40$  are resistant (see Clinical results).

Viruses isolated from patients on darunavir/ritonavir 600/100 mg twice daily experiencing virologic failure by rebound that were susceptible to tipranavir at baseline remained susceptible to tipranavir after treatment in the vast majority of cases.

The lowest rates of developing resistant HIV virus are observed in ART-naïve patients who are treated for the first time with darunavir in combination with other ART.

The table below shows the development of HIV 1 protease mutations and loss of susceptibility to PIs in virologic failures at endpoint in the *ARTEMIS*, *ODIN* and *TITAN* trials.

	ARTEMIS Week 192	ODIN Week 48		TITAN Week 48
	Darunavir/ ritonavir 800/100 mg once daily N=343	Darunavir/ ritonavir 800/100 mg once daily N=294	Darunavir/ ritonavir 600/100 mg twice daily N=296	Darunavir/ ritonavir 600/100 mg twice daily N=298
Total number of virologic failures <sup>a</sup> , n (%)	55 (16.0%)	65 (22.1%)	54 (18.2%)	31 (10.4%)
Rebounders				
Never suppressed subjects	39 (11.4%) 16 (4.7%)	11 (3.7%) 54 (18.4%)	11 (3.7%) 43 (14.5%)	16 (5.4%) 15 (5.0%)
Number of subjects with virologic failure and paired baseline/endpoint genotypes, developing mutations <sup>b</sup> at endpoint, n/N				
Primary (major) PI	0/43	1/60	0/42	6/28
Mutations				
PI RAMs	4/43	7/60	4/42	10/28
Number of subjects with virologic failure and paired baseline/endpoint phenotypes, showing loss of susceptibility to PIs at endpoint compared to baseline, n/N				
PI				
darunavir	0/39	1/58	0/41	3/26
amprenavir	0/39	1/58	0/40	0/22
atazanavir	0/39	2/56	0/40	0/22
indinavir	0/39	2/57	0/40	1/24
lopinavir	0/39	1/58	0/40	0/23
saquinavir	0/39	0/56	0/40	0/22
tipranavir	0/39	0/58	0/41	1/25

<sup>a</sup> TLOVR non-VF censored algorithm based on HIV-1 RNA  $< 50$  copies/ml, except for *TITAN* (HIV-1 RNA  $< 400$  copies/ml)

<sup>b</sup> IAS-USA lists

## Cross-resistance

Darunavir FC was less than 10 for 90% of 3,309 clinical isolates resistant to amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir, saquinavir and/or tipranavir showing that viruses resistant to most PIs remain susceptible to darunavir.

In the virologic failures of the *ARTEMIS* trial no cross-resistance with other PIs was observed.

## Clinical results

### Adult patients

For clinical trial results in ART-naïve adult patients, refer to the Summary of Product Characteristics for Darunavir Mylan 400 mg and 800 mg tablets.

### Efficacy of darunavir 600 mg twice daily co-administered with 100 mg ritonavir twice daily in ART-experienced patients

The evidence of efficacy of darunavir co-administered with ritonavir (600/100 mg twice daily) in ART-experienced patients is based on the 96 weeks analysis of the Phase III trial *TITAN* in ART-experienced lopinavir naïve patients, on the 48 week analysis of the Phase III trial *ODIN* in ART-experienced patients with no DRV-RAMs, and on the analyses of 96 weeks data from the Phase IIb trials *POWER* 1 and 2 in ART-experienced patients with high level of PI resistance.

*TITAN* is a randomised, controlled, open-label Phase III trial comparing darunavir co-administered with ritonavir (600/100 mg twice daily) versus lopinavir/ritonavir (400/100 mg twice daily) in ART-experienced, lopinavir naïve HIV-1 infected adult patients. Both arms used an Optimised Background Regimen (OBR) consisting of at least 2 antiretrovirals (NRTIs with or without NNRTIs).

The table below shows the efficacy data of the 48 week analysis from the *TITAN* trial.

TITAN			
Outcomes	Darunavir/ritonavir 600/100 mg twice daily + OBR N=298	Lopinavir/ ritonavir 400/100 mg twice daily + OBR N=297	Treatment difference (95% CI of difference)
HIV-1 RNA < 50 copies/ml <sup>a</sup>	70.8% (211)	60.3% (179)	10.5% (2.9; 18.1) <sup>b</sup>
median CD4+ cell count change from baseline (x 10 <sup>6</sup> /L) <sup>c</sup>	88	81	

<sup>a</sup> Imputations according to the TLOVR algorithm

<sup>b</sup> Based on a normal approximation of the difference in % response

<sup>c</sup> NC=F

At 48 weeks non-inferiority in virologic response to the darunavir/ritonavir treatment, defined as the percentage of patients with plasma HIV-1 RNA level < 400 and < 50 copies/ml, was demonstrated (at the pre-defined 12% non-inferiority margin) for both ITT and OP populations. These results were confirmed in the analysis of data at 96 weeks of treatment in the *TITAN* trial, with 60.4% of patients in the darunavir/ritonavir arm having HIV-1 RNA < 50 copies/ml at week 96 compared to 55.2% in the lopinavir/ritonavir arm [difference: 5.2%, 95% CI (-2.8; 13.1)].

*ODIN* is a Phase III, randomised, open-label trial comparing darunavir/ritonavir 800/100 mg once daily versus darunavir/ritonavir 600/100 mg twice daily in ART-experienced HIV-1 infected patients with screening genotype resistance testing showing no darunavir RAMs (i.e. V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V, L89V) and a screening HIV-1 RNA > 1,000 copies/ml. Efficacy analysis is based on 48 weeks of treatment (see table below). Both arms used an optimised background regimen (OBR) of ≥ 2 NRTIs.

ODIN			
<i>Outcomes</i>	Darunavir/ritonavir 800/100 mg once daily + OBR N=294	Darunavir/ritonavir 600/100 mg twice daily + OBR N=296	Treatment difference (95% CI of difference)
HIV-1 RNA < 50 copies/ml <sup>a</sup>	72.1% (212)	70.9% (210)	1.2% (-6.1; 8.5) <sup>b</sup>
With Baseline HIV-1 RNA (copies/ml)			
< 100,000	77.6% (198/255)	73.2% (194/265)	4.4% (-3.0; 11.9)
≥ 100,000	35.9% (14/39)	51.6% (16/31)	-15.7% (-39.2; 7.7)
With Baseline CD4+ cell count (x 10 <sup>6</sup> /L)			
≥ 100	75.1% (184/245)	72.5% (187/258)	2.6% (-5.1; 10.3)
< 100	57.1% (28/49)	60.5% (23/38)	-3.4% (-24.5; 17.8)
With HIV-1 clade			
Type B	70.4% (126/179)	64.3% (128/199)	6.1% (-3.4; 15.6)
Type AE	90.5% (38/42)	91.2% (31/34)	-0.7% (-14.0; 12.6)
Type C	72.7% (32/44)	78.8% (26/33)	-6.1% (-2.6; 13.7)
Other <sup>c</sup>	55.2% (16/29)	83.3% (25/30)	-28.2% (-51.0; -5.3)
mean CD4+ cell count change from baseline (x 10 <sup>6</sup> /L) <sup>e</sup>	108	112	-5d (-25; 16)

<sup>a</sup> Imputations according to the TLOVR algorithm

<sup>b</sup> Based on a normal approximation of the difference in % response

<sup>c</sup> Clades A1, D, F1, G, K, CRF02\_AG, CRF12\_BF, and CRF06\_CPX

<sup>d</sup> Difference in means

<sup>e</sup> Last Observation Carried Forward imputation

At 48 weeks, virologic response, defined as the percentage of patients with plasma HIV-1 RNA level < 50 copies/ml, with darunavir/ritonavir 800/100 mg once daily treatment was demonstrated to be non-inferior (at the pre-defined 12% non-inferiority margin) compared to darunavir/ritonavir 600/100 mg twice daily for both ITT and OP populations.

Darunavir/ritonavir 800/100 mg once daily in ART-experienced patients should not be used in patients with one or more darunavir resistance associated mutations (DRV-RAMs) or HIV-1 RNA ≥ 100,000 copies/ml or CD4+ cell count < 100 cells x 10<sup>6</sup>/L (see sections 4.2 and 4.4). Limited data is available in patients with HIV-1 clades other than B.

**POWER 1** and **POWER 2** are randomised, controlled trials comparing darunavir co-administered with ritonavir (600/100 mg twice daily) with a control group receiving an investigator-selected PI(s) regimen in HIV-1 infected patients who had previously failed more than 1 PI containing regimen. An OBR consisting of at least 2 NRTIs with or without enfuvirtide (ENF) was used in both trials.

The table below shows the efficacy data of the 48-week and 96-week analyses from the pooled **POWER 1** and **POWER 2** trials.

POWER 1 and POWER 2 pooled data						
Outcomes	Week 48			Week 96		
	Darunavir/ ritonavir 600/100 mg twice daily n=131	Control n=124	Treatment difference	Darunavir/ ritonavir 600/100 mg twice daily n=131	Control n=124	Treatment difference
HIV RNA < 50 copies/ml <sup>a</sup>	45.0% (59)	11.3% (14)	33.7% (23.4%; 44.1%) <sup>c</sup>	38.9% (51)	8.9% (11)	30.1% (20.1; 40.0) <sup>c</sup>
CD4+ cell count mean change from baseline (x 10 <sup>6</sup> /L) <sup>b</sup>	103	17	86 (57; 114) <sup>c</sup>	133	15	118 (83.9; 153.4) <sup>c</sup>

<sup>a</sup> Imputations according to the TLOVR algorithm

<sup>b</sup> Last Observation Carried Forward imputation

<sup>c</sup> 95% confidence intervals.

Analyses of data through 96 weeks of treatment in the *POWER* trials demonstrated sustained antiretroviral efficacy and immunologic benefit.

Out of the 59 patients who responded with complete viral suppression (< 50 copies/ml) at week 48, 47 patients (80% of the responders at week 48) remained responders at week 96.

#### *Baseline genotype or phenotype and virologic outcome*

Baseline genotype and darunavir FC (shift in susceptibility relative to reference) were shown to be a predictive factor of virologic outcome.

*Proportion (%) of patients with response (HIV-1 RNA < 50 copies/ml at week 24) to darunavir co-administered with ritonavir (600/100 mg twice daily) by baseline genotype<sup>a</sup>, and baseline darunavir FC and by use of enfuvirtide (ENF): As treated analysis of the POWER and DUET trials.*

Response (HIV-1 RNA < 50 copies/ml at week 24) %, n/N	Number of baseline mutations <sup>a</sup>				Baseline DRV FC <sup>b</sup>			
	All ranges	0-2	3	≥ 4	All ranges	≤ 10	10-40	> 40
All patients	45% 455/1,014	54% 359/660	39% 67/172	12% 20/171	45% 455/1,014	55% 364/659	29% 59/203	8% 9/118
Patients with no/non-naïve use of ENF <sup>c</sup>	39% 290/741	50% 238/477	29% 35/120	7% 10/135	39% 290/741	51% 244/477	17% 25/147	5% 5/94
Patients with naïve use of ENF <sup>d</sup>	60% 165/273	66% 121/183	62% 32/52	28% 10/36	60% 165/273	66% 120/182	61% 34/56	17% 4/24

<sup>a</sup> Number of mutations from the list of mutations associated with a diminished response to darunavir/ritonavir (V11I, V32I, L33F, I47V, I50V, I54L or M, T74P, L76V, I84V or L89V)

<sup>b</sup> fold change in EC<sub>50</sub>

<sup>c</sup> “Patients with no/non-naïve use of ENF” are patients who did not use ENF or who used ENF but not for the first time

<sup>d</sup> “Patients with naïve use of ENF” are patients who used ENF for the first time

#### Paediatric patients

For clinical trial results in ART-naïve paediatric patients aged 12 to 17 years, refer to the Summary of Product Characteristics for Darunavir Mylan 400 mg and 800 mg tablets.

#### ART-experienced paediatric patients from the age of 6 to < 18 years, and weighing at least 20 kg

**DELPHI** is an open-label, Phase II trial evaluating the pharmacokinetics, safety, tolerability, and efficacy of darunavir with low dose ritonavir in 80 ART-experienced HIV-1 infected paediatric patients aged 6 to 17 years and weighing at least 20 kg. These patients received darunavir/ritonavir twice daily in combination with other antiretroviral agents (see section 4.2 for dosage



recommendations per body weight). Virologic response was defined as a decrease in plasma HIV-1 RNA viral load of at least 1.0 log<sub>10</sub> versus baseline.

In the study, patients who were at risk of discontinuing therapy due to intolerance of ritonavir oral solution (e.g. taste aversion) were allowed to switch to the capsule formulation. Of the 44 patients taking ritonavir oral solution, 27 switched to the 100 mg capsule formulation and exceeded the weight-based ritonavir dose without changes in observed safety.

DELPHI	
<i>Outcomes at week 48</i>	Darunavir/ritonavir N=80
HIV-1 RNA < 50 copies/ml <sup>a</sup>	47.5% (38)
CD4+ cell count mean change from baseline <sup>b</sup>	147

<sup>a</sup> Imputations according to the TLOVR algorithm.

<sup>b</sup> Non-completer is failure imputation: patients who discontinued prematurely are imputed with a change equal to 0.

According to the TLOVR non-virologic failure censored algorithm 24 (30.0%) patients experienced virological failure, of which 17 (21.3%) patients were rebounders and 7 (8.8%) patients were non-responders.

#### ART-experienced paediatric patients from the age of 3 to < 6 years

The pharmacokinetics, safety, tolerability and efficacy of darunavir/ritonavir twice daily, in combination with other antiretroviral agents in 21 ART-experienced HIV-1 infected paediatric patients aged 3 to < 6 years and weighing 10 kg to < 20 kg was evaluated in an open-label, Phase II trial, **ARIEL**. Patients received a weight-based twice daily treatment regimen, patients weighing 10 kg to < 15 kg received darunavir/ritonavir 25/3 mg/kg twice daily, and patients weighing 15 kg to < 20 kg received darunavir/ritonavir 375/50 mg twice daily. At week 48, the virologic response, defined as the percentage of patients with confirmed plasma viral load < 50 HIV-1 RNA copies/ml, was evaluated in 16 paediatric patients 15 kg to < 20 kg and 5 paediatric patients 10 kg to < 15 kg receiving darunavir/ritonavir in combination with other antiretroviral agents (see section 4.2 for dosage recommendations per body weight).

ARIEL		
<i>Outcomes at week 48</i>	Darunavir/ritonavir	
	10 kg to < 15 kg N=5	15 kg to < 20 kg N=16
HIV-1 RNA < 50 copies/ml <sup>a</sup>	80.0% (4)	81.3% (13)
CD4+ percent change from baseline <sup>b</sup>	4	4
CD4+ cell count mean change from baseline <sup>b</sup>	16	241

<sup>a</sup> Imputations according to the TLOVR algorithm.

<sup>b</sup> NC=F

Limited efficacy data are available in paediatric patients below 15 kg and no recommendation on a posology can be made.

#### *Pregnancy and postpartum*

Darunavir/ritonavir (600/100 mg twice daily or 800/100 mg once daily) in combination with a background regimen was evaluated in a clinical trial of 36 pregnant women (18 in each arm) during the second and third trimesters, and postpartum. Virologic response was preserved throughout the study period in both arms. No mother to child transmission occurred in the infants born to the 31 subjects who stayed on the antiretroviral treatment through delivery. There were no new clinically relevant safety findings compared with the known safety profile of darunavir/ritonavir in HIV-1 infected adults (see sections 4.2, 4.4 and 5.2).

## 5.2 Pharmacokinetic properties

The pharmacokinetic properties of darunavir, co-administered with ritonavir, have been evaluated in healthy adult volunteers and in HIV-1 infected patients. Exposure to darunavir was higher in HIV-1 infected patients than in healthy subjects. The increased exposure to darunavir in HIV-1 infected patients compared to healthy subjects may be explained by the higher concentrations of  $\alpha_1$ -acid glycoprotein (AAG) in HIV-1 infected patients, resulting in higher darunavir binding to plasma AAG and, therefore, higher plasma concentrations.

Darunavir is primarily metabolised by CYP3A. Ritonavir inhibits CYP3A, thereby increasing the plasma concentrations of darunavir considerably.

### Absorption

Darunavir was rapidly absorbed following oral administration. Maximum plasma concentration of darunavir in the presence of low dose ritonavir is generally achieved within 2.5-4.0 hours.

The absolute oral bioavailability of a single 600 mg dose of darunavir alone was approximately 37% and increased to approximately 82% in the presence of 100 mg twice daily ritonavir. The overall pharmacokinetic enhancement effect by ritonavir was an approximate 14-fold increase in the systemic exposure of darunavir when a single dose of 600 mg darunavir was given orally in combination with ritonavir at 100 mg twice daily (see section 4.4).

When administered without food, the relative bioavailability of darunavir in the presence of low dose ritonavir is 30% lower as compared to intake with food. Therefore, darunavir tablets should be taken with ritonavir and with food. The type of food does not affect exposure to darunavir.

### Distribution

Darunavir is approximately 95% bound to plasma protein. Darunavir binds primarily to plasma  $\alpha_1$ -acid glycoprotein.

Following intravenous administration, the volume of distribution of darunavir alone was  $88.1 \pm 59.0$  l (Mean  $\pm$  SD) and increased to  $131 \pm 49.9$  l (Mean  $\pm$  SD) in the presence of 100 mg twice-daily ritonavir.

### Biotransformation

*In vitro* experiments with human liver microsomes (HLMs) indicate that darunavir primarily undergoes oxidative metabolism. Darunavir is extensively metabolised by the hepatic CYP system and almost exclusively by isozyme CYP3A4. A  $^{14}\text{C}$ -darunavir trial in healthy volunteers showed that a majority of the radioactivity in plasma after a single 400/100 mg darunavir with ritonavir dose was due to the parent active substance. At least 3 oxidative metabolites of darunavir have been identified in humans; all showed activity that was at least 10-fold less than the activity of darunavir against wild type HIV.

### Elimination

After a 400/100 mg  $^{14}\text{C}$ -darunavir with ritonavir dose, approximately 79.5% and 13.9% of the administered dose of  $^{14}\text{C}$ -darunavir could be retrieved in faeces and urine, respectively. Unchanged darunavir accounted for approximately 41.2% and 7.7% of the administered dose in faeces and urine, respectively. The terminal elimination half-life of darunavir was approximately 15 hours when combined with ritonavir.

The intravenous clearance of darunavir alone (150 mg) and in the presence of low dose ritonavir was 32.8 l/h and 5.9 l/h, respectively.

## Special populations

### *Paediatric population*

The pharmacokinetics of darunavir in combination with ritonavir taken twice daily in 74 treatment-experienced paediatric patients, aged 6 to 17 years and weighing at least 20 kg, showed that the administered weight-based doses of darunavir/ritonavir resulted in darunavir exposure comparable to that in adults receiving darunavir/ritonavir 600/100 mg twice daily (see section 4.2).

The pharmacokinetics of darunavir in combination with ritonavir taken twice daily in 14 treatment-experienced paediatric patients, aged 3 to < 6 years and weighing at least 15 kg to < 20 kg, showed that weight-based dosages resulted in darunavir exposure that was comparable to that achieved in adults receiving darunavir/ritonavir 600/100 mg twice daily (see section 4.2).

The pharmacokinetics of darunavir in combination with ritonavir taken once daily in 12 ART-naïve paediatric patients, aged 12 to < 18 years and weighing at least 40 kg, showed that darunavir/ritonavir 800/100 mg once daily results in darunavir exposure that was comparable to that achieved in adults receiving darunavir/ritonavir 800/100 mg once daily. Therefore the same once daily dosage may be used in treatment-experienced adolescents aged 12 to < 18 years and weighing at least 40 kg without darunavir resistance associated mutations (DRV-RAMs)\* and who have plasma HIV-1 RNA < 100,000 copies/ml and CD4+ cell count  $\geq 100$  cells  $\times 10^6/L$  (see section 4.2).

\* DRV-RAMs: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

The pharmacokinetics of darunavir in combination with ritonavir taken once daily in 10 treatment-experienced paediatric patients, aged 3 to < 6 years and weighing at least 14 kg to < 20 kg, showed that weight-based dosages resulted in darunavir exposure that was comparable to that achieved in adults receiving darunavir/ritonavir 800/100 mg once daily (see section 4.2). In addition, pharmacokinetic modeling and simulation of darunavir exposures in paediatric patients across the ages of 3 to < 18 years confirmed the darunavir exposures as observed in the clinical studies and allowed the identification of weight-based darunavir/ritonavir once daily dosing regimens for paediatric patients weighing at least 15 kg that are either ART-naïve or treatment-experienced paediatric patients without DRV-RAMs\* and who have plasma HIV-1 RNA < 100,000 copies/ml and CD4+ cell count  $\geq 100$  cells  $\times 10^6/L$  (see section 4.2).

\* DRV-RAMs: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

### *Elderly*

Population pharmacokinetic analysis in HIV infected patients showed that darunavir pharmacokinetics are not considerably different in the age range (18 to 75 years) evaluated in HIV infected patients (n=12, age  $\geq 65$ ) (see section 4.4). However, only limited data were available in patients above the age of 65 year.

### *Gender*

Population pharmacokinetic analysis showed a slightly higher darunavir exposure (16.8%) in HIV infected females compared to males. This difference is not clinically relevant.

### *Renal impairment*

Results from a mass balance study with  $^{14}C$ -darunavir with ritonavir showed that approximately 7.7% of the administered dose of darunavir is excreted in the urine unchanged.

Although darunavir has not been studied in patients with renal impairment, population pharmacokinetic analysis showed that the pharmacokinetics of darunavir were not significantly affected in HIV infected patients with moderate renal impairment (CrCl between 30-60 ml/min, n=20) (see sections 4.2 and 4.4).

### *Hepatic impairment*

Darunavir is primarily metabolised and eliminated by the liver. In a multiple dose study with darunavir co-administered with ritonavir (600/100 mg) twice daily, it was demonstrated that the total plasma concentrations of darunavir in subjects with mild (Child-Pugh Class A, n=8) and moderate (Child-Pugh Class B, n=8) hepatic impairment were comparable with those in healthy subjects. However,

unbound darunavir concentrations were approximately 55% (Child-Pugh Class A) and 100% (Child-Pugh Class B) higher, respectively. The clinical relevance of this increase is unknown therefore, darunavir should be used with caution. The effect of severe hepatic impairment on the pharmacokinetics of darunavir has not been studied (see sections 4.2, 4.3 and 4.4).

#### *Pregnancy and postpartum*

The exposure to total darunavir and ritonavir after intake of darunavir/ritonavir 600/100 mg twice daily and darunavir/ritonavir 800/100 mg once daily as part of an antiretroviral regimen was generally lower during pregnancy compared with postpartum. However, for unbound (i.e. active) darunavir, the pharmacokinetic parameters were less reduced during pregnancy compared to postpartum, due to an increase in the unbound fraction of darunavir during pregnancy compared to postpartum.

<b>Pharmacokinetic results of total darunavir after administration of darunavir/ritonavir at 600/100 mg twice daily as part of an antiretroviral regimen, during the second trimester of pregnancy, the third trimester of pregnancy and postpartum</b>			
<b>Pharmacokinetics of total darunavir (mean ± SD)</b>	<b>Second trimester of pregnancy (n=12)<sup>a</sup></b>	<b>Third trimester of pregnancy (n=12)</b>	<b>Postpartum (6-12 weeks) (n=12)</b>
$C_{max}$ , ng/ml	4,668 ± 1,097	5,328 ± 1,631	6,659 ± 2,364
$AUC_{12h}$ , ng.h/ml	39,370 ± 9,597	45,880 ± 17,360	56,890 ± 26,340
$C_{min}$ , ng/ml	1,922 ± 825	2,661 ± 1,269	2,851 ± 2,216

<sup>a</sup> n=11 for  $AUC_{12h}$

<b>Pharmacokinetic results of total darunavir after administration of darunavir/ritonavir at 800/100 mg once daily as part of an antiretroviral regimen, during the second trimester of pregnancy, the third trimester of pregnancy and postpartum</b>			
<b>Pharmacokinetics of total darunavir (mean ± SD)</b>	<b>Second trimester of pregnancy (n=17)</b>	<b>Third Trimester of pregnancy (n=15)</b>	<b>Postpartum (6-12 weeks) (n=16)</b>
$C_{max}$ , ng/ml	4,964 ± 1,505	5,132 ± 1,198	7,310 ± 1,704
$AUC_{24h}$ , ng.h/ml	62,289 ± 16,234	61,112 ± 13,790	92,116 ± 29,241
$C_{min}$ , ng/ml	1,248 ± 542	1,075 ± 594	1,473 ± 1,141

In women receiving darunavir/ritonavir 600/100 mg twice daily during the second trimester of pregnancy, mean intra-individual values for total darunavir  $C_{max}$ ,  $AUC_{12h}$  and  $C_{min}$  were 28%, 26% and 26% lower, respectively, as compared with postpartum; during the third trimester of pregnancy, total darunavir  $C_{max}$ ,  $AUC_{12h}$  and  $C_{min}$  values were 18%, 16% lower and 2% higher, respectively, as compared with postpartum.

In women receiving darunavir/ritonavir 800/100 mg once daily during the second trimester of pregnancy, mean intra-individual values for total darunavir  $C_{max}$ ,  $AUC_{24h}$  and  $C_{min}$  were 33%, 31% and 30% lower, respectively, as compared with postpartum; during the third trimester of pregnancy, total darunavir  $C_{max}$ ,  $AUC_{24h}$  and  $C_{min}$  values were 29%, 32% and 50% lower, respectively, as compared with postpartum.

### **5.3 Preclinical safety data**

Animal toxicology studies have been conducted at exposures up to clinical exposure levels with darunavir alone, in mice, rats and dogs and in combination with ritonavir in rats and dogs.

In repeated-dose toxicology studies in mice, rats and dogs, there were only limited effects of treatment with darunavir. In rodents the target organs identified were the haematopoietic system, the blood coagulation system, liver and thyroid. A variable but limited decrease in red blood cell-related parameters was observed, together with increases in activated partial thromboplastin time.

Changes were observed in liver (hepatocyte hypertrophy, vacuolation, increased liver enzymes) and thyroid (follicular hypertrophy). In the rat, the combination of darunavir with ritonavir lead to a small increase in effect on RBC parameters, liver and thyroid and increased incidence of islet fibrosis in the pancreas (in male rats only) compared to treatment with darunavir alone. In the dog, no major toxicity findings or target organs were identified up to exposures equivalent to clinical exposure at the recommended dose.

In a study conducted in rats, the number of corpora lutea and implantations were decreased in the presence of maternal toxicity. Otherwise, there were no effects on mating or fertility with darunavir treatment up to 1,000 mg/kg/day and exposure levels below (AUC-0.5 fold) of that in human at the clinically recommended dose. Up to same dose levels, there was no teratogenicity with darunavir in rats and rabbits when treated alone nor in mice when treated in combination with ritonavir. The exposure levels were lower than those with the recommended clinical dose in humans. In a pre- and postnatal development assessment in rats, darunavir with and without ritonavir, caused a transient reduction in body weight gain of the offspring pre-weaning and there was a slight delay in the opening of eyes and ears. Darunavir in combination with ritonavir caused a reduction in the number of pups that exhibited the startle response on day 15 of lactation and a reduced pup survival during lactation. These effects may be secondary to pup exposure to the active substance via the milk and/or maternal toxicity. No post weaning functions were affected with darunavir alone or in combination with ritonavir. In juvenile rats receiving darunavir up to days 23-26, increased mortality was observed with convulsions in some animals. Exposure in plasma, liver and brain was considerably higher than in adult rats after comparable doses in mg/kg between days 5 and 11 of age. After day 23 of life, the exposure was comparable to that in adult rats. The increased exposure was likely at least partly due to immaturity of the drug-metabolising enzymes in juvenile animals. No treatment related mortalities were noted in juvenile rats dosed at 1,000 mg/kg darunavir (single dose) on day 26 of age or at 500 mg/kg (repeated dose) from day 23 to 50 of age, and the exposures and toxicity profile were comparable to those observed in adult rats.

Due to uncertainties regarding the rate of development of the human blood brain barrier and liver enzymes, darunavir with low dose ritonavir should not be used in paediatric patients below 3 years of age.

Darunavir was evaluated for carcinogenic potential by oral gavage administration to mice and rats up to 104 weeks. Daily doses of 150, 450 and 1,000 mg/kg were administered to mice and doses of 50, 150 and 500 mg/kg were administered to rats. Dose-related increases in the incidences of hepatocellular adenomas and carcinomas were observed in males and females of both species. Thyroid follicular cell adenomas were noted in male rats. Administration of darunavir did not cause a statistically significant increase in the incidence of any other benign or malignant neoplasm in mice or rats. The observed hepatocellular and thyroid tumours in rodents are considered to be of limited relevance to humans. Repeated administration of darunavir to rats caused hepatic microsomal enzyme induction and increased thyroid hormone elimination, which predispose rats, but not humans, to thyroid neoplasms. At the highest tested doses, the systemic exposures (based on AUC) to darunavir were between 0.4- and 0.7-fold (mice) and 0.7- and 1-fold (rats), relative to those observed in humans at the recommended therapeutic doses.

After 2 years administration of darunavir at exposures at or below the human exposure, kidney changes were observed in mice (nephrosis) and rats (chronic progressive nephropathy).

Darunavir was not mutagenic or genotoxic in a battery of *in vitro* and *in vivo* assays including bacterial reverse mutation (Ames), chromosomal aberration in human lymphocytes and *in vivo* micronucleus test in mice.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Silica, colloidal anhydrous  
Cellulose microcrystalline  
Crospovidone  
Sodium starch glycolate  
Hypromellose  
Magnesium stearate

#### Tablet film-coat

Polyvinyl alcohol, partially hydrolysed  
Titanium dioxide (E171)  
Macrogol  
Talc

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years  
In-use shelf life after first opening HDPE bottle: 100 days

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

#### Darunavir Mylan 75 mg film-coated tablets

PVC/PE/PVDC-Al blister pack containing 480 tablets and 480x1 tablets.  
Cold form PVC/Al/OPA-Al blister pack containing 480 tablets and 480x1 tablets.  
HDPE bottle pack with a PP screw cap containing 480 tablets.

#### Darunavir Mylan 150 mg film-coated tablets

PVC/PE/PVDC-Al blister pack containing 240 tablets and 240x1 tablets.  
Cold form PVC/Al/OPA-Al blister pack containing 240 tablets and 240x1 tablets.  
HDPE bottle pack with a PP screw cap containing 60 and 240 tablets.

#### Darunavir Mylan 300 mg film-coated tablets

PVC/PE/PVDC-Al blister pack containing 30, 60 and 120 tablets and 120x1 tablets.  
Cold form PVC/Al/OPA-Al blister pack containing 30, 60 and 120 tablets and 120x1 tablets.  
HDPE bottle pack with a PP screw cap containing 30 and 120 tablets.

#### Darunavir Mylan 600 mg film-coated tablets

PVC/PE/PVDC-Al blister pack containing 30 and 60 tablets and 60x1 tablets.  
Cold form PVC/Al/OPA-Al blister pack containing 30 and 60 tablets and 60x1 tablets.

HDPE bottle pack with a PP screw cap containing 30, 60, 90 tablets.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

No special requirements for disposal.

## **7. MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

## **8. MARKETING AUTHORISATION NUMBER(S)**

### Darunavir Mylan 75 mg film-coated tablets

EU/1/16/1140/001  
EU/1/16/1140/002  
EU/1/16/1140/003  
EU/1/16/1140/004  
EU/1/16/1140/005

### Darunavir Mylan 150 mg film-coated tablets

EU/1/16/1140/006  
EU/1/16/1140/007  
EU/1/16/1140/008  
EU/1/16/1140/009  
EU/1/16/1140/010  
EU/1/16/1140/011

### Darunavir Mylan 300 mg film-coated tablets

EU/1/16/1140/012  
EU/1/16/1140/013  
EU/1/16/1140/014  
EU/1/16/1140/015  
EU/1/16/1140/016  
EU/1/16/1140/017  
EU/1/16/1140/018  
EU/1/16/1140/019  
EU/1/16/1140/020  
EU/1/16/1140/021

### Darunavir Mylan 600 mg film-coated tablets

EU/1/16/1140/030  
EU/1/16/1140/031  
EU/1/16/1140/032  
EU/1/16/1140/033

EU/1/16/1140/034  
EU/1/16/1140/035  
EU/1/16/1140/036  
EU/1/16/1140/037  
EU/1/16/1140/038

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 04 January 2017

Date of latest renewal: 16 September 2021

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.



## **1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 400 mg film-coated tablets  
Darunavir Mylan 800 mg film-coated tablets

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

### Darunavir Mylan 400 mg film-coated tablets

Each film-coated tablet contains 400 mg of darunavir.

### Darunavir Mylan 800 mg film-coated tablets

Each film-coated tablet contains 800 mg of darunavir.

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

### Darunavir Mylan 400 mg film-coated tablets

Film-coated tablet

White to off-white, oval shaped, biconvex film-coated tablets approximately 19.2 mm by 9.6 mm, debossed with 'M' on one side and 'DV4' on the other side.

### Darunavir Mylan 800 mg film-coated tablets

Film-coated tablet.

White to off-white, oval shaped, biconvex film-coated tablets approximately 21.2 mm by 10.6 mm, debossed with 'M' on one side and 'DV8' on the other side.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Darunavir co-administered with low dose ritonavir is indicated in combination with other antiretroviral medicinal products for the treatment of patients with human immunodeficiency virus (HIV-1) infection.

Darunavir co-administered with cobicistat is indicated in combination with other antiretroviral medicinal products for the treatment of human immunodeficiency virus (HIV-1) infection in adults and adolescents (aged 12 years and older, weighing at least 40 kg) (see section 4.2).

Darunavir Mylan 400 mg and 800 mg tablets may be used to provide suitable dose regimens for the treatment of HIV-1 infection in adult and paediatric patients from the age of 3 years and at least 40 kg body weight who are:

- antiretroviral therapy (ART)-naïve (see section 4.2).
- ART-experienced with no darunavir resistance associated mutations (DRV-RAMs) and who have plasma HIV-1 RNA < 100,000 copies/ml and CD4<sup>+</sup> cell count ≥ 100 cells x 10<sup>6</sup>/L. In deciding to initiate treatment with darunavir in such ART-experienced patients, genotypic testing should guide the use of darunavir (see sections 4.2, 4.3, 4.4 and 5.1).

## 4.2 Posology and method of administration

Therapy should be initiated by a healthcare provider experienced in the management of HIV infection. After therapy with darunavir has been initiated, patients should be advised not to alter the dosage, dose form or discontinue therapy without discussing with their healthcare provider.

The interaction profile of darunavir depends on whether ritonavir or cobicistat is used as pharmacokinetic enhancer. Darunavir may therefore have different contraindications and recommendations for concomitant medications depending on whether the compound is boosted with ritonavir or cobicistat (see sections 4.3, 4.4 and 4.5).

### Posology

Darunavir must always be given orally with cobicistat or low dose ritonavir as a pharmacokinetic enhancer and in combination with other antiretroviral medicinal products. The Summary of Product Characteristics of cobicistat or ritonavir as appropriate, must therefore be consulted prior to initiation of therapy with darunavir. Cobicistat is not indicated for use in twice daily regimens or for use in the paediatric population less than 12 years of age weighing less than 40 kg.

#### *ART-naïve adult patients*

The recommended dose regimen is 800 mg once daily taken with cobicistat 150 mg once daily or ritonavir 100 mg once daily taken with food. Darunavir Mylan 400 mg and 800 mg tablets can be used to construct the once daily 800 mg regimen.

#### *ART-experienced adult patients*

The recommended dose regimens are as follows:

- In ART-experienced patients with no darunavir resistance associated mutations (DRV-RAMs)\* and who have plasma HIV-1 RNA < 100,000 copies/ml and CD4+ cell count  $\geq 100$  cells  $\times 10^6/L$  (see section 4.1) a regimen of 800 mg once daily with cobicistat 150 mg once daily or ritonavir 100 mg once daily taken with food may be used. Darunavir 400 mg and 800 mg tablets can be used to construct the once daily 800 mg regimen.
- In all other ART-experienced patients or if HIV-1 genotype testing is not available, the recommended dose regimen is 600 mg twice daily taken with ritonavir 100 mg twice daily taken with food. See the Summary of Product Characteristics for Darunavir Mylan 75 mg, 150 mg, 300 mg or 600 mg tablets.

\* DRV-RAMs: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

#### *ART-naïve paediatric patients (3 to 17 years of age and weighing at least 40 kg)*

The recommended dose regimen is 800 mg once daily with ritonavir 100 mg once daily taken with food or 800 mg once daily with cobicistat 150 mg once daily taken with food (in adolescent patients 12 years of age or older). Darunavir Mylan 400 mg and 800 mg tablets can be used to construct the once daily 800 mg regimen. The dose of cobicistat to be used with darunavir in children less than 12 years of age has not been established.

#### *ART-experienced paediatric patients (3 to 17 years of age and weighing at least 40 kg)*

The dose of cobicistat to be used with darunavir in children less than 12 years of age has not been established.

The recommended dose regimens are as follows:

- In ART-experienced patients without DRV-RAMs\* and who have plasma HIV-1 RNA < 100,000 copies/ml and CD4+ cell count  $\geq 100$  cells  $\times 10^6/L$  (see section 4.1) a regimen of 800 mg once daily with ritonavir 100 mg once daily taken with food or 800 mg once daily with cobicistat 150 mg once daily taken with food (in adolescent patients 12 years of age or older) may be used. Darunavir Mylan 400 mg and 800 mg tablets can be used to construct the once daily 800 mg regimen. The dose of cobicistat to be used with darunavir in children less than 12 years of age has not been established.

- In all other ART-experienced patients or if HIV-1 genotype testing is not available, the recommended dose regimen described in the Summary of Product Characteristics for Darunavir Mylan 75 mg, 150 mg, 300 mg and 600 mg tablets.

\* DRV-RAMs: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

#### *Advice on missed doses*

If a once daily dose of darunavir and/or cobicistat or ritonavir is missed within 12 hours of the time it is usually taken, patients should be instructed to take the prescribed dose of darunavir and cobicistat or ritonavir with food as soon as possible. If this is noticed later than 12 hours after the time it is usually taken, the missed dose should not be taken and the patient should resume the usual dosing schedule.

This guidance is based on the half-life of darunavir in the presence of cobicistat or ritonavir and the recommended dosing interval of approximately 24 hours.

If a patient vomits within 4 hours of taking the medicine, another dose of Darunavir Mylan with cobicistat or ritonavir should be taken with food as soon as possible. If a patient vomits more than 4 hours after taking the medicine, the patient does not need to take another dose of Darunavir Mylan with cobicistat or ritonavir until the next regularly scheduled time.

#### Special populations

##### *Elderly*

Limited information is available in this population, and therefore, darunavir should be used with caution in this age group (see sections 4.4 and 5.2).

##### *Hepatic impairment*

Darunavir is metabolised by the hepatic system. No dose adjustment is recommended in patients with mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment, however, darunavir should be used with caution in these patients. No pharmacokinetic data are available in patients with severe hepatic impairment. Severe hepatic impairment could result in an increase of darunavir exposure and a worsening of its safety profile. Therefore, darunavir must not be used in patients with severe hepatic impairment (Child-Pugh Class C) (see sections 4.3, 4.4 and 5.2).

##### *Renal impairment*

No dose adjustment is required for darunavir/ritonavir in patients with renal impairment (see sections 4.4 and 5.2). Cobicistat has not been studied in patients receiving dialysis, and, therefore, no recommendation can be made for the use of darunavir/cobicistat in these patients.

Cobicistat inhibits the tubular secretion of creatinine and may cause modest increases in serum creatinine and modest declines in creatinine clearance. Hence, the use of creatinine clearance as an estimate of renal elimination capacity may be misleading. Cobicistat as a pharmacokinetic enhancer of darunavir should, therefore, not be initiated in patients with creatine clearance less than 70 ml/min if any co-administered agent requires dose adjustment based on creatinine clearance: e.g. emtricitabine, lamivudine, tenofovir disoproxil (as fumarate, phosphate or succinate) or adefovir dipivoxil. For information on cobicistat, consult the cobicistat Summary of Product Characteristics.

##### *Paediatric population*

Darunavir should not be used in children

- below 3 years of age, because of safety concerns (see sections 4.4 and 5.3), or,
- less than 15 kg body weight, as the dose for this population has not been established in a sufficient number of patients (see section 5.1).

Darunavir taken with cobicistat should not be used in children aged 3 to 11 years of age weighing < 40 kg as the dose of cobicistat to be used in these children has not been established (see sections 4.4 and 5.3).

Darunavir Mylan 400 mg and 800 mg tablets are not suitable for this patient population. Other formulations are available, see the Summary of Product Characteristics for Darunavir Mylan 75 mg, 150 mg, 300 mg, 600 mg tablets.

#### *Pregnancy and postpartum*

No dose adjustment is required for darunavir/ritonavir during pregnancy and postpartum.

Darunavir/ritonavir should be used during pregnancy only if the potential benefit justifies the potential risk (see sections 4.4, 4.6 and 5.2).

Treatment with darunavir/cobicistat 800/150 mg during pregnancy results in low darunavir exposure (see sections 4.4 and 5.2). Therefore, therapy with Darunavir/cobicistat should not be initiated during pregnancy, and women who become pregnant during therapy with Darunavir/cobicistat should be switched to an alternative regimen (see sections 4.4 and 4.6). Darunavir/ritonavir may be considered as an alternative.

#### Method of administration

Patients should be instructed to take Darunavir Mylan with cobicistat or low dose ritonavir within 30 minutes after completion of a meal. The type of food does not affect the exposure to darunavir (see sections 4.4, 4.5 and 5.2).

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Patients with severe (Child-Pugh Class C) hepatic impairment.

Concomitant treatment with any of the following medicinal products given the expected decrease in plasma concentrations of darunavir, ritonavir and cobicistat and the potential for loss of therapeutic effect (see sections 4.4 and 4.5).

Applicable to darunavir boosted with either ritonavir or cobicistat:

- The combination product lopinavir/ritonavir (see section 4.5).
- The strong CYP3A inducers rifampicin and herbal preparations containing St John's wort (*Hypericum perforatum*). Co-administration is expected to reduce plasma concentrations of darunavir, ritonavir and cobicistat, which could lead to loss of therapeutic effect and possible development of resistance (see sections 4.4 and 4.5).

Applicable to darunavir boosted with cobicistat, not when boosted with ritonavir:

- Darunavir boosted with cobicistat is more sensitive for CYP3A induction than darunavir boosted with ritonavir. Concomitant use with strong CYP3A inducers is contraindicated, since these may reduce the exposure to cobicistat and darunavir leading to loss of therapeutic effect. Strong CYP3A inducers include e.g. carbamazepine, phenobarbital and phenytoin (see sections 4.4 and 4.5).

Darunavir boosted with either ritonavir or cobicistat inhibits the elimination of active substances that are highly dependent on CYP3A for clearance, which results in increased exposure to the co-administered medicinal product. Therefore, concomitant treatment with such medicinal products for which elevated plasma concentrations are associated with serious and/or life-threatening events is contraindicated (applies to darunavir boosted with either ritonavir or cobicistat). These active substances include e.g.:

- alfuzosin
- amiodarone, bepridil, dronedarone, ivabradine, quinidine, ranolazine
- astemizole, terfenadine
- colchicine when used in patients with renal and/or hepatic impairment (see section 4.5)
- ergot derivatives (e.g. dihydroergotamine, ergometrine, ergotamine, methylergonovine)
- elbasvir/grazoprevir

- cisapride
- dapoxetine
- domperidone
- naloxegol
- lurasidone, pimozide, quetiapine, sertindole (see section 4.5)
- triazolam, midazolam administered orally (for caution on parenterally administered midazolam, see section 4.5)
- sildenafil - when used for the treatment of pulmonary arterial hypertension, avanafil
- simvastatin, lovastatin and lomitapide (see section 4.5)
- ticagrelor (see section 4.5).

#### **4.4 Special warnings and precautions for use**

Regular assessment of virological response is advised. In the setting of lack or loss of virological response, resistance testing should be performed.

Darunavir 400 mg or 800 mg must always be given orally with cobicistat or low dose ritonavir as a pharmacokinetic enhancer and in combination with other antiretroviral medicinal products (see section 5.2). The Summary of Product Characteristics of cobicistat or ritonavir as appropriate, must therefore be consulted prior to initiation of therapy with darunavir.

Increasing the dose of ritonavir from that recommended in section 4.2 did not significantly affect darunavir concentrations. It is not recommended to alter the dose of cobicistat or ritonavir. Darunavir binds predominantly to  $\alpha_1$ -acid glycoprotein. This protein binding is concentration-dependent indicative for saturation of binding. Therefore, protein displacement of medicinal products highly bound to  $\alpha_1$ -acid glycoprotein cannot be ruled out (see section 4.5).

##### ART-experienced patients – once daily dosing

Darunavir used in combination with cobicistat or low dose ritonavir once daily in ART-experienced patients should not be used in patients with one or more darunavir resistance associated mutations (DRV-RAMs) or HIV-1 RNA  $\geq 100,000$  copies/ml or CD4+ cell count  $< 100$  cells  $\times 10^6/L$  (see section 4.2). Combinations with optimised background regimen (OBRs) other than  $\geq 2$  NRTIs have not been studied in this population. Limited data are available in patients with HIV-1 clades other than B (see section 5.1).

##### Paediatric population

Darunavir is not recommended for use in paediatric patients below 3 years of age or less than 15 kg body weight (see sections 4.2 and 5.3).

##### Pregnancy

Darunavir/ritonavir should be used during pregnancy only if the potential benefit justifies the potential risk. Caution should be used in pregnant women with concomitant medications which may further decrease darunavir exposure (see sections 4.5 and 5.2).

Treatment with darunavir/cobicistat 800/150 mg once daily during the second and third trimester has been shown to result in low darunavir exposure, with a reduction of around 90% in  $C_{min}$  levels (see section 5.2). Cobicistat levels decrease and may not provide sufficient boosting. The substantial reduction in darunavir exposure may result in virological failure and an increased risk of mother to child transmission of HIV infection. Therefore, therapy with Darunavir/cobicistat should not be initiated during pregnancy, and women who become pregnant during therapy with Darunavir/cobicistat should be switched to an alternative regimen (see sections 4.2 and 4.6). Darunavir given with low dose ritonavir may be considered as an alternative.

## Elderly

As limited information is available on the use of darunavir in patients aged 65 and over, caution should be exercised in the administration of darunavir in elderly patients, reflecting the greater frequency of decreased hepatic function and of concomitant disease or other therapy (see sections 4.2 and 5.2).

## Severe skin reactions

During the darunavir/ritonavir clinical development program (N=3,063), severe skin reactions, which may be accompanied with fever and/or elevations of transaminases, have been reported in 0.4% of patients. DRESS (Drug Rash with Eosinophilia and Systemic Symptoms) and Stevens-Johnson syndrome has been rarely (< 0.1%) reported, and during post-marketing experience toxic epidermal necrolysis and acute generalised exanthematous pustulosis have been reported. Darunavir should be discontinued immediately if signs or symptoms of severe skin reactions develop. These can include, but are not limited to, severe rash or rash accompanied by fever, general malaise, fatigue, muscle or joint aches, blisters, oral lesions, conjunctivitis, hepatitis and/or eosinophilia.

Rash occurred more commonly in treatment-experienced patients receiving regimens containing darunavir/ritonavir + raltegravir compared to patients receiving darunavir/ritonavir without raltegravir or raltegravir without darunavir (see section 4.8).

Darunavir contains a sulphonamide moiety. Darunavir should be used with caution in patients with a known sulphonamide allergy.

## Hepatotoxicity

Drug-induced hepatitis (e.g. acute hepatitis, cytolytic hepatitis) has been reported with darunavir. During the darunavir/ritonavir clinical development program (N=3,063), hepatitis was reported in 0.5% of patients receiving combination antiretroviral therapy with darunavir/ritonavir. Patients with pre-existing liver dysfunction, including chronic active hepatitis B or C, have an increased risk for liver function abnormalities including severe and potentially fatal hepatic adverse reactions. In case of concomitant antiviral therapy for hepatitis B or C, please refer to the relevant product information for these medicinal products.

Appropriate laboratory testing should be conducted prior to initiating therapy with darunavir used in combination with cobicistat or low dose ritonavir and patients should be monitored during treatment. Increased AST/ALT monitoring should be considered in patients with underlying chronic hepatitis, cirrhosis, or in patients who have pre-treatment elevations of transaminases, especially during the first several months of darunavir used in combination with cobicistat or low dose ritonavir treatment.

If there is evidence of new or worsening liver dysfunction (including clinically significant elevation of liver enzymes and/or symptoms such as fatigue, anorexia, nausea, jaundice, dark urine, liver tenderness, hepatomegaly) in patients using darunavir used in combination with cobicistat or low dose ritonavir, interruption or discontinuation of treatment should be considered promptly.

## Patients with coexisting conditions

### *Hepatic impairment*

The safety and efficacy of darunavir have not been established in patients with severe underlying liver disorders and darunavir is therefore contraindicated in patients with severe hepatic impairment. Due to an increase in the unbound darunavir plasma concentrations, darunavir should be used with caution in patients with mild or moderate hepatic impairment (see sections 4.2, 4.3 and 5.2).

### *Renal impairment*

No special precautions or dose adjustments for darunavir/ritonavir are required in patients with renal impairment. As darunavir and ritonavir are highly bound to plasma proteins, it is unlikely that they

will be significantly removed by haemodialysis or peritoneal dialysis. Therefore, no special precautions or dose adjustments are required in these patients (see sections 4.2 and 5.2). Cobicistat has not been studied in patients receiving dialysis, therefore, no recommendation can be made for the use of darunavir/cobicistat in these patients (see section 4.2).

Cobicistat decreases the estimated creatinine clearance due to inhibition of tubular secretion of creatinine. This should be taken into consideration if darunavir with cobicistat is administered to patients in whom the estimated creatinine clearance is used to adjust doses of co-administered medicinal products (see section 4.2 and cobicistat SmPC).

There are currently inadequate data to determine whether co-administration of tenofovir disoproxil and cobicistat is associated with a greater risk of renal adverse reactions compared with regimens that include tenofovir disoproxil without cobicistat.

#### *Haemophilic patients*

There have been reports of increased bleeding, including spontaneous skin haematomas and haemarthrosis in patients with haemophilia type A and B treated with PIs. In some patients additional factor VIII was given. In more than half of the reported cases, treatment with PIs was continued or reintroduced if treatment had been discontinued. A causal relationship has been suggested, although the mechanism of action has not been elucidated. Haemophilic patients should, therefore, be made aware of the possibility of increased bleeding.

#### *Weight and metabolic parameters*

An increase in weight and in levels of blood lipids and glucose may occur during antiretroviral therapy. Such changes may in part be linked to disease control and life style. For lipids, there is in some cases evidence for a treatment effect, while for weight gain there is no strong evidence relating this to any particular treatment. For monitoring of blood lipids and glucose reference is made to established HIV treatment guidelines. Lipid disorders should be managed as clinically appropriate.

#### Osteonecrosis

Although the aetiology is considered to be multifactorial (including corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index), cases of osteonecrosis have been reported particularly in patients with advanced HIV disease and/or long-term exposure to combination antiretroviral therapy (CART). Patients should be advised to seek medical advice if they experience joint aches and pain, joint stiffness or difficulty in movement.

#### Immune reconstitution inflammatory syndrome

In HIV infected patients with severe immune deficiency at the time of initiation of combination antiretroviral therapy (CART), an inflammatory reaction to asymptomatic or residual opportunistic pathogens may arise and cause serious clinical conditions, or aggravation of symptoms. Typically, such reactions have been observed within the first weeks or months of initiation of CART. Relevant examples are cytomegalovirus retinitis, generalised and/or focal mycobacterial infections and pneumonia caused by *Pneumocystis jirovecii* (formerly known as *Pneumocystis carinii*). Any inflammatory symptoms should be evaluated and treatment instituted when necessary. In addition, reactivation of herpes simplex and herpes zoster has been observed in clinical studies with darunavir co-administered with low dose ritonavir.

Autoimmune disorders (such as Graves' disease and autoimmune hepatitis) have also been reported to occur in the setting of immune reactivation; however, the reported time to onset is more variable and these events can occur many months after initiation of treatment (see section 4.8).

#### Interactions with medicinal products

Several of the interaction studies have been performed with darunavir at lower than recommended doses. The effects on co-administered medicinal products may thus be underestimated and clinical

monitoring of safety may be indicated. For full information on interactions with other medicinal products see section 4.5.

#### *Pharmacokinetic enhancer and concomitant medications*

Darunavir has different interaction profiles depending on whether the compound is boosted with ritonavir or cobicistat:

- Darunavir boosted with cobicistat is more sensitive for CYP3A induction: concomitant use of darunavir/cobicistat and strong CYP3A inducers is therefore contraindicated (see section 4.3), and concomitant use with weak to moderate CYP3A inducers is not recommended (see section 4.5). Concomitant use of darunavir/ritonavir and darunavir/cobicistat with lopinavir/ritonavir, rifampicin and herbal products containing St John's wort, *Hypericum perforatum*, is contraindicated (see section 4.5).
- Unlike ritonavir, cobicistat does not have inducing effects on enzymes or transport proteins (see section 4.5). If switching the pharmacoenhancer from ritonavir to cobicistat, caution is required during the first two weeks of treatment with darunavir/cobicistat, particularly if doses of any concomitantly administered medicinal products have been titrated or adjusted during use of ritonavir as a pharmacoenhancer. A dose reduction of the co-administered drug may be needed in these cases.

Efavirenz in combination with boosted darunavir may result in sub-optimal darunavir  $C_{min}$ . If efavirenz is to be used in combination with darunavir, the darunavir/ritonavir 600/100 mg twice daily regimen should be used. See the Summary of Product Characteristics for Darunavir Mylan 75 mg, 150 mg, 300 mg and 600 mg tablets (see section 4.5).

Life-threatening and fatal drug interactions have been reported in patients treated with colchicine and strong inhibitors of CYP3A and P-glycoprotein (P-gp; see sections 4.3 and 4.5).

#### Darunavir Mylan contains sodium

Darunavir Mylan 400 mg and 800 mg film-coated tablets contain less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

### **4.5 Interaction with other medicinal products and other forms of interaction**

The interaction profile of darunavir may differ depending on whether ritonavir or cobicistat is used as pharmacoenhancer. The recommendations given for concomitant use of darunavir and other medicinal products may therefore differ depending on whether darunavir is boosted with ritonavir or cobicistat (see sections 4.3 and 4.4), and caution is also required during the first time of treatment if switching the pharmacoenhancer from ritonavir to cobicistat (see section 4.4).

#### Medicinal products that affect darunavir exposure (ritonavir as pharmacoenhancer)

Darunavir and ritonavir are metabolised by CYP3A. Medicinal products that induce CYP3A activity would be expected to increase the clearance of darunavir and ritonavir, resulting in lowered plasma concentrations of these compounds and consequently that of darunavir, leading to loss of therapeutic effect and possible development of resistance (see sections 4.3 and 4.4). CYP3A inducers that are contraindicated include rifampicin, St John's wort and lopinavir.

Co-administration of darunavir and ritonavir with other medicinal products that inhibit CYP3A may decrease the clearance of darunavir and ritonavir, which may result in increased plasma concentrations of darunavir and ritonavir. Co-administration with strong CYP3A4 inhibitors is not recommended and caution is warranted, these interactions are described in the interaction table below (e.g. indinavir, azole antifungals like clotrimazole).



### Medicinal products that affect darunavir exposure (cobicistat as pharmacoenhancer)

Darunavir and cobicistat are metabolised by CYP3A, and co-administration with CYP3A inducers may therefore result in subtherapeutic plasma exposure to darunavir. Darunavir boosted with cobicistat is more sensitive to CYP3A induction than ritonavir-boosted darunavir: co-administration of darunavir/cobicistat with medicinal products that are strong inducers of CYP3A (e.g. St John's wort, rifampicin, carbamazepine, phenobarbital, and phenytoin) is contraindicated (see section 4.3). Co-administration of darunavir/cobicistat with weak to moderate CYP3A inducers (e.g. efavirenz, etravirine, nevirapine, fluticasone, and bosentan) is not recommended (see interaction table below).

For co-administration with strong CYP3A4 inhibitors, the same recommendations apply independent of whether darunavir is boosted with ritonavir or with cobicistat (see section above).

### Medicinal products that may be affected by darunavir boosted with ritonavir

Darunavir and ritonavir are inhibitors of CYP3A, CYP2D6 and P-gp. Co-administration of darunavir/ritonavir with medicinal products primarily metabolised by CYP3A and/or CYP2D6 or transported by P-gp may result in increased systemic exposure to such medicinal products, which could increase or prolong their therapeutic effect and adverse reactions.

Darunavir co-administered with low dose ritonavir must not be combined with medicinal products that are highly dependent on CYP3A for clearance and for which increased systemic exposure is associated with serious and/or life-threatening events (narrow therapeutic index) (see section 4.3).

Co-administration of boosted darunavir with drugs that have active metabolite(s) formed by CYP3A may result in reduced plasma concentrations of these active metabolite(s), potentially leading to loss of their therapeutic effect (see the Interaction table below).

The overall pharmacokinetic enhancement effect by ritonavir was an approximate 14-fold increase in the systemic exposure of darunavir when a single dose of 600 mg darunavir was given orally in combination with ritonavir at 100 mg twice daily. Therefore, darunavir must only be used in combination with a pharmacokinetic enhancer (see sections 4.4 and 5.2).

A clinical study utilising a cocktail of medicinal products that are metabolised by cytochromes CYP2C9, CYP2C19 and CYP2D6 demonstrated an increase in CYP2C9 and CYP2C19 activity and inhibition of CYP2D6 activity in the presence of darunavir/ritonavir, which may be attributed to the presence of low dose ritonavir. Co-administration of darunavir and ritonavir with medicinal products which are primarily metabolised by CYP2D6 (such as flecainide, propafenone, metoprolol) may result in increased plasma concentrations of these medicinal products, which could increase or prolong their therapeutic effect and adverse reactions. Co-administration of darunavir and ritonavir with medicinal products primarily metabolised by CYP2C9 (such as warfarin) and CYP2C19 (such as methadone) may result in decreased systemic exposure to such medicinal products, which could decrease or shorten their therapeutic effect.

Although the effect on CYP2C8 has only been studied *in vitro*, co-administration of darunavir and ritonavir and medicinal products primarily metabolised by CYP2C8 (such as paclitaxel, rosiglitazone, repaglinide) may result in decreased systemic exposure to such medicinal products, which could decrease or shorten their therapeutic effect.

Ritonavir inhibits the transporters P-glycoprotein, OATP1B1 and OATP1B3, and co-administration with substrates of these transporters can result in increased plasma concentrations of these compounds (e.g. dabigatran etexilate, digoxin, statins and bosentan; see the Interaction table below).

### Medicinal products that may be affected by darunavir boosted with cobicistat

The recommendations for darunavir boosted with ritonavir are similar to the recommendation for darunavir boosted with cobicistat with regard to substrates of CYP3A4, CYP2D6, P-glycoprotein,

OATP1B1 and OATP1B3 (see contraindications and recommendations presented in the section above). Cobicistat 150 mg given with darunavir 800 mg once daily enhances darunavir pharmacokinetic parameters in a comparable way to ritonavir (see section 5.2).

Unlike ritonavir, cobicistat does not induce CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19 or UGT1A1. For further information on cobicistat, consult the cobicistat Summary of Product Characteristics.

### Interaction table

Interaction studies have only been performed in adults.

Several of the interaction studies (indicated by <sup>#</sup> in the table below) have been performed at lower than recommended doses of darunavir or with a different dosing regimen (see section 4.2 Posology). The effects on co-administered medicinal products may thus be underestimated and clinical monitoring of safety may be indicated.

The interaction profile of darunavir depends on whether ritonavir or cobicistat is used as pharmacokinetic enhancer. Darunavir may therefore have different recommendations for concomitant medications depending on whether the compound is boosted with ritonavir or cobicistat. No interaction studies presented in the table have been performed with darunavir boosted with cobicistat. The same recommendations apply, unless specifically indicated. For further information on cobicistat, consult the cobicistat Summary of Product Characteristics.

Interactions between darunavir/ritonavir and antiretroviral and non-antiretroviral medicinal products are listed in the table below. The direction of the arrow for each pharmacokinetic parameter is based on the 90% confidence interval of the geometric mean ratio being within ( $\leftrightarrow$ ), below ( $\downarrow$ ) or above ( $\uparrow$ ) the 80-125% range (not determined as “ND”).

In the table below the specific pharmacokinetic enhancer is specified when recommendations differ. When the recommendation is the same for darunavir when co-administered with a low dose ritonavir or cobicistat, the term “boosted darunavir” is used.

The below list of examples of drug -drug interactions is not comprehensive and therefore the label of each drug that is co-administered with darunavir should be consulted for information related to the route of metabolism, interaction pathways, potential risks, and specific actions to be taken with regards to co-administration.

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
<b>HIV ANTIRETROVIRALS</b>		
<b><i>Integrase strand transfer inhibitors</i></b>		
Dolutegravir	dolutegravir AUC $\downarrow$ 22% dolutegravir C <sub>24h</sub> $\downarrow$ 38% dolutegravir C <sub>max</sub> $\downarrow$ 11% darunavir $\leftrightarrow$ * * Using cross-study comparisons to historical pharmacokinetic data	Boosted darunavir and dolutegravir can be used without dose adjustment.
Raltegravir	Some clinical studies suggest raltegravir may cause a modest decrease in darunavir plasma concentrations.	At present the effect of raltegravir on darunavir plasma concentrations does not appear to be clinically relevant. Boosted darunavir and raltegravir can be used without dose adjustments.

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
<i>Nucleo(s)ide reverse transcriptase inhibitors (NRTIs)</i>		
Didanosine 400 mg once daily	didanosine AUC ↓ 9% didanosine C <sub>min</sub> ND didanosine C <sub>max</sub> ↓ 16% darunavir AUC ↔ darunavir C <sub>min</sub> ↔ darunavir C <sub>max</sub> ↔	Boosted darunavir and didanosine can be used without dose adjustments. Didanosine is to be administered on an empty stomach, thus it should be administered 1 hour before or 2 hours after boosted darunavir given with food.
Tenofovir disoproxil 245 mg once daily <sup>‡</sup>	tenofovir AUC ↑ 22% tenofovir C <sub>min</sub> ↑ 37% tenofovir C <sub>max</sub> ↑ 24% #darunavir AUC ↑ 21% #darunavir C <sub>min</sub> ↑ 24% #darunavir C <sub>max</sub> ↑ 16% (↑ tenofovir from effect on MDR-1 transport in the renal tubules)	Monitoring of renal function may be indicated when boosted darunavir is given in combination with tenofovir disoproxil, particularly in patients with underlying systemic or renal disease, or in patients taking nephrotoxic agents.  Darunavir co-administered with cobicistat lowers the creatinine clearance. Refer to section 4.4 if creatinine clearance is used for dose adjustment of tenofovir disoproxil.
Emtricitabine/tenofovir alafenamide	Tenofovir alafenamide ↔ Tenofovir ↑	The recommended dose of emtricitabine/tenofovir alafenamide is 200/10 mg once daily when used with boosted darunavir.
Abacavir Emtricitabine Lamivudine Stavudine Zidovudine	Not studied. Based on the different elimination pathways of the other NRTIs zidovudine, emtricitabine, stavudine, lamivudine, that are primarily renally excreted, and abacavir for which metabolism is not mediated by CYP450, no interactions are expected for these medicinal compounds and boosted darunavir.	Boosted darunavir can be used with these NRTIs without dose adjustment.  Darunavir co-administered with cobicistat lowers the creatinine clearance. Refer to section 4.4 if creatinine clearance is used for dose adjustment of emtricitabine or lamivudine.

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
<b><i>Non-nucleo(s)tide reverse transcriptase inhibitors (NNRTIs)</i></b>		
Efavirenz 600 mg once daily	efavirenz AUC ↑ 21% efavirenz C <sub>min</sub> ↑ 17% efavirenz C <sub>max</sub> ↑ 15% #darunavir AUC ↓ 13% #darunavir C <sub>min</sub> ↓ 31% #darunavir C <sub>max</sub> ↓ 15% (↑ efavirenz from CYP3A inhibition) (↓ darunavir from CYP3A induction)	Clinical monitoring for central nervous system toxicity associated with increased exposure to efavirenz may be indicated when darunavir co-administered with low dose ritonavir is given in combination with efavirenz.  Efavirenz in combination with darunavir/ritonavir 800/100 mg once daily may result in sub-optimal darunavir C <sub>min</sub> . If efavirenz is to be used in combination with darunavir/ritonavir, the darunavir/ritonavir 600/100 mg twice daily regimen should be used (see section 4.4).  Co-administration with darunavir co-administered with cobicistat is not recommended (see section 4.4).
Etravirine 100 mg twice daily	etravirine AUC ↓ 37% etravirine C <sub>min</sub> ↓ 49% etravirine C <sub>max</sub> ↓ 32% darunavir AUC ↑ 15% darunavir C <sub>min</sub> ↔ darunavir C <sub>max</sub> ↔	Darunavir co-administered with low dose ritonavir and etravirine <b>200 mg twice daily</b> can be used without dose adjustments.  Co-administration with darunavir co-administered with cobicistat is not recommended (see section 4.4).
Nevirapine 200 mg twice daily	nevirapine AUC ↑ 27% nevirapine C <sub>min</sub> ↑ 47% nevirapine C <sub>max</sub> ↑ 18% #darunavir: concentrations were consistent with historical data (↑ nevirapine from CYP3A inhibition)	Darunavir co-administered with low dose ritonavir and nevirapine can be used without dose adjustments.  Co-administration with darunavir co-administered with cobicistat is not recommended (see section 4.4).
Rilpivirine 150 mg once daily	rilpivirine AUC ↑ 130% rilpivirine C <sub>min</sub> ↑ 178% rilpivirine C <sub>max</sub> ↑ 79% darunavir AUC ↔ darunavir C <sub>min</sub> ↓ 11% darunavir C <sub>max</sub> ↔	Boosted darunavir and rilpivirine can be used without dose adjustments.

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
<b><i>HIV Protease inhibitors (PIs) - without additional co-administration of low dose ritonavir†</i></b>		
Atazanavir 300 mg once daily	<p>atazanavir AUC ↔            atazanavir C<sub>min</sub> ↑ 52%            atazanavir C<sub>max</sub> ↓ 11%            #darunavir AUC ↔            #darunavir C<sub>min</sub> ↔            #darunavir C<sub>max</sub> ↔</p> <p>Atazanavir: comparison of atazanavir/ritonavir 300/100 mg once daily vs. atazanavir 300 mg once daily in combination with darunavir/ritonavir 400/100 mg twice daily.            Darunavir: comparison of darunavir/ritonavir 400/100 mg twice daily vs. darunavir/ritonavir 400/100 mg twice daily in combination with atazanavir 300 mg once daily.</p>	<p>Darunavir co-administered with low dose ritonavir and atazanavir can be used without dose adjustments.</p> <p>Darunavir co-administered with cobicistat should not be used in combination with another antiretroviral agent that requires pharmacoenhancement by means of co- administration with an inhibitor of CYP3A4 (see section 4.5).</p>
Indinavir 800 mg twice daily	<p>indinavir AUC ↑ 23%            indinavir C<sub>min</sub> ↑ 125%            indinavir C<sub>max</sub> ↔            #darunavir AUC ↑ 24%            #darunavir C<sub>min</sub> ↑ 44%            #darunavir C<sub>max</sub> ↑ 11%</p> <p>Indinavir: comparison of indinavir/ritonavir 800/100 mg twice daily vs. indinavir/darunavir/ritonavir 800/400/100 mg twice daily.            Darunavir: comparison of darunavir/ritonavir 400/100 mg twice daily vs. darunavir/ritonavir 400/100 mg in combination with indinavir 800 mg twice daily.</p>	<p>When used in combination with darunavir co-administered with low dose ritonavir, dose adjustment of indinavir from 800 mg twice daily to 600 mg twice daily may be warranted in case of intolerance.</p> <p>Darunavir co-administered with cobicistat should not be used in combination with another antiretroviral agent that requires pharmacoenhancement by means of co- administration with an inhibitor of CYP3A4 (see section 4.5).</p>
Saquinavir 1,000 mg twice daily	<p>#darunavir AUC ↓ 26%            #darunavir C<sub>min</sub> ↓ 42%            #darunavir C<sub>max</sub> ↓ 17%            saquinavir AUC ↓ 6%            saquinavir C<sub>min</sub> ↓ 18%            saquinavir C<sub>max</sub> ↓ 6%</p> <p>Saquinavir: comparison of saquinavir/ritonavir 1,000/100 mg twice daily vs. saquinavir/darunavir/ritonavir 1,000/400/100 mg twice daily            Darunavir: comparison of darunavir/ritonavir 400/100 mg twice daily vs. darunavir/ritonavir 400/100 mg in combination with saquinavir 1,000 mg twice daily.</p>	<p>It is not recommended to combine darunavir co-administered with low dose ritonavir with saquinavir.</p> <p>Darunavir co-administered with cobicistat should not be used in combination with another antiretroviral agent that requires pharmacoenhancement by means of co- administration with an inhibitor of CYP3A4 (see section 4.5).</p>

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
<b>HIV Protease inhibitors (PIs) - with co-administration of low dose ritonavir<sup>‡</sup></b>		
Lopinavir/ritonavir 400/100 mg twice daily	lopinavir AUC ↑ 9% lopinavir C <sub>min</sub> ↑ 23% lopinavir C <sub>max</sub> ↓ 2% darunavir AUC ↓ 38% <sup>‡</sup> darunavir C <sub>min</sub> ↓ 51% <sup>‡</sup> darunavir C <sub>max</sub> ↓ 21% <sup>‡</sup>	Due to a decrease in the exposure (AUC) of darunavir by 40%, appropriate doses of the combination have not been established. Hence, concomitant use of boosted darunavir and the combination product lopinavir/ritonavir is contraindicated (see section 4.3).
Lopinavir/ritonavir 533/133.3 mg twice daily	lopinavir AUC ↔ lopinavir C <sub>min</sub> ↑ 13% lopinavir C <sub>max</sub> ↑ 11% darunavir AUC ↓ 41% darunavir C <sub>min</sub> ↓ 55% darunavir C <sub>max</sub> ↓ 21% <sup>‡</sup> based upon non dose normalised values	
<b>CCR5 ANTAGONIST</b>		
Maraviroc 150 mg twice daily	maraviroc AUC ↑ 305% maraviroc C <sub>min</sub> ND maraviroc C <sub>max</sub> ↑ 129% darunavir, ritonavir concentrations were consistent with historical data	The maraviroc dose should be 150 mg twice daily when co-administered with boosted darunavir.
<b>α1-ADRENORECEPTOR ANTAGONIST</b>		
Alfuzosin	Based on theoretical considerations darunavir is expected to increase alfuzosin plasma concentrations. (CYP3A inhibition)	Co-administration of boosted darunavir and alfuzosin is contraindicated (see section 4.3).
<b>ANAESTHETIC</b>		
Alfentanil	Not studied. The metabolism of alfentanil is mediated via CYP3A, and may as such be inhibited by boosted darunavir.	The concomitant use with boosted darunavir may require to lower the dose of alfentanil and requires monitoring for risks of prolonged or delayed respiratory depression.
<b>ANTIANGINA/ANTIARRHYTHMIC</b>		
Disopyramide Flecainide Lidocaine (systemic) Mexiletine Propafenone	Not studied. Boosted darunavir is expected to increase these antiarrhythmic plasma concentrations. (CYP3A and/or CYP2D6 inhibition)	Caution is warranted and therapeutic concentration monitoring, if available, is recommended for these antiarrhythmics when co-administered with boosted darunavir.
Amiodarone Bepidil Dronedarone Ivabradine Quinidine Ranolazine		Co-administration of boosted darunavir and amiodarone, bepridil, dronedarone, ivabradine, quinidine, or ranolazine is contraindicated (see section 4.3).
Digoxin 0.4 mg single dose	digoxin AUC ↑ 61% digoxin C <sub>min</sub> ND digoxin C <sub>max</sub> ↑ 29% (↑ digoxin from probable inhibition of P-gp)	Given that digoxin has a narrow therapeutic index, it is recommended that the lowest possible dose of digoxin should initially be prescribed in case digoxin is given to patients on boosted darunavir therapy. The digoxin dose should be carefully titrated to obtain the desired clinical effect while assessing the overall clinical state of the subject.

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
<b>ANTIBIOTIC</b>		
Clarithromycin 500 mg twice daily	clarithromycin AUC ↑ 57% clarithromycin C <sub>min</sub> ↑ 174% clarithromycin C <sub>max</sub> ↑ 26% #darunavir AUC ↓ 13% #darunavir C <sub>min</sub> ↑ 1% #darunavir C <sub>max</sub> ↓ 17% 14-OH-clarithromycin concentrations were not detectable when combined with darunavir/ritonavir. (↑ clarithromycin from CYP3A inhibition and possible P-gp inhibition)	Caution should be exercised when clarithromycin is combined with boosted darunavir.  For patients with renal impairment the Summary of Product Characteristics for clarithromycin should be consulted for the recommended dose.
<b>ANTICOAGULANT/PLATELET AGGREGATION INHIBITOR</b>		
Apixaban Rivaroxaban	Not studied. Co-administration of boosted darunavir with these anticoagulants may increase concentrations of the anticoagulant. (CYP3A and/or P-gp inhibition)	The use of boosted darunavir with a direct oral anticoagulant (DOAC) that is metabolised by CYP3A4 and transported by P-gp is not recommended as this may lead to an increased bleeding risk.
Dabigatran etexilate Edoxaban	dabigatran etexilate (150 mg): darunavir/ritonavir 800/100 mg single dose: dabigatran AUC ↑ 72% dabigatran C <sub>max</sub> ↑ 64%  darunavir/ritonavir 800/100 mg once daily: dabigatran AUC ↑ 18% dabigatran C <sub>max</sub> ↑ 22%  darunavir/cobicistat 800/150 mg single dose: dabigatran AUC ↑ 164% dabigatran C <sub>max</sub> ↑ 164%  darunavir/cobicistat 800/150 mg once daily: dabigatran AUC ↑ 88% dabigatran C <sub>max</sub> ↑ 99%	Darunavir/ritonavir: Clinical monitoring and/or dose reduction of the DOAC should be considered when a DOAC transported by P-gp but not metabolised by CYP3A4, including dabigatran etexilate and edoxaban, is co-administered with darunavir/rtv.  Darunavir/cobicistat: Clinical monitoring and dose reduction is required when a DOAC transported by P-gp but not metabolised by CYP3A4, including dabigatran etexilate and edoxaban, is co-administered with darunavir/cobi.
Ticagrelor	Based on theoretical considerations, co-administration of boosted darunavir with ticagrelor may increase concentrations of ticagrelor (CYP3A and/or P-glycoprotein inhibition).	Concomitant administration of boosted darunavir with ticagrelor is contraindicated (see section 4.3).
Clopidogrel	Not studied. Co-administration of clopidogrel with boosted darunavir is expected to decrease clopidogrel active metabolite plasma concentration, which may reduce the antiplatelet activity of clopidogrel.	Co-administration of clopidogrel with boosted darunavir is not recommended.  Use of other antiplatelets not affected by CYP inhibition or induction (e.g. prasugrel) is recommended.

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
Warfarin	Not studied. Warfarin concentrations may be affected when co-administered with boosted darunavir.	It is recommended that the international normalised ratio (INR) be monitored when warfarin is combined with boosted darunavir.
<b>ANTICONVULSANTS</b>		
Phenobarbital Phenytoin	Not studied. Phenobarbital and phenytoin are expected to decrease plasma concentrations of darunavir and its pharmacoenhancer. (induction of CYP450 enzymes)	Darunavir co-administered with low dose ritonavir should not be used in combination with these medicines.  The use of these medicines with darunavir/cobicistat is contraindicated (see section 4.3).
Carbamazepine 200 mg twice daily	carbamazepine AUC ↑ 45% carbamazepine C <sub>min</sub> ↑ 54% carbamazepine C <sub>max</sub> ↑ 43% darunavir AUC ↔ darunavir C <sub>min</sub> ↓ 15% darunavir C <sub>max</sub> ↔	No dose adjustment for darunavir/ritonavir is recommended. If there is a need to combine darunavir/ritonavir and carbamazepine, patients should be monitored for potential carbamazepine-related adverse events. Carbamazepine concentrations should be monitored and its dose should be titrated for adequate response. Based upon the findings, the carbamazepine dose may need to be reduced by 25% to 50% in the presence of darunavir/ritonavir.  The use of carbamazepine with darunavir co-administered with cobicistat is contraindicated (see section 4.3).
Clonazepam	Not studied. Co-administration of boosted darunavir with clonazepam may increase concentrations of clonazepam. (CYP3A inhibition)	Clinical monitoring is recommended when co-administering boosted darunavir with clonazepam.



INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
<b>ANTIDEPRESSANTS</b>		
Paroxetine 20 mg once daily	paroxetine AUC ↓ 39% paroxetine C <sub>min</sub> ↓ 37% paroxetine C <sub>max</sub> ↓ 36% #darunavir AUC ↔ #darunavir C <sub>min</sub> ↔ #darunavir C <sub>max</sub> ↔	If antidepressants are co-administered with boosted darunavir, the recommended approach is a dose titration of the antidepressant based on a clinical assessment of antidepressant response. In addition, patients on a stable dose of these antidepressants who start treatment with boosted darunavir should be monitored for antidepressant response.
Sertraline 50 mg once daily	sertraline AUC ↓ 49% sertraline C <sub>min</sub> ↓ 49% sertraline C <sub>max</sub> ↓ 44% #darunavir AUC ↔ #darunavir C <sub>min</sub> ↓ 6% #darunavir C <sub>max</sub> ↔  In contrast to these data with darunavir/ritonavir, darunavir/cobicistat may increase these antidepressant plasma concentrations (CYP2D6 and/or CYP3A inhibition).	
Amitriptyline Desipramine Imipramine Nortriptyline Trazodone	Concomitant use of boosted darunavir and these antidepressants may increase concentrations of the antidepressant. (CYP2D6 and/or CYP3A inhibition)	Clinical monitoring is recommended when co-administering boosted darunavir with these antidepressants and a dose adjustment of the antidepressant may be needed.
<b>ANTI-DIABETICS</b>		
Metformin	Not studied. Based on theoretical considerations darunavir co-administered with cobicistat is expected to increase metformin plasma concentrations. (MATE1 inhibition)	Careful patient monitoring and dose adjustment of metformin is recommended in patients who are taking darunavir co-administered with cobicistat. (not applicable for darunavir co-administered with ritonavir)
<b>ANTIEMETICS</b>		
Domperidone	Not studied.	Co-administration of domperidone with boosted darunavir is contraindicated.
<b>ANTIFUNGALS</b>		
Voriconazole	Not studied. Ritonavir may decrease plasma concentrations of voriconazole. (induction of CYP450 enzymes)  Concentrations of voriconazole may increase or decrease when co-administered with darunavir co-administered with cobicistat. (inhibition of CYP450 enzymes)	Voriconazole should not be combined with boosted darunavir unless an assessment of the benefit/risk ratio justifies the use of voriconazole.

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
Fluconazole Isavuconazole Itraconazole Posaconazole	Not studied. Boosted darunavir may increase antifungal plasma concentrations and posaconazole, isavuconazole, itraconazole or fluconazole may increase darunavir concentrations. (CYP3A and/or P-gp inhibition)	Caution is warranted and clinical monitoring is recommended. When co-administration is required the daily dose of itraconazole should not exceed 200 mg.
Clotrimazole	Not studied. Concomitant systemic use of clotrimazole and boosted darunavir may increase plasma concentrations of darunavir and/or clotrimazole. Darunavir AUC <sub>24h</sub> ↑ 33% (based on population pharmacokinetic model)	
ANTIGOUT MEDICINES		
Colchicine	Not studied. Concomitant use of colchicine and boosted darunavir may increase the exposure to colchicine. (CYP3A and/ or P-gp inhibition)	A reduction in colchicine dosage or an interruption of colchicine treatment is recommended in patients with normal renal or hepatic function if treatment with boosted darunavir is required. For patients with renal or hepatic impairment colchicine with boosted darunavir is contraindicated (see sections 4.3 and 4.4).
ANTIMALARIALS		
Artemether/Lumefantrine 80/480 mg, 6 doses at 0, 8, 24, 36, 48, and 60 hours	artemether AUC ↓ 16% artemether C <sub>min</sub> ↔ artemether C <sub>max</sub> ↓ 18% dihydroartemisinin AUC ↓ 18% dihydroartemisinin C <sub>min</sub> ↔ dihydroartemisinin C <sub>max</sub> ↓ 18% lumefantrine AUC ↑ 175% lumefantrine C <sub>min</sub> ↑ 126% lumefantrine C <sub>max</sub> ↑ 65% darunavir AUC ↔ darunavir C <sub>min</sub> ↓ 13% darunavir C <sub>max</sub> ↔	The combination of boosted darunavir and artemether/lumefantrine can be used without dose adjustments; however, due to the increase in lumefantrine exposure, the combination should be used with caution.
ANTIMYCOBACTERIALS		
Rifampicin Rifapentine	Not studied. Rifapentine and rifampicin are strong CYP3A inducers and have been shown to cause profound decreases in concentrations of other protease inhibitors, which can result in virological failure and resistance development (CYP450 enzyme induction). During attempts to overcome the decreased exposure by increasing the dose of other protease inhibitors with low dose ritonavir, a high frequency of liver reactions was seen with rifampicin.	The combination of rifapentine and boosted darunavir is not recommended.  The combination of rifampicin and boosted darunavir is contraindicated (see section 4.3).

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
Rifabutin 150 mg once every other day	<p>rifabutin AUC** ↑ 55%</p> <p>rifabutin C<sub>min</sub>** ↑ ND</p> <p>rifabutin C<sub>max</sub>** ↔</p> <p>darunavir AUC ↑ 53%</p> <p>darunavir C<sub>min</sub> ↑ 68%</p> <p>darunavir C<sub>max</sub> ↑ 39%</p> <p>** sum of active moieties of rifabutin (parent drug + 25-O-desacetyl metabolite)</p> <p>The interaction trial showed a comparable daily systemic exposure for rifabutin between treatment at 300 mg once daily alone and 150 mg once every other day in combination with darunavir/ritonavir (600/100 mg twice daily) with an about 10-fold increase in the daily exposure to the active metabolite 25- O-desacetyl rifabutin. Furthermore, AUC of the sum of active moieties of rifabutin (parent drug + 25-O- desacetyl metabolite) was increased 1.6-fold, while C<sub>max</sub> remained comparable. Data on comparison with a 150 mg once daily reference dose is lacking.</p> <p>(Rifabutin is an inducer and substrate of CYP3A.) An increase of systemic exposure to darunavir was observed when darunavir co-administered with 100 mg ritonavir was co-administered with rifabutin (150 mg once every other day).</p>	<p>A dosage reduction of rifabutin by 75% of the usual dose of 300 mg/day (i.e. rifabutin 150 mg once every other day) and increased monitoring for rifabutin related adverse events is warranted in patients receiving the combination with darunavir co-administered with ritonavir. In case of safety issues, a further increase of the dosing interval for rifabutin and/or monitoring of rifabutin levels should be considered. Consideration should be given to official guidance on the appropriate treatment of tuberculosis in HIV infected patients. Based upon the safety profile of darunavir/ritonavir, the increase in darunavir exposure in the presence of rifabutin does not warrant a dose adjustment for darunavir/ritonavir. Based on pharmacokinetic modeling, this dosage reduction of 75% is also applicable if patients receive rifabutin at doses other than 300 mg/day.</p> <p>Co-administration of darunavir co-administered with cobicistat and rifabutin is not recommended.</p>
<b>ANTINEOPLASTICS</b>		
Dasatinib Nilotinib Vinblastine Vincristine	Not studied. Boosted darunavir is expected to increase these antineoplastic plasma concentrations. (CYP3A inhibition)	Concentrations of these medicinal products may be increased when co-administered with boosted darunavir resulting in the potential for increased adverse events usually associated with these agents. Caution should be exercised when combining one of these antineoplastic agents with boosted darunavir.
Everolimus Irinotecan		Concomitant use of everolimus or Irinotecan and boosted darunavir is not recommended.
<b>ANTIPSYCHOTICS/NEUROLEPTICS</b>		
Quetiapine	Not studied. Boosted darunavir is expected to increase these antipsychotic plasma concentrations. (CYP3A inhibition)	Concomitant administration of boosted darunavir and quetiapine is contraindicated as it may increase quetiapine-related toxicity. Increased concentrations of quetiapine may lead to coma (see section 4.3).

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
Perphenazine Risperidone Thioridazine  Lurasidone Pimozide Sertindole	Not studied. Boosted darunavir is expected to increase these antipsychotic plasma concentrations. (CYP3A, CYP2D6 and/or P-gp inhibition)	A dose decrease may be needed for these drugs when co-administered with boosted darunavir.  Concomitant administration of boosted darunavir and lurasidone, pimozide or sertindole is contraindicated (see section 4.3).
<b>β-BLOCKERS</b>		
Carvedilol Metoprolol Timolol	Not studied. Boosted darunavir is expected to increase these β-blocker plasma concentrations. (CYP2D6 inhibition)	Clinical monitoring is recommended when co-administering boosted darunavir with β-blockers. A lower dose of the β-blocker should be considered.
<b>CALCIUM CHANNEL BLOCKERS</b>		
Amlodipine Diltiazem Felodipine Nicardipine Nifedipine Verapamil	Not studied. Boosted darunavir can be expected to increase the plasma concentrations of calcium channel blockers. (CYP3A and/or CYP2D6 inhibition)	Clinical monitoring of therapeutic and adverse effects is recommended when these medicines are concomitantly administered with boosted darunavir.
<b>CORTICOSTEROIDS</b>		
Corticosteroids primarily metabolised by CYP3A (including betamethasone, budesonide, fluticasone, mometasone, prednisone, triamcinolone)	<p>Fluticasone: in a clinical study where ritonavir 100 mg capsules twice daily were co-administered with 50 µg intranasal fluticasone propionate (4 times daily) for 7 days in healthy subjects, fluticasone propionate plasma concentrations increased significantly, whereas the intrinsic cortisol levels decreased by approximately 86% (90% CI 82-89%). Greater effects may be expected when fluticasone is inhaled. Systemic corticosteroid effects including Cushing's syndrome and adrenal suppression have been reported in patients receiving ritonavir and inhaled or intranasally administered fluticasone. The effects of high fluticasone systemic exposure on ritonavir plasma levels are unknown.</p> <p>Other corticosteroids: interaction not studied. Plasma concentrations of these medicinal products may be increased when co-administered with boosted darunavir, resulting in reduced serum cortisol concentrations.</p>	<p>Concomitant use of boosted darunavir and corticosteroids (all routes of administration) that are metabolised by CYP3A may increase the risk of development of systemic corticosteroid effects, including Cushing's syndrome and adrenal suppression.</p> <p>Co-administration with CYP3A-metabolised corticosteroids is not recommended unless the potential benefit to the patient outweighs the risk, in which case patients should be monitored for systemic corticosteroid effects.</p> <p>Alternative corticosteroids which are less dependent on CYP3A metabolism e.g. beclomethasone should be considered, particularly for long term use.</p>
Dexamethasone (systemic)	Not studied. Dexamethasone may decrease plasma concentrations of darunavir. (CYP3A induction)	Systemic dexamethasone should be used with caution when combined with boosted darunavir.

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
<b>ENDOTHELIN RECEPTOR ANTAGONISTS</b>		
Bosentan	Not studied. Concomitant use of bosentan and boosted darunavir may increase plasma concentrations of bosentan. Bosentan is expected to decrease plasma concentrations of darunavir and/or its pharmacoenhancer. (CYP3A induction)	When administered concomitantly with darunavir and low dose ritonavir, the patient's tolerability of bosentan should be monitored.  Co-administration of darunavir co-administered with cobicistat and bosentan is not recommended.
<b>HEPATITIS C VIRUS (HCV) DIRECT-ACTING ANTIVIRALS</b>		
<b>NS3-4A protease inhibitors</b>		
Elbasvir/grazoprevir	Boosted darunavir may increase the exposure to grazoprevir. (CYP3A and OATP1B inhibition).	Concomitant use of boosted darunavir and elbasvir/grazoprevir is contraindicated (see section 4.3).
Glecaprevir/pibrentasvir	Based on theoretical considerations boosted darunavir may increase the exposure to glecaprevir and pibrentasvir. (P-gp, BCRP and/or OATP1B1/3 inhibition)	It is not recommended to co-administer boosted darunavir with glecaprevir/pibrentasvir.
<b>HERBAL PRODUCTS</b>		
St John's wort ( <i>Hypericum perforatum</i> )	Not studied. St John's wort is expected to decrease the plasma concentrations of darunavir or its pharmacoenhancers. (CYP450 induction)	Boosted darunavir must not be used concomitantly with products containing St John's wort ( <i>Hypericum perforatum</i> ) (see section 4.3). If a patient is already taking St John's wort, stop St John's wort and if possible check viral levels. Darunavir exposure (and also ritonavir exposure) may increase on stopping St John's wort. The inducing effect may persist for at least 2 weeks after cessation of treatment with St John's wort.
<b>HMG CO-A REDUCTASE INHIBITORS</b>		
Lovastatin Simvastatin	Not studied. Lovastatin and simvastatin are expected to have markedly increased plasma concentrations when co-administered with boosted darunavir. (CYP3A inhibition)	Increased plasma concentrations of lovastatin or simvastatin may cause myopathy, including rhabdomyolysis. Concomitant use of boosted darunavir with lovastatin and simvastatin is therefore contraindicated (see section 4.3).
Atorvastatin 10 mg once daily	atorvastatin AUC ↑ 3-4 fold atorvastatin C <sub>min</sub> ↑ ≈5.5-10 fold atorvastatin C <sub>max</sub> ↑ ≈2 fold #darunavir/ritonavir  atorvastatin AUC ↑ 290% <sup>Ω</sup> atorvastatin C <sub>max</sub> ↑ 319% <sup>Ω</sup> atorvastatin C <sub>min</sub> ND <sup>Ω</sup> <sup>Ω</sup> with darunavir/cobicistat 800/150 mg	When administration of atorvastatin and boosted darunavir is desired, it is recommended to start with an atorvastatin dose of 10 mg once daily. A gradual dose increase of atorvastatin may be tailored to the clinical response.

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
Pravastatin 40 mg single dose	pravastatin AUC ↑ 81% <sup>†</sup> pravastatin C <sub>min</sub> ND pravastatin C <sub>max</sub> ↑ 63% <sup>†</sup> an up to five-fold increase was seen in a limited subset of subjects	When administration of pravastatin and boosted darunavir is required, it is recommended to start with the lowest possible dose of pravastatin and titrate up to the desired clinical effect while monitoring for safety.
Rosuvastatin 10 mg once daily	rosuvastatin AUC ↑ 48% <sup>‡</sup> rosuvastatin C <sub>max</sub> ↑ 144% <sup>‡</sup> <sup>‡</sup> based on published data with darunavir/ritonavir  rosuvastatin AUC ↑ 93% <sup>§</sup> rosuvastatin C <sub>max</sub> ↑ 277% <sup>§</sup> rosuvastatin C <sub>min</sub> ND <sup>§</sup> <sup>§</sup> with darunavir/cobicistat 800/150 mg	When administration of rosuvastatin and boosted darunavir is required, it is recommended to start with the lowest possible dose of rosuvastatin and titrate up to the desired clinical effect while monitoring for safety.
<b>OTHER LIPID MODIFYING AGENTS</b>		
Lomitapide	Based on theoretical considerations boosted darunavir is expected to increase the exposure of lomitapide when co-administered. (CYP3A inhibition)	Co-administration is contraindicated (see section 4.3).
<b>H2-RECEPTOR ANTAGONISTS</b>		
Ranitidine 150 mg twice daily	#darunavir AUC ↔ #darunavir C <sub>min</sub> ↔ #darunavir C <sub>max</sub> ↔	Boosted darunavir can be co-administered with H2-receptor antagonists without dose adjustments.
<b>IMMUNOSUPPRESSANTS</b>		
Ciclosporin Sirolimus Tacrolimus Everolimus	Not studied. Exposure to these immunosuppressants will be increased when co-administered with boosted darunavir. (CYP3A inhibition)	Therapeutic drug monitoring of the immunosuppressive agent must be done when co-administration occurs. Concomitant use of everolimus and boosted darunavir is not recommended.
<b>INHALED BETA AGONISTS</b>		
Salmeterol	Not studied. Concomitant use of salmeterol and boosted darunavir may increase plasma concentrations of salmeterol.	Concomitant use of salmeterol and boosted darunavir is not recommended. The combination may result in increased risk of cardiovascular adverse event with salmeterol, including QT prolongation, palpitations and sinus tachycardia.
<b>NARCOTIC ANALGESICS / TREATMENT OF OPIOID DEPENDENCE</b>		
Methadone individual dose ranging from 55 mg to 150 mg once daily	R(-) methadone AUC ↓ 16% R(-) methadone C <sub>min</sub> ↓ 15% R(-) methadone C <sub>max</sub> ↓ 24%  Darunavir/cobicistat may, in contrast, increase methadone plasma concentrations (see cobicistat SmPC).	No adjustment of methadone dosage is required when initiating co-administration with boosted darunavir. However, adjustment of the methadone dose may be necessary when concomitantly administered for a longer period of time. Therefore, clinical monitoring is recommended, as maintenance therapy may need to be adjusted in some patients.

<b>INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS</b>		
<b>Medicinal products by therapeutic areas</b>	<b>Interaction Geometric mean change (%)</b>	<b>Recommendations concerning co-administration</b>
Buprenorphine/naloxone 8/2 mg–16/4 mg once daily	buprenorphine AUC ↓ 11% buprenorphine C <sub>min</sub> ↔ buprenorphine C <sub>max</sub> ↓ 8% norbuprenorphine AUC ↑ 46% norbuprenorphine C <sub>min</sub> ↑ 71% norbuprenorphine C <sub>max</sub> ↑ 36% naloxone AUC ↔ naloxone C <sub>min</sub> ND naloxone C <sub>max</sub> ↔	The clinical relevance of the increase in norbuprenorphine pharmacokinetic parameters has not been established. Dose adjustment for buprenorphine may not be necessary when co-administered with boosted darunavir but a careful clinical monitoring for signs of opiate toxicity is recommended.
Fentanyl Oxycodone Tramadol	Based on theoretical considerations boosted darunavir may increase plasma concentrations of these analgesics. (CYP2D6 and/or CYP3A inhibition)	Clinical monitoring is recommended when co-administering boosted darunavir with these analgesics.
<b>OESTROGEN-BASED CONTRACEPTIVES</b>		
Drospirenone Ethinylestradiol (3 mg/0.02 mg once daily)	drospirenone AUC ↑ 58% <sup>ε</sup> drospirenone C <sub>min</sub> ND <sup>ε</sup> drospirenone C <sub>max</sub> ↑ 15% <sup>ε</sup> ethinylestradiol AUC ↓ 30% <sup>ε</sup> ethinylestradiol C <sub>min</sub> ND <sup>ε</sup> ethinylestradiol C <sub>max</sub> ↓ 14% <sup>ε</sup> <sup>ε</sup> with darunavir/cobicistat	When darunavir is coadministered with a drospirenone-containing product, clinical monitoring is recommended due to the potential for hyperkalaemia.  Alternative or additional contraceptive measures are recommended when oestrogen-based contraceptives are co-administered with boosted darunavir. Patients using oestrogens as hormone replacement therapy should be clinically monitored for signs of oestrogen deficiency.
Ethinylestradiol Norethindrone 35 µg/1 mg once daily	ethinylestradiol AUC ↓ 44% ethinylestradiol C <sub>min</sub> ↓ 62% ethinylestradiol C <sub>max</sub> ↓ 32% norethindrone AUC ↓ 14% norethindrone C <sub>min</sub> ↓ 30% norethindrone C <sub>max</sub> ↔ <sup>β</sup> with darunavir/ritonavir	
<b>OPIOID ANTAGONIST</b>		
Naloxegol	Not studied.	Co-administration of boosted darunavir and naloxegol is contraindicated.
<b>PHOSPHODIESTERASE, TYPE 5 (PDE-5) INHIBITORS</b>		
For the treatment of erectile dysfunction Avanafil Sildenafil Tadalafil Vardenafil	In an interaction study <sup>#</sup> , a comparable systemic exposure to sildenafil was observed for a single intake of 100 mg sildenafil alone and a single intake of 25 mg sildenafil co-administered with darunavir and low dose ritonavir.	The combination of avanafil and boosted darunavir is contraindicated (see section 4.3). Concomitant use of other PDE-5 inhibitors for the treatment of erectile dysfunction with boosted darunavir should be done with caution. If concomitant use of boosted darunavir with sildenafil, vardenafil or tadalafil is indicated, sildenafil at a single dose not exceeding 25 mg in 48 hours, vardenafil at a single dose not exceeding 2.5 mg in 72 hours or tadalafil at a single dose not exceeding 10 mg in 72 hours is recommended.

INTERACTIONS AND DOSE RECOMMENDATIONS WITH OTHER MEDICINAL PRODUCTS		
Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
For the treatment of pulmonary arterial hypertension Sildenafil Tadalafil	Not studied. Concomitant use of sildenafil or tadalafil for the treatment of pulmonary arterial hypertension and boosted darunavir may increase plasma concentrations of sildenafil or tadalafil. (CYP3A inhibition)	A safe and effective dose of sildenafil for the treatment of pulmonary arterial hypertension co-administered with boosted darunavir has not been established. There is an increased potential for sildenafil-associated adverse events (including visual disturbances, hypotension, prolonged erection and syncope). Therefore, co-administration of boosted darunavir and sildenafil when used for the treatment of pulmonary arterial hypertension is contraindicated (see section 4.3). Co-administration of tadalafil for the treatment of pulmonary arterial hypertension with boosted darunavir is not recommended.
<b>PROTON PUMP INHIBITORS</b>		
Omeprazole 20 mg once daily	#darunavir AUC ↔ #darunavir C <sub>min</sub> ↔ #darunavir C <sub>max</sub> ↔	Boosted darunavir can be co-administered with proton pump inhibitors without dose adjustments.
<b>SEDATIVES/HYPNOTICS</b>		
Buspirone Clorazepate Diazepam Estazolam Flurazepam Midazolam (parenteral) Zolpidem          Midazolam (oral) Triazolam	Not studied. Sedative/hypnotics are extensively metabolised by CYP3A. Co-administration with boosted darunavir may cause a large increase in the concentration of these medicines.  If parenteral midazolam is co-administered with boosted darunavir it may cause a large increase in the concentration of this benzodiazepine. Data from concomitant use of parenteral midazolam with other protease inhibitors suggest a possible 3-4 fold increase in midazolam plasma levels.	Clinical monitoring is recommended when co-administering boosted darunavir with these sedatives/hypnotics and a lower dose of the sedatives/hypnotics should be considered.  If parenteral midazolam is co-administered with boosted darunavir, it should be done in an intensive care unit (ICU) or similar setting, which ensures close clinical monitoring and appropriate medical management in case of respiratory depression and/or prolonged sedation. Dose adjustment for midazolam should be considered, especially if more than a single dose of midazolam is administered.  Boosted darunavir with triazolam or oral midazolam is contraindicated (see section 4.3)
<b>TREATMENT FOR PREMATURE EJACULATION</b>		
Dapoxetine	Not studied.	Co-administration of boosted darunavir with dapoxetine is contraindicated.
<b>UROLOGICAL DRUGS</b>		
Fesoterodine Solifenacin	Not studied.	Use with caution. Monitor for fesoterodine or solifenacin adverse reactions, dose reduction of fesoterodine or solifenacin may be necessary.

# Studies have been performed at lower than recommended doses of darunavir or with a different dosing regimen (see section 4.2 Posology).



- † The efficacy and safety of the use of darunavir with 100 mg ritonavir and any other HIV PI (e.g. (fos)amprenavir and tipranavir) has not been established in HIV patients. According to current treatment guidelines, dual therapy with protease inhibitors is generally not recommended.
- ‡ Study was conducted with tenofovir disoproxil fumarate 300 mg once daily.

## 4.6 Fertility, pregnancy and lactation

### Pregnancy

As a general rule, when deciding to use antiretroviral agents for the treatment of HIV infection in pregnant women and consequently for reducing the risk of HIV vertical transmission to the newborn, the animal data as well as the clinical experience in pregnant women should be taken into account.

There are no adequate and well controlled studies on pregnancy outcome with darunavir in pregnant women. Studies in animals do not indicate direct harmful effects with respect to pregnancy, embryonal/foetal development, parturition or postnatal development (see section 5.3).

Treatment with darunavir/cobicistat 800/150 mg during pregnancy results in low darunavir exposure (see section 5.2), which may be associated with an increased risk of treatment failure and an increased risk of HIV transmission to the child. Therapy with Darunavir/cobicistat should not be initiated during pregnancy, and women who become pregnant during therapy with Darunavir/cobicistat should be switched to an alternative regimen (see sections 4.2 and 4.4).

Darunavir co-administered with low dose ritonavir should be used during pregnancy only if the potential benefit justifies the potential risk.

### Breast-feeding

It is not known whether darunavir is excreted in human milk. Studies in rats have demonstrated that darunavir is excreted in milk and at high levels (1,000 mg/kg/day) resulted in toxicity of the offspring.

Because of the potential for adverse reactions in breast-fed infants, women should be instructed not to breast-feed if they are receiving darunavir.

In order to avoid transmission of HIV to the infant it is recommended that women living with HIV do not breast-feed.

### Fertility

No human data on the effect of darunavir on fertility are available. There was no effect on mating or fertility with darunavir treatment in rats (see section 5.3).

## 4.7 Effects on ability to drive and use machines

Darunavir in combination with cobicistat or ritonavir has no or negligible influence on the ability to drive and use machines. However, dizziness has been reported in some patients during treatment with regimens containing darunavir co-administered with cobicistat or low dose ritonavir and should be borne in mind when considering a patient's ability to drive or operate machinery (see section 4.8).

## 4.8 Undesirable effects

### Summary of the safety profile

During the clinical development program (N=2,613 treatment-experienced subjects who initiated therapy with darunavir/ritonavir 600/100 mg twice daily), 51.3% of subjects experienced at least one adverse reaction. The total mean treatment duration for subjects was 95.3 weeks. The most frequent adverse reactions reported in clinical trials and as spontaneous reports are diarrhoea, nausea, rash, headache and vomiting. The most frequent serious reactions are acute renal failure, myocardial

infarction, immune reconstitution inflammatory syndrome, thrombocytopenia, osteonecrosis, diarrhoea, hepatitis and pyrexia.

In the 96 week analysis, the safety profile of darunavir/ritonavir 800/100 mg once daily in treatment-naïve subjects was similar to that seen with darunavir/ritonavir 600/100 mg twice daily in treatment-experienced subjects except for nausea which was observed more frequently in treatment-naïve subjects. This was driven by mild intensity nausea. No new safety findings were identified in the 192 week analysis of the treatment-naïve subjects in which the mean treatment duration of darunavir/ritonavir 800/100 mg once daily was 162.5 weeks.

During the Phase III clinical trial GS-US-216-130 with darunavir/cobicistat (N=313 treatment-naïve and treatment-experienced subjects), 66.5% of subjects experienced at least one adverse reaction. The mean treatment duration was 58.4 weeks. The most frequent adverse reactions reported were diarrhoea (28%), nausea (23%), and rash (16%). Serious adverse reactions are diabetes mellitus, (drug) hypersensitivity, immune reconstitution inflammatory syndrome, rash and vomiting.

For information on cobicistat, consult the cobicistat Summary of Product Characteristics.

#### Tabulated list of adverse reactions

Adverse reactions are listed by system organ class (SOC) and frequency category. Within each frequency category, adverse reactions are presented in order of decreasing seriousness. Frequency categories are defined as follows: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ) and not known (frequency cannot be estimated from the available data).

#### *Adverse reactions observed with darunavir/ritonavir in clinical trials and post-marketing*

MedDRA system organ class Frequency category	Adverse reaction
<i>Infections and infestations</i>	
Uncommon	herpes simplex
<i>Blood and lymphatic system disorders</i>	
Uncommon	thrombocytopenia, neutropenia, anaemia, leucopenia
Rare	increased eosinophil count
<i>Immune system disorders</i>	
Uncommon	immune reconstitution inflammatory syndrome, (drug) hypersensitivity
<i>Endocrine disorders</i>	
Uncommon	hypothyroidism, increased blood thyroid stimulating hormone
<i>Metabolism and nutrition disorders</i>	
Common	diabetes mellitus, hypertriglyceridaemia, hypercholesterolaemia, hyperlipidaemia
Uncommon	gout, anorexia, decreased appetite, decreased weight, increased weight, hyperglycaemia, insulin resistance, decreased high density lipoprotein, increased appetite, polydipsia, increased blood lactate dehydrogenase

MedDRA system organ class Frequency category	Adverse reaction
<i>Psychiatric disorders</i>	
Common	insomnia
Uncommon	depression, disorientation, anxiety, sleep disorder, abnormal dreams, nightmare, decreased libido
Rare	confusional state, altered mood, restlessness
<i>Nervous system disorders</i>	
Common	headache, peripheral neuropathy, dizziness
Uncommon	lethargy, paraesthesia, hypoaesthesia, dysgeusia, disturbance in attention, memory impairment, somnolence
Rare	syncope, convulsion, ageusia, sleep phase rhythm disturbance
<i>Eye disorders</i>	
Uncommon	conjunctival hyperaemia, dry eye
Rare	visual disturbance
<i>Ear and labyrinth disorders</i>	
Uncommon	vertigo
<i>Cardiac disorders</i>	
Uncommon	myocardial infarction, angina pectoris, prolonged electrocardiogram QT, tachycardia
Rare	acute myocardial infarction, sinus bradycardia, palpitations
<i>Vascular disorders</i>	
Uncommon	hypertension, flushing
<i>Respiratory, thoracic and mediastinal disorders</i>	
Uncommon	dyspnoea, cough, epistaxis, throat irritation
Rare	rhinorrhoea
<i>Gastrointestinal disorders</i>	
Very common	diarrhoea
Common	vomiting, nausea, abdominal pain, increased blood amylase, dyspepsia, abdominal distension, flatulence
Uncommon	pancreatitis, gastritis, gastrooesophageal reflux disease, aphthous stomatitis, retching, dry mouth, abdominal discomfort, constipation, increased lipase, eructation, oral dysaesthesia
Rare	stomatitis, haematemesis, cheilitis, dry lip, coated tongue
<i>Hepatobiliary disorders</i>	
Common	increased alanine aminotransferase
Uncommon	hepatitis, cytolytic hepatitis, hepatic steatosis, hepatomegaly, increased transaminase, increased aspartate aminotransferase, increased blood bilirubin, increased blood alkaline phosphatase, increased gamma-glutamyltransferase

MedDRA system organ class Frequency category	Adverse reaction
<i>Skin and subcutaneous tissue disorders</i>	
Common	rash (including macular, maculopapular, papular, erythematous and pruritic rash), pruritus
Uncommon	angioedema, generalised rash, allergic dermatitis, urticaria, eczema, erythema, hyperhidrosis, night sweats, alopecia, acne, dry skin, nail pigmentation
Rare	DRESS, Stevens-Johnson syndrome, erythema multiforme, dermatitis, seborrhoeic dermatitis, skin lesion, xeroderma
Not known	toxic epidermal necrolysis, acute generalised exanthematous pustulosis
<i>Musculoskeletal and connective tissue disorders</i>	
Uncommon	myalgia, osteonecrosis, muscle spasms, muscular weakness, arthralgia, pain in extremity, osteoporosis, increased blood creatine phosphokinase
Rare	musculoskeletal stiffness, arthritis, joint stiffness
<i>Renal and urinary disorders</i>	
Uncommon	acute renal failure, renal failure, nephrolithiasis, increased blood creatinine, proteinuria, bilirubinuria, dysuria, nocturia, pollakiuria
Rare	decreased creatinine renal clearance, crystal nephropathy <sup>§</sup>
<i>Reproductive system and breast disorders</i>	
Uncommon	erectile dysfunction, gynaecomastia
<i>General disorders and administration site conditions</i>	
Common	asthenia, fatigue
Uncommon	pyrexia, chest pain, peripheral oedema, malaise, feeling hot, irritability, pain
Rare	chills, abnormal feeling, xerosis

<sup>§</sup> adverse reaction identified in the post-marketing setting. Per the guideline on Summary of Product Characteristics (Revision 2, September 2009), the frequency of this adverse reaction in the post-marketing setting was determined using the "Rule of 3".

#### *Adverse reactions observed with darunavir/cobicistat in adult patients*

MedDRA system organ class Frequency category	Adverse reaction
<i>Immune system disorders</i>	
Common	(drug) hypersensitivity
Uncommon	immune reconstitution inflammatory syndrome
<i>Metabolism and nutrition disorders</i>	
Common	anorexia, diabetes mellitus, hypercholesterolaemia, hypertriglyceridaemia, hyperlipidaemia
<i>Psychiatric disorders</i>	
Common	abnormal dreams

MedDRA system organ class Frequency category	Adverse reaction
<i>Nervous system disorders</i>	
Very common	headache
<i>Gastrointestinal disorders</i>	
Very common	diarrhoea, nausea
Common	vomiting, abdominal pain, abdominal distension, dyspepsia, flatulence, pancreatic enzymes increased
Uncommon	pancreatitis acute
<i>Hepatobiliary disorders</i>	
Common	hepatic enzyme increased
Uncommon	hepatitis*, cytolytic hepatitis*
<i>Skin and subcutaneous tissue disorders</i>	
Very common	rash (including macular, maculopapular, papular, erythematous, pruritic rash, generalised rash, and allergic dermatitis)
Common	angioedema, pruritus, urticaria
Rare	drug reaction with eosinophilia and systemic symptoms*, Stevens-Johnson syndrome*
Not known	toxic epidermal necrolysis*, acute generalised exanthematous pustulosis*
<i>Musculoskeletal and connective tissue disorders</i>	
Common	myalgia
Uncommon	osteonecrosis*
<i>Renal and urinary disorders</i>	
Rare	crystal nephropathy*§
<i>Reproductive system and breast disorders</i>	
Uncommon	gynaecomastia*
<i>General disorders and administration site conditions</i>	
Common	fatigue
Uncommon	asthenia
<i>Investigations</i>	
Common	increased blood creatinine

\* these adverse drug reactions have not been reported in clinical trial experience with darunavir/cobicistat but have been noted with darunavir/ritonavir treatment and could be expected with darunavir/cobicistat too.

§ adverse reaction identified in the post-marketing setting. Per the guideline on Summary of Product Characteristics (Revision 2, September 2009), the frequency of this adverse reaction in the post-marketing setting was determined using the "Rule of 3".

## Description of selected adverse reactions

### *Rash*

In clinical trials, rash was mostly mild to moderate, often occurring within the first four weeks of treatment and resolving with continued dosing. In cases of severe skin reaction see the warning in section 4.4. In a single arm trial investigating darunavir 800 mg once daily in combination with cobicistat 150 mg once daily and other antiretrovirals 2.2% of patients discontinued treatment due to rash.

During the clinical development program of raltegravir in treatment-experienced patients, rash, irrespective of causality, was more commonly observed with regimens containing darunavir/ritonavir + raltegravir compared to those containing darunavir/ritonavir without raltegravir or raltegravir without darunavir/ritonavir. Rash considered by the investigator to be drug-related occurred at similar rates. The exposure-adjusted rates of rash (all causality) were 10.9, 4.2, and 3.8 per 100 patient-years (PYR), respectively; and for drug-related rash were 2.4, 1.1, and 2.3 per 100 PYR, respectively. The rashes observed in clinical studies were mild to moderate in severity and did not result in discontinuation of therapy (see section 4.4).

### *Metabolic parameters*

Weight and levels of blood lipids and glucose may increase during antiretroviral therapy (see section 4.4).

### *Musculoskeletal abnormalities*

Increased CPK, myalgia, myositis and rarely, rhabdomyolysis have been reported with the use of protease inhibitors, particularly in combination with NRTIs.

Cases of osteonecrosis have been reported, particularly in patients with generally acknowledged risk factors, advanced HIV disease or long-term exposure to combination antiretroviral therapy (CART). The frequency of this is unknown (see section 4.4).

### *Immune reconstitution inflammatory syndrome*

In HIV infected patients with severe immune deficiency at the time of initiation of combination antiretroviral therapy (CART), an inflammatory reaction to asymptomatic or residual opportunistic infections may arise. Autoimmune disorders (such as Graves' disease and autoimmune hepatitis) have also been reported; however, the reported time to onset is more variable and these events can occur many months after initiation of treatment (see section 4.4).

### *Bleeding in haemophiliac patients*

There have been reports of increased spontaneous bleeding in haemophiliac patients receiving antiretroviral protease inhibitors (see section 4.4).

## Paediatric population

The safety assessment of darunavir with ritonavir in paediatric patients is based on the 48-week analysis of safety data from three Phase II trials. The following patient populations were evaluated (see section 5.1):

- 80 ART-experienced HIV-1 infected paediatric patients aged from 6 to 17 years and weighing at least 20 kg who received darunavir tablets with low dose ritonavir twice daily in combination with other antiretroviral agents.
- 21 ART-experienced HIV-1 infected paediatric patients aged from 3 to < 6 years and weighing 10 kg to < 20 kg (16 participants from 15 kg to < 20 kg) who received darunavir oral suspension with low dose ritonavir twice daily in combination with other antiretroviral agents.
- 12 ART-naïve HIV-1 infected paediatric patients aged from 12 to 17 years and weighing at least 40 kg who received darunavir tablets with low dose ritonavir once daily in combination with other antiretroviral agents (see section 5.1).

Overall, the safety profile in these paediatric patients was similar to that observed in the adult population.

The safety assessment of darunavir with cobicistat in paediatric patients was evaluated in adolescents aged 12 to less than 18 years, weighing at least 40 kg through the clinical trial GS-US-216-0128 (treatment experienced, virologically suppressed, N=7). Safety analyses of this study in adolescent subjects did not identify new safety concerns compared to the known safety profile of darunavir and cobicistat in adult subjects.

#### Other special populations

##### *Patients co-infected with hepatitis B and/or hepatitis C virus*

Among 1,968 treatment-experienced patients receiving darunavir co-administered with ritonavir 600/100 mg twice daily, 236 patients were co-infected with hepatitis B or C. Co-infected patients were more likely to have baseline and treatment emergent hepatic transaminase elevations than those without chronic viral hepatitis (see section 4.4).

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

## **4.9 Overdose**

Human experience of acute overdose with darunavir co-administered with cobicistat or low dose ritonavir is limited. Single doses up to 3,200 mg of darunavir as oral solution alone and up to 1,600 mg of the tablet formulation of darunavir in combination with ritonavir have been administered to healthy volunteers without untoward symptomatic effects.

There is no specific antidote for overdose with darunavir. Treatment of overdose with darunavir consists of general supportive measures including monitoring of vital signs and observation of the clinical status of the patient. Since darunavir is highly protein bound, dialysis is unlikely to be beneficial in significant removal of the active substance.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Antivirals for systemic use, protease inhibitors, ATC code: J05AE10.

#### Mechanism of action

Darunavir is an inhibitor of the dimerisation and of the catalytic activity of the HIV-1 protease (KD of  $4.5 \times 10^{-12}$  M). It selectively inhibits the cleavage of HIV encoded Gag-Pol polyproteins in virus infected cells, thereby preventing the formation of mature infectious virus particles.

#### Antiviral activity *in vitro*

Darunavir exhibits activity against laboratory strains and clinical isolates of HIV-1 and laboratory strains of HIV-2 in acutely infected T-cell lines, human peripheral blood mononuclear cells and human monocytes/macrophages with median EC<sub>50</sub> values ranging from 1.2 to 8.5 nM (0.7 to 5.0 ng/ml). Darunavir demonstrates antiviral activity *in vitro* against a broad panel of HIV-1 group M (A, B, C, D, E, F, G) and group O primary isolates with EC<sub>50</sub> values ranging from < 0.1 to 4.3 nM.

These EC<sub>50</sub> values are well below the 50% cellular toxicity concentration range of 87 µM to > 100 µM.

### Resistance

*In vitro* selection of darunavir-resistant virus from wild type HIV-1 was lengthy (> 3 years). The selected viruses were unable to grow in the presence of darunavir concentrations above 400 nM. Viruses selected in these conditions and showing decreased susceptibility to darunavir (range: 23-50-fold) harboured 2 to 4 amino acid substitutions in the protease gene. The decreased susceptibility to darunavir of the emerging viruses in the selection experiment could not be explained by the emergence of these protease mutations.

The clinical trial data from ART-experienced patients (TITAN trial and the pooled analysis of the POWER 1, 2 and 3 and DUET 1 and 2 trials) showed that virologic response to darunavir co-administered with low dose ritonavir was decreased when 3 or more darunavir RAMs (V11I, V32I, L33F, I47V, I50V, I54L or M, T74P, L76V, I84V and L89V) were present at baseline or when these mutations developed during treatment.

Increasing baseline darunavir fold change in EC<sub>50</sub> (FC) was associated with decreasing virologic response. A lower and upper clinical cut-off of 10 and 40 were identified. Isolates with baseline FC ≤ 10 are susceptible; isolates with FC > 10 to 40 have decreased susceptibility; isolates with FC > 40 are resistant (see Clinical results).

Viruses isolated from patients on darunavir/ritonavir 600/100 mg twice daily experiencing virologic failure by rebound that were susceptible to tipranavir at baseline remained susceptible to tipranavir after treatment in the vast majority of cases.

The lowest rates of developing resistant HIV virus are observed in ART-naïve patients who are treated for the first time with darunavir in combination with other ART.

The table below shows the development of HIV-1 protease mutations and loss of susceptibility to PIs in virologic failures at endpoint in the ARTEMIS, ODIN and TITAN trials.

	ARTEMIS Week 192	ODIN Week 48		TITAN Week 48
	darunavir/ ritonavir 800/100 mg once daily N=343	darunavir/ ritonavir 800/100 mg once daily N=294	darunavir/ ritonavir 600/100 mg twice daily N=296	darunavir/ ritonavir 600/100 mg twice daily N=298
Total number of virologic failures <sup>a</sup> , n (%)	55 (16.0%)	65 (22.1%)	54 (18.2%)	31 (10.4%)
Rebounders	39 (11.4%)	11 (3.7%)	11 (3.7%)	16 (5.4%)
Never suppressed subjects	16 (4.7%)	54 (18.4%)	43 (14.5%)	15 (5.0%)
Number of subjects with virologic failure and paired baseline/endpoint genotypes, developing mutations <sup>b</sup> at endpoint, n/N				
Primary (major) PI mutations	0/43	1/60	0/42	6/28
PI RAMs	4/43	7/60	4/42	10/28



	ARTEMIS Week 192	ODIN Week 48		TITAN Week 48
	darunavir/ ritonavir 800/100 mg once daily N=343	darunavir/ ritonavir 800/100 mg once daily N=294	darunavir/ ritonavir 600/100 mg twice daily N=296	darunavir/ ritonavir 600/100 mg twice daily N=298
Number of subjects with virologic failure and paired baseline/endpoint phenotypes, showing loss of susceptibility to PIs at endpoint compared to baseline, n/N				
PI				
darunavir	0/39	1/58	0/41	3/26
amprenavir	0/39	1/58	0/40	0/22
atazanavir	0/39	2/56	0/40	0/22
indinavir	0/39	2/57	0/40	1/24
lopinavir	0/39	1/58	0/40	0/23
saquinavir	0/39	0/56	0/40	0/22
tipranavir	0/39	0/58	0/41	1/25

<sup>a</sup> TLOVR non-VF censored algorithm based on HIV-1 RNA < 50 copies/ml, except for TITAN (HIV-1 RNA < 400 copies/ml)

<sup>b</sup> IAS-USA lists

Low rates of developing resistant HIV-1 virus were observed in ART-naïve patients who are treated for the first time with darunavir/cobicistat once daily in combination with other ART, and in ART-experienced patients with no darunavir RAMs receiving darunavir/cobicistat in combination with other ART. The table below shows the development of HIV-1 protease mutations and resistance to PIs in virologic failures at endpoint in the GS-US-216-130 trial.

	GS-US-216-130 Week 48	
	Treatment-naïve darunavir/cobicistat 800/150 mg once daily N=295	Treatment-experienced darunavir/cobicistat 800/150 mg once daily N=18
Number of subjects with virologic failure <sup>a</sup> and genotype data that develop mutations <sup>b</sup> at endpoint, n/N		
Primary (major) PI mutations	0/8	1/7
PI RAMs	2/8	1/7
Number of subjects with virologic failure <sup>a</sup> and phenotype data that show resistance to PIs at endpoint <sup>c</sup> , n/N		
HIV PI		
darunavir	0/8	0/7
amprenavir	0/8	0/7
atazanavir	0/8	0/7
indinavir	0/8	0/7
lopinavir	0/8	0/7
saquinavir	0/8	0/7
tipranavir	0/8	0/7

<sup>a</sup> Virologic failures were defined as: never suppressed: confirmed HIV-1 RNA < 1 log<sub>10</sub> reduction from baseline and ≥ 50 copies/ml at the week-8; rebound: HIV-1 RNA < 50 copies/ml followed by confirmed HIV-1 RNA to ≥ 400 copies/ml or confirmed > 1 log<sub>10</sub> HIV-1 RNA increase from the nadir; discontinuations with HIV-1 RNA ≥ 400 copies/ml at last visit

<sup>b</sup> IAS-USA lists

<sup>c</sup> In GS-US216-130 baseline phenotype was not available

## Cross-resistance

Darunavir FC was less than 10 for 90% of 3,309 clinical isolates resistant to amprenavir, atazanavir, indinavir, lopinavir, nelfinavir, ritonavir, saquinavir and/or tipranavir showing that viruses resistant to most PIs remain susceptible to darunavir.

In the virologic failures of the ARTEMIS trial no cross-resistance with other PIs was observed. In the virologic failures of the GS-US-216-130 trial no cross-resistance with other HIV PIs was observed.

## Clinical results

The pharmacokinetic enhancing effect of cobicistat on darunavir was evaluated in a Phase I study in healthy subjects that were administered darunavir 800 mg with either cobicistat at 150 mg or ritonavir at 100 mg once daily. The steady-state pharmacokinetic parameters of darunavir were comparable when boosted with cobicistat versus ritonavir. For information on cobicistat, consult the cobicistat Summary of Product Characteristics.

## Adult patients

### *Efficacy of darunavir 800 mg once daily co-administered with 150 mg cobicistat once daily in ART-naïve and ART-experienced patients*

GS-US-216-130 is a single arm, open-label, Phase III trial evaluating the pharmacokinetics, safety, tolerability, and efficacy of darunavir with cobicistat in 313 HIV-1 infected adult patients (295 treatment-naïve and 18 treatment-experienced). These patients received darunavir 800 mg once daily in combination with cobicistat 150 mg once daily with an investigator selected background regimen consisting of 2 active NRTIs.

HIV-1 infected patients who were eligible for this trial had a screening genotype showing no darunavir RAMs and plasma HIV-1 RNA  $\geq 1,000$  copies/ml. The table below shows the efficacy data of the 48 week analyses from the GS-US-216-130 trial:

	GS-US-216-130		
Outcomes at Week 48	Treatment-naïve darunavir/cobicistat 800/150 mg once daily + OBR N=295	Treatment-experienced darunavir/cobicistat 800/150 mg once daily + OBR N=18	All subjects darunavir/cobicistat 800/150 mg once daily + OBR N=313
HIV-1 RNA < 50 copies/ml <sup>a</sup>	245 (83.1%)	8 (44.4%)	253 (80.8%)
mean HIV-1 RNA log change from baseline (log <sub>10</sub> copies/ml)	-3.01	-2.39	-2.97
CD4+ cell count mean change from baseline <sup>b</sup>	+174	+102	+170

<sup>a</sup> Imputations according to the TLOVR algorithm

<sup>b</sup> Last Observation Carried Forward imputation

### *Efficacy of darunavir 800 mg once daily co-administered with 100 mg ritonavir once daily in ART-naïve patients*

The evidence of efficacy of darunavir/ritonavir 800/100 mg once daily is based on the analyses of 192 week data from the randomised, controlled, open-label Phase III trial ARTEMIS in antiretroviral treatment-naïve HIV-1 infected patients comparing darunavir/ritonavir 800/100 mg once daily with lopinavir/ritonavir 800/200 mg per day (given as a twice-daily or as a once-daily regimen). Both arms used a fixed background regimen consisting of tenofovir disoproxil fumarate 300 mg once daily and emtricitabine 200 mg once daily.

The table below shows the efficacy data of the 48 week and 96 week analyses from the ARTEMIS trial:

ARTEMIS						
	Week 48 <sup>a</sup>			Week 96 <sup>b</sup>		
Outcomes	Darunavir/ ritonavir 800/100 mg once daily N=343	Lopinavir/ ritonavir 800/200 mg per day N=346	Treatment difference (95% CI of difference)	Darunavir/ ritonavir 800/100 mg once daily N=343	Lopinavir/ ritonavir 800/200 mg per day N=346	Treatment difference (95% CI of difference)
HIV-1 RNA < 50 copies/ml <sup>c</sup> All patients	83.7% (287)	78.3% (271)	5.3% (-0.5; 11.2) <sup>d</sup>	79.0% (271)	70.8% (245)	8.2% (1.7; 14.7) <sup>d</sup>
With baseline HIV-RNA < 100,000	85.8% (194/226)	84.5% (191/226)	1.3% (-5.2; 7.9) <sup>d</sup>	80.5% (182/226)	75.2% (170/226)	5.3% (-2.3; 13.0) <sup>d</sup>
With baseline HIV-RNA ≥ 100,000	79.5% (93/117)	66.7% (80/120)	12.8% (1.6; 24.1) <sup>d</sup>	76.1% (89/117)	62.5% (75/120)	13.6% (1.9; 25.3) <sup>d</sup>
With baseline CD4+ cell count < 200	79.4% (112/141)	70.3% (104/148)	9.2% (-0.8; 19.2) <sup>d</sup>	78.7% (111/141)	64.9% (96/148)	13.9% (3.5; 24.2) <sup>d</sup>
With baseline CD4+ cell count ≥ 200	86.6% (175/202)	84.3% (167/198)	2.3% (-4.6; 9.2) <sup>d</sup>	79.2% (160/202)	75.3% (149/198)	4.0% (-4.3; 12.2) <sup>d</sup>
median CD4+ cell count change from baseline (x 10 <sup>6</sup> /L) <sup>e</sup>	137	141		171	188	

<sup>a</sup> Data based on analyses at week 48

<sup>b</sup> Data based on analyses at week 96

<sup>c</sup> Imputations according to the TLOVR algorithm

<sup>d</sup> Based on normal approximation to the difference in % response

<sup>e</sup> Non-completer is failure imputation: patients who discontinued prematurely are imputed with a change equal to 0

Non-inferiority in virologic response to the darunavir/ritonavir treatment, defined as the percentage of patients with plasma HIV-1 RNA level < 50 copies/ml, was demonstrated (at the pre-defined 12% non-inferiority margin) for both Intent-To-Treat (ITT) and On Protocol (OP) populations in the 48 week analysis. These results were confirmed in the analyses of data at 96 weeks of treatment in the ARTEMIS trial. These results were sustained up to 192 weeks of treatment in the ARTEMIS trial.

#### Efficacy of darunavir 800 mg once daily co-administered with 100 mg ritonavir once daily in ART-experienced patients

ODIN is a Phase III, randomised, open-label trial comparing darunavir/ritonavir 800/100 mg once daily versus darunavir/ritonavir 600/100 mg twice daily in ART-experienced HIV-1 infected patients with screening genotype resistance testing showing no darunavir RAMs (i.e. V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V, L89V) and a screening HIV-1 RNA > 1,000 copies/ml.

Efficacy analysis is based on 48 weeks of treatment (see table below). Both arms used an optimised background regimen (OBR) of ≥ 2 NRTIs.

ODIN			
Outcomes	Darunavir/ritonavir 800/100 mg once daily + OBR N=294	Darunavir/ritonavir 600/100 mg twice daily + OBR N=296	Treatment difference (95% CI of difference)
HIV-1 RNA < 50 copies/ml <sup>a</sup>	72.1% (212)	70.9% (210)	1.2% (-6.1; 8.5) <sup>b</sup>
With Baseline HIV-1 RNA (copies/ml)			
< 100,000	77.6% (198/255)	73.2% (194/265)	4.4% (-3.0; 11.9)
≥ 100,000	35.9% (14/39)	51.6% (16/31)	-15.7% (-39.2; 7.7)
With Baseline CD4+ cell count (x 10 <sup>6</sup> /L)			
≥ 100	75.1% (184/245)	72.5% (187/258)	2.6% (-5.1; 10.3)
< 100	57.1% (28/49)	60.5% (23/38)	-3.4% (-24.5; 17.8)
With HIV-1 clade			
Type B	70.4% (126/179)	64.3% (128/199)	6.1% (-3.4; 15.6)
Type AE	90.5% (38/42)	91.2% (31/34)	-0.7% (-14.0; 12.6)
Type C	72.7% (32/44)	78.8% (26/33)	-6.1% (-2.6; 13.7)
Other <sup>c</sup>	55.2% (16/29)	83.3% (25/30)	-28.2% (-51.0; -5.3)
mean CD4+ cell count change from baseline (x 10 <sup>6</sup> /L) <sup>e</sup>	108	112	-5 <sup>d</sup> (-25; 16)

<sup>a</sup> Imputations according to the TLOVR algorithm

<sup>b</sup> Based on a normal approximation of the difference in % response

<sup>c</sup> Clades A1, D, F1, G, K, CRF02\_AG, CRF12\_BF, and CRF06\_CPX

<sup>d</sup> Difference in means

<sup>e</sup> Last Observation Carried Forward imputation

At 48 weeks, virologic response, defined as the percentage of patients with plasma HIV-1 RNA level < 50 copies/ml, with darunavir/ritonavir 800/100 mg once daily treatment was demonstrated to be non-inferior (at the pre-defined 12% non-inferiority margin) compared to darunavir/ritonavir 600/100 mg twice daily for both ITT and OP populations.

Darunavir/ritonavir 800/100 mg once daily in ART-experienced patients should not be used in patients with one or more darunavir resistance associated mutations (DRV-RAMs) or HIV-1 RNA ≥ 100,000 copies/ml or CD4+ cell count < 100 cells x 10<sup>6</sup>/L (see section 4.2 and 4.4). Limited data is available in patients with HIV-1 clades other than B.

### Paediatric patients

ART-naïve paediatric patients from the age of 12 years to < 18 years, and weighing at least 40 kg DIONE is an open-label, Phase II trial evaluating the pharmacokinetics, safety, tolerability, and efficacy of darunavir with low dose ritonavir in 12 ART-naïve HIV-1 infected paediatric patients aged 12 to less than 18 years and weighing at least 40 kg. These patients received darunavir/ritonavir 800/100 mg once daily in combination with other antiretroviral agents. Virologic response was defined as a decrease in plasma HIV-1 RNA viral load of at least 1.0 log<sub>10</sub> versus baseline.

DIONE	
Outcomes at week 48	Darunavir/ritonavir N=12
HIV-1 RNA < 50 copies/ml <sup>a</sup>	83.3% (10)
CD4+ percent change from baseline <sup>b</sup>	14

DIONE	
<i>Outcomes at week 48</i>	Darunavir/ritonavir N=12
CD4+ cell count mean change from baseline <sup>b</sup>	221
≥ 1.0 log <sub>10</sub> decrease from baseline in plasma viral load	100%

<sup>a</sup> Imputations according to the TLOVR algorithm.

<sup>b</sup> Non-completer is failure imputation: patients who discontinued prematurely are imputed with a change equal to 0.

In the open-label, Phase II/III trial GS-US-216-0128, the efficacy, safety, and pharmacokinetics of darunavir 800 mg and cobicistat 150 mg (administered as separate tablets) and at least 2 NRTIs were evaluated in 7 HIV-1 infected, treatment-experienced, virologically suppressed adolescents weighing at least 40 kg. Patients were on a stable antiretroviral regimen (for at least 3 months), consisting of darunavir administered with ritonavir, combined with 2 NRTIs. They were switched from ritonavir to cobicistat 150 mg once daily and continued darunavir (N=7) and 2 NRTIs.

Virologic outcome in ART-experienced, virologically suppressed adolescents at week 48	
GS-US-216-0128	
Outcomes at Week 48	Darunavir/cobicistat + at least 2 NRTIs (N=7)
HIV-1 RNA < 50 copies/mL per FDA Snapshot Approach	85.7% (6)
CD4+ percent median change from baseline <sup>a</sup>	-6.1%
CD4+ cell count median change from baseline <sup>a</sup>	-342 cells/mm <sup>3</sup>

<sup>a</sup> No imputation (observed data).

For additional clinical study results in ART-experienced adults and paediatric patients, refer to the Summary of Product Characteristics for Darunavir Mylan 75 mg, 150 mg, 300 mg or 600 mg tablets.

### *Pregnancy and postpartum*

Darunavir/ritonavir (600/100 mg twice daily or 800/100 mg once daily) in combination with a background regimen was evaluated in a clinical trial of 36 pregnant women (18 in each arm) during the second and third trimesters, and postpartum. Virologic response was preserved throughout the study period in both arms. No mother to child transmission occurred in the infants born to the 31 subjects who stayed on the antiretroviral treatment through delivery. There were no new clinically relevant safety findings compared with the known safety profile of darunavir/ritonavir in HIV-1 infected adults (see sections 4.2, 4.4 and 5.2).

## **5.2 Pharmacokinetic properties**

The pharmacokinetic properties of darunavir, co-administered with cobicistat or ritonavir, have been evaluated in healthy adult volunteers and in HIV-1 infected patients. Exposure to darunavir was higher in HIV-1 infected patients than in healthy subjects. The increased exposure to darunavir in HIV-1 infected patients compared to healthy subjects may be explained by the higher concentrations of  $\alpha_1$ -acid glycoprotein (AAG) in HIV-1 infected patients, resulting in higher darunavir binding to plasma AAG and, therefore, higher plasma concentrations.

Darunavir is primarily metabolised by CYP3A. Cobicistat and ritonavir inhibit CYP3A, thereby increasing the plasma concentrations of darunavir considerably.

For information on cobicistat pharmacokinetic properties, consult the cobicistat Summary of Product Characteristics.

## Absorption

Darunavir was rapidly absorbed following oral administration. Maximum plasma concentration of darunavir in the presence of low dose ritonavir is generally achieved within 2.5-4.0 hours.

The absolute oral bioavailability of a single 600 mg dose of darunavir alone was approximately 37% and increased to approximately 82% in the presence of 100 mg twice daily ritonavir. The overall pharmacokinetic enhancement effect by ritonavir was an approximate 14-fold increase in the systemic exposure of darunavir when a single dose of 600 mg darunavir was given orally in combination with ritonavir at 100 mg twice daily (see section 4.4).

When administered without food, the relative bioavailability of darunavir in the presence of cobicistat or low dose ritonavir is lower as compared to intake with food. Therefore, darunavir tablets should be taken with cobicistat or ritonavir and with food. The type of food does not affect exposure to darunavir.

## Distribution

Darunavir is approximately 95% bound to plasma protein. Darunavir binds primarily to plasma  $\alpha_1$ -acid glycoprotein.

Following intravenous administration, the volume of distribution of darunavir alone was  $88.1 \pm 59.0$  l (Mean  $\pm$  SD) and increased to  $131 \pm 49.9$  l (Mean  $\pm$  SD) in the presence of 100 mg twice-daily ritonavir.

## Biotransformation

*In vitro* experiments with human liver microsomes (HLMs) indicate that darunavir primarily undergoes oxidative metabolism. Darunavir is extensively metabolised by the hepatic CYP system and almost exclusively by isozyme CYP3A4. A  $^{14}\text{C}$ -darunavir trial in healthy volunteers showed that a majority of the radioactivity in plasma after a single 400/100 mg darunavir with ritonavir dose was due to the parent active substance. At least 3 oxidative metabolites of darunavir have been identified in humans; all showed activity that was at least 10-fold less than the activity of darunavir against wild type HIV.

## Elimination

After a 400/100 mg  $^{14}\text{C}$ -darunavir with ritonavir dose, approximately 79.5% and 13.9% of the administered dose of  $^{14}\text{C}$ -darunavir could be retrieved in faeces and urine, respectively. Unchanged darunavir accounted for approximately 41.2% and 7.7% of the administered dose in faeces and urine, respectively. The terminal elimination half-life of darunavir was approximately 15 hours when combined with ritonavir.

The intravenous clearance of darunavir alone (150 mg) and in the presence of low dose ritonavir was 32.8 l/h and 5.9 l/h, respectively.

## Special populations

### *Paediatric population*

The pharmacokinetics of darunavir in combination with ritonavir taken twice daily in 74 treatment-experienced paediatric patients, aged 6 to 17 years and weighing at least 20 kg, showed that the administered weight-based doses of darunavir/ritonavir resulted in darunavir exposure comparable to that in adults receiving darunavir/ritonavir 600/100 mg twice daily (see section 4.2).

The pharmacokinetics of darunavir in combination with ritonavir taken twice daily in 14 treatment-experienced paediatric patients, aged 3 to < 6 years and weighing at least 15 kg to < 20 kg, showed that weight-based dosages resulted in darunavir exposure that was comparable to that achieved in adults receiving darunavir/ritonavir 600/100 mg twice daily (see section 4.2).

The pharmacokinetics of darunavir in combination with ritonavir taken once daily in 12 ART-naïve paediatric patients, aged 12 to < 18 years and weighing at least 40 kg, showed that darunavir/ritonavir 800/100 mg once daily results in darunavir exposure that was comparable to that achieved in adults receiving darunavir/ritonavir 800/100 mg once daily. Therefore the same once daily dosage may be used in treatment-experienced adolescents aged 12 to < 18 years and weighing at least 40 kg without darunavir resistance associated mutations (DRV-RAMs)\* and who have plasma HIV-1 RNA < 100,000 copies/ml and CD4+ cell count  $\geq 100$  cells  $\times 10^6/L$  (see section 4.2).

\* DRV-RAMs: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

The pharmacokinetics of darunavir in combination with ritonavir taken once daily in 10 treatment-experienced paediatric patients, aged 3 to < 6 years and weighing at least 14 kg to < 20 kg, showed that weight-based dosages resulted in darunavir exposure that was comparable to that achieved in adults receiving darunavir/ritonavir 800/100 mg once daily (see section 4.2). In addition, pharmacokinetic modeling and simulation of darunavir exposures in paediatric patients across the ages of 3 to < 18 years confirmed the darunavir exposures as observed in the clinical studies and allowed the identification of weight-based darunavir/ritonavir once daily dosing regimens for paediatric patients weighing at least 15 kg that are either ART-naïve or treatment-experienced paediatric patients without DRV-RAMs\* and who have plasma HIV-1 RNA < 100,000 copies/ml and CD4+ cell count  $\geq 100$  cells  $\times 10^6/L$  (see section 4.2).

\* DRV-RAMs: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V and L89V

The pharmacokinetics of darunavir 800 mg co-administered with cobicistat 150 mg in paediatric patients have been studied in 7 adolescents aged 12 to less than 18 years, weighing at least 40 kg in Study GS-US-216-0128. The geometric mean adolescent exposure ( $AUC_{tau}$ ) was similar for darunavir and increased 19% for cobicistat compared to exposures achieved in adults who received darunavir 800 mg co-administered with cobicistat 150 mg in Study GS-US-216-0130. The difference observed for cobicistat was not considered clinically relevant.

	<b>Adults in Study GS-US-216-0130, week 24 (Reference)<sup>a</sup> Mean (%CV) GLSM</b>	<b>Adolescents in Study GS-US-216-0128, day 10 (Test)<sup>b</sup> Mean (%CV) GLSM</b>	<b>GLSM Ratio (90% CI) (Test/Reference)</b>
N	60 <sup>c</sup>	7	
<b>DRV PK Parameter</b>			
$AUC_{tau}$ (h.ng/mL) <sup>d</sup>	81,646 (32.2) 77,534	80,877 (29.5) 77,217	1.00 (0.79-1.26)
$C_{max}$ (ng/mL)	7,663 (25.1) 7,422	7,506 (21.7) 7,319	0.99 (0.83-1.17)
$C_{tau}$ (ng/mL) <sup>d</sup>	1,311 (74.0) 947	1,087 (91.6) 676	0.71 (0.34-1.48)
<b>COBI PK Parameter</b>			
$AUC_{tau}$ (h.ng/mL) <sup>d</sup>	7,596 (48.1) 7,022	8,741 (34.9) 8,330	1.19 (0.95-1.48)
$C_{max}$ (ng/mL)	991 (33.4) 945	1,116 (20.0) 1,095	1.16 (1.00-1.35)
$C_{tau}$ (ng/mL) <sup>d</sup>	32.8 (289.4) 17.2 <sup>e</sup>	28.3 (157.2) 22.0 <sup>e</sup>	1.28 (0.51-3.22)

<sup>a</sup> Week 24 intensive PK data from subjects who received DRV 800 mg + COBI 150 mg.

<sup>b</sup> Day 10 intensive PK data from subjects who received DRV 800 mg + COBI 150 mg.

<sup>c</sup> N=59 for  $AUC_{tau}$  and  $C_{tau}$ .

<sup>d</sup> Concentration at predose (0 hours) was used as surrogate for concentration at 24 hours for the purposes of estimating  $AUC_{tau}$  and  $C_{tau}$  in Study GS-US-216-0128.

<sup>e</sup> N=57 and N=5 for GLSM of  $C_{\text{tau}}$  in Study GS-US-216-0130 and Study GS-US-216-0128, respectively.

### *Elderly*

Population pharmacokinetic analysis in HIV infected patients showed that darunavir pharmacokinetics are not considerably different in the age range (18 to 75 years) evaluated in HIV infected patients (n=12, age  $\geq 65$ ) (see section 4.4). However, only limited data were available in patients above the age of 65 year.

### *Gender*

Population pharmacokinetic analysis showed a slightly higher darunavir exposure (16.8%) in HIV infected females compared to males. This difference is not clinically relevant.

### *Renal impairment*

Results from a mass balance study with  $^{14}\text{C}$ -darunavir with ritonavir showed that approximately 7.7% of the administered dose of darunavir is excreted in the urine unchanged.

Although darunavir has not been studied in patients with renal impairment, population pharmacokinetic analysis showed that the pharmacokinetics of darunavir were not significantly affected in HIV infected patients with moderate renal impairment (CrCl between 30-60 ml/min, n=20) (see sections 4.2 and 4.4).

### *Hepatic impairment*

Darunavir is primarily metabolised and eliminated by the liver. In a multiple dose study with darunavir co-administered with ritonavir (600/100 mg) twice daily, it was demonstrated that the total plasma concentrations of darunavir in subjects with mild (Child-Pugh Class A, n=8) and moderate (Child-Pugh Class B, n=8) hepatic impairment were comparable with those in healthy subjects. However, unbound darunavir concentrations were approximately 55% (Child-Pugh Class A) and 100% (Child-Pugh Class B) higher, respectively. The clinical relevance of this increase is unknown therefore, darunavir should be used with caution. The effect of severe hepatic impairment on the pharmacokinetics of darunavir has not been studied (see sections 4.2, 4.3 and 4.4).

### *Pregnancy and postpartum*

The exposure to total darunavir and ritonavir after intake of darunavir/ritonavir 600/100 mg twice daily and darunavir/ritonavir 800/100 mg once daily as part of an antiretroviral regimen was generally lower during pregnancy compared with postpartum. However, for unbound (i.e. active) darunavir, the pharmacokinetic parameters were less reduced during pregnancy compared to postpartum, due to an increase in the unbound fraction of darunavir during pregnancy compared to postpartum.

<b>Pharmacokinetic results of total darunavir after administration of darunavir/ritonavir at 600/100 mg twice daily as part of an antiretroviral regimen, during the second trimester of pregnancy, the third trimester of pregnancy and postpartum</b>			
<b>Pharmacokinetics of total darunavir (mean <math>\pm</math> SD)</b>	<b>Second trimester of pregnancy (n=12)<sup>a</sup></b>	<b>Third trimester of pregnancy (n=12)</b>	<b>Postpartum (6-12 weeks) (n=12)</b>
$C_{\text{max}}$ , ng/ml	4,668 $\pm$ 1,097	5,328 $\pm$ 1,631	6,659 $\pm$ 2,364
AUC <sub>12h</sub> , ng.h/ml	39,370 $\pm$ 9,597	45,880 $\pm$ 17,360	56,890 $\pm$ 26,340
$C_{\text{min}}$ , ng/ml	1,922 $\pm$ 825	2,661 $\pm$ 1,269	2,851 $\pm$ 2,216

<sup>a</sup> n=11 for AUC<sub>12h</sub>



<b>Pharmacokinetic results of total darunavir after administration of darunavir/ritonavir at 800/100 mg once daily as part of an antiretroviral regimen, during the second trimester of pregnancy, the third trimester of pregnancy and postpartum</b>			
<b>Pharmacokinetics of total darunavir (mean ± SD)</b>	<b>Second trimester of pregnancy (n=17)</b>	<b>Third Trimester of pregnancy (n=15)</b>	<b>Postpartum (6-12 weeks) (n=16)</b>
$C_{max}$ , ng/ml	4,964 ± 1,505	5,132 ± 1,198	7,310 ± 1,704
$AUC_{24h}$ , ng.h/ml	62,289 ± 16,234	61,112 ± 13,790	92,116 ± 29,241
$C_{min}$ , ng/ml	1,248 ± 542	1,075 ± 594	1,473 ± 1,141

In women receiving darunavir/ritonavir 600/100 mg twice daily during the second trimester of pregnancy, mean intra-individual values for total darunavir  $C_{max}$ ,  $AUC_{12h}$  and  $C_{min}$  were 28%, 26% and 26% lower, respectively, as compared with postpartum; during the third trimester of pregnancy, total darunavir  $C_{max}$ ,  $AUC_{12h}$  and  $C_{min}$  values were 18%, 16% lower and 2% higher, respectively, as compared with postpartum.

In women receiving darunavir/ritonavir 800/100 mg once daily during the second trimester of pregnancy, mean intra-individual values for total darunavir  $C_{max}$ ,  $AUC_{24h}$  and  $C_{min}$  were 33%, 31% and 30% lower, respectively, as compared with postpartum; during the third trimester of pregnancy, total darunavir  $C_{max}$ ,  $AUC_{24h}$  and  $C_{min}$  values were 29%, 32% and 50% lower, respectively, as compared with postpartum.

Treatment with darunavir/cobicistat 800/150 mg once daily during pregnancy results in low darunavir exposure. In women receiving darunavir/cobicistat during the second trimester of pregnancy, mean intra-individual values for total darunavir  $C_{max}$ ,  $AUC_{24h}$  and  $C_{min}$  were 49%, 56% and 92% lower, respectively, as compared with postpartum; during the third trimester of pregnancy, total darunavir  $C_{max}$ ,  $AUC_{24h}$  and  $C_{min}$  values were 37%, 50% and 89% lower, respectively, as compared with postpartum. The unbound fraction was also substantially reduced, including around 90% reductions of  $C_{min}$  levels. The main cause of these low exposures is a marked reduction in cobicistat exposure as a consequence of pregnancy-associated enzyme induction (see below).

<b>Pharmacokinetic results of total darunavir after administration of darunavir/cobicistat 800/150 mg once daily as part of an antiretroviral regimen, during the second trimester of pregnancy, the third trimester of pregnancy, and postpartum</b>			
<b>Pharmacokinetics of total darunavir (mean ± SD)</b>	<b>Second trimester of pregnancy (n=7)</b>	<b>Third trimester of pregnancy (n=6)</b>	<b>Postpartum (6-12 weeks) (n=6)</b>
$C_{max}$ , ng/mL	4,340 ± 1,616	4,910 ± 970	7,918 ± 2,199
$AUC_{24h}$ , ng.h/mL	47,293 ± 19,058	47,991 ± 9,879	99,613 ± 34,862
$C_{min}$ , ng/mL	168 ± 149	184 ± 99	1,538 ± 1,344

The exposure to cobicistat was lower during pregnancy, potentially leading to suboptimal boosting of darunavir. During the second trimester of pregnancy, cobicistat  $C_{max}$ ,  $AUC_{24h}$ , and  $C_{min}$  were 50%, 63%, and 83% lower, respectively, as compared with postpartum. During the third trimester of pregnancy, cobicistat  $C_{max}$ ,  $AUC_{24h}$ , and  $C_{min}$ , were 27%, 49%, and 83% lower, respectively, as compared with postpartum.

### 5.3 Preclinical safety data

Animal toxicology studies have been conducted at exposures up to clinical exposure levels with darunavir alone, in mice, rats and dogs and in combination with ritonavir in rats and dogs.

In repeated-dose toxicology studies in mice, rats and dogs, there were only limited effects of treatment with darunavir. In rodents the target organs identified were the haematopoietic system, the blood coagulation system, liver and thyroid. A variable but limited decrease in red blood cell-related parameters was observed, together with increases in activated partial thromboplastin time.

Changes were observed in liver (hepatocyte hypertrophy, vacuolation, increased liver enzymes) and thyroid (follicular hypertrophy). In the rat, the combination of darunavir with ritonavir lead to a small increase in effect on RBC parameters, liver and thyroid and increased incidence of islet fibrosis in the pancreas (in male rats only) compared to treatment with darunavir alone. In the dog, no major toxicity findings or target organs were identified up to exposures equivalent to clinical exposure at the recommended dose.

In a study conducted in rats, the number of corpora lutea and implantations were decreased in the presence of maternal toxicity. Otherwise, there were no effects on mating or fertility with darunavir treatment up to 1,000 mg/kg/day and exposure levels below (AUC-0.5 fold) of that in human at the clinically recommended dose. Up to same dose levels, there was no teratogenicity with darunavir in rats and rabbits when treated alone nor in mice when treated in combination with ritonavir. The exposure levels were lower than those with the recommended clinical dose in humans. In a pre- and postnatal development assessment in rats, darunavir with and without ritonavir, caused a transient reduction in body weight gain of the offspring pre-weaning and there was a slight delay in the opening of eyes and ears. Darunavir in combination with ritonavir caused a reduction in the number of pups that exhibited the startle response on day 15 of lactation and a reduced pup survival during lactation. These effects may be secondary to pup exposure to the active substance via the milk and/or maternal toxicity. No post weaning functions were affected with darunavir alone or in combination with ritonavir. In juvenile rats receiving darunavir up to days 23-26, increased mortality was observed with convulsions in some animals. Exposure in plasma, liver and brain was considerably higher than in adult rats after comparable doses in mg/kg between days 5 and 11 of age. After day 23 of life, the exposure was comparable to that in adult rats. The increased exposure was likely at least partly due to immaturity of the drug-metabolising enzymes in juvenile animals. No treatment related mortalities were noted in juvenile rats dosed at 1,000 mg/kg darunavir (single dose) on day 26 of age or at 500 mg/kg (repeated dose) from day 23 to 50 of age, and the exposures and toxicity profile were comparable to those observed in adult rats.

Due to uncertainties regarding the rate of development of the human blood brain barrier and liver enzymes, darunavir with low dose ritonavir should not be used in paediatric patients below 3 years of age.

Darunavir was evaluated for carcinogenic potential by oral gavage administration to mice and rats up to 104 weeks. Daily doses of 150, 450 and 1,000 mg/kg were administered to mice and doses of 50, 150 and 500 mg/kg were administered to rats. Dose-related increases in the incidences of hepatocellular adenomas and carcinomas were observed in males and females of both species. Thyroid follicular cell adenomas were noted in male rats. Administration of darunavir did not cause a statistically significant increase in the incidence of any other benign or malignant neoplasm in mice or rats. The observed hepatocellular and thyroid tumours in rodents are considered to be of limited relevance to humans. Repeated administration of darunavir to rats caused hepatic microsomal enzyme induction and increased thyroid hormone elimination, which predispose rats, but not humans, to thyroid neoplasms. At the highest tested doses, the systemic exposures (based on AUC) to darunavir were between 0.4- and 0.7-fold (mice) and 0.7- and 1-fold (rats), relative to those observed in humans at the recommended therapeutic doses.

After 2 years administration of darunavir at exposures at or below the human exposure, kidney changes were observed in mice (nephrosis) and rats (chronic progressive nephropathy).

Darunavir was not mutagenic or genotoxic in a battery of *in vitro* and *in vivo* assays including bacterial reverse mutation (Ames), chromosomal aberration in human lymphocytes and *in vivo* micronucleus test in mice.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Silica, colloidal anhydrous  
Cellulose microcrystalline  
Crospovidone  
Sodium starch glycolate  
Hypromellose  
Magnesium stearate

#### Tablet film-coat

Polyvinyl alcohol partially hydrolysed  
Titanium dioxide (E171)  
Macrogol  
Talc

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

#### Darunavir Mylan 400 mg film-coated tablets

3 years  
In-use shelf life after first opening HDPE bottle: 100 days.

#### Darunavir Mylan 800 mg film-coated tablets

3 years  
In-use shelf life after first opening HDPE bottle: 90 days.

### **6.4 Special precautions for storage**

#### Darunavir Mylan 400 mg film-coated tablets

This medicinal product does not require any special storage conditions.

#### Darunavir Mylan 800 mg film-coated tablets

PVC/PE/PVDC-Al blister pack  
Do not store above 25°C.

Cold form PVC/Al/OPA-Al blister pack  
This medicinal product does not require any special storage conditions.

HDPE bottle pack  
This medicinal product does not require any special storage conditions.

## **6.5 Nature and contents of container**

### Darunavir Mylan 400 mg film-coated tablets

PVC/PE/PVDC-Al blister pack containing 30 and 60 tablets and 60x1 tablets.  
Cold form PVC/Al/OPA-Al blister pack containing 30 and 60 tablets and 60x1 tablets.  
HDPE bottle pack with a PP screw cap containing 60 and 100 tablets.

### Darunavir Mylan 800 mg film-coated tablets

PVC/PE/PVDC-Al blister pack containing 30 tablets and 30x1 tablets.  
Cold form PVC/Al/OPA-Al blister pack containing 30 tablets and 30x1 tablets.  
HDPE bottle pack with a PP screw cap containing 30, 60, 90 tablets.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal and other handling**

No special requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

## **8. MARKETING AUTHORISATION NUMBER(S)**

### Darunavir Mylan 400 mg film-coated tablets

EU/1/16/1140/022  
EU/1/16/1140/023  
EU/1/16/1140/024  
EU/1/16/1140/025  
EU/1/16/1140/026  
EU/1/16/1140/027  
EU/1/16/1140/028  
EU/1/16/1140/029

### Darunavir Mylan 800 mg film-coated tablets

EU/1/16/1140/039  
EU/1/16/1140/040  
EU/1/16/1140/041  
EU/1/16/1140/042  
EU/1/16/1140/043  
EU/1/16/1140/044  
EU/1/16/1140/045

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 04 January 2017

Date of latest renewal: 16 September 2021

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## **ANNEX II**

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**

### Name and address of the manufacturers responsible for batch release

Mylan Hungary Kft  
Mylan utca 1, Komárom, 2900,  
Hungary

McDermott Laboratories Limited trading as Gerard Laboratories  
35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13  
Ireland

Mylan Germany GmbH  
Zweigniederlassung Bad Homburg v. d. Hoehe, Benzstrasse 1  
Bad Homburg v. d. Hoehe  
Hessen, 61352  
Germany

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription.

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**



## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON (BLISTERS AND BOTTLES)**

**1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 75 mg film-coated tablets  
darunavir

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 75 mg of darunavir.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet

Blister carton

480 film-coated tablets

480 x 1 film-coated tablets

Bottle carton

480 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

For oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

&lt;Bottles Only&gt; Once opened, use within 100 days

[On carton only:]

Open date: \_\_\_\_\_

**9. SPECIAL STORAGE CONDITIONS****10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1140/001  
EU/1/16/1140/002  
EU/1/16/1140/003  
EU/1/16/1140/004  
EU/1/16/1140/005

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Darunavir Mylan 75 mg

<b>17. UNIQUE IDENTIFIER – 2D BARCODE</b>
-------------------------------------------

2D barcode carrying the unique identifier included.

<b>18. UNIQUE IDENTIFIER – HUMAN READABLE DATA</b>
----------------------------------------------------

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING****BOTTLE LABEL****1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 75 mg film-coated tablets  
darunavir

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 75 mg of darunavir.

**3. LIST OF EXCIPIENTS****4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet  
480 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

For oral use.  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP  
Once opened, use within 100 days

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1140/005

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

<b>MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS</b>
------------------------------------------------------------

<b>BLISTERS</b>
-----------------

<b>1. NAME OF THE MEDICINAL PRODUCT</b>
-----------------------------------------

Darunavir Mylan 75 mg film-coated tablets  
darunavir

<b>2. NAME OF THE MARKETING AUTHORISATION HOLDER</b>
------------------------------------------------------

Mylan Pharmaceuticals Limited

<b>3. EXPIRY DATE</b>
-----------------------

EXP

<b>4. BATCH NUMBER</b>
------------------------

Lot

<b>5. OTHER</b>
-----------------

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON (BLISTERS AND BOTTLES)**

**1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 150 mg film-coated tablets  
darunavir

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 150 mg of darunavir.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet

Blister carton

240 film-coated tablets

240 x 1 film-coated tablets

Bottle carton

60 film-coated tablets

240 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

For oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**



**8. EXPIRY DATE**

EXP

&lt;Bottles Only&gt; Once opened, use within 100 days

[On carton only:]

Open date: \_\_\_\_\_

**9. SPECIAL STORAGE CONDITIONS****10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1140/006  
EU/1/16/1140/007  
EU/1/16/1140/008  
EU/1/16/1140/009  
EU/1/16/1140/010  
EU/1/16/1140/011

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Darunavir Mylan 150 mg

<b>17. UNIQUE IDENTIFIER – 2D BARCODE</b>
-------------------------------------------

2D barcode carrying the unique identifier included.

<b>18. UNIQUE IDENTIFIER – HUMAN READABLE DATA</b>
----------------------------------------------------

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING****BOTTLE LABEL****1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 150 mg film-coated tablets  
darunavir

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 150 mg of darunavir.

**3. LIST OF EXCIPIENTS****4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet

60 film-coated tablets

240 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

For oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

Once opened, use within 100 days

**9. SPECIAL STORAGE CONDITIONS**

<b>10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE</b>
--------------------------------------------------------------------------------------------------------------------------------------------------

<b>11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER</b>
-------------------------------------------------------------------

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

<b>12. MARKETING AUTHORISATION NUMBERS</b>
--------------------------------------------

EU/1/16/1140/010  
EU/1/16/1140/011

<b>13. BATCH NUMBER</b>
-------------------------

Lot

<b>14. GENERAL CLASSIFICATION FOR SUPPLY</b>
----------------------------------------------

<b>15. INSTRUCTIONS ON USE</b>
--------------------------------

<b>16. INFORMATION IN BRAILLE</b>
-----------------------------------

<b>17. UNIQUE IDENTIFIER – 2D BARCODE</b>
-------------------------------------------

<b>18. UNIQUE IDENTIFIER – HUMAN READABLE DATA</b>
----------------------------------------------------

<b>MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS</b>
------------------------------------------------------------

<b>BLISTERS</b>
-----------------

<b>1. NAME OF THE MEDICINAL PRODUCT</b>
-----------------------------------------

Darunavir Mylan 150 mg film-coated tablets  
darunavir

<b>2. NAME OF THE MARKETING AUTHORISATION HOLDER</b>
------------------------------------------------------

Mylan Pharmaceuticals Limited

<b>3. EXPIRY DATE</b>
-----------------------

EXP

<b>4. BATCH NUMBER</b>
------------------------

Lot

<b>5. OTHER</b>
-----------------

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON (BLISTERS AND BOTTLES)**

**1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 300 mg film-coated tablets  
darunavir

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 300 mg of darunavir.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet

Blister carton

30 film-coated tablets

60 film-coated tablets

120 film-coated tablets

120 x 1 film-coated tablets

Bottle carton

30 film-coated tablets

120 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

For oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

&lt;Bottles Only&gt; Once opened, use within 100 days

[On carton only:]

Open date: \_\_\_\_\_

**9. SPECIAL STORAGE CONDITIONS****10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1140/012  
EU/1/16/1140/013  
EU/1/16/1140/014  
EU/1/16/1140/015  
EU/1/16/1140/016  
EU/1/16/1140/017  
EU/1/16/1140/018  
EU/1/16/1140/019  
EU/1/16/1140/020  
EU/1/16/1140/021

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE**

<b>16. INFORMATION IN BRAILLE</b>
-----------------------------------

Darunavir Mylan 300 mg

<b>17. UNIQUE IDENTIFIER – 2D BARCODE</b>
-------------------------------------------

2D barcode carrying the unique identifier included.

<b>18. UNIQUE IDENTIFIER – HUMAN READABLE DATA</b>
----------------------------------------------------

PC  
SN  
NN



**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING****BOTTLE LABEL****1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 300 mg film-coated tablets  
darunavir

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 300 mg of darunavir.

**3. LIST OF EXCIPIENTS****4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet

30 film-coated tablets

120 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

For oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

Once opened, use within 100 days

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1140/020  
EU/1/16/1140/021

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

<b>MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS</b>
------------------------------------------------------------

<b>BLISTERS</b>
-----------------

<b>1. NAME OF THE MEDICINAL PRODUCT</b>
-----------------------------------------

Darunavir Mylan 300 mg film-coated tablets  
darunavir

<b>2. NAME OF THE MARKETING AUTHORISATION HOLDER</b>
------------------------------------------------------

Mylan Pharmaceuticals Limited

<b>3. EXPIRY DATE</b>
-----------------------

EXP

<b>4. BATCH NUMBER</b>
------------------------

Lot

<b>5. OTHER</b>
-----------------

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON (BLISTERS AND BOTTLES)**

**1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 400 mg film-coated tablets  
darunavir

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 400 mg of darunavir.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablets

Blister carton

30 film-coated tablets

60 film-coated tablets

60 x 1 film-coated tablets

Bottle carton

60 film-coated tablets

100 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

For oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

&lt;Bottles Only&gt; Once opened, use within 100 days

[On carton only:]

Open date: \_\_\_\_\_

**9. SPECIAL STORAGE CONDITIONS****10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1140/022  
EU/1/16/1140/023  
EU/1/16/1140/024  
EU/1/16/1140/025  
EU/1/16/1140/026  
EU/1/16/1140/027  
EU/1/16/1140/028  
EU/1/16/1140/029

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Darunavir Mylan 400 mg

<b>17. UNIQUE IDENTIFIER – 2D BARCODE</b>
-------------------------------------------

2D barcode carrying the unique identifier included.

<b>18. UNIQUE IDENTIFIER – HUMAN READABLE DATA</b>
----------------------------------------------------

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING****BOTTLE LABEL****1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 400 mg film-coated tablets  
darunavir

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 400 mg of darunavir.

**3. LIST OF EXCIPIENTS****4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet

60 film-coated tablets

100 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

For oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

Once opened, use within 100 days

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1140/028  
EU/1/16/1140/029

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**



<b>MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS</b>
------------------------------------------------------------

<b>BLISTERS</b>
-----------------

<b>1. NAME OF THE MEDICINAL PRODUCT</b>
-----------------------------------------

Darunavir Mylan 400 mg film-coated tablets  
darunavir

<b>2. NAME OF THE MARKETING AUTHORISATION HOLDER</b>
------------------------------------------------------

Mylan Pharmaceuticals Limited

<b>3. EXPIRY DATE</b>
-----------------------

EXP

<b>4. BATCH NUMBER</b>
------------------------

Lot

<b>5. OTHER</b>
-----------------

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****CARTON (BLISTERS AND BOTTLES)****1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 600 mg film-coated tablets  
darunavir

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 600 mg of darunavir.

**3. LIST OF EXCIPIENTS****4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet

Blister carton

30 film-coated tablets

60 film-coated tablets

60 x 1 film-coated tablets

Bottle carton

30 film-coated tablets

60 film-coated tablets

90 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

For oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

&lt;Bottles Only&gt; Once opened, use within 100 days

[On carton only:]

Open date: \_\_\_\_\_

**9. SPECIAL STORAGE CONDITIONS****10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1140/030  
EU/1/16/1140/031  
EU/1/16/1140/032  
EU/1/16/1140/033  
EU/1/16/1140/034  
EU/1/16/1140/035  
EU/1/16/1140/036  
EU/1/16/1140/037  
EU/1/16/1140/038

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Darunavir Mylan 600 mg

<b>17. UNIQUE IDENTIFIER – 2D BARCODE</b>
-------------------------------------------

2D barcode carrying the unique identifier included.

<b>18. UNIQUE IDENTIFIER – HUMAN READABLE DATA</b>
----------------------------------------------------

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING****BOTTLE LABEL****1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 600 mg film-coated tablets  
darunavir

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 600 mg of darunavir.

**3. LIST OF EXCIPIENTS****4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet

30 film-coated tablets

60 film-coated tablets

90 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

For oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

Once opened, use within 100 days

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1140/036  
EU/1/16/1140/037  
EU/1/16/1140/038

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

<b>MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS</b>
------------------------------------------------------------

<b>BLISTERS</b>
-----------------

<b>1. NAME OF THE MEDICINAL PRODUCT</b>
-----------------------------------------

Darunavir Mylan 600 mg film-coated tablets  
darunavir

<b>2. NAME OF THE MARKETING AUTHORISATION HOLDER</b>
------------------------------------------------------

Mylan Pharmaceuticals Limited

<b>3. EXPIRY DATE</b>
-----------------------

EXP

<b>4. BATCH NUMBER</b>
------------------------

Lot

<b>5. OTHER</b>
-----------------

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON (BLISTERS AND BOTTLES)**

**1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 800 mg film-coated tablets  
darunavir

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 800 mg of darunavir.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet

Blister carton

30 film-coated tablets

30 x 1 film-coated tablets

Bottle carton

30 film-coated tablets

60 film-coated tablets

90 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

For oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**



**8. EXPIRY DATE**

EXP

&lt;Bottles Only&gt; Once opened, use within 90 days

[On carton only:]

Open date: \_\_\_\_\_

**9. SPECIAL STORAGE CONDITIONS**

&lt;PVC/PE/PVDC-Al blister pack only&gt;

Do not store above 25°C.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1140/039

EU/1/16/1140/040

EU/1/16/1140/041

EU/1/16/1140/042

EU/1/16/1140/043

EU/1/16/1140/044

EU/1/16/1140/045

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Darunavir Mylan 800 mg

<b>17. UNIQUE IDENTIFIER – 2D BARCODE</b>
-------------------------------------------

2D barcode carrying the unique identifier included.

<b>18. UNIQUE IDENTIFIER – HUMAN READABLE DATA</b>
----------------------------------------------------

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING****BOTTLE LABEL****1. NAME OF THE MEDICINAL PRODUCT**

Darunavir Mylan 800 mg film-coated tablets  
darunavir

**2. STATEMENT OF ACTIVE SUBSTANCE**

Each film-coated tablet contains 800 mg of darunavir.

**3. LIST OF EXCIPIENTS****4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet

30 film-coated tablets

60 film-coated tablets

90 film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

For oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

Once opened, use within 90 days

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1140/043  
EU/1/16/1140/044  
EU/1/16/1140/045

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

<b>MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS</b>
------------------------------------------------------------

<b>BLISTERS</b>
-----------------

<b>1. NAME OF THE MEDICINAL PRODUCT</b>
-----------------------------------------

Darunavir Mylan 800 mg film-coated tablets  
darunavir

<b>2. NAME OF THE MARKETING AUTHORISATION HOLDER</b>
------------------------------------------------------

Mylan Pharmaceuticals Limited

<b>3. EXPIRY DATE</b>
-----------------------

EXP

<b>4. BATCH NUMBER</b>
------------------------

Lot

<b>5. OTHER</b>
-----------------

## **B. PACKAGE LEAFLET**

## Package leaflet: Information for the user

### Darunavir Mylan 75 mg film-coated tablets

darunavir

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Darunavir Mylan is and what it is used for
2. What you need to know before you take Darunavir Mylan
3. How to take Darunavir Mylan
4. Possible side effects
5. How to store Darunavir Mylan
6. Contents of the pack and other information

#### 1. What Darunavir Mylan is and what it is used for

##### What is Darunavir Mylan?

Darunavir Mylan contains the active substance darunavir. Darunavir is an antiretroviral medicine used in the treatment of Human Immunodeficiency Virus (HIV) infection. It belongs to a group of medicines called protease inhibitors. Darunavir works by reducing the amount of HIV in your body. This will improve your immune system and reduces the risk of developing illnesses linked to HIV infection.

##### What it is used for?

Darunavir is used to treat adults and children of 3 years of age and above, and at least 15 kilogram body weight who are infected by HIV and who have already used other antiretroviral medicines.

Darunavir must be taken in combination with a low dose of ritonavir and other anti-HIV medicines. Your doctor will discuss with you which combination of medicines is best for you.

#### 2. What you need to know before you take Darunavir Mylan

##### Do not take Darunavir Mylan

- if you are **allergic** to darunavir or any of the other ingredients of this medicine (listed in section 6) or to ritonavir.
- if you have **severe liver problems**. Ask your doctor if you are unsure about the severity of your liver disease. Some additional tests might be necessary.

##### Do not combine Darunavir Mylan with any of the following medicines

If you are taking any of these, ask your doctor about switching to another medicine.

Medicine	Purpose of the medicine
<i>Avanafil</i>	to treat erectile dysfunction
<i>Astemizole</i> or <i>terfenadine</i>	to treat allergy symptoms
<i>Triazolam</i> and <i>oral</i> (taken by mouth) <i>midazolam</i>	to help you sleep and/or relieve anxiety

Medicine	Purpose of the medicine
<i>Cisapride</i>	to treat some stomach conditions
<i>Colchicine</i> (if you have kidney and/or liver problems)	to treat gout or familial Mediterranean fever
<i>Lurasidone, pimozide, quetiapine</i> or <i>sertindole</i>	to treat psychiatric conditions
<i>Ergot alkaloids</i> like <i>ergotamine</i> , <i>dihydroergotamine</i> , <i>ergometrine</i> and <i>methylergonovine</i>	to treat migraine headaches
<i>Amiodarone</i> , <i>bepiridil</i> , <i>dronedarone</i> , <i>ivabradine</i> , <i>quinidine</i> , <i>ranolazine</i>	to treat certain heart disorders e.g. abnormal heart beat
<i>Lovastatin</i> <i>simvastatin</i> and <i>lomitapide</i>	to lower cholesterol levels
<i>Rifampicin</i>	to treat some infections such as tuberculosis
The combination product <i>lopinavir/ritonavir</i>	this anti-HIV medicine belongs to the same class as darunavir
<i>Elbasvir/grazoprevir</i>	to treat hepatitis C infection
<i>Alfuzosin</i>	to treat enlarged prostate
<i>Sildenafil</i>	to treat high blood pressure in the pulmonary circulation
<i>Ticagrelor</i>	to help stop the clumping of platelets in the treatment of patients with a history of a heart attack
<i>Naloxegol</i>	to treat opioid induced constipation
<i>Dapoxetine</i>	to treat premature ejaculation
<i>Domperidone</i>	to treat nausea and vomiting

Do not combine darunavir with products that contain St John's wort (*Hypericum perforatum*).

### Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking Darunavir Mylan.

Darunavir is not a cure for HIV infection.

People taking darunavir may still develop infections or other illnesses associated with HIV infection. You must keep in regular contact with your doctor.

People taking darunavir may develop a skin rash. Infrequently a rash may become severe or potentially life-threatening. Please contact your doctor whenever you develop a rash.

In patients taking darunavir and raltegravir (for HIV infection), rashes (generally mild or moderate) may occur more frequently than in patients taking either medicine separately.

### Tell your doctor about your situation BEFORE and DURING your treatment

Make sure that you check the following points and tell your doctor if any of these apply to you.

- Tell your doctor if you have had **problems with your liver** before, including hepatitis B or C infection. Your doctor may evaluate how severe your liver disease is before deciding if you can take darunavir.
- Tell your doctor if you have **diabetes**. Darunavir might increase sugar levels in the blood.
- Tell your doctor immediately if you notice any **symptoms of infection** (for example enlarged lymph nodes and fever). In some patients with advanced HIV infection and a history of opportunistic infection, signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started. It is believed that these symptoms are due to an improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms.
- In addition to the opportunistic infections, autoimmune disorders (a condition that occurs when the immune system attacks healthy body tissue) may also occur after you start taking medicines for the treatment of your HIV infection. Autoimmune disorders may occur many months after the start of treatment. If you notice any symptoms of infection or other symptoms such as



muscle weakness, weakness beginning in the hands and feet and moving up towards the trunk of the body, palpitations, tremor or hyperactivity, please inform your doctor immediately to seek necessary treatment.

- Tell your doctor if you have **haemophilia**. Darunavir might increase the risk of bleeding.
- Tell your doctor if you are **allergic to sulphonamides** (e.g. used to treat certain infections).
- Tell your doctor if you notice any **musculoskeletal problems**. Some patients taking combination antiretroviral therapy may develop a bone disease called osteonecrosis (death of bone tissue caused by loss of blood supply to the bone). The length of combination antiretroviral therapy, corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index, among others, may be some of the many risk factors for developing this disease. Signs of osteonecrosis are joint stiffness, aches and pains (especially of the hip, knee and shoulder) and difficulty in movement. If you notice any of these symptoms please inform your doctor.

### Elderly

Darunavir has only been used in limited numbers of patients 65 years or older. If you belong to this age group, please discuss with your doctor if you can use Darunavir Mylan.

### Children

Darunavir is not for use in children younger than 3 years of age or weighing less than 15 kilograms.

### Other medicines and Darunavir Mylan

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines.

There are some medicines that **you must not combine** with darunavir. These are mentioned above under the heading '**Do not combine Darunavir Mylan with any of the following medicines:**'

In most cases, darunavir can be combined with anti-HIV medicines belonging to another class [e.g. NRTIs (nucleoside reverse transcriptase inhibitors), NNRTIs (non-nucleoside reverse transcriptase inhibitors), CCR5 antagonists and FIs (fusion inhibitors)]. Darunavir with ritonavir has not been tested with all PIs (protease inhibitors) and must not be used with other HIV PIs. In some cases dosage of other medicines might need to be changed. Therefore always tell your doctor if you take other anti-HIV medicines and follow your doctor's instruction carefully on which medicines can be combined.

The effects of darunavir might be reduced if you take any of the following products. Tell your doctor if you take:

- *Phenobarbital, phenytoin* (to prevent seizures)
- *Dexamethasone* (corticosteroid)
- *Efavirenz* (HIV infection)
- *Rifapentine, rifabutin* (medicines to treat some infections such as tuberculosis)
- *Saquinavir* (HIV infection).

The effects of other medicines might be influenced if you take darunavir and your doctor might want to do some additional blood tests. Tell your doctor if you take:

- *Amlodipine, diltiazem, disopyramide, carvedilol, felodipine, flecainide, lidocaine, metoprolol, mexiletine, nifedipine, nicardipine, propafenone, timolol, verapamil* (for heart disease) as the therapeutic effect or side effects of these medicines may be increased.
- *Apixaban, dabigatran etexilate, edoxaban, rivaroxaban, warfarin, clopidogrel* (to reduce clotting of the blood) as their therapeutic effect or side effects may be altered.
- Oestrogen-based hormonal contraceptives and hormonal replacement therapy. Darunavir might reduce its effectiveness. When used for birth control, alternative methods of non-hormonal contraception are recommended.
- Ethinylestradiol/drospirenone. Darunavir might increase the risk for elevated potassium levels by drospirenone.
- *Atorvastatin, pravastatin, rosuvastatin* (to lower cholesterol levels). The risk of muscle damage might be increased. Your doctor will evaluate which cholesterol lowering regimen is best for your specific situation.
- *Clarithromycin* (antibiotic)

- *Ciclosporin, everolimus, tacrolimus, sirolimus* (for dampening down your immune system) as the therapeutic effect or side effects of these medicines might be increased.
- *Corticosteroids including betamethasone, budesonide, fluticasone, mometasone, prednisone, triamcinolone*. These medicines are used to treat allergies, asthma, inflammatory bowel diseases, inflammatory conditions of the skin, eyes, joints and muscles and other inflammatory conditions. These medicines are generally taken orally, inhaled, injected or applied to the skin. If alternatives cannot be used, its use should only take place after medical evaluation and under close monitoring by your doctor for corticosteroid side effects.
- *Buprenorphine/naloxone* (medicines to treat opioid dependence)
- *Salmeterol* (medicine to treat asthma)
- *Artemether/lumefantrine* (a combination medicine to treat malaria)
- *Dasatinib, everolimus, irinotecan, nilotinib, vinblastine, vincristine* (to treat cancer)
- *Sildenafil, tadalafil, vardenafil* (for erectile dysfunction or to treat a heart and lung disorder called pulmonary arterial hypertension)
- *Glecaprevir/pibrentasvir* (to treat hepatitis C infection)
- *Fentanyl, oxycodone, tramadol* (to treat pain)
- *Fesoterodine, solifenacin* (to treat urologic disorders).
- Your doctor might want to do some additional blood tests and the dosage of other medicines might need to be changed since either their own or darunavir's therapeutic effect or side effects may be influenced when combined.

Tell your doctor if you take:

- *Dabigatran etexilate, edoxaban, warfarin* (to reduce clotting of the blood)
- *Alfentanil* (injectable strong and short-acting painkiller that is used for surgical procedures)
- *Digoxin* (to treat certain heart disorders)
- *Clarithromycin* (antibiotic)
- *Itraconazole, isavuconazole, fluconazole, posaconazole, clotrimazole* (to treat fungal infections). Voriconazole should only be taken after medical evaluation.
- *Rifabutin* (against bacterial infections)
- *Sildenafil, vardenafil, tadalafil* (for erectile dysfunction or high blood pressure in the pulmonary circulation)
- *Amitriptyline, desipramine, imipramine, nortriptyline, paroxetine, sertraline, trazodone* (to treat depression and anxiety)
- *Maraviroc* (to treat HIV infection)
- *Methadone* (to treat opiate dependence)
- *Carbamazepine, clonazepam* (to prevent seizures or to treat certain types of nerve pain)
- *Colchicine* (to treat gout or familial Mediterranean fever)
- *Bosentan* (to treat high blood pressure in the pulmonary circulation)
- *Buspirone, clorazepate, diazepam, estazolam, flurazepam, midazolam when used as injection, zolpidem* (sedative agents)
- *Perphenazine, risperidone, thioridazine* (to treat psychiatric conditions).

This is **not** a complete list of medicines. Tell your healthcare provider about **all** medicines that you are taking.

### **Darunavir Mylan with food and drink**

See section 3 'How to take Darunavir Mylan'.

### **Pregnancy and breast-feeding**

Tell your doctor immediately if you are pregnant or planning to become pregnant. Pregnant women must not take darunavir with ritonavir unless specifically directed by the doctor. Pregnant women should not take darunavir with cobicistat.

Because of the potential for side effects in breast-fed infants, women should not breast-feed if they are receiving Darunavir Mylan.

Breast-feeding is not recommended in women living with HIV because HIV infection can be passed on to the baby in breast milk. If you are breast-feeding, or thinking about breast-feeding, you should discuss it with your doctor as soon as possible.

### **Driving and using machines**

Do not operate machines or drive if you feel dizzy after taking darunavir.

### **Darunavir Mylan contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially ‘sodium-free’.

## **3. How to take Darunavir Mylan**

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure.

Even if you feel better, do not stop taking Darunavir Mylan and ritonavir without talking to your doctor.

After therapy has been initiated, the dose or dosage form must not be changed or therapy must not be stopped without instruction of the doctor.

### **Dose for children of 3 years of age and above, weighing at least 15 kilograms who have not taken antiretroviral medicines before (your child’s doctor will determine this)**

The doctor will work out the right once daily dose based on the weight of the child (see table below).

This dose must not exceed the recommended adult dose, which is 800 milligram darunavir together with 100 milligram ritonavir once a day.

The doctor will inform you on how much Darunavir Mylan tablets and how much ritonavir (capsules, tablets or solution) the child must take.

<b>Weight</b>	<b>One darunavir dose is</b>	<b>One ritonavir<sup>a</sup> dose is</b>
between 15 and 30 kilograms	600 milligram	100 milligram
between 30 and 40 kilograms	675 milligram	100 milligram
more than 40 kilograms	800 milligram	100 milligram

<sup>a</sup> ritonavir oral solution: 80 milligram per milliliter

### **Dose for children of 3 years of age and above, weighing at least 15 kilograms who have taken antiretroviral medicines before (your child’s doctor will determine this)**

The doctor will work out the right dose based on the weight of the child (see table below). The doctor will determine if once daily dosing or twice daily dosing is appropriate for the child. This dose must not exceed the recommended adult dose, which is 600 milligram darunavir together with 100 milligram ritonavir two times per day or 800 milligram darunavir together with 100 milligram ritonavir once a day.

The doctor will inform you on how many Darunavir Mylan tablets and how much ritonavir (capsules, tablets or solution) the child must take. Tablets of other strengths are available and your doctor may have prescribed a certain combination of tablets to construct the appropriate dosing regimen.

Other forms of this medicine may be more suitable for children: ask your doctor or pharmacist.

#### **Twice daily dosing**

<b>Weight</b>	<b>One dose is</b>
between 15 and 30 kilograms	375 milligram darunavir + 50 milligram ritonavir twice a day
between 30 and 40 kilograms	450 milligram darunavir + 60 milligram ritonavir twice a day

Weight	One dose is
more than 40 kilograms*	600 milligram darunavir + 100 milligram ritonavir twice a day

\* For children aged 12 or more and weighing at least 40 kilograms, your child's doctor will determine if Darunavir Mylan 800 milligram once daily dosing may be used. This cannot be administered with these 75 milligram tablets. Other strengths of Darunavir Mylan are available.

Once daily dosing

Weight	One darunavir dose is	One ritonavir <sup>a</sup> dose is
between 15 and 30 kilograms	600 milligram	100 milligram
between 30 and 40 kilograms	675 milligram	100 milligram
more than 40 kilograms	800 milligram	100 milligram

### **Instructions for children**

- The child must take darunavir always together with ritonavir. Darunavir cannot work properly without ritonavir.
- The child must take the appropriate doses of darunavir and ritonavir two times per day or once a day. If prescribed darunavir twice daily the child must take one dose in the morning, and one dose in the evening. Your child's doctor will determine the appropriate dosing regimen for your child.
- The child must take darunavir with food. Darunavir cannot work properly without food. The type of food is not important.
- The child must swallow the tablets with a drink such as water or milk.

### **Dose for adults who have not taken antiretroviral medicines before (your doctor will determine this)**

You will require a different dose of darunavir which cannot be administered with these 75 milligram tablets. Other strengths of Darunavir Mylan are available.

### **Dose for adults who have taken antiretroviral medicines before (your doctor will determine this)**

The dose is either:

- 600 milligram darunavir (2 tablets containing 300 milligram of darunavir or 1 tablet containing 600 milligram of darunavir) together with 100 milligram ritonavir twice daily.
- OR
- 800 milligram darunavir (2 tablets containing 400 milligram of darunavir or 1 tablet containing 800 milligram of darunavir) together with 100 milligram ritonavir once daily. Darunavir Mylan 400 milligram and 800 milligram tablets are only to be used to construct the once daily 800 milligram regimen.

Please discuss with your doctor which dose is right for you.

### **Instructions for adults**

- Take darunavir always together with ritonavir. Darunavir cannot work properly without ritonavir.
- In the morning, take 600 milligram darunavir together with 100 milligram ritonavir.
- In the evening, take 600 milligram darunavir together with 100 milligram ritonavir.
- Take darunavir with food. Darunavir cannot work properly without food. The type of food is not important.
- Swallow the tablets with a drink such as water or milk.
- Darunavir Mylan 75 milligram and 150 milligram tablets have been developed for use in children, but can also be used in adults in some cases.

### **If you take more Darunavir Mylan than you should**

Contact your doctor, pharmacist or nurse immediately.

**If you forget to take Darunavir Mylan**

If you notice **within 6 hours**, you must take the tablets immediately. Always take with ritonavir and food. If you notice **after 6 hours**, then skip the intake and take the next doses as usual. Do not take a double dose to make up for a forgotten dose.

**If you vomit after taking Darunavir Mylan and ritonavir**

If you vomit **within 4 hours** of taking the medicine, another dose of Darunavir Mylan and ritonavir should be taken with food as soon as possible. If you vomit **more than 4 hours** after taking the medicine, then you do not need to take another dose of Darunavir Mylan and ritonavir until the next regularly scheduled time.

Contact your doctor **if you are uncertain** about what to do if you miss a dose or vomit.

**Do not stop taking Darunavir Mylan without talking to your doctor first**

Anti-HIV medicines may make you feel better. Even when you feel better, do not stop taking darunavir. Talk to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

**4. Possible side effects**

During HIV therapy there may be an increase in weight and in levels of blood lipids and glucose. This is partly linked to restored health and life style, and in the case of blood lipids sometimes to the HIV medicines themselves. Your doctor will test for these changes.

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Tell your doctor if you develop any of the following side effects**

Liver problems that may occasionally be severe have been reported. Your doctor should do blood tests before you start darunavir. If you have chronic hepatitis B or C infection, your doctor should check your blood tests more often because you have an increased chance of developing liver problems. Talk to your doctor about the signs and symptoms of liver problems. These may include yellowing of your skin or whites of your eyes, dark (tea coloured) urine, pale coloured stools (bowel movements), nausea, vomiting, loss of appetite, or pain, aching, or pain and discomfort on your right side below your ribs.

Skin rash (more often when used in combination with raltegravir), itching. The rash is usually mild to moderate. A skin rash might also be a symptom of a rare severe situation. It is therefore important to talk to your doctor if you develop a rash. Your doctor will advise you how to deal with your symptoms or whether darunavir must be stopped.

Other severe side effects were diabetes (common), and inflammation of the pancreas (uncommon).

Very common side effects (may affect more than 1 in 10 people)

- diarrhoea.

Common side effects (may affect up to 1 in 10 people)

- vomiting, nausea, abdominal pain or distension, dyspepsia, flatulence
- headache, tiredness, dizziness, drowsiness, numbness, tingling or pain in hands or feet, loss of strength, difficulty falling asleep.

Uncommon side effects (may affect up to 1 in 100 people)

- chest pain, changes in electrocardiogram, rapid heart beating
- decreased or abnormal skin sensitivity, pins and needles, attention disturbance, loss of memory, problems with your balance
- difficulty breathing, cough, nosebleed, throat irritation
- inflammation of the stomach or mouth, heartburn, retching, dry mouth, discomfort of the abdomen, constipation, belching

- kidney failure, kidney stones, difficult discharge of urine, frequent or excessive passage of urine, sometimes at night
- urticaria, severe swelling of the skin and other tissues (most often the lips or the eyes), eczema, excessive sweating, night sweats, hair loss, acne, scaly skin, colouration of nails
- muscle pain, muscle cramps or weakness, pain in extremity, osteoporosis
- slowing down of the thyroid gland function. This can be seen in a blood test.
- high blood pressure, flushing
- red or dry eyes
- fever, swelling of lower limbs due to fluids, malaise, irritability, pain
- symptoms of infection, herpes simplex
- erectile dysfunction, enlargement of breasts
- sleeping problems, sleepiness, depression, anxiety, abnormal dreams, decrease in sexual drive

Rare side effects (may affect up to 1 in 1,000 people)

- a reaction called DRESS [severe rash, which may be accompanied by fever, fatigue, swelling of the face or lymph glands, increase of eosinophils (type of white blood cells), effects on liver, kidney or lung]
- heart attack, slow heart beating, palpitations
- visual disturbance
- chills, feeling abnormal
- a feeling of confusion or disorientation, altered mood, restlessness
- fainting, epileptic fits, changes or loss of taste
- mouth sores, vomiting blood, inflammation of the lips, dry lips, coated tongue
- running nose
- skin lesions, dry skin
- stiffness of muscles or joints, joint pain with or without inflammation
- changes in some values of your blood cells or chemistry. These can be seen in the results of blood and/or urine tests. Your doctor will explain these to you. Examples are: increase in some white blood cells
- darunavir crystals in the kidney causing kidney disease.

Some side effects are typical for anti-HIV medicines in the same family as darunavir. These are:

- muscle pain, tenderness or weakness. On rare occasions, these muscle disorders have been serious.

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Darunavir Mylan**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, blister and on the bottle after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

For bottles: Once opened, use within 100 days.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away any medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Darunavir Mylan contains

- The active substance is darunavir. Each tablet contains 75 milligrams of darunavir.
- The other ingredients are colloidal anhydrous silica, cellulose microcrystalline, crospovidone, sodium starch glycolate, hypromellose, magnesium stearate. The film-coating contains polyvinyl alcohol partially hydrolysed, titanium dioxide (E171), macrogol and talc.

### What Darunavir Mylan looks like and contents of the pack

Darunavir Mylan 75 mg film-coated tablets are white to off-white, oval shaped with two curved sides, marked with M on one side and DV1 on the other side.

Darunavir Mylan 75 mg film-coated tablets are available in blister packs containing 480 tablets and in plastic bottles containing 480 tablets.

Not all pack sizes may be marketed.

### Marketing Authorisation Holder

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

### Manufacturer

Mylan Hungary Kft  
H-2900 Komárom, Mylan utca 1  
Hungary

McDermott Laboratories Limited trading as Gerard Laboratories  
35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13  
Ireland

Mylan Germany GmbH  
Zweigniederlassung Bad Homburg v. d. Hoehe, Benzstrasse 1  
Bad Homburg v. d. Hoehe  
Hessen, 61352  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### **België/Belgique/Belgien**

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00

#### **Lietuva**

Mylan Healthcare UAB  
Tel: +370 5 205 1288

#### **България**

Майлан ЕООД  
Тел: +359 2 44 55 400

#### **Luxembourg/Luxemburg**

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00  
(Belgique/Belgien)

#### **Česká republika**

Viatis CZ s.r.o.  
Tel: + 420 222 004 400

#### **Magyarország**

Mylan EPD Kft.  
Tel.: + 36 1 465 2100

**Danmark**

Viatriis ApS  
Tlf: +45 28 11 69 32

**Deutschland**

Viatriis Healthcare GmbH  
Tel: +49 800 0700 800

**Eesti**

BGP Products Switzerland GmbH Eesti  
filiaal  
Tel: + 372 6363 052

**Ελλάδα**

Generics Pharma Hellas EΠE  
Τηλ: +30 210 993 6410

**España**

Viatriis Pharmaceuticals, S.L.U.  
Tel: + 34 900 102 712

**France**

Viatriis Santé  
Tél: +33 4 37 25 75 00

**Hrvatska**

Viatriis Hrvatska d.o.o.  
Tel: +385 1 23 50 599

**Ireland**

Mylan Ireland Limited  
Tel: +353 1 8711600

**Ísland**

Icepharma hf.  
Sími: +354 540 8000

**Italia**

Mylan Italia S.r.l.  
Tel: + 39 02 612 46921

**Κύπρος**

Varnavas Hadjipanayis Ltd  
Τηλ: + 357 2220 7700

**Latvija**

Mylan Healthcare SIA  
Tel: +371 676 055 80

**Malta**

V.J. Salomone Pharma Ltd  
Tel: + 356 21 22 01 74

**Nederland**

Mylan BV  
Tel: +31 (0)20 426 3300

**Norge**

Viatriis AS  
Tlf: + 47 66 75 33 00

**Österreich**

Arcana Arzneimittel GmbH  
Tel: +43 1 416 2418

**Polska**

Mylan Healthcare Sp. z o.o.  
Tel.: + 48 22 546 64 00

**Portugal**

Mylan, Lda.  
Tel: + 351 214 127 200

**România**

BGP Products SRL  
Tel: +40 372 579 000

**Slovenija**

Viatriis d.o.o.  
Tel: + 386 1 23 63 180

**Slovenská republika**

Viatriis Slovakia s.r.o.  
Tel: +421 2 32 199 100

**Suomi/Finland**

Viatriis Oy  
Puh/Tel: +358 20 720 9555

**Sverige**

Viatriis AB  
Tel: +46 (0)8 630 19 00

**United Kingdom (Northern Ireland)**

Mylan IRE Healthcare Limited  
Tel: +353 18711600

**This leaflet was last revised in {MM/YYYY}.**



**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu>.

## Package leaflet: Information for the user

### Darunavir Mylan 150 mg film-coated tablets

darunavir

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Darunavir Mylan is and what it is used for
2. What you need to know before you take Darunavir Mylan
3. How to take Darunavir Mylan
4. Possible side effects
5. How to store Darunavir Mylan
6. Contents of the pack and other information

#### 1. What Darunavir Mylan is and what it is used for

##### What is Darunavir Mylan?

Darunavir Mylan contains the active substance darunavir. Darunavir is an antiretroviral medicine used in the treatment of Human Immunodeficiency Virus (HIV) infection. It belongs to a group of medicines called protease inhibitors. Darunavir works by reducing the amount of HIV in your body. This will improve your immune system and reduces the risk of developing illnesses linked to HIV infection.

##### What it is used for?

Darunavir is used to treat adults and children of 3 years of age and above, and at least 15 kilogram body weight who are infected by HIV and who have already used other antiretroviral medicines.

Darunavir must be taken in combination with a low dose of ritonavir and other anti-HIV medicines. Your doctor will discuss with you which combination of medicines is best for you.

#### 2. What you need to know before you take Darunavir Mylan

##### Do not take Darunavir Mylan

- if you are **allergic** to darunavir or any of the other ingredients of this medicine (listed in section 6) or to ritonavir.
- if you have **severe liver problems**. Ask your doctor if you are unsure about the severity of your liver disease. Some additional tests might be necessary.

##### Do not combine Darunavir Mylan with any of the following medicines

If you are taking any of these, ask your doctor about switching to another medicine.

Medicine	Purpose of the medicine
<i>Avanafil</i>	to treat erectile dysfunction
<i>Astemizole</i> or <i>terfenadine</i>	to treat allergy symptoms
<i>Triazolam</i> and <i>oral</i> (taken by mouth) <i>midazolam</i>	to help you sleep and/or relieve anxiety
<i>Cisapride</i>	to treat some stomach conditions

<b>Medicine</b>	<b>Purpose of the medicine</b>
<i>Colchicine</i> (if you have kidney and/or liver problems)	to treat gout or familial Mediterranean fever
<i>Lurasidone, pimozide, quetiapine or sertindole</i>	to treat psychiatric conditions
<i>Ergot alkaloids</i> like <i>ergotamine</i> , <i>dihydroergotamine</i> , <i>ergometrine</i> and <i>methylergonovine</i>	to treat migraine headaches
<i>Amiodarone, bepridil, dronedarone, ivabradine, quinidine, ranolazine</i>	to treat certain heart disorders e.g. abnormal heart beat
<i>Lovastatin, simvastatin and lomitapide</i>	to lower cholesterol levels
<i>Rifampicin</i>	to treat some infections such as tuberculosis
The combination product <i>lopinavir/ritonavir</i>	this anti-HIV medicine belongs to the same class as darunavir
<i>Elbasvir/grazoprevir</i>	to treat hepatitis C infection
<i>Alfuzosin</i>	to treat enlarged prostate
<i>Sildenafil</i>	to treat high blood pressure in the pulmonary circulation
<i>Ticagrelor</i>	to help stop the clumping of platelets in the treatment of patients with a history of a heart attack
<i>Naloxegol</i>	to treat opioid induced constipation
<i>Dapoxetine</i>	to treat premature ejaculation
<i>Domperidone</i>	to treat nausea and vomiting

Do not combine darunavir with products that contain St John's wort (*Hypericum perforatum*).

### **Warnings and precautions**

Talk to your doctor, pharmacist or nurse before taking Darunavir Mylan.

Darunavir is not a cure for HIV infection.

People taking darunavir may still develop infections or other illnesses associated with HIV infection. You must keep in regular contact with your doctor.

People taking darunavir may develop a skin rash. Infrequently a rash may become severe or potentially life-threatening. Please contact your doctor whenever you develop a rash.

In patients taking darunavir and raltegravir (for HIV infection), rashes (generally mild or moderate) may occur more frequently than in patients taking either medicine separately.

### **Tell your doctor about your situation BEFORE and DURING your treatment**

Make sure that you check the following points and tell your doctor if any of these apply to you.

- Tell your doctor if you have had **problems with your liver** before, including hepatitis B or C infection. Your doctor may evaluate how severe your liver disease is before deciding if you can take darunavir.
- Tell your doctor if you have **diabetes**. Darunavir might increase sugar levels in the blood.
- Tell your doctor immediately if you notice any **symptoms of infection** (for example enlarged lymph nodes and fever). In some patients with advanced HIV infection and a history of opportunistic infection, signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started. It is believed that these symptoms are due to an improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms.
- In addition to the opportunistic infections, autoimmune disorders (a condition that occurs when the immune system attacks healthy body tissue) may also occur after you start taking medicines for the treatment of your HIV infection. Autoimmune disorders may occur many months after the start of treatment. If you notice any symptoms of infection or other symptoms such as muscle weakness, weakness beginning in the hands and feet and moving up towards the trunk of

the body, palpitations, tremor or hyperactivity, please inform your doctor immediately to seek necessary treatment.

- Tell your doctor if you have **haemophilia**. Darunavir might increase the risk of bleeding.
- Tell your doctor if you are **allergic to sulphonamides** (e.g. used to treat certain infections).
- Tell your doctor if you notice any **musculoskeletal problems**. Some patients taking combination antiretroviral therapy may develop a bone disease called osteonecrosis (death of bone tissue caused by loss of blood supply to the bone). The length of combination antiretroviral therapy, corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index, among others, may be some of the many risk factors for developing this disease. Signs of osteonecrosis are joint stiffness, aches and pains (especially of the hip, knee and shoulder) and difficulty in movement. If you notice any of these symptoms please inform your doctor.

### Elderly

Darunavir has only been used in limited numbers of patients 65 years or older. If you belong to this age group, please discuss with your doctor if you can use Darunavir Mylan.

### Children

Darunavir is not for use in children younger than 3 years of age or weighing less than 15 kilograms.

### Other medicines and Darunavir Mylan

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines.

There are some medicines that **you must not combine** with darunavir. These are mentioned above under the heading '**Do not combine Darunavir Mylan with any of the following medicines:**'

In most cases, darunavir can be combined with anti-HIV medicines belonging to another class [e.g. NRTIs (nucleoside reverse transcriptase inhibitors), NNRTIs (non-nucleoside reverse transcriptase inhibitors), CCR5 antagonists and FIs (fusion inhibitors)]. Darunavir with ritonavir has not been tested with all PIs (protease inhibitors) and must not be used with other HIV PIs. In some cases dosage of other medicines might need to be changed. Therefore always tell your doctor if you take other anti-HIV medicines and follow your doctor's instruction carefully on which medicines can be combined.

The effects of darunavir might be reduced if you take any of the following products. Tell your doctor if you take:

- *Phenobarbital, phenytoin* (to prevent seizures)
- *Dexamethasone* (corticosteroid)
- *Efavirenz* (HIV infection)
- *Rifapentine, rifabutin* (medicines to treat some infections such as tuberculosis)
- *Saquinavir* (HIV infection).

The effects of other medicines might be influenced if you take darunavir and your doctor might want to do some additional blood tests. Tell your doctor if you take:

- *Amlodipine, diltiazem, disopyramide, carvedilol, felodipine, flecainide, lidocaine, metoprolol, mexiletine, nifedipine, nicardipine, propafenone, timolol, verapamil* (for heart disease) as the therapeutic effect or side effects of these medicines may be increased.
- *Apixaban dabigatran etexilate, edoxaban, rivaroxaban, warfarin, clopidogrel* (to reduce clotting of the blood) as their therapeutic effect or side effects may be altered.
- Oestrogen-based hormonal contraceptives and hormonal replacement therapy. Darunavir might reduce its effectiveness. When used for birth control, alternative methods of non-hormonal contraception are recommended.
- *Ethinylestradiol/drospirenone*. Darunavir might increase the risk for elevated potassium levels by drospirenone.
- *Atorvastatin, pravastatin, rosuvastatin* (to lower cholesterol levels). The risk of muscle damage might be increased. Your doctor will evaluate which cholesterol lowering regimen is best for your specific situation.
- *Clarithromycin* (antibiotic)

- *Ciclosporin, everolimus, tacrolimus, sirolimus* (for dampening down your immune system) as the therapeutic effect or side effects of these medicines might be increased.
- *Corticosteroids including betamethasone, budesonide, fluticasone, mometasone, prednisone, triamcinolone*. These medicines are used to treat allergies, asthma, inflammatory bowel diseases, inflammatory conditions of the skin, eyes, joints and muscles and other inflammatory conditions. These medicines are generally taken orally, inhaled, injected or applied to the skin. If alternatives cannot be used, its use should only take place after medical evaluation and under close monitoring by your doctor for corticosteroid side effects.
- *Buprenorphine/naloxone* (medicines to treat opioide dependence)
- *Salmeterol* (medicine to treat asthma)
- *Artemether/lumefantrine* (a combination medicine to treat malaria)
- *Dasatinib, everolimus, irinotecan, nilotinib, vinblastine, vincristine* (to treat cancer)
- *Sildenafil, tadalafil, vardenafil* (for erectile dysfunction or to treat a heart and lung disorder called pulmonary arterial hypertension)
- *Glecaprevir/pibrentasvir* (to treat hepatitis C infection)
- *Fentanyl, oxycodone, tramadol* (to treat pain)
- *Fesoterodine, solifenacin* (to treat urologic disorders).

Your doctor might want to do some additional blood tests and the dosage of other medicines might need to be changed since either their own or darunavir's therapeutic effect or side effects may be influenced when combined.

Tell your doctor if you take:

- *Dabigatran etexilate, edoxaban, warfarin* (to reduce clotting of the blood)
- *Alfentanil* (injectable strong and short-acting painkiller that is used for surgical procedures)
- *Digoxin* (to treat certain heart disorders)
- *Clarithromycin* (antibiotic)
- *Itraconazole, isavuconazole, fluconazole, posaconazole, clotrimazole* (to treat fungal infections). Voriconazole should only be taken after medical evaluation.
- *Rifabutin* (against bacterial infections)
- *Sildenafil, vardenafil, tadalafil* (for erectile dysfunction or high blood pressure in the pulmonary circulation)
- *Amitriptyline, desipramine, imipramine, nortriptyline, paroxetine, sertraline, trazodone* (to treat depression and anxiety)
- *Maraviroc* (to treat HIV infection)
- *Methadone* (to treat opiate dependence)
- *Carbamazepine, clonazepam* (to prevent seizures or to treat certain types of nerve pain)
- *Colchicine* (to treat gout or familial Mediterranean fever)
- *Bosentan* (to treat high blood pressure in the pulmonary circulation)
- *Buspirone, clorazepate, diazepam, estazolam, flurazepam, midazolam when used as injection, zolpidem* (sedative agents)
- *Perphenazine, risperidone, thioridazine* (to treat psychiatric conditions).

This is **not** a complete list of medicines. Tell your healthcare provider about **all** medicines that you are taking.

### **Darunavir Mylan with food and drink**

See section 3 'How to take Darunavir Mylan.

### **Pregnancy and breast-feeding**

Tell your doctor immediately if you are pregnant or planning to become pregnant. Pregnant women must not take darunavir with ritonavir unless specifically directed by the doctor. Pregnant women should not take darunavir with cobicistat.

Because of the potential for side effects in breast-fed infants, women should not breast-feed if they are receiving Darunavir Mylan.

Breast-feeding is not recommended in women living with HIV because HIV infections can be passed on to the baby in breast milk. If you are breast-feeding, or thinking about breast-feeding, you should discuss it with your doctor as soon as possible.

### **Driving and using machines**

Do not operate machines or drive if you feel dizzy after taking darunavir.

### **Darunavir Mylan contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially ‘sodium-free’.

## **3. How to take Darunavir Mylan**

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure.

Even if you feel better, do not stop taking Darunavir Mylan and ritonavir without talking to your doctor.

After therapy has been initiated, the dose or dosage form must not be changed or therapy must not be stopped without instruction of the doctor.

### **Dose for children of 3 years of age and above, weighing at least 15 kilograms who have not taken antiretroviral medicines before (your child’s doctor will determine this)**

The doctor will work out the right once daily dose based on the weight of the child (see table below). This dose must not exceed the recommended adult dose, which is 800 milligram darunavir together with 100 milligram ritonavir once a day.

The doctor will inform you on how much Darunavir Mylan tablets and how much ritonavir (capsules, tablets or solution) the child must take.

<b>Weight</b>	<b>One darunavir dose is</b>	<b>One ritonavir<sup>a</sup> dose is</b>
between 15 and 30 kilograms	600 milligram	100 milligram
between 30 and 40 kilograms	675 milligram	100 milligram
more than 40 kilograms	800 milligram	100 milligram

<sup>a</sup> ritonavir oral solution: 80 milligram per milliliter

### **Dose for children of 3 years of age and above, weighing at least 15 kilograms who have taken antiretroviral medicines before (your child’s doctor will determine this)**

The doctor will work out the right dose based on the weight of the child (see table below). The doctor will determine if once daily dosing or twice daily dosing is appropriate for the child. This dose must not exceed the recommended adult dose, which is 600 milligram darunavir together with 100 milligram ritonavir two times per day or 800 milligram darunavir together with 100 milligram ritonavir once a day.

The doctor will inform you on how many Darunavir Mylan tablets and how much ritonavir (capsules, tablets or solution) the child must take. Tablets of other strengths are available and your doctor may have prescribed a certain combination of tablets to construct the appropriate dosing regimen. Other forms of this medicine may be more suitable for children: ask your doctor or pharmacist.

#### **Twice daily dosing**

<b>Weight</b>	<b>One dose is</b>
between 15 and 30 kilograms	375 milligram darunavir + 50 milligram ritonavir twice a day
between 30 and 40 kilograms	450 milligram darunavir + 60 milligram ritonavir twice a day

<b>Weight</b>	<b>One dose is</b>
more than 40 kilograms*	600 milligram darunavir + 100 milligramritonavir twice a day

\* For children aged 12 or more and weighing at least 40 kilograms, your child's doctor will determine if Darunavir Mylan 800 milligram once daily dosing may be used. This cannot be administered with these 150 milligram tablets. Other strengths of Darunavir Mylan are available.

Once daily dosing

<b>Weight</b>	<b>One darunavir dose is</b>	<b>One ritonavir<sup>a</sup> dose is</b>
between 15 and 30 kilograms	600 milligram	100 milligram
between 30 and 40 kilograms	675 milligram	100 milligram
more than 40 kilograms	800 milligram	100 milligram

<sup>a</sup> ritonavir oral solution: 80 milligram per milliliter

### ***Instructions for children***

- The child must take darunavir always together with ritonavir. Darunavir cannot work properly without ritonavir.
- The child must take the appropriate doses of darunavir and ritonavir two times per day or once a day. If prescribed darunavir twice daily the child must take one dose in the morning, and one dose in the evening. Your child's doctor will determine the appropriate dosing regimen for your child.
- The child must take darunavir with food. Darunavir cannot work properly without food. The type of food is not important.
- The child must swallow the tablets with a drink such as water or milk.

### **Dose for adults who have not taken antiretroviral medicines before (your doctor will determine this)**

You will require a different dose of darunavir which cannot be administered with these 150 milligram tablets. Other strengths of Darunavir Mylan are available.

### **Dose for adults who have taken antiretroviral medicines before (your doctor will determine this)**

The dose is either:

- 600 milligram darunavir (2 tablets containing 300 milligram of darunavir or 1 tablet containing 600 milligram of darunavir) together with 100 milligram ritonavir twice daily.
- OR
- 800 milligram darunavir (2 tablets containing 400 milligram of darunavir or 1 tablet containing 800 milligram of darunavir) together with 100 milligram ritonavir once daily. Darunavir Mylan 400 milligram and 800 milligram tablets are only to be used to construct the once daily 800 milligram regimen.

Please discuss with your doctor which dose is right for you.

### ***Instructions for adults***

- Take darunavir always together with ritonavir. Darunavir cannot work properly without ritonavir.
- In the morning, take 600 milligram darunavir together with 100 milligram ritonavir.
- In the evening, take 600 milligram darunavir together with 100 milligram ritonavir.
- Take darunavir with food. Darunavir cannot work properly without food. The type of food is not important.
- Swallow the tablets with a drink such as water or milk.
- Darunavir Mylan 75 milligram and 150 milligram tablets have been developed for use in children, but can also be used in adults in some cases.

### **If you take more Darunavir Mylan than you should**

Contact your doctor, pharmacist or nurse immediately.

**If you forget to take Darunavir Mylan**

If you notice **within 6 hours**, you must take the tablets immediately. Always take with ritonavir and food. If you notice **after 6 hours**, then skip the intake and take the next doses as usual. Do not take a double dose to make up for a forgotten dose.

**If you vomit after taking Darunavir Mylan and ritonavir**

If you vomit **within 4 hours** of taking the medicine, another dose of Darunavir Mylan and ritonavir should be taken with food as soon as possible. If you vomit **more than 4 hours** after taking the medicine, then you do not need to take another dose of Darunavir Mylan and ritonavir until the next regularly scheduled time.

Contact your doctor **if you are uncertain** about what to do if you miss a dose or vomit.

**Do not stop taking Darunavir Mylan without talking to your doctor first**

Anti-HIV medicines make you feel better. Even when you feel better, do not stop taking darunavir. Talk to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

**4. Possible side effects**

During HIV therapy there may be an increase in weight and in levels of blood lipids and glucose. This is partly linked to restored health and life style, and in the case of blood lipids sometimes to the HIV medicines themselves. Your doctor will test for these changes.

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Tell your doctor if you develop any of the following side effects**

Liver problems that may occasionally be severe have been reported. Your doctor should do blood tests before you start darunavir. If you have chronic hepatitis B or C infection, your doctor should check your blood tests more often because you have an increased chance of developing liver problems. Talk to your doctor about the signs and symptoms of liver problems. These may include yellowing of your skin or whites of your eyes, dark (tea coloured) urine, pale coloured stools (bowel movements), nausea, vomiting, loss of appetite, or pain, aching, or pain and discomfort on your right side below your ribs.

Skin rash (more often when used in combination with raltegravir), itching. The rash is usually mild to moderate. A skin rash might also be a symptom of a rare severe situation. It is therefore important to talk to your doctor if you develop a rash. Your doctor will advise you how to deal with your symptoms or whether darunavir must be stopped.

Other severe side effects were diabetes (common), and inflammation of the pancreas (uncommon).

Very common side effects (may affect more than 1 in 10 people)

- diarrhoea.

Common side effects (may affect up to 1 in 10 people)

- vomiting, nausea, abdominal pain or distension, dyspepsia, flatulence
- headache, tiredness, dizziness, drowsiness, numbness, tingling or pain in hands or feet, loss of strength, difficulty falling asleep.

Uncommon side effects (may affect up to 1 in 100 people)

- chest pain, changes in electrocardiogram, rapid heart beating
- decreased or abnormal skin sensitivity, pins and needles, attention disturbance, loss of memory, problems with your balance
- difficulty breathing, cough, nosebleed, throat irritation
- inflammation of the stomach or mouth, heartburn, retching, dry mouth, discomfort of the abdomen, constipation, belching



- kidney failure, kidney stones, difficult discharge of urine, frequent or excessive passage of urine, sometimes at night
- urticaria, severe swelling of the skin and other tissues (most often the lips or the eyes), eczema, excessive sweating, night sweats, hair loss, acne, scaly skin, colouration of nails
- muscle pain, muscle cramps or weakness, pain in extremity, osteoporosis
- slowing down of the thyroid gland function. This can be seen in a blood test.
- high blood pressure, flushing
- red or dry eyes
- fever, swelling of lower limbs due to fluids, malaise, irritability, pain
- symptoms of infection, herpes simplex
- erectile dysfunction, enlargement of breasts
- sleeping problems, sleepiness, depression, anxiety, abnormal dreams, decrease in sexual drive

Rare side effects (may affect up to 1 in 1,000 people)

- a reaction called DRESS [severe rash, which may be accompanied by fever, fatigue, swelling of the face or lymph glands, increase of eosinophils (type of white blood cells), effects on liver, kidney or lung]
- heart attack, slow heart beating, palpitations
- visual disturbance
- chills, feeling abnormal
- a feeling of confusion or disorientation, altered mood, restlessness
- fainting, epileptic fits, changes or loss of taste
- mouth sores, vomiting blood, inflammation of the lips, dry lips, coated tongue
- running nose
- skin lesions, dry skin
- stiffness of muscles or joints, joint pain with or without inflammation
- changes in some values of your blood cells or chemistry. These can be seen in the results of blood and/or urine tests. Your doctor will explain these to you. Examples are: increase in some white blood cells
- darunavir crystals in the kidney causing kidney disease.

Some side effects are typical for anti-HIV medicines in the same family as darunavir. These are:

- muscle pain, tenderness or weakness. On rare occasions, these muscle disorders have been serious.

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Darunavir Mylan**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, blister and on the bottle after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

For bottles: Once opened, use within 100 days.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away any medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Darunavir Mylan contains

- The active substance is darunavir. Each tablet contains 150 milligram of darunavir.
- The other ingredients are colloidal anhydrous silica, cellulose microcrystalline, crospovidone, sodium starch glycolate, hypromellose, magnesium stearate. The film-coating contains polyvinyl alcohol partially hydrolysed, titanium dioxide (E171), macrogol and talc.

### What Darunavir Mylan looks like and contents of the pack

Darunavir Mylan 150 mg film-coated tablets are white to off-white, capsule shaped, with two curved sides, marked with M on one side and DV2 on the other side.

Darunavir Mylan 150 mg film-coated tablets are available in blister packs containing 240 tablets and in plastic bottles containing 60 and 240 tablets.

Not all pack sizes may be marketed.

### Marketing Authorisation Holder

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

### Manufacturers

Mylan Hungary Kft  
H-2900 Komárom, Mylan utca 1  
Hungary

McDermott Laboratories Limited trading as Gerard Laboratories  
35/36 Baldoye Industrial Estate, Grange Road, Dublin 13  
Ireland

Mylan Germany GmbH  
Zweigniederlassung Bad Homburg v. d. Hoehe, Benzstrasse 1  
Bad Homburg v. d. Hoehe  
Hessen, 61352  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### **België/Belgique/Belgien**

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00

#### **Lietuva**

Mylan Healthcare UAB  
Tel: +370 5 205 1288

#### **България**

Майлан ЕООД  
Тел: +359 2 44 55 400

#### **Luxembourg/Luxemburg**

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00  
(Belgique/Belgien)

#### **Česká republika**

Viatrix CZ s.r.o.  
Tel: + 420 222 004 400

#### **Magyarország**

Mylan EPD Kft.  
Tel.: + 36 1 465 2100

**Danmark**

Viatriis ApS  
Tlf: +45 28 11 69 32

**Malta**

V.J. Salomone Pharma Ltd  
Tel: + 356 21 22 01 74

**Deutschland**

Viatriis Healthcare GmbH  
Tel: +49 800 0700 800

**Nederland**

Mylan BV  
Tel: +31 (0)20 426 3300

**Eesti**

BGP Products Switzerland GmbH  
Eesti filiaal  
Tel: + 372 6363 052

**Norge**

Viatriis AS  
Tlf: + 47 66 75 33 00

**Ελλάδα**

Generics Pharma Hellas EΠE  
Τηλ: +30 210 993 6410

**Österreich**

Arcana Arzneimittel GmbH  
Tel: +43 1 416 2418

**España**

Viatriis Pharmaceuticals, S.L.U.  
Tel: + 34 900 102 712

**Polska**

Mylan Healthcare Sp. z o.o.  
Tel.: + 48 22 546 64 00

**France**

Viatriis Santé  
Tél: +33 4 37 25 75 00

**Portugal**

Mylan, Lda.  
Tel: + 351 214 127 200

**Hrvatska**

Viatriis Hrvatska d.o.o.  
Tel: +385 1 23 50 599

**România**

BGP Products SRL  
Tel: +40 372 579 000

**Ireland**

Mylan Ireland Limited  
Tel: +353 1 8711600

**Slovenija**

Viatriis d.o.o.  
Tel: + 386 1 23 63 180

**Ísland**

Icepharma hf.  
Sími: +354 540 8000

**Slovenská republika**

Viatriis Slovakia s.r.o.  
Tel: +421 2 32 199 100

**Italia**

Mylan Italia S.r.l.  
Tel: + 39 02 612 46921

**Suomi/Finland**

Viatriis Oy  
Puh/Tel: +358 20 720 9555

**Κύπρος**

Varnavas Hadjipanayis Ltd  
Τηλ: + 357 2220 7700

**Sverige**

Viatriis AB  
Tel: +46 (0)8 630 19 00

**Latvija**

Mylan Healthcare SIA  
Tel: +371 676 055 80

**United Kingdom (Northern Ireland)**

Mylan IRE Healthcare Limited  
Tel: +353 18711600

**This leaflet was last revised in {MM/YYYY}.**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu>.

## Package leaflet: Information for the user

### Darunavir Mylan 300 mg film-coated tablets

darunavir

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Darunavir Mylan is and what it is used for
2. What you need to know before you take Darunavir Mylan
3. How to take Darunavir Mylan
4. Possible side effects
5. How to store Darunavir Mylan
6. Contents of the pack and other information

#### 1. What Darunavir Mylan is and what it is used for

##### What is Darunavir Mylan?

Darunavir Mylan contains the active substance darunavir. Darunavir is an antiretroviral medicine used in the treatment of Human Immunodeficiency Virus (HIV) infection. It belongs to a group of medicines called protease inhibitors. Darunavir works by reducing the amount of HIV in your body. This will improve your immune system and reduces the risk of developing illnesses linked to HIV infection.

##### What it is used for?

Darunavir is used to treat adults and children of 3 years of age and above, and at least 15 kilogram body weight who are infected by HIV and who have already used other antiretroviral medicines.

Darunavir must be taken in combination with a low dose of ritonavir and other anti-HIV medicines. Your doctor will discuss with you which combination of medicines is best for you.

#### 2. What you need to know before you take Darunavir Mylan

##### Do not take Darunavir Mylan

- if you are **allergic** to darunavir or any of the other ingredients of this medicine (listed in section 6) or to ritonavir.
- if you have **severe liver problems**. Ask your doctor if you are unsure about the severity of your liver disease. Some additional tests might be necessary.

##### Do not combine Darunavir Mylan with any of the following medicines

If you are taking any of these, ask your doctor about switching to another medicine.

Medicine	Purpose of the medicine
<i>Avanafil</i>	to treat erectile dysfunction
<i>Astemizole</i> or <i>terfenadine</i>	to treat allergy symptoms
<i>Triazolam</i> and <i>oral</i> (taken by mouth) <i>midazolam</i>	to help you sleep and/or relieve anxiety

Medicine	Purpose of the medicine
<i>Cisapride</i>	to treat some stomach conditions
<i>Colchicine</i> (if you have kidney and/or liver problems)	to treat gout or familial Mediterranean fever
<i>Lurasidone, pimozide, quetiapine</i> or <i>sertindole</i>	to treat psychiatric conditions
<i>Ergot alkaloids</i> like <i>ergotamine</i> , <i>dihydroergotamine</i> , <i>ergometrine</i> and <i>methylergonovine</i>	to treat migraine headaches
<i>Amiodarone</i> , <i>bepiridil</i> , <i>dronedarone</i> , <i>ivabradine</i> , <i>quinidine</i> , <i>ranolazine</i>	to treat certain heart disorders e.g. abnormal heart beat
<i>Lovastatin</i> , <i>simvastatin</i> and <i>lomitapide</i>	to lower cholesterol levels
<i>Rifampicin</i>	to treat some infections such as tuberculosis
The combination product <i>lopinavir/ritonavir</i>	this anti-HIV medicine belongs to the same class as Darunavir Mylan
<i>Elbasvir/grazoprevir</i>	to treat hepatitis C infection
<i>Alfuzosin</i>	to treat enlarged prostate
<i>Sildenafil</i>	to treat high blood pressure in the pulmonary circulation
<i>Ticagrelor</i>	to help stop the clumping of platelets in the treatment of patients with a history of a heart attack
<i>Naloxegol</i>	to treat opioid induced constipation
<i>Dapoxetine</i>	to treat premature ejaculation
<i>Domperidone</i>	to treat nausea and vomiting

Do not combine darunavir with products that contain St John's wort (*Hypericum perforatum*).

### Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking Darunavir Mylan.

Darunavir is not a cure for HIV infection.

People taking darunavir may still develop infections or other illnesses associated with HIV infection. You must keep in regular contact with your doctor.

People taking darunavir may develop a skin rash. Infrequently a rash may become severe or potentially life-threatening. Please contact your doctor whenever you develop a rash.

In patients taking darunavir and raltegravir (for HIV infection), rashes (generally mild or moderate) may occur more frequently than in patients taking either medicine separately.

### Tell your doctor about your situation BEFORE and DURING your treatment

Make sure that you check the following points and tell your doctor if any of these apply to you.

- Tell your doctor if you have had **problems with your liver** before, including hepatitis B or C infection. Your doctor may evaluate how severe your liver disease is before deciding if you can take darunavir.
- Tell your doctor if you have **diabetes**. Darunavir might increase sugar levels in the blood.
- Tell your doctor immediately if you notice any **symptoms of infection** (for example enlarged lymph nodes and fever). In some patients with advanced HIV infection and a history of opportunistic infection, signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started. It is believed that these symptoms are due to an improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms.
- In addition to the opportunistic infections, autoimmune disorders (a condition that occurs when the immune system attacks healthy body tissue) may also occur after you start taking medicines for the treatment of your HIV infection. Autoimmune disorders may occur many months after

the start of treatment. If you notice any symptoms of infection or other symptoms such as muscle weakness, weakness beginning in the hands and feet and moving up towards the trunk of the body, palpitations, tremor or hyperactivity, please inform your doctor immediately to seek necessary treatment.

- Tell your doctor if you have **haemophilia**. Darunavir might increase the risk of bleeding.
- Tell your doctor if you are **allergic to sulphonamides** (e.g. used to treat certain infections).
- Tell your doctor if you notice any **musculoskeletal problems**. Some patients taking combination antiretroviral therapy may develop a bone disease called osteonecrosis (death of bone tissue caused by loss of blood supply to the bone). The length of combination antiretroviral therapy, corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index, among others, may be some of the many risk factors for developing this disease. Signs of osteonecrosis are joint stiffness, aches and pains (especially of the hip, knee and shoulder) and difficulty in movement. If you notice any of these symptoms please inform your doctor.

### Elderly

Darunavir has only been used in limited numbers of patients 65 years or older. If you belong to this age group, please discuss with your doctor if you can use Darunavir Mylan.

### Children

Darunavir is not for use in children younger than 3 years of age or weighing less than 15 kilograms.

### Other medicines and Darunavir Mylan

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines.

There are some medicines that **you must not combine** with darunavir. These are mentioned above under the heading '**Do not combine Darunavir Mylan with any of the following medicines:**'

In most cases, darunavir can be combined with anti-HIV medicines belonging to another class [e.g. NRTIs (nucleoside reverse transcriptase inhibitors), NNRTIs (non-nucleoside reverse transcriptase inhibitors), CCR5 antagonists and FIs (fusion inhibitors)]. Darunavir with ritonavir has not been tested with all PIs (protease inhibitors) and must not be used with other HIV PIs. In some cases dosage of other medicines might need to be changed. Therefore always tell your doctor if you take other anti-HIV medicines and follow your doctor's instruction carefully on which medicines can be combined.

The effects of darunavir might be reduced if you take any of the following products. Tell your doctor if you take:

- *Phenobarbital, phenytoin* (to prevent seizures)
- *Dexamethasone* (corticosteroid)
- *Efavirenz* (HIV infection)
- *Rifapentine, rifabutin* (medicines to treat some infections such as tuberculosis)
- *Saquinavir* (HIV infection).

The effects of other medicines might be influenced if you take darunavir and your doctor might want to do some additional blood tests. Tell your doctor if you take:

- *Amlodipine, diltiazem, disopyramide, carvedilol, felodipine, flecainide, lidocaine, metoprolol, mexiletine, nifedipine, nicardipine, propafenone, timolol, verapamil* (for heart disease) as the therapeutic effect or side effects of these medicines may be increased.
- *Apixaban, dabigatran etexilate, edoxaban, rivaroxaban, warfarin, clopidogrel* (to reduce clotting of the blood) as their therapeutic effect or side effects may be altered.
- Oestrogen-based hormonal contraceptives and hormonal replacement therapy. Darunavir might reduce its effectiveness. When used for birth control, alternative methods of non-hormonal contraception are recommended.
- *Ethinylestradiol/drospirenone*. Darunavir might increase the risk for elevated potassium levels by drospirenone.
- *Atorvastatin, pravastatin, rosuvastatin* (to lower cholesterol levels). The risk of muscle damage might be increased. Your doctor will evaluate which cholesterol lowering regimen is best for your specific situation.

- *Clarithromycin* (antibiotic)
- *Ciclosporin, everolimus, tacrolimus, sirolimus* (for dampening down your immune system) as the therapeutic effect or side effects of these medicines might be increased.
- *Corticosteroids including betamethasone, budesonide, fluticasone, mometasone, prednisone, triamcinolone*. These medicines are used to treat allergies, asthma, inflammatory bowel diseases, inflammatory conditions of the skin, eyes, joints and muscles and other inflammatory conditions. These medicines are generally taken orally, inhaled, injected or applied to the skin. If alternatives cannot be used, its use should only take place after medical evaluation and under close monitoring by your doctor for corticosteroid side effects.
- *Buprenorphine/naloxone* (medicines to treat opioide dependence)
- *Salmeterol* (medicine to treat asthma)
- *Artemether/lumefantrine* (a combination medicine to treat malaria)
- *Dasatinib, everolimus, irinotecan, nilotinib, vinblastine, vincristine* (to treat cancer)
- *Sildenafil, tadalafil, vardenafil* (for erectile dysfunction or to treat a heart and lung disorder called pulmonary arterial hypertension)
- *Glecaprevir/pibrentasvir* (to treat hepatitis C infection)
- *Fentanyl, oxycodone, tramadol* (to treat pain)
- *Fesoterodine, solifenacin* (to treat urologic disorders).

Your doctor might want to do some additional blood tests and the dosage of other medicines might need to be changed since either their own or darunavir's therapeutic effect or side effects may be influenced when combined.

Tell your doctor if you take:

- *Dabigatran etexilate, edoxaban, warfarin* (to reduce clotting of the blood)
- *Alfentanil* (injectable strong and short-acting painkiller that is used for surgical procedures)
- *Digoxin* (to treat certain heart disorders)
- *Clarithromycin* (antibiotic)
- *Itraconazole, isavuconazole, fluconazole, posaconazole, clotrimazole* (to treat fungal infections). Voriconazole should only be taken after medical evaluation.
- *Rifabutin* (against bacterial infections)
- *Sildenafil, vardenafil, tadalafil* (for erectile dysfunction or high blood pressure in the pulmonary circulation)
- *Amitriptyline, desipramine, imipramine, nortriptyline, paroxetine, sertraline, trazodone* (to treat depression and anxiety)
- *Maraviroc* (to treat HIV infection)
- *Methadone* (to treat opiate dependence)
- *Carbamazepine, clonazepam* (to prevent seizures or to treat certain types of nerve pain)
- *Colchicine* (to treat gout or familial Mediterranean fever)
- *Bosentan* (to treat high blood pressure in the pulmonary circulation)
- *Buspirone, clorazepate, diazepam, estazolam, flurazepam, midazolam when used as injection, zolpidem* (sedative agents)
- *Perphenazine, risperidone, thioridazine* (to treat psychiatric conditions).

This is **not** a complete list of medicines. Tell your healthcare provider about **all** medicines that you are taking.

### **Darunavir Mylan with food and drink**

See section 3 'How to take Darunavir Mylan'.

### **Pregnancy and breast-feeding**

Tell your doctor immediately if you are pregnant or planning to become pregnant. Pregnant women must not take darunavir with ritonavir unless specifically directed by the doctor. Pregnant women should not take darunavir with cobicistat.

Because of the potential for side effects in breast-fed infants, women should not breast-feed if they are receiving Darunavir Mylan.



Breast-feeding is not recommended in women living with HIV because HIV infection can be passed on to the baby in breast milk. If you are breast-feeding, or thinking about breast-feeding, you should discuss it with your doctor as soon as possible.

### **Driving and using machines**

Do not operate machines or drive if you feel dizzy after taking Darunavir Mylan.

### **Darunavir Mylan contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially ‘sodium-free’.

## **3. How to take Darunavir Mylan**

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure.

Even if you feel better, do not stop taking Darunavir Mylan and ritonavir without talking to your doctor.

After therapy has been initiated, the dose or dosage form must not be changed or therapy must not be stopped without instruction of the doctor.

### **Dose for adults who have not taken antiretroviral medicines before (your doctor will determine this)**

You will require a different dose of darunavir which cannot be administered with these 300 milligram tablets. Other strengths of Darunavir Mylan are available.

### **Dose for adults who have taken antiretroviral medicines before (your doctor will determine this)**

The dose is either:

- 600 milligram darunavir (2 tablets containing 300 milligram of darunavir or 1 tablet containing 600 milligram of darunavir) together with 100 milligram ritonavir twice daily.
- OR
- 800 milligram darunavir (2 tablets containing 400 milligram of darunavir or 1 tablet containing 800 milligram of darunavir) together with 100 milligram ritonavir once daily. Darunavir Mylan 400 milligram and 800 milligram tablets are only to be used to construct the once daily 800 milligram regimen.

Please discuss with your doctor which dose is right for you.

### **Instructions for adults**

- Take darunavir always together with ritonavir. Darunavir cannot work properly without ritonavir.
- In the morning, take two 300 milligram darunavir tablets together with 100 milligram ritonavir.
- In the evening, take two 300 milligram darunavir tablets together with 100 milligram ritonavir.
- Take darunavir with food. Darunavir cannot work properly without food. The type of food is not important.
- Swallow the tablets with a drink such as water or milk.
- Darunavir 75 milligram and 150 milligram tablets have been developed for use in children, but can also be used in adults in some cases.

### **Dose for children of 3 years of age and above, weighing at least 15 kilograms who have not taken antiretroviral medicines before (your child's doctor will determine this)**

The doctor will work out the right once daily dose based on the weight of the child (see table below). This dose must not exceed the recommended adult dose, which is 800 milligram darunavir together with 100 milligram ritonavir once a day.

The doctor will inform you on how much Darunavir Mylan tablets and how much ritonavir (capsules, tablets or solution) the child must take.

Weight	One darunavir dose is	One ritonavir <sup>a</sup> dose is
between 15 and 30 kilograms	600 milligram	100 milligram
between 30 and 40 kilograms	675 milligram	100 milligram
more than 40 kilograms	800 milligram	100 milligram

<sup>a</sup> ritonavir oral solution: 80 milligram per milliliter

**Dose for children of 3 years of age and above, weighing at least 15 kilograms who have taken antiretroviral medicines before (your child's doctor will determine this)**

The doctor will work out the right dose based on the weight of the child (see table below). The doctor will determine if once daily dosing or twice daily dosing is appropriate for the child. This dose must not exceed the recommended adult dose, which is 600 milligram darunavir together with 100 milligram ritonavir two times per day or 800 milligram darunavir together with 100 milligram ritonavir once a day.

The doctor will inform you on how many Darunavir Mylan tablets and how much ritonavir (capsules, tablets or solution) the child must take. Tablets of lower strengths are available and your doctor may have prescribed a certain combination of tablets to construct the appropriate dosing regimen.

Other forms of this medicine may be more suitable for children: ask your doctor or pharmacist.

**Twice daily dosing**

Weight	One dose is
between 15 and 30 kilograms	375 milligram darunavir + 50 milligram ritonavir twice a day
between 30 and 40 kilograms	450 milligram darunavir + 60 milligram ritonavir twice a day
more than 40 kilograms*	600 milligram darunavir + 100 milligram ritonavir twice a day

\* For children aged 12 or more and weighing at least 40 kilograms, your child's doctor will determine if Darunavir Mylan 800 milligram once daily dosing may be used. This cannot be administered with these 300 milligram tablets. Other strengths of Darunavir Mylan are available.

**Once daily dosing**

Weight	One darunavir dose is	One ritonavir <sup>a</sup> dose is
between 15 and 30 kilograms	600 milligram	100 milligram
between 30 and 40 kilograms	675 milligram	100 milligram
more than 40 kilograms	800 milligram	100 milligram

<sup>a</sup> ritonavir oral solution: 80 milligram per milliliter

**Instructions for children**

- The child must take darunavir always together with ritonavir. Darunavir cannot work properly without ritonavir.
- The child must take the appropriate doses of darunavir and ritonavir two times per day or once a day. If prescribed darunavir twice daily the child must take one dose in the morning, and one dose in the evening. Your child's doctor will determine the appropriate dosing regimen for your child.
- The child must take darunavir with food. Darunavir cannot work properly without food. The type of food is not important.
- The child must swallow the tablets with a drink such as water or milk.

**If you take more Darunavir Mylan than you should**

Contact your doctor, pharmacist or nurse immediately.

**If you forget to take Darunavir Mylan**

If you notice **within 6 hours**, you must take the tablets immediately. Always take with ritonavir and food. If you notice **after 6 hours**, then skip the intake and take the next doses as usual. Do not take a double dose to make up for a forgotten dose.

**If you vomit after taking Darunavir Mylan and ritonavir**

If you vomit **within 4 hours** of taking the medicine, another dose of Darunavir Mylan and ritonavir should be taken with food as soon as possible. If you vomit **more than 4 hours** after taking the medicine, then you do not need to take another dose of Darunavir Mylan and ritonavir until the next regularly scheduled time.

Contact your doctor **if you are uncertain** about what to do if you miss a dose or vomit.

**Do not stop taking Darunavir Mylan without talking to your doctor first**

Anti-HIV medicines may make you feel better. Even when you feel better, do not stop taking Darunavir Mylan. Talk to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

**4. Possible side effects**

During HIV therapy there may be an increase in weight and in levels of blood lipids and glucose. This is partly linked to restored health and life style, and in the case of blood lipids sometimes to the HIV medicines themselves. Your doctor will test for these changes.

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Tell your doctor if you develop any of the following side effects**

Liver problems that may occasionally be severe have been reported. Your doctor should do blood tests before you start darunavir. If you have chronic hepatitis B or C infection, your doctor should check your blood tests more often because you have an increased chance of developing liver problems. Talk to your doctor about the signs and symptoms of liver problems. These may include yellowing of your skin or whites of your eyes, dark (tea coloured) urine, pale coloured stools (bowel movements), nausea, vomiting, loss of appetite, or pain, aching, or pain and discomfort on your right side below your ribs.

Skin rash (more often when used in combination with raltegravir), itching. The rash is usually mild to moderate. A skin rash might also be a symptom of a rare severe situation. It is therefore important to talk to your doctor if you develop a rash. Your doctor will advise you how to deal with your symptoms or whether darunavir must be stopped.

Other severe side effects were diabetes (common), and inflammation of the pancreas (uncommon).  
Very common side effects (may affect more than 1 in 10 people)

- diarrhoea.

Common side effects (may affect up to 1 in 10 people)

- vomiting, nausea, abdominal pain or distension, dyspepsia, flatulence
- headache, tiredness, dizziness, drowsiness, numbness, tingling or pain in hands or feet, loss of strength, difficulty falling asleep.

Uncommon side effects (may affect up to 1 in 100 people)

- chest pain, changes in electrocardiogram, rapid heart beating
- decreased or abnormal skin sensitivity, pins and needles, attention disturbance, loss of memory, problems with your balance
- difficulty breathing, cough, nosebleed, throat irritation
- inflammation of the stomach or mouth, heartburn, retching, dry mouth, discomfort of the abdomen, constipation, belching
- kidney failure, kidney stones, difficult discharge of urine, frequent or excessive passage of urine, sometimes at night
- urticaria, severe swelling of the skin and other tissues (most often the lips or the eyes), eczema, excessive sweating, night sweats, hair loss, acne, scaly skin, colouration of nails

- muscle pain, muscle cramps or weakness, pain in extremity, osteoporosis
- slowing down of the thyroid gland function. This can be seen in a blood test.
- high blood pressure, flushing
- red or dry eyes
- fever, swelling of lower limbs due to fluids, malaise, irritability, pain
- symptoms of infection, herpes simplex
- erectile dysfunction, enlargement of breasts
- sleeping problems, sleepiness, depression, anxiety, abnormal dreams, decrease in sexual drive

Rare side effects (may affect up to 1 in 1,000 people)

- a reaction called DRESS [severe rash, which may be accompanied by fever, fatigue, swelling of the face or lymph glands, increase of eosinophils (type of white blood cells), effects on liver, kidney or lung]
- heart attack, slow heart beating, palpitations
- visual disturbance
- chills, feeling abnormal
- a feeling of confusion or disorientation, altered mood, restlessness
- fainting, epileptic fits, changes or loss of taste
- mouth sores, vomiting blood, inflammation of the lips, dry lips, coated tongue
- running nose
- skin lesions, dry skin
- stiffness of muscles or joints, joint pain with or without inflammation
- changes in some values of your blood cells or chemistry. These can be seen in the results of blood and/or urine tests. Your doctor will explain these to you. Examples are: increase in some white blood cells
- darunavir crystals in the kidney causing kidney disease.

Some side effects are typical for anti-HIV medicines in the same family as darunavir. These are:

- muscle pain, tenderness or weakness. On rare occasions, these muscle disorders have been serious.

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Darunavir Mylan**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, blister and on the bottle after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

For bottles: Once opened, use within 100 days.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away any medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Darunavir Mylan contains

- The active substance is darunavir. Each tablet contains 300 milligram of darunavir.
- The other ingredients are colloidal anhydrous silica, cellulose microcrystalline, crospovidone (Type A), sodium starch glycolate, hypromellose, magnesium stearate. The film-coating contains polyvinyl alcohol partially hydrolysed, titanium dioxide (E171), macrogol and talc.

### What Darunavir Mylan looks like and contents of the pack

Darunavir Mylan 300 mg film-coated tablets are white to off-white, oval shaped, with two curved sides, marked with M on one side and DV3 on the other side.

Darunavir Mylan 300 mg film-coated tablets are available in blister packs containing 30, 60 and 120 tablets and in plastic bottles containing 30 and 120 tablets.

Not all pack sizes may be marketed.

### Marketing Authorisation Holder

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

### Manufacturer

Mylan Hungary Kft  
H-2900 Komárom, Mylan utca 1  
Hungary

McDermott Laboratories Limited trading as Gerard Laboratories  
35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13  
Ireland

Mylan Germany GmbH  
Zweigniederlassung Bad Homburg v. d. Hoehe, Benzstrasse 1  
Bad Homburg v. d. Hoehe  
Hessen, 61352  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### België/Belgique/Belgien

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00

#### Lietuva

Mylan Healthcare UAB  
Tel: +370 5 205 1288

#### България

Майлан ЕООД  
Тел: +359 2 44 55 400

#### Luxembourg/Luxemburg

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00  
(Belgique/Belgien)

#### Česká republika

Viatrix CZ s.r.o.  
Tel: + 420 222 004 400

#### Magyarország

Mylan EPD Kft.  
Tel.: + 36 1 465 2100

**Danmark**

Viatriis ApS  
Tlf: +45 28 11 69 32

**Deutschland**

Viatriis Healthcare GmbH  
Tel: +49 800 0700 800

**Eesti**

BGP Products Switzerland GmbH  
Eesti filiaal  
Tel: + 372 6363 052

**Ελλάδα**

Generics Pharma Hellas EΠE  
Τηλ: +30 210 993 6410

**España**

Viatriis Pharmaceuticals, S.L.U.  
Tel: + 34 900 102 712

**France**

Viatriis Santé  
Tél: +33 4 37 25 75 00

**Hrvatska**

Viatriis Hrvatska d.o.o.  
Tel: +385 1 23 50 599

**Ireland**

Mylan Ireland Limited  
Tel: +353 1 8711600

**Ísland**

Icepharma hf.  
Sími: +354 540 8000

**Italia**

Mylan Italia S.r.l.  
Tel: + 39 02 612 46921

**Κύπρος**

Varnavas Hadjipanayis Ltd  
Τηλ: + 357 2220 7700

**Latvija**

Mylan Healthcare SIA  
Tel: +371 676 055 80

**Malta**

V.J. Salomone Pharma Ltd  
Tel: + 356 21 22 01 74

**Nederland**

Mylan BV  
Tel: +31 (0)20 426 3300

**Norge**

Viatriis AS  
Tlf: + 47 66 75 33 00

**Österreich**

Arcana Arzneimittel GmbH  
Tel: +43 1 416 2418

**Polska**

Mylan Healthcare Sp. z o.o.  
Tel.: + 48 22 546 64 00

**Portugal**

Mylan, Lda.  
Tel: + 351 214 127 200

**România**

BGP Products SRL  
Tel: +40 372 579 000

**Slovenija**

Viatriis d.o.o.  
Tel: + 386 1 23 63 180

**Slovenská republika**

Viatriis Slovakia s.r.o.  
Tel: +421 2 32 199 100

**Suomi/Finland**

Viatriis Oy  
Puh/Tel: +358 20 720 9555

**Sverige**

Viatriis AB  
Tel: +46 (0)8 630 19 00

**United Kingdom (Northern Ireland)**

Mylan IRE Healthcare Limited  
Tel: +353 18711600

**This leaflet was last revised in {MM/YYYY}.**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu>.

## **Package leaflet: Information for the user**

### **Darunavir Mylan 400 mg film-coated tablets**

darunavir

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Darunavir Mylan is and what it is used for
2. What you need to know before you take Darunavir Mylan
3. How to take Darunavir Mylan
4. Possible side effects
5. How to store Darunavir Mylan
6. Contents of the pack and other information

#### **1. What Darunavir Mylan is and what it is used for**

##### **What is Darunavir Mylan?**

Darunavir Mylan contains the active substance darunavir. Darunavir is an antiretroviral medicine used in the treatment of Human Immunodeficiency Virus (HIV) infection. It belongs to a group of medicines called protease inhibitors. Darunavir works by reducing the amount of HIV in your body. This will improve your immune system and reduces the risk of developing illnesses linked to HIV infection.

##### **What it is used for?**

Darunavir is used to treat adults and children 3 years of age and above, at least 40 kilograms body weight) who are infected by HIV and

- who have not used antiretroviral medicines before.
- in certain patients who have used antiretroviral medicines before (your doctor will determine this).

Darunavir must be taken in combination with a low dose of cobicistat or ritonavir and other anti-HIV medicines. Your doctor will discuss with you which combination of medicines is best for you.

#### **2. What you need to know before you take Darunavir Mylan**

##### **Do not take Darunavir Mylan**

- if you are **allergic** to darunavir or any of the other ingredients of this medicine (listed in section 6) or to cobicistat or ritonavir.
- if you have **severe liver problems**. Ask your doctor if you are unsure about the severity of your liver disease. Some additional tests might be necessary.



**Do not combine Darunavir Mylan with any of the following medicines**

If you are taking any of these, ask your doctor about switching to another medicine.

Medicine	Purpose of the medicine
<i>Avanafil</i>	to treat erectile dysfunction
<i>Astemizole</i> or <i>terfenadine</i>	to treat allergy symptoms
<i>Triazolam</i> and <i>oral</i> (taken by mouth) <i>midazolam</i>	to help you sleep and/or relieve anxiety
<i>Cisapride</i>	to treat some stomach conditions
<i>Colchicine</i> (if you have kidney and/or liver problems)	to treat gout or familial Mediterranean fever
<i>Lurasidone</i> , <i>pimozide</i> , <i>quetiapine</i> or <i>sertindole</i>	to treat psychiatric conditions
<i>Ergot alkaloids</i> like <i>ergotamine</i> , <i>dihydroergotamine</i> , <i>ergometrine</i> and <i>methylergonovine</i>	to treat migraine headaches
<i>Amiodarone</i> , <i>bepiridil</i> , <i>dronedarone</i> , <i>ivabradine</i> , <i>quinidine</i> , <i>ranolazine</i>	to treat certain heart disorders e.g. abnormal heart beat
<i>Lovastatin</i> , <i>simvastatin</i> and <i>lomitapide</i>	to lower cholesterol levels
<i>Rifampicin</i>	to treat some infections such as tuberculosis
The combination product <i>lopinavir/ritonavir</i>	this anti-HIV medicine belongs to the same class as darunavir
<i>Elbasvir/grazoprevir</i>	to treat hepatitis C infection
<i>Alfuzosin</i>	to treat enlarged prostate
<i>Sildenafil</i>	to treat high blood pressure in the pulmonary circulation
<i>Ticagrelor</i>	to help stop the clumping of platelets in the treatment of patients with a history of a heart attack
<i>Naloxegol</i>	to treat opioid induced constipation
<i>Dapoxetine</i>	to treat premature ejaculation
<i>Domperidone</i>	to treat nausea and vomiting

Do not combine darunavir with products that contain St John's wort (*Hypericum perforatum*).

**Warnings and precautions**

Talk to your doctor, pharmacist or nurse before taking Darunavir Mylan.

Darunavir is not a cure for HIV infection.

People taking darunavir may still develop infections or other illnesses associated with HIV infection. You must keep in regular contact with your doctor.

People taking darunavir may develop a skin rash. Infrequently a rash may become severe or potentially life-threatening. Please contact your doctor whenever you develop a rash.

In patients taking darunavir and raltegravir (for HIV infection), rashes (generally mild or moderate) may occur more frequently than in patients taking either medicine separately.

**Tell your doctor about your situation BEFORE and DURING your treatment**

Make sure that you check the following points and tell your doctor if any of these apply to you.

- Tell your doctor if you have had **problems with your liver** before, including hepatitis B or C infection. Your doctor may evaluate how severe your liver disease is before deciding if you can take darunavir.
- Tell your doctor if you have **diabetes**. Darunavir might increase sugar levels in the blood.
- Tell your doctor immediately if you notice any **symptoms of infection** (for example enlarged lymph nodes and fever). In some patients with advanced HIV infection and a history of opportunistic infection, signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started. It is believed that these symptoms are due to an

- improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms.
- In addition to the opportunistic infections, autoimmune disorders (a condition that occurs when the immune system attacks healthy body tissue) may also occur after you start taking medicines for the treatment of your HIV infection. Autoimmune disorders may occur many months after the start of treatment. If you notice any symptoms of infection or other symptoms such as muscle weakness, weakness beginning in the hands and feet and moving up towards the trunk of the body, palpitations, tremor or hyperactivity, please inform your doctor immediately to seek necessary treatment.
  - Tell your doctor if you have **haemophilia**. Darunavir might increase the risk of bleeding.
  - Tell your doctor if you are **allergic to sulphonamides** (e.g. used to treat certain infections).
  - Tell your doctor if you notice any **musculoskeletal problems**. Some patients taking combination antiretroviral therapy may develop a bone disease called osteonecrosis (death of bone tissue caused by loss of blood supply to the bone). The length of combination antiretroviral therapy, corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index, among others, may be some of the many risk factors for developing this disease. Signs of osteonecrosis are joint stiffness, aches and pains (especially of the hip, knee and shoulder) and difficulty in movement. If you notice any of these symptoms please inform your doctor.

### **Elderly**

Darunavir has only been used in limited numbers of patients 65 years or older. If you belong to this age group, please discuss with your doctor if you can use Darunavir Mylan.

### **Children and adolescents**

Darunavir is not for use in children younger than 3 years of age or weighing less than 15 kilograms.

### **Other medicines and Darunavir Mylan**

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines.

There are some medicines that **you must not combine** with darunavir. These are mentioned above under the heading '**Do not combine Darunavir Mylan with any of the following medicines:**'

In most cases, darunavir can be combined with anti-HIV medicines belonging to another class [e.g. NRTIs (nucleoside reverse transcriptase inhibitors), NNRTIs (non-nucleoside reverse transcriptase inhibitors), CCR5 antagonists and FIs (fusion inhibitors)]. Darunavir with cobicistat or ritonavir has not been tested with all PIs (protease inhibitors) and must not be used with other HIV PIs. In some cases dosage of other medicines might need to be changed. Therefore always tell your doctor if you take other anti-HIV medicines and follow your doctor's instruction carefully on which medicines can be combined.

The effects of darunavir might be reduced if you take any of the following products. Tell your doctor if you take:

- *Phenobarbital, phenytoin* (to prevent seizures)
- *Dexamethasone* (corticosteroid)
- *Efavirenz* (HIV infection)
- *Rifapentine, rifabutin* (medicines to treat some infections such as tuberculosis)
- *Saquinavir* (HIV infection).

The effects of other medicines might be influenced if you take darunavir and your doctor might want to do some additional blood tests. Tell your doctor if you take:

- *Amlodipine, diltiazem, disopyramide, carvedilol, felodipine, flecainide, lidocaine, metoprolol, mexiletine, nifedipine, nicardipine, propafenone, timolol, verapamil* (for heart disease) as the therapeutic effect or side effects of these medicines may be increased.
- *Apixaban, dabigatran etexilate, edoxaban, rivaroxaban, warfarin, clopidogrel* (to reduce clotting of the blood) as their therapeutic effect or side effects may be altered.

- Oestrogen-based hormonal contraceptives and hormonal replacement therapy. Darunavir might reduce its effectiveness. When used for birth control, alternative methods of non-hormonal contraception are recommended.
- *Ethinylestradiol/drospirenone*. Darunavir might increase the risk for elevated potassium levels by drospirenone.
- *Atorvastatin, pravastatin, rosuvastatin* (to lower cholesterol levels). The risk of muscle damage might be increased. Your doctor will evaluate which cholesterol lowering regimen is best for your specific situation.
- *Clarithromycin* (antibiotic)
- *Ciclosporin, everolimus, tacrolimus, sirolimus* (for dampening down your immune system) as the therapeutic effect or side effects of these medicines might be increased.
- *Corticosteroids including betamethasone, budesonide, fluticasone, mometasone, prednisone, triamcinolone*. These medicines are used to treat allergies, asthma, inflammatory bowel diseases, inflammatory conditions of the skin, eyes, joints and muscles and other inflammatory conditions. These medicines are generally taken orally, inhaled, injected or applied to the skin. If alternatives cannot be used, its use should only take place after medical evaluation and under close monitoring by your doctor for corticosteroid side effects.
- *Buprenorphine/naloxone* (medicines to treat opioid dependence)
- *Salmeterol* (medicine to treat asthma)
- *Artemether/lumefantrine* (a combination medicine to treat malaria)
- *Dasatinib, everolimus, irinotecan, nilotinib, vinblastine, vincristine* (to treat cancer)
- *Sildenafil, tadalafil, vardenafil* (for erectile dysfunction or to treat a heart and lung disorder called pulmonary arterial hypertension)
- *Glecaprevir/pibrentasvir* (to treat hepatitis C infection)
- *Fentanyl, oxycodone, tramadol* (to treat pain)
- *Fesoterodine, solifenacin* (to treat urologic disorders).

Your doctor might want to do some additional blood tests and the dosage of other medicines might need to be changed since either their own or darunavir's therapeutic effect or side effects may be influenced when combined.

Tell your doctor if you take:

- *Dabigatran etexilate, edoxaban, warfarin* (to reduce clotting of the blood)
- *Alfentanil* (injectable strong and short-acting painkiller that is used for surgical procedures)
- *Digoxin* (to treat certain heart disorders)
- *Clarithromycin* (antibiotic)
- *Itraconazole, isavuconazole, fluconazole, posaconazole, clotrimazole* (to treat fungal infections). Voriconazole should only be taken after medical evaluation.
- *Rifabutin* (against bacterial infections)
- *Sildenafil, vardenafil, tadalafil* (for erectile dysfunction or high blood pressure in the pulmonary circulation)
- *Amitriptyline, desipramine, imipramine, nortriptyline, paroxetine, sertraline, trazodone* (to treat depression and anxiety)
- *Maraviroc* (to treat HIV infection)
- *Methadone* (to treat opiate dependence)
- *Carbamazepine, clonazepam* (to prevent seizures or to treat certain types of nerve pain)
- *Colchicine* (to treat gout or familial Mediterranean fever)
- *Bosentan* (to treat high blood pressure in the pulmonary circulation)
- *Buspirone, clorazepate, diazepam, estazolam, flurazepam, midazolam when used as injection, zolpidem* (sedative agents)
- *Perphenazine, risperidone, thioridazine* (to treat psychiatric conditions)
- *Metformin* (to treat type 2 diabetes).

This is **not** a complete list of medicines. Tell your healthcare provider about **all** medicines that you are taking.

### **Darunavir Mylan with food and drink**

See section 3 'How to take Darunavir Mylan.

### **Pregnancy and breast-feeding**

Tell your doctor immediately if you are pregnant or planning to become pregnant. Pregnant women must not take darunavir with ritonavir unless specifically directed by the doctor. Pregnant women should not take darunavir with cobicistat.

Because of the potential for side effects in breast-fed infants, women should not breast-feed if they are receiving Darunavir Mylan.

Breast-feeding is not recommended in women living with HIV because HIV infection can be passed on to the baby in breast milk. If you are breast-feeding or thinking about breast-feeding, you should discuss it with your doctor as soon as possible.

### **Driving and using machines**

Do not operate machines or drive if you feel dizzy after taking darunavir.

### **Darunavir Mylan contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

## **3. How to take Darunavir Mylan**

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure.

Even if you feel better, do not stop taking darunavir and cobicistat or ritonavir without talking to your doctor.

After therapy has been initiated, the dose or dosage form must not be changed or therapy must not be stopped without instruction of the doctor.

Darunavir 400 mg tablets are only to be used to construct the once daily 800 mg regimen.

### **Dose for adults who have not taken antiretroviral medicines before (your doctor will determine this)**

The usual dose of darunavir is 800 milligram (2 tablets containing 400 milligram of Darunavir Mylan or 1 tablet containing 800 milligram of Darunavir Mylan) once daily.

You must take darunavir every day and always in combination with 150 milligram of cobicistat or 100 milligram of ritonavir and with food. Darunavir cannot work properly without cobicistat or ritonavir and food. You must eat a meal or a snack within 30 minutes prior to taking your darunavir and cobicistat or ritonavir. The type of food is not important. Even if you feel better, do not stop taking darunavir and cobicistat or ritonavir without talking to your doctor.

### **Instructions for adults**

- Take two 400 milligram tablets at the same time, once a day, every day.
- Take darunavir always together with 150 milligram of cobicistat or 100 milligram of ritonavir.
- Take darunavir with food.
- Swallow the tablets with a drink such as water or milk.
- Take your other HIV medicines used in combination with darunavir and cobicistat or ritonavir as recommended by your doctor.

### **Dose for adults who have taken antiretroviral medicines before (your doctor will determine this)**

The dose is either:

- 800 milligram darunavir (2 tablets containing 400 milligram of darunavir or 1 tablet containing 800 milligram of darunavir) together with 150 milligram cobicistat or 100 milligram ritonavir once daily.

OR

- 600 milligram darunavir (2 tablets containing 300 milligram of darunavir or 1 tablet containing 600 milligram of darunavir) together with 100 milligram ritonavir twice daily.

Please discuss with your doctor which dose is right for you.

**Dose for children 3 years of age and above with ritonavir, and 12 years of age and above with cobicistat, weighing more than 40 kilograms who have not taken antiretroviral medicines before (your child's doctor will determine this)**

- The usual dose of darunavir is 800 milligram (2 tablets containing 400 milligram of darunavir or 1 tablet containing 800 milligram of darunavir) together with 100 milligram ritonavir or 150 milligram of cobicistat once daily.

**Dose for children 3 years of age and above with ritonavir, and 12 years of age and above with cobicistat, weighing more than 40 kilograms who have taken antiretroviral medicines before (your child's doctor will determine this)**

The dose is either:

- 800 milligram darunavir (2 tablets containing 400 milligram of darunavir or 1 tablet containing 800 milligram of darunavir) together with 100 milligram ritonavir or 150 milligram of cobicistat once daily.
- OR
- 600 milligram darunavir (2 tablets containing 300 milligram of darunavir or 1 tablet containing 600 milligram of darunavir) together with 100 milligram ritonavir twice daily.

Please discuss with your doctor which dose is right for you.

***Instructions for children 3 years of age and above with ritonavir, and 12 years of age and above with cobicistat, weighing more than 40 kilograms***

- Take 800 milligram darunavir (2 tablets containing 400 milligram of darunavir or 1 tablet containing 800 milligram of darunavir) at the same time, once a day, every day.
- Take darunavir always together with 100 milligram of ritonavir or 150 milligram of cobicistat.
- Take darunavir with food.
- Swallow the tablets with a drink such as water or milk.
- Take your other HIV medicines used in combination with darunavir and ritonavir or cobicistat as recommended by your doctor.

**If you take more Darunavir Mylan than you should**

Contact your doctor, pharmacist or nurse immediately.

**If you forget to take Darunavir Mylan**

If you notice **within 12 hours**, you must take the tablets immediately. Always take with cobicistat or ritonavir and food. If you notice **after 12 hours**, then skip the intake and take the next doses as usual. Do not take a double dose to make up for a forgotten dose.

**If you vomit after taking Darunavir Mylan and cobicistat or ritonavir**

If you vomit **within 4 hours** of taking the medicine, another dose of Darunavir Mylan and cobicistat or ritonavir should be taken with food as soon as possible. If you vomit **more than 4 hours** after taking the medicine, then you do not need to take another dose of Darunavir Mylan and cobicistat or ritonavir until the next regularly scheduled time.

Contact your doctor **if you are uncertain** about what to do if you miss a dose or vomit.

**Do not stop taking Darunavir Mylan without talking to your doctor first**

Anti-HIV medicines may make you feel better. Even when you feel better, do not stop taking Darunavir Mylan. Talk to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

**4. Possible side effects**

During HIV therapy there may be an increase in weight and in levels of blood lipids and glucose. This is partly linked to restored health and life style, and in the case of blood lipids sometimes to the HIV medicines themselves. Your doctor will test for these changes.

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Tell your doctor if you develop any of the following side effects**

Liver problems that may occasionally be severe have been reported. Your doctor should do blood tests before you start darunavir. If you have chronic hepatitis B or C infection, your doctor should check your blood tests more often because you have an increased chance of developing liver problems. Talk to your doctor about the signs and symptoms of liver problems. These may include yellowing of your skin or whites of your eyes, dark (tea coloured) urine, pale coloured stools (bowel movements), nausea, vomiting, loss of appetite, or pain, aching, or pain and discomfort on your right side below your ribs.

Skin rash (more often when used in combination with raltegravir), itching. The rash is usually mild to moderate. A skin rash might also be a symptom of a rare severe situation. It is therefore important to talk to your doctor if you develop a rash. Your doctor will advise you how to deal with your symptoms or whether darunavir must be stopped.

Other severe side effects were diabetes (common), and inflammation of the pancreas (uncommon).

Very common side effects (may affect more than 1 in 10 people)

- diarrhoea.

Common side effects (may affect up to 1 in 10 people)

- vomiting, nausea, abdominal pain or distension, dyspepsia, flatulence
- headache, tiredness, dizziness, drowsiness, numbness, tingling or pain in hands or feet, loss of strength, difficulty falling asleep.

Uncommon side effects (may affect up to 1 in 100 people)

- chest pain, changes in electrocardiogram, rapid heart beating
- decreased or abnormal skin sensitivity, pins and needles, attention disturbance, loss of memory, problems with your balance
- difficulty breathing, cough, nosebleed, throat irritation
- inflammation of the stomach or mouth, heartburn, retching, dry mouth, discomfort of the abdomen, constipation, belching
- kidney failure, kidney stones, difficult discharge of urine, frequent or excessive passage of urine, sometimes at night
- urticaria, severe swelling of the skin and other tissues (most often the lips or the eyes), eczema, excessive sweating, night sweats, hair loss, acne, scaly skin, colouration of nails
- muscle pain, muscle cramps or weakness, pain in extremity, osteoporosis
- slowing down of the thyroid gland function. This can be seen in a blood test.
- high blood pressure, flushing
- red or dry eyes
- fever, swelling of lower limbs due to fluids, malaise, irritability, pain
- symptoms of infection, herpes simplex
- erectile dysfunction, enlargement of breasts

- sleeping problems, sleepiness, depression, anxiety, abnormal dreams, decrease in sexual drive

Rare side effects (may affect up to 1 in 1,000 people)

- a reaction called DRESS [severe rash, which may be accompanied by fever, fatigue, swelling of the face or lymph glands, increase of eosinophils (type of white blood cells), effects on liver, kidney or lung]
- heart attack, slow heart beating, palpitations
- visual disturbance
- chills, feeling abnormal
- a feeling of confusion or disorientation, altered mood, restlessness
- fainting, epileptic fits, changes or loss of taste
- mouth sores, vomiting blood, inflammation of the lips, dry lips, coated tongue
- running nose
- skin lesions, dry skin
- stiffness of muscles or joints, joint pain with or without inflammation
- changes in some values of your blood cells or chemistry. These can be seen in the results of blood and/or urine tests. Your doctor will explain these to you. Examples are: increase in some white blood cells
- darunavir crystals in the kidney causing kidney disease.

Some side effects are typical for anti-HIV medicines in the same family as darunavir. These are:

- muscle pain, tenderness or weakness. On rare occasions, these muscle disorders have been serious.

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Darunavir Mylan**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, blister and on the bottle after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

For bottles: Once opened, use within 100 days.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away any medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Darunavir Mylan contains**

- The active substance is darunavir. Each tablet contains 400 milligrams of darunavir.
- The other ingredients are colloidal anhydrous silica, cellulose microcrystalline, crospovidone, sodium starch glycolate, hypromellose, magnesium stearate. The film-coating contains polyvinyl alcohol partially hydrolysed, titanium dioxide (E171), macrogol and talc.

## What Darunavir Mylan looks like and contents of the pack

Darunavir Mylan 400 mg film-coated tablets are white to off-white, oval shaped with two curved sides, marked with M on one side and DV4 on the other side.

Darunavir Mylan 400 mg film-coated tablets are available blister packs containing 30 and 60 tablets and in the HDPE bottles of 60 and 100 tablets.

Not all pack sizes may be marketed.

### Marketing Authorisation Holder

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

### Manufacturer

Mylan Hungary Kft  
H-2900 Komárom, Mylan utca 1  
Hungary

McDermott Laboratories Limited trading as Gerard Laboratories  
35/36 Baldoye Industrial Estate, Grange Road, Dublin 13  
Ireland

Mylan Germany GmbH  
Zweigniederlassung Bad Homburg v. d. Hoehe, Benzstrasse 1  
Bad Homburg v. d. Hoehe  
Hessen, 61352  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### **België/Belgique/Belgien**

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00

#### **Lietuva**

Mylan Healthcare UAB  
Tel: +370 5 205 1288

#### **България**

Майлан ЕООД  
Тел: +359 2 44 55 400

#### **Luxembourg/Luxemburg**

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00  
(Belgique/Belgien)

#### **Česká republika**

Viatrix CZ s.r.o.  
Tel: + 420 222 004 400

#### **Magyarország**

Mylan EPD Kft.  
Tel.: + 36 1 465 2100

#### **Danmark**

Viatrix ApS  
Tlf: +45 28 11 69 32

#### **Malta**

V.J. Salomone Pharma Ltd  
Tel: + 356 21 22 01 74

#### **Deutschland**

Viatrix Healthcare GmbH  
Tel: +49 800 0700 800

#### **Nederland**

Mylan BV  
Tel: +31 (0)20 426 3300



**Eesti**

BGP Products Switzerland GmbH Eesti filiaal  
Tel: + 372 6363 052

**Ελλάδα**

Generics Pharma Hellas EΠE  
Τηλ: +30 210 993 6410

**España**

Viatriis Pharmaceuticals, S.L.U.  
Tel: + 34 900 102 712

**France**

Viatriis Santé  
Tél: +33 4 37 25 75 00

**Hrvatska**

Viatriis Hrvatska d.o.o.  
Tel: +385 1 23 50 599

**Ireland**

Mylan Ireland Limited  
Tel: +353 1 8711600

**Ísland**

Icepharma hf.  
Sími: +354 540 8000

**Italia**

Mylan Italia S.r.l.  
Tel: + 39 02 612 46921

**Κύπρος**

Varnavas Hadjipanayis Ltd  
Τηλ: + 357 2220 7700

**Latvija**

Mylan Healthcare SIA  
Tel: +371 676 055 80

**Norge**

Viatriis AS  
Tlf: + 47 66 75 33 00

**Österreich**

Arcana Arzneimittel GmbH  
Tel: +43 1 416 2418

**Polska**

Mylan Healthcare Sp. z o.o.  
Tel.: + 48 22 546 64 00

**Portugal**

Mylan, Lda.  
Tel: + 351 214 127 200

**România**

BGP Products SRL  
Tel: +40 372 579 000

**Slovenija**

Viatriis d.o.o.  
Tel: + 386 1 23 63 180

**Slovenská republika**

Viatriis Slovakia s.r.o.  
Tel: +421 2 32 199 100

**Suomi/Finland**

Viatriis Oy  
Puh/Tel: +358 20 720 9555

**Sverige**

Viatriis AB  
Tel: +46 (0)8 630 19 00

**United Kingdom (Northern Ireland)**

Mylan IRE Healthcare Limited Tel: +353  
18711600

**This leaflet was last revised in {MM/YYYY}.**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu>.

## Package leaflet: Information for the user

### Darunavir Mylan 600 mg film-coated tablets

darunavir

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Darunavir Mylan is and what it is used for
2. What you need to know before you take Darunavir Mylan
3. How to take Darunavir Mylan
4. Possible side effects
5. How to store Darunavir Mylan
6. Contents of the pack and other information

#### 1. What Darunavir Mylan is and what it is used for

##### What is Darunavir Mylan?

Darunavir Mylan contains the active substance darunavir. Darunavir is an antiretroviral medicine used in the treatment of Human Immunodeficiency Virus (HIV) infection. It belongs to a group of medicines called protease inhibitors. Darunavir works by reducing the amount of HIV in your body. This will improve your immune system and reduces the risk of developing illnesses linked to HIV infection.

##### What it is used for?

Darunavir is used to treat adults and children of 3 years of age and above, and at least 15 kilogram body weight who are infected by HIV and who have already used other antiretroviral medicines. Darunavir must be taken in combination with a low dose of ritonavir and other anti-HIV medicines. Your doctor will discuss with you which combination of medicines is best for you.

#### 2. What you need to know before you take Darunavir Mylan

##### Do not take Darunavir Mylan

- if you are **allergic** to darunavir or any of the other ingredients of this medicine (listed in section 6) or to ritonavir.
- if you have **severe liver problems**. Ask your doctor if you are unsure about the severity of your liver disease. Some additional tests might be necessary.

##### Do not combine Darunavir Mylan with any of the following medicines

If you are taking any of these, ask your doctor about switching to another medicine.

Medicine	Purpose of the medicine
<i>Avanafil</i>	to treat erectile dysfunction
<i>Astemizole</i> or <i>terfenadine</i>	to treat allergy symptoms
<i>Triazolam</i> and <i>oral</i> (taken by mouth) <i>midazolam</i>	to help you sleep and/or relieve anxiety
<i>Cisapride</i>	to treat some stomach conditions

<b>Medicine</b>	<b>Purpose of the medicine</b>
<i>Colchicine</i> (if you have kidney and/or liver problems)	to treat gout or familial Mediterranean fever
<i>Lurasidone, pimozide, quetiapine or sertindole</i>	to treat psychiatric conditions
<i>Ergot alkaloids</i> like <i>ergotamine</i> , <i>dihydroergotamine</i> , <i>ergometrine</i> and <i>methylergonovine</i>	to treat migraine headaches
<i>Amiodarone, bepridil, dronedarone, ivabradine, quinidine, ranolazine</i>	to treat certain heart disorders e.g. abnormal heart beat
<i>Lovastatin, simvastatin and lomitapide</i>	to lower cholesterol levels
<i>Rifampicin</i>	to treat some infections such as tuberculosis
The combination product <i>lopinavir/ritonavir</i>	this anti-HIV medicine belongs to the same class as Darunavir Mylan
<i>Elbasvir/grazoprevir</i>	to treat hepatitis C infection
<i>Alfuzosin</i>	to treat enlarged prostate
<i>Sildenafil</i>	to treat high blood pressure in the pulmonary circulation
<i>Ticagrelor</i>	to help stop the clumping of platelets in the treatment of patients with a history of a heart attack
<i>Naloxegol</i>	to treat opioid induced constipation
<i>Dapoxetine</i>	to treat premature ejaculation
<i>Domperidone</i>	to treat nausea and vomiting

Do not combine darunavir with products that contain St John's wort (*Hypericum perforatum*).

### **Warnings and precautions**

Talk to your doctor, pharmacist or nurse before taking Darunavir Mylan.

Darunavir is not a cure for HIV infection.

People taking darunavir may still develop infections or other illnesses associated with HIV infection. You must keep in regular contact with your doctor.

People taking darunavir may develop a skin rash. Infrequently a rash may become severe or potentially life-threatening. Please contact your doctor whenever you develop a rash.

In patients taking darunavir and raltegravir (for HIV infection), rashes (generally mild or moderate) may occur more frequently than in patients taking either medicine separately.

### **Tell your doctor about your situation BEFORE and DURING your treatment**

Make sure that you check the following points and tell your doctor if any of these apply to you.

- Tell your doctor if you have had **problems with your liver** before, including hepatitis B or C infection. Your doctor may evaluate how severe your liver disease is before deciding if you can take darunavir.
- Tell your doctor if you have **diabetes**. Darunavir might increase sugar levels in the blood.
- Tell your doctor immediately if you notice any **symptoms of infection** (for example enlarged lymph nodes and fever). In some patients with advanced HIV infection and a history of opportunistic infection, signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started. It is believed that these symptoms are due to an improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms.
- In addition to the opportunistic infections, autoimmune disorders (a condition that occurs when the immune system attacks healthy body tissue) may also occur after you start taking medicines for the treatment of your HIV infection. Autoimmune disorders may occur many months after the start of treatment. If you notice any symptoms of infection or other symptoms such as muscle weakness, weakness beginning in the hands and feet and moving up towards the trunk of

the body, palpitations, tremor or hyperactivity, please inform your doctor immediately to seek necessary treatment.

- Tell your doctor if you have **haemophilia**. Darunavir might increase the risk of bleeding.
- Tell your doctor if you are **allergic to sulphonamides** (e.g. used to treat certain infections).
- Tell your doctor if you notice any **musculoskeletal problems**. Some patients taking combination antiretroviral therapy may develop a bone disease called osteonecrosis (death of bone tissue caused by loss of blood supply to the bone). The length of combination antiretroviral therapy, corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index, among others, may be some of the many risk factors for developing this disease. Signs of osteonecrosis are joint stiffness, aches and pains (especially of the hip, knee and shoulder) and difficulty in movement. If you notice any of these symptoms please inform your doctor.

### **Elderly**

Darunavir has only been used in limited numbers of patients 65 years or older. If you belong to this age group, please discuss with your doctor if you can use Darunavir Mylan.

### **Children**

Darunavir is not for use in children younger than 3 years of age or weighing less than 15 kilograms.

### **Other medicines and Darunavir Mylan**

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines.

There are some medicines that **you must not combine** with darunavir. These are mentioned above under the heading '**Do not combine Darunavir Mylan with any of the following medicines:**'

In most cases, darunavir can be combined with anti-HIV medicines belonging to another class [e.g. NRTIs (nucleoside reverse transcriptase inhibitors), NNRTIs (non-nucleoside reverse transcriptase inhibitors), CCR5 antagonists and FIs (fusion inhibitors)]. Darunavir with ritonavir has not been tested with all PIs (protease inhibitors) and must not be used with other HIV PIs. In some cases dosage of other medicines might need to be changed. Therefore always tell your doctor if you take other anti-HIV medicines and follow your doctor's instruction carefully on which medicines can be combined.

The effects of darunavir might be reduced if you take any of the following products. Tell your doctor if you take:

- *Phenobarbital, phenytoin* (to prevent seizures)
- *Dexamethasone* (corticosteroid)
- *Efavirenz* (HIV infection)
- *Rifapentine, rifabutin* (medicines to treat some infections such as tuberculosis)
- *Saquinavir* (HIV infection).

The effects of other medicines might be influenced if you take darunavir and your doctor might want to do some additional blood tests. Tell your doctor if you take:

- *Amlodipine, diltiazem, disopyramide, carvedilol, felodipine, flecainide, lidocaine, metoprolol, mexiletine, nifedipine, nicardipine, propafenone, timolol, verapamil* (for heart disease) as the therapeutic effect or side effects of these medicines may be increased.
- *Apixaban, dabigatran etexilate, edoxaban, rivaroxaban, warfarin, clopidogrel* (to reduce clotting of the blood) as their therapeutic effect or side effects may be altered.
- Oestrogen-based hormonal contraceptives and hormonal replacement therapy. Darunavir might reduce its effectiveness. When used for birth control, alternative methods of non-hormonal contraception are recommended.
- *Ethinylestradiol/drospirenone*. Darunavir might increase the risk for elevated potassium levels by drospirenone.
- *Atorvastatin, pravastatin, rosuvastatin* (to lower cholesterol levels). The risk of muscle damage might be increased. Your doctor will evaluate which cholesterol lowering regimen is best for your specific situation.
- *Clarithromycin* (antibiotic)

- *Ciclosporin, everolimus, tacrolimus, sirolimus* (for dampening down your immune system) as the therapeutic effect or side effects of these medicines might be increased.
- *Corticosteroids including betamethasone, budesonide, fluticasone, mometasone, prednisone, triamcinolone*. These medicines are used to treat allergies, asthma, inflammatory bowel diseases, inflammatory conditions of the skin, eyes, joints and muscles and other inflammatory conditions. These medicines are generally taken orally, inhaled, injected or applied to the skin. If alternatives cannot be used, its use should only take place after medical evaluation and under close monitoring by your doctor for corticosteroid side effects.
- *Buprenorphine/naloxone* (medicines to treat opioid dependence)
- *Salmeterol* (medicine to treat asthma)
- *Artemether/lumefantrine* (a combination medicine to treat malaria)
- *Dasatinib, everolimus, irinotecan, nilotinib, vinblastine, vincristine* (to treat cancer)
- *Sildenafil, tadalafil, vardenafil* (for erectile dysfunction or to treat a heart and lung disorder called pulmonary arterial hypertension)
- *Glecaprevir/pibrentasvir* (to treat hepatitis C infection)
- *Fentanyl, oxycodone, tramadol* (to treat pain)
- *Fesoterodine, solifenacin* (to treat urologic disorders).

Your doctor might want to do some blood tests and the dosage of other medicines might need to be changed since either their own or darunavir's therapeutic effect or side effects may be influenced when combined.

Tell your doctor if you take:

- *Dabigatran etexilate, edoxaban, warfarin* (to reduce clotting of the blood)
- *Alfentanil* (injectable strong and short-acting painkiller that is used for surgical procedures)
- *Digoxin* (to treat certain heart disorders)
- *Clarithromycin* (antibiotic)
- *Itraconazole, isavuconazole, fluconazole, posaconazole, clotrimazole* (to treat fungal infections). Voriconazole should only be taken after medical evaluation.
- *Rifabutin* (against bacterial infections)
- *Sildenafil, vardenafil, tadalafil* (for erectile dysfunction or high blood pressure in the pulmonary circulation)
- *Amitriptyline, desipramine, imipramine, nortriptyline, paroxetine, sertraline, trazodone* (to treat depression and anxiety)
- *Maraviroc* (to treat HIV infection)
- *Methadone* (to treat opiate dependence)
- *Carbamazepine, clonazepam* (to prevent seizures or to treat certain types of nerve pain)
- *Colchicine* (to treat gout or familial Mediterranean fever)
- *Bosentan* (to treat high blood pressure in the pulmonary circulation)
- *Buspirone, clorazepate, diazepam, estazolam, flurazepam, midazolam when used as injection, zolpidem* (sedative agents)
- *Perphenazine, risperidone, thioridazine* (to treat psychiatric conditions).

This is **not** a complete list of medicines. Tell your healthcare provider about **all** medicines that you are taking.

### **Darunavir Mylan with food and drink**

See section 3 'How to take Darunavir Mylan.

### **Pregnancy and breast-feeding**

Tell your doctor immediately if you are pregnant or planning to become pregnant. Pregnant women must not take darunavir with ritonavir unless specifically directed by the doctor. Pregnant women should not take darunavir with cobicistat.

Because of the potential for side effects in breast-fed infants, women should not breast-feed if they are receiving Darunavir Mylan.

Breast-feeding is not recommended in women living with HIV because HIV infection can be passed on to the baby in breast milk. If you are breast-feeding, or thinking about breast-feeding, you should discuss it with your doctor as soon as possible.

### **Driving and using machines**

Do not operate machines or drive if you feel dizzy after taking Darunavir Mylan.

### **Darunavir Mylan contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially ‘sodium-free’.

## **3. How to take Darunavir Mylan**

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure.

Even if you feel better, do not stop taking Darunavir Mylan and ritonavir without talking to your doctor.

After therapy has been initiated, the dose or dosage form must not be changed or therapy must not be stopped without instruction of the doctor.

### **Dose for adults who have not taken antiretroviral medicines before (your doctor will determine this)**

You will require a different dose of darunavir which cannot be administered with these 600 milligram tablets. Other strengths of Darunavir Mylan are available.

### **Dose for adults who have taken antiretroviral medicines before (your doctor will determine this)**

The dose is either:

- 600 milligram darunavir (2 tablets containing 300 milligram of darunavir or 1 tablet containing 600 milligram of darunavir) together with 100 milligram ritonavir twice daily.
- OR
- 800 milligram darunavir (2 tablets containing 400 milligram of darunavir or 1 tablet containing 800 milligram of darunavir) together with 100 milligram ritonavir once daily. Darunavir Mylan 400 milligram and 800 milligram tablets are only to be used to construct the once daily 800 milligram regimen.

Please discuss with your doctor which dose is right for you.

### **Instructions for adults**

- Take darunavir always together with ritonavir. Darunavir cannot work properly without ritonavir.
- In the morning, take one 600 milligram darunavir tablet together with 100 milligram ritonavir.
- In the evening, take one 600 milligram darunavir tablet together with 100 milligram ritonavir.
- Take darunavir with food. Darunavir cannot work properly without food. The type of food is not important.
- Swallow the tablets with a drink such as water or milk.
- Darunavir Mylan 75 milligram and 150 milligram tablets have been developed for use in children, but can also be used in adults in some cases.

### **Dose for children of 3 years of age and above, weighing at least 15 kilograms who have not taken antiretroviral medicines before (your child's doctor will determine this)**

The doctor will work out the right once daily dose based on the weight of the child (see table below). This dose must not exceed the recommended adult dose, which is 800 milligram darunavir together with 100 milligram ritonavir once a day.

The doctor will inform you on how much Darunavir Mylan tablets and how much ritonavir (capsules, tablets or solution) the child must take.

Weight	One darunavir dose is	One ritonavir <sup>a</sup> dose is
between 15 and 30 kilograms	600 milligram	100 milligram
between 30 and 40 kilograms	675 milligram	100 milligram
more than 40 kilograms	800 milligram	100 milligram

<sup>a</sup> ritonavir oral solution: 80 milligram per milliliter

**Dose for children of 3 years of age and above, weighing at least 15 kilograms who have taken antiretroviral medicines before (your child's doctor will determine this)**

The doctor will work out the right dose based on the weight of the child (see table below). The doctor will determine if once daily dosing or twice daily dosing is appropriate for the child. This dose must not exceed the recommended adult dose, which is 600 milligram darunavir together with 100 milligram of ritonavir two times per day or 800 milligram darunavir together with 100 milligram ritonavir once a day.

The doctor will inform you on how many Darunavir Mylan tablets and how much ritonavir (capsules, tablets or solution) the child must take. Tablets of lower strengths are available to construct the appropriate dosing regimen.

Other forms of this medicine may be more suitable for children: ask your doctor or pharmacist.

**Twice daily dosing**

Weight	One dose is
between 15 and 30 kilograms	375 milligram darunavir + 50 milligram ritonavir twice a day
between 30 and 40 kilograms	450 milligram darunavir + 60 milligram ritonavir twice a day
more than 40 kilograms*	600 milligram darunavir + 100 milligram ritonavir twice a day

\* For children aged 12 or more and weighing at least 40 kilograms, your child's doctor will determine if Darunavir Mylan 800 milligram once daily dosing may be used. This cannot be administered with these 600 milligram tablets. Other strengths of Darunavir Mylan are available.

**Once daily dosing**

Weight	One darunavir dose is	One ritonavir <sup>a</sup> dose is
between 15 and 30 kilograms	600 milligram	100 milligram
between 30 and 40 kilograms	675 milligram	100 milligram
more than 40 kilograms	800 milligram	100 milligram

<sup>a</sup> ritonavir oral solution: 80 milligram per milliliter

**Instructions for children**

- The child must take darunavir always together with ritonavir. Darunavir cannot work properly without ritonavir.
- The child must take the appropriate doses of darunavir and ritonavir two times per day or once a day. If prescribed darunavir twice daily the child must take one dose in the morning, and one dose in the evening. Your child's doctor will determine the appropriate dosing regimen for your child.
- The child must take darunavir with food. Darunavir cannot work properly without food. The type of food is not important.
- The child must swallow the tablets with a drink such as water or milk.
- Darunavir Mylan 75 mg and 150 mg tablets have been developed for use in children weighing less than 40 kilograms, but can also be used in adults in some cases.

**If you take more Darunavir Mylan than you should**

Contact your doctor, pharmacist or nurse immediately.

**If you forget to take Darunavir Mylan**

If you notice **within 6 hours**, you must take your missed dose immediately. Always take with ritonavir and food. If you notice **after 6 hours**, then skip the intake and take the next doses as usual. Do not take a double dose to make up for a forgotten dose.

**If you vomit after taking Darunavir Mylan and ritonavir**

If you vomit **within 4 hours** of taking the medicine, another dose of Darunavir Mylan and ritonavir should be taken with food as soon as possible. If you vomit **more than 4 hours** after taking the medicine, then you do not need to take another dose of Darunavir Mylan and ritonavir until the next regularly scheduled time.

Contact your doctor **if you are uncertain** about what to do if you miss a dose or vomit.

**Do not stop taking Darunavir Mylan without talking to your doctor first**

Anti-HIV medicine may make you feel better. Even when you feel better, do not stop taking Darunavir Mylan. Talk to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

**4. Possible side effects**

During HIV therapy there may be an increase in weight and in levels of blood lipids and glucose. This is partly linked to restored health and life style, and in the case of blood lipids sometimes to the HIV medicines themselves. Your doctor will test for these changes.

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Tell your doctor if you develop any of the following side effects**

Liver problems that may occasionally be severe have been reported. Your doctor should do blood tests before you start darunavir. If you have chronic hepatitis B or C infection, your doctor should check your blood tests more often because you have an increased chance of developing liver problems. Talk to your doctor about the signs and symptoms of liver problems. These may include yellowing of your skin or whites of your eyes, dark (tea coloured) urine, pale coloured stools (bowel movements), nausea, vomiting, loss of appetite, or pain, aching, or pain and discomfort on your right side below your ribs.

Skin rash (more often when used in combination with raltegravir), itching. The rash is usually mild to moderate. A skin rash might also be a symptom of a rare severe situation. It is therefore important to talk to your doctor if you develop a rash. Your doctor will advise you how to deal with your symptoms or whether darunavir must be stopped.

Other severe side effects were diabetes (common), and inflammation of the pancreas (uncommon).

Very common side effects (may affect more than 1 in 10 people)

- diarrhoea.

Common side effects (may affect up to 1 in 10 people)

- vomiting, nausea, abdominal pain or distension, dyspepsia, flatulence
- headache, tiredness, dizziness, drowsiness, numbness, tingling or pain in hands or feet, loss of strength, difficulty falling asleep.

Uncommon side effects (may affect up to 1 in 100 people)

- chest pain, changes in electrocardiogram, rapid heart beating
- decreased or abnormal skin sensitivity, pins and needles, attention disturbance, loss of memory, problems with your balance
- difficulty breathing, cough, nosebleed, throat irritation
- inflammation of the stomach or mouth, heartburn, retching, dry mouth, discomfort of the abdomen, constipation, belching



- kidney failure, kidney stones, difficult discharge of urine, frequent or excessive passage of urine, sometimes at night
- urticaria, severe swelling of the skin and other tissues (most often the lips or the eyes), eczema, excessive sweating, night sweats, hair loss, acne, scaly skin, colouration of nails
- muscle pain, muscle cramps or weakness, pain in extremity, osteoporosis
- slowing down of the thyroid gland function. This can be seen in a blood test.
- high blood pressure, flushing
- red or dry eyes
- fever, swelling of lower limbs due to fluids, malaise, irritability, pain
- symptoms of infection, herpes simplex
- erectile dysfunction, enlargement of breasts
- sleeping problems, sleepiness, depression, anxiety, abnormal dreams, decrease in sexual drive

Rare side effects (may affect up to 1 in 1,000 people)

- a reaction called DRESS [severe rash, which may be accompanied by fever, fatigue, swelling of the face or lymph glands, increase of eosinophils (type of white blood cells), effects on liver, kidney or lung]
- heart attack, slow heart beating, palpitations
- visual disturbance
- chills, feeling abnormal
- a feeling of confusion or disorientation, altered mood, restlessness
- fainting, epileptic fits, changes or loss of taste
- mouth sores, vomiting blood, inflammation of the lips, dry lips, coated tongue
- running nose
- skin lesions, dry skin
- stiffness of muscles or joints, joint pain with or without inflammation
- changes in some values of your blood cells or chemistry. These can be seen in the results of blood and/or urine tests. Your doctor will explain these to you. Examples are: increase in some white blood cells
- darunavir crystals in the kidney causing kidney disease.

Some side effects are typical for anti-HIV medicines in the same family as darunavir. These are:

- muscle pain, tenderness or weakness. On rare occasions, these muscle disorders have been serious.

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Darunavir Mylan**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, blister and on the bottle after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

For bottles: Once opened, use within 100 days.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away any medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Darunavir Mylan contains

- The active substance is darunavir. Each tablet contains 600 milligram of darunavir.
- The other ingredients are colloidal anhydrous silica, cellulose microcrystalline, crospovidone, sodium starch glycolate, hypromellose, magnesium stearate. The film-coating contains polyvinyl alcohol partially hydrolysed, titanium dioxide (E171), macrogol and talc.

### What Darunavir Mylan looks like and contents of the pack

Darunavir Mylan 600 mg film-coated tablets are white to off-white, oval shaped, with two curved sides, marked with M on one side and DV5 on the other side.

Darunavir Mylan 600 mg film-coated tablets are available in blister packs containing 30 and 60 tablets and in plastic bottles containing 30, 60 and 90 tablets.

Not all pack sizes may be marketed.

### Marketing Authorisation Holder

Mylan Pharmaceuticals Limited  
Damastown Industrial Park,  
Mulhuddart, Dublin 15,  
DUBLIN  
Ireland

### Manufacturers

Mylan Hungary Kft  
H-2900 Komárom, Mylan utca 1  
Hungary

McDermott Laboratories Limited trading as Gerard Laboratories  
35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13  
Ireland

Mylan Germany GmbH  
Zweigniederlassung Bad Homburg v. d. Hoehe, Benzstrasse 1  
Bad Homburg v. d. Hoehe  
Hessen, 61352  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

### België/Belgique/Belgien

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00

### Lietuva

Mylan Healthcare UAB  
Tel: +370 5 205 1288

### България

Майлан ЕООД  
Тел: +359 2 44 55 400

### Luxembourg/Luxemburg

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00  
(Belgique/Belgien)

### Česká republika

Viatrix CZ s.r.o.  
Tel: + 420 222 004 400

### Magyarország

Mylan EPD Kft.  
Tel.: + 36 1 465 2100

**Danmark**

Viatriis ApS  
Tlf: +45 28 11 69 32

**Deutschland**

Viatriis Healthcare GmbH  
Tel: +49 800 0700 800

**Eesti**

BGP Products Switzerland GmbH Eesti filiaal  
Tel: + 372 6363 052

**Ελλάδα**

Generics Pharma Hellas EΠE  
Τηλ: +30 210 993 6410

**España**

Viatriis Pharmaceuticals, S.L.U.  
Tel: + 34 900 102 712

**France**

Viatriis Santé  
Tél: +33 4 37 25 75 00

**Hrvatska**

Viatriis Hrvatska d.o.o.  
Tel: +385 1 23 50 599

**Ireland**

Mylan Ireland Limited  
Tel: +353 1 8711600

**Ísland**

Icepharma hf.  
Sími: +354 540 8000

**Italia**

Mylan Italia S.r.l.  
Tel: + 39 02 612 46921

**Κύπρος**

Varnavas Hadjipanayis Ltd  
Τηλ: + 357 2220 7700

**Latvija**

Mylan Healthcare SIA  
Tel: +371 676 055 80

**Malta**

V.J. Salomone Pharma Ltd  
Tel: + 356 21 22 01 74

**Nederland**

Mylan BV  
Tel: +31 (0)20 426 3300

**Norge**

Viatriis AS  
Tlf: + 47 66 75 33 00

**Österreich**

Arcana Arzneimittel GmbH  
Tel: +43 1 416 2418

**Polska**

Mylan Healthcare Sp. z o.o.  
Tel.: + 48 22 546 64 00

**Portugal**

Mylan, Lda.  
Tel: + 351 214 127 200

**România**

BGP Products SRL  
Tel: +40 372 579 000

**Slovenija**

Viatriis d.o.o.  
Tel: + 386 1 23 63 180

**Slovenská republika**

Viatriis Slovakia s.r.o.  
Tel: +421 2 32 199 100

**Suomi/Finland**

Viatriis Oy  
Puh/Tel: +358 20 720 9555

**Sverige**

Viatriis AB  
Tel: +46 (0)8 630 19 00

**United Kingdom (Northern Ireland)**

Mylan IRE Healthcare Limited  
Tel: +353 18711600

**This leaflet was last revised in {MM/YYYY}.**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.

## **Package leaflet: Information for the user**

### **Darunavir Mylan 800 mg film-coated tablets**

darunavir

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Darunavir Mylan is and what it is used for
2. What you need to know before you take Darunavir Mylan
3. How to take Darunavir Mylan
4. Possible side effects
5. How to store Darunavir Mylan
6. Contents of the pack and other information

#### **1. What Darunavir Mylan is and what it is used for**

##### **What is Darunavir Mylan?**

Darunavir Mylan contains the active substance darunavir. Darunavir is an antiretroviral medicine used in the treatment of Human Immunodeficiency Virus (HIV) infection. It belongs to a group of medicines called protease inhibitors. Darunavir works by reducing the amount of HIV in your body. This will improve your immune system and reduces the risk of developing illnesses linked to HIV infection.

##### **What it is used for?**

Darunavir is used to treat adults and children (3 years of age and above, at least 40 kilograms body weight) who are infected by HIV and

- who have not used antiretroviral medicines before.
- in certain patients who have used antiretroviral medicines before (your doctor will determine this).

Darunavir must be taken in combination with a low dose of cobicistat or ritonavir and other anti-HIV medicines. Your doctor will discuss with you which combination of medicines is best for you.

#### **2. What you need to know before you take Darunavir Mylan**

##### **Do not take Darunavir Mylan**

- if you are **allergic** to darunavir or any of the other ingredients of this medicine (listed in section 6) or to cobicistat or ritonavir.
- if you have **severe liver problems**. Ask your doctor if you are unsure about the severity of your liver disease. Some additional tests might be necessary.

**Do not combine Darunavir Mylan with any of the following medicines**

If you are taking any of these, ask your doctor about switching to another medicine.

Medicine	Purpose of the medicine
<i>Avanafil</i>	to treat erectile dysfunction
<i>Astemizole</i> or <i>terfenadine</i>	to treat allergy symptoms
<i>Triazolam</i> and <i>oral</i> (taken by mouth) <i>midazolam</i>	to help you sleep and/or relieve anxiety
<i>Cisapride</i>	to treat some stomach conditions
<i>Colchicine</i> (if you have kidney and/or liver problems)	to treat gout or familial Mediterranean fever
<i>Lurasidone</i> , <i>pimozide</i> , <i>quetiapine</i> or <i>sertindole</i>	to treat psychiatric conditions
<i>Ergot alkaloids</i> like <i>ergotamine</i> , <i>dihydroergotamine</i> , <i>ergometrine</i> and <i>methylergonovine</i>	to treat migraine headaches
<i>Amiodarone</i> , <i>bepiridil</i> , <i>dronedarone</i> , <i>ivabradine</i> , <i>quinidine</i> , <i>ranolazine</i>	to treat certain heart disorders e.g. abnormal heart beat
<i>Lovastatin</i> , <i>simvastatin</i> and <i>lomitapide</i>	to lower cholesterol levels
<i>Rifampicin</i>	to treat some infections such as tuberculosis
The combination product <i>lopinavir/ritonavir</i>	this anti-HIV medicine belongs to the same class as Darunavir Mylan
<i>Elbasvir/grazoprevir</i>	to treat hepatitis C infection
<i>Alfuzosin</i>	to treat enlarged prostate
<i>Sildenafil</i>	to treat high blood pressure in the pulmonary circulation
<i>Ticagrelor</i>	to help stop the clumping of platelets in the treatment of patients with a history of a heart attack
<i>Naloxegol</i>	to treat opioid induced constipation
<i>Dapoxetine</i>	to treat premature ejaculation
<i>Domperidone</i>	to treat nausea and vomiting

Do not combine darunavir with products that contain St John's wort (*Hypericum perforatum*).

**Warnings and precautions**

Talk to your doctor, pharmacist or nurse before taking Darunavir Mylan.

Darunavir is not a cure for HIV infection.

People taking darunavir may still develop infections or other illnesses associated with HIV infection. You must keep in regular contact with your doctor.

People taking darunavir may develop a skin rash. Infrequently a rash may become severe or potentially life-threatening. Please contact your doctor whenever you develop a rash.

In patients taking darunavir and raltegravir (for HIV infection), rashes (generally mild or moderate) may occur more frequently than in patients taking either medicine separately.

**Tell your doctor about your situation BEFORE and DURING your treatment**

Make sure that you check the following points and tell your doctor if any of these apply to you.

- Tell your doctor if you have had **problems with your liver** before, including hepatitis B or C infection. Your doctor may evaluate how severe your liver disease is before deciding if you can take darunavir.
- Tell your doctor if you have **diabetes**. Darunavir might increase sugar levels in the blood.
- Tell your doctor immediately if you notice any **symptoms of infection** (for example enlarged lymph nodes and fever). In some patients with advanced HIV infection and a history of opportunistic infection, signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started. It is believed that these symptoms are due to an

improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms.

- In addition to the opportunistic infections, autoimmune disorders (a condition that occurs when the immune system attacks healthy body tissue) may also occur after you start taking medicines for the treatment of your HIV infection. Autoimmune disorders may occur many months after the start of treatment. If you notice any symptoms of infection or other symptoms such as muscle weakness, weakness beginning in the hands and feet and moving up towards the trunk of the body, palpitations, tremor or hyperactivity, please inform your doctor immediately to seek necessary treatment.
- Tell your doctor if you have **haemophilia**. Darunavir might increase the risk of bleeding.
- Tell your doctor if you are **allergic to sulphonamides** (e.g. used to treat certain infections).
- Tell your doctor if you notice any **musculoskeletal problems**. Some patients taking combination antiretroviral therapy may develop a bone disease called osteonecrosis (death of bone tissue caused by loss of blood supply to the bone). The length of combination antiretroviral therapy, corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index, among others, may be some of the many risk factors for developing this disease. Signs of osteonecrosis are joint stiffness, aches and pains (especially of the hip, knee and shoulder) and difficulty in movement. If you notice any of these symptoms please inform your doctor.

### Elderly

Darunavir has only been used in limited numbers of patients 65 years or older. If you belong to this age group, please discuss with your doctor if you can use Darunavir Mylan.

### Children and adolescents

Darunavir Mylan 800 mg tablets is not for use in children younger than 3 years of age or weighing less than 40 kilograms.

### Other medicines and Darunavir Mylan

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines.

There are some medicines that **you must not combine** with darunavir. These are mentioned above under the heading '**Do not combine Darunavir Mylan with any of the following medicines:**'

In most cases, darunavir can be combined with anti-HIV medicines belonging to another class [e.g. NRTIs (nucleoside reverse transcriptase inhibitors), NNRTIs (non-nucleoside reverse transcriptase inhibitors), CCR5 antagonists and FIs (fusion inhibitors)]. Darunavir with cobicistat or ritonavir has not been tested with all PIs (protease inhibitors) and must not be used with other HIV PIs. In some cases dosage of other medicines might need to be changed. Therefore always tell your doctor if you take other anti-HIV medicines and follow your doctor's instruction carefully on which medicines can be combined.

The effects of darunavir might be reduced if you take any of the following products. Tell your doctor if you take:

- *Phenobarbital, phenytoin* (to prevent seizures)
- *Dexamethasone* (corticosteroid)
- *Efavirenz* (HIV infection)
- *Rifapentine, rifabutin* (medicines to treat some infections such as tuberculosis)
- *Saquinavir* (HIV infection).

The effects of other medicines might be influenced if you take darunavir and your doctor might want to do some additional blood tests. Tell your doctor if you take:

- *Amlodipine, diltiazem, disopyramide, carvedilol, felodipine, flecainide, lidocaine, metoprolol, mexiletine, nifedipine, nicardipine, propafenone, timolol, verapamil* (for heart disease) as the therapeutic effect or side effects of these medicines may be increased.
- *Apixaban, dabigatran etexilate, edoxaban, rivaroxaban, warfarin, clopidogrel* (to reduce clotting of the blood) as their therapeutic effect or side effects may be altered.

- Oestrogen-based hormonal contraceptives and hormonal replacement therapy. Darunavir might reduce its effectiveness. When used for birth control, alternative methods of non-hormonal contraception are recommended.
- *Ethinylestradiol/drospirenone*. Darunavir might increase the risk for elevated potassium levels by drospirenone.
- *Atorvastatin, pravastatin, rosuvastatin* (to lower cholesterol levels). The risk of muscle damage might be increased. Your doctor will evaluate which cholesterol lowering regimen is best for your specific situation.
- *Clarithromycin* (antibiotic)
- *Ciclosporin, everolimus, tacrolimus, sirolimus* (for dampening down your immune system) as the therapeutic effect or side effects of these medicines might be increased.
- *Corticosteroids including betamethasone, budesonide, fluticasone, mometasone, prednisone, triamcinolone*. These medicines are used to treat allergies, asthma, inflammatory bowel diseases, inflammatory conditions of the skin, eyes, joints and muscles and other inflammatory conditions. These medicines are generally taken orally, inhaled, injected or applied to the skin. If alternatives cannot be used, its use should only take place after medical evaluation and under close monitoring by your doctor for corticosteroid side effects.
- *Buprenorphine/naloxone* (medicines to treat opioid dependence)
- *Salmeterol* (medicine to treat asthma)
- *Artemether/lumefantrine* (a combination medicine to treat malaria)
- *Dasatinib, everolimus, irinotecan, nilotinib, vinblastine, vincristine* (to treat cancer)
- *Sildenafil, tadalafil, vardenafil* (for erectile dysfunction or to treat a heart and lung disorder called pulmonary arterial hypertension)
- *Glecaprevir/pibrentasvir* (to treat hepatitis C infection)
- *Fentanyl, oxycodone, tramadol* (to treat pain)
- *Fesoterodine, solifenacin* (to treat urologic disorders).

Your doctor might want to do some additional tests and the dosage of other medicines might need to be changed since either their own or darunavir's therapeutic effect or side effects may be influenced when combined.

Tell your doctor if you take:

- *Dabigatran etexilate, edoxaban, warfarin* (to reduce clotting of the blood)
- *Alfentanil* (injectable strong and short-acting painkiller that is used for surgical procedures)
- *Digoxin* (to treat certain heart disorders)
- *Clarithromycin* (antibiotic)
- *Itraconazole, isavuconazole, fluconazole, posaconazole, clotrimazole* (to treat fungal infections). Voriconazole should only be taken after medical evaluation.
- *Rifabutin* (against bacterial infections)
- *Sildenafil, vardenafil, tadalafil* (for erectile dysfunction or high blood pressure in the pulmonary circulation)
- *Amitriptyline, desipramine, imipramine, nortriptyline, paroxetine, sertraline, trazodone* (to treat depression and anxiety)
- *Maraviroc* (to treat HIV infection)
- *Methadone* (to treat opiate dependence)
- *Carbamazepine, clonazepam* (to prevent seizures or to treat certain types of nerve pain)
- *Colchicine* (to treat gout or familial Mediterranean fever)
- *Bosentan* (to treat high blood pressure in the pulmonary circulation)
- *Buspirone, clorazepate, diazepam, estazolam, flurazepam, midazolam when used as injection, zolpidem* (sedative agents)
- *Perphenazine, risperidone, thioridazine* (to treat psychiatric conditions)
- *Metformin* (to treat type 2 diabetes).

This is **not** a complete list of medicines. Tell your healthcare provider about **all** medicines that you are taking.

### **Darunavir Mylan with food and drink**

See section 3 'How to take Darunavir Mylan.

### **Pregnancy and breast-feeding**

Tell your doctor immediately if you are pregnant or planning to become pregnant. Pregnant women must not take darunavir with ritonavir unless specifically directed by the doctor. Pregnant women should not take darunavir with cobicistat.

Because of the potential for side effects in breast-fed infants, women should not breast-feed if they are receiving Darunavir Mylan.

Breast-feeding is not recommended in women living with HIV because HIV infection can be passed on to the baby in breast milk. If you are breast-feeding, or thinking about breast-feeding, you should discuss it with your doctor as soon as possible.

### **Driving and using machines**

Do not operate machines or drive if you feel dizzy after taking darunavir.

### **Darunavir Mylan contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially ‘sodium-free’.

## **3. How to take Darunavir Mylan**

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure.

Even if you feel better, do not stop taking darunavir and cobicistat or ritonavir without talking to your doctor.

After therapy has been initiated, the dose or dosage form must not be changed or therapy must not be stopped without instruction of the doctor.

Darunavir Mylan 800 mg tablets are intended for once daily use only.

### **Dose for adults who have not taken antiretroviral medicines before (your doctor will determine this)**

The usual dose of darunavir is 800 milligram (2 tablets containing 400 milligram of darunavir or 1 tablet containing 800 milligram of darunavir) once daily.

You must take darunavir every day and always in combination with 150 milligram of cobicistat or 100 milligram of ritonavir and with food. Darunavir cannot work properly without cobicistat or ritonavir and food. You must eat a meal or a snack within 30 minutes prior to taking your darunavir and cobicistat or ritonavir. The type of food is not important. Even if you feel better, do not stop taking darunavir and cobicistat or ritonavir without talking to your doctor.

### **Instructions for adults**

- Take one 800 milligram tablet at the same time, once a day, every day.
- Take darunavir always together with 150 milligram of cobicistat or 100 milligram of ritonavir.
- Take darunavir with food.
- Swallow the tablet with a drink such as water or milk.
- Take your other HIV medicines used in combination with darunavir and cobicistat or ritonavir as recommended by your doctor.

### **Dose for adults who have taken antiretroviral medicines before (your doctor will determine this)**

The dose is either:

- 800 milligram darunavir (2 tablets containing 400 milligram of darunavir or 1 tablet containing 800 milligram of darunavir) together with 150 milligram cobicistat or 100 milligram ritonavir once daily.
- OR



- 600 milligram darunavir (2 tablets containing 300 milligram of darunavir or 1 tablet containing 600 milligram of darunavir) together with 100 milligram ritonavir twice daily.

Please discuss with your doctor which dose is right for you.

**Dose for children 3 years of age and above with ritonavir, and 12 years of age and above with cobicistat, weighing more than 40 kilograms who have not taken antiretroviral medicines before (your child's doctor will determine this)**

- The usual dose of darunavir is 800 milligram (2 tablets containing 400 milligram of darunavir or 1 tablet containing 800 milligram of darunavir) together with 100 milligram ritonavir or 150 milligram of cobicistat once daily.

**Dose for children 3 years of age and above with ritonavir, and 12 years of age and above with cobicistat, weighing more than 40 kilograms who have taken antiretroviral medicines before (your child's doctor will determine this)**

The dose is either:

- 800 milligram darunavir (2 tablets containing 400 milligram of darunavir or 1 tablet containing 800 milligram of darunavir) together with 100 milligram ritonavir or 150 milligram of cobicistat once daily.  
OR
- 600 milligram darunavir (2 tablets containing 300 milligram of darunavir or 1 tablet containing 600 milligram of darunavir) together with 100 milligram ritonavir twice daily.

Please discuss with your doctor which dose is right for you.

***Instructions for children 3 years of age and above with ritonavir, and 12 years of age and above with cobicistat, weighing more than 40 kilograms***

- Take 800 milligram darunavir (2 tablets containing 400 milligram of Darunavir Mylan or 1 tablet containing 800 milligram of Darunavir Mylan) at the same time, once a day, every day.
- Take darunavir always together with 100 milligram of ritonavir or 150 milligram of cobicistat.
- Take darunavir with food.
- Swallow the tablets with a drink such as water or milk.
- Take your other HIV medicines used in combination with darunavir and ritonavir or cobicistat as recommended by your doctor.

**If you take more Darunavir Mylan than you should**

Contact your doctor, pharmacist or nurse immediately.

**If you forget to take Darunavir Mylan**

If you notice **within 12 hours**, you must take the tablets immediately. Always take with cobicistat or ritonavir and food. If you notice **after 12 hours**, then skip the intake and take the next doses as usual. Do not take a double dose to make up for a forgotten dose.

**If you vomit after taking Darunavir Mylan and cobicistat or ritonavir**

If you vomit **within 4 hours** of taking the medicine, another dose of Darunavir Mylan and cobicistat or ritonavir should be taken with food as soon as possible. If you vomit **more than 4 hours** after taking the medicine, then you do not need to take another dose of Darunavir Mylan and cobicistat or ritonavir until the next regularly scheduled time.

Contact your doctor **if you are uncertain** about what to do if you miss a dose or vomit.

**Do not stop taking Darunavir Mylan without talking to your doctor first**

Anti-HIV medicines may make you feel better. Even when you feel better, do not stop taking Darunavir Mylan. Talk to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

#### **4. Possible side effects**

During HIV therapy there may be an increase in weight and in levels of blood lipids and glucose. This is partly linked to restored health and life style, and in the case of blood lipids sometimes to the HIV medicines themselves. Your doctor will test for these changes.

Like all medicines, this medicine can cause side effects, although not everybody gets them.

##### **Tell your doctor if you develop any of the following side effects**

Liver problems that may occasionally be severe have been reported. Your doctor should do blood tests before you start darunavir. If you have chronic hepatitis B or C infection, your doctor should check your blood tests more often because you have an increased chance of developing liver problems. Talk to your doctor about the signs and symptoms of liver problems. These may include yellowing of your skin or whites of your eyes, dark (tea coloured) urine, pale coloured stools (bowel movements), nausea, vomiting, loss of appetite, or pain, aching, or pain and discomfort on your right side below your ribs.

Skin rash (more often when used in combination with raltegravir), itching. The rash is usually mild to moderate. A skin rash might also be a symptom of a rare severe situation. It is therefore important to talk to your doctor if you develop a rash. Your doctor will advise you how to deal with your symptoms or whether darunavir must be stopped.

Other severe side effects were diabetes (common), and inflammation of the pancreas (uncommon).

Very common side effects (may affect more than 1 in 10 people)

- diarrhoea.

Common side effects (may affect up to 1 in 10 people)

- vomiting, nausea, abdominal pain or distension, dyspepsia, flatulence
- headache, tiredness, dizziness, drowsiness, numbness, tingling or pain in hands or feet, loss of strength, difficulty falling asleep.

Uncommon side effects (may affect up to 1 in 100 people)

- chest pain, changes in electrocardiogram, rapid heart beating
- decreased or abnormal skin sensitivity, pins and needles, attention disturbance, loss of memory, problems with your balance
- difficulty breathing, cough, nosebleed, throat irritation
- inflammation of the stomach or mouth, heartburn, retching, dry mouth, discomfort of the abdomen, constipation, belching
- kidney failure, kidney stones, difficult discharge of urine, frequent or excessive passage of urine, sometimes at night
- urticaria, severe swelling of the skin and other tissues (most often the lips or the eyes), eczema, excessive sweating, night sweats, hair loss, acne, scaly skin, colouration of nails
- muscle pain, muscle cramps or weakness, pain in extremity, osteoporosis
- slowing down of the thyroid gland function. This can be seen in a blood test.
- high blood pressure, flushing
- red or dry eyes
- fever, swelling of lower limbs due to fluids, malaise, irritability, pain
- symptoms of infection, herpes simplex
- erectile dysfunction, enlargement of breasts
- sleeping problems, sleepiness, depression, anxiety, abnormal dreams, decrease in sexual drive

Rare side effects (may affect up to 1 in 1,000 people)

- a reaction called DRESS [severe rash, which may be accompanied by fever, fatigue, swelling of the face or lymph glands, increase of eosinophils (type of white blood cells), effects on liver, kidney or lung]

- heart attack, slow heart beating, palpitations
- visual disturbance
- chills, feeling abnormal
- a feeling of confusion or disorientation, altered mood, restlessness
- fainting, epileptic fits, changes or loss of taste
- mouth sores, vomiting blood, inflammation of the lips, dry lips, coated tongue
- running nose
- skin lesions, dry skin
- stiffness of muscles or joints, joint pain with or without inflammation
- changes in some values of your blood cells or chemistry. These can be seen in the results of blood and/or urine tests. Your doctor will explain these to you. Examples are: increase in some white blood cells
- darunavir crystals in the kidney causing kidney disease.

Some side effects are typical for anti-HIV medicines in the same family as darunavir. These are:

- muscle pain, tenderness or weakness. On rare occasions, these muscle disorders have been serious.

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Darunavir Mylan**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, blister and on the bottle after EXP. The expiry date refers to the last day of that month.

For blisters with plastic on one side and aluminium on the other: Do not store above 25°C.

For blisters with aluminium on both sides: This medicine does not require any special storage conditions.

For bottles: Once opened, use within 90 days. This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away any medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Darunavir Mylan contains**

- The active substance is darunavir. Each tablet contains 800 milligram of darunavir.
- The other ingredients are colloidal anhydrous silica, cellulose microcrystalline, crospovidone, sodium starch glycolate, hypromellose, magnesium stearate. The film-coating contains polyvinyl alcohol partially hydrolysed, titanium dioxide (E171), macrogol and talc.

### **What Darunavir Mylan looks like and contents of the pack**

Darunavir Mylan 800 mg film-coated tablets are white to off-white, oval shaped, with two curved sides, marked with M on one side and DV8 on the other side.

Darunavir Mylan 800 mg film-coated tablets are available in blister packs containing 30 tablets and in plastic bottles containing 30, 60 and 90 tablets.

Not all pack sizes may be marketed.

**Marketing Authorisation Holder**

Mylan Pharmaceuticals Limited  
Damastown Industrial Park, Mulhuddart,  
Dublin 15,  
DUBLIN  
Ireland

**Manufacturers**

Mylan Hungary Kft  
H-2900 Komárom, Mylan utca 1  
Hungary

McDermott Laboratories Limited trading as Gerard Laboratories  
35/36 Baldoye Industrial Estate, Grange Road, Dublin 13  
Ireland

Mylan Germany GmbH  
Zweigniederlassung Bad Homburg v. d. Hoehe, Benzstrasse 1  
Bad Homburg v. d. Hoehe  
Hessen, 61352  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00

**Lietuva**

Mylan Healthcare UAB  
Tel: +370 5 205 1288

**България**

Майлан ЕООД  
Тел: +359 2 44 55 400

**Luxembourg/Luxemburg**

Mylan bvba/sprl  
Tél/Tel: + 32 (0)2 658 61 00  
(Belgique/Belgien)

**Česká republika**

Viatrix CZ s.r.o.  
Tel: + 420 222 004 400

**Magyarország**

Mylan EPD Kft.  
Tel.: + 36 1 465 2100

**Danmark**

Viatrix ApS  
Tlf: +45 28 11 69 32

**Malta**

V.J. Salomone Pharma Ltd  
Tel: + 356 21 22 01 74

**Deutschland**

Viatrix Healthcare GmbH  
Tel: +49 800 0700 800

**Nederland**

Mylan BV  
Tel: +31 (0)20 426 3300

**Eesti**

BGP Products Switzerland GmbH Eesti filiaal  
Tel: + 372 6363 052

**Norge**

Viatrix AS  
Tlf: + 47 66 75 33 00

**Ελλάδα**

Generics Pharma Hellas ΕΠΕ  
Τηλ: +30 210 993 6410

**España**

Viatrix Pharmaceuticals, S.L.U.  
Tel: + 34 900 102 712

**France**

Viatrix Santé  
Tél: +33 4 37 25 75 00

**Hrvatska**

Viatrix Hrvatska d.o.o.  
Tel: +385 1 23 50 599

**Ireland**

Mylan Ireland Limited  
Tel: +353 1 8711600

**Ísland**

Icepharma hf.  
Sími: +354 540 8000

**Italia**

Mylan Italia S.r.l.  
Tel: + 39 02 612 46921

**Κύπρος**

Varnavas Hadjipanayis Ltd  
Τηλ: + 357 2220 7700

**Latvija**

Mylan Healthcare SIA  
Tel: +371 676 055 80

**Österreich**

Arcana Arzneimittel GmbH  
Tel: +43 1 416 2418

**Polska**

Mylan Healthcare Sp. z o.o.  
Tel.: + 48 22 546 64 00

**Portugal**

Mylan, Lda.  
Tel: + 351 214 127 200

**România**

BGP Products SRL  
Tel: +40 372 579 000

**Slovenija**

Viatrix d.o.o.  
Tel: + 386 1 23 63 180

**Slovenská republika**

Viatrix Slovakia s.r.o.  
Tel: +421 2 32 199 100

**Suomi/Finland**

Viatrix Oy  
Puh/Tel: +358 20 720 9555

**Sverige**

Viatrix AB  
Tel: +46 (0)8 630 19 00

**United Kingdom (Northern Ireland)**

Mylan IRE Healthcare Limited  
Tel: +353 18711600

**This leaflet was last revised in {MM/YYYY}.**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu>.