

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Dexdor 100 micrograms/ml concentrate for solution for infusion

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 1 ml of concentrate contains dexmedetomidine hydrochloride equivalent to 100 micrograms dexmedetomidine.

Each 2 ml ampoule contains 200 micrograms of dexmedetomidine.

Each 2 ml vial contains 200 micrograms of dexmedetomidine.

Each 4 ml vial contains 400 micrograms of dexmedetomidine.

Each 10 ml vial contains 1 000 micrograms of dexmedetomidine.

The concentration of the final solution after dilution should be either 4 micrograms/ml or 8 micrograms/ml.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Concentrate for solution for infusion (sterile concentrate).

The concentrate is a clear, colourless solution, pH 4.5–7.0

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

For sedation of adult ICU (Intensive Care Unit) patients requiring a sedation level not deeper than arousal in response to verbal stimulation (corresponding to Richmond Agitation-Sedation Scale (RASS) 0 to -3).

For sedation of non-intubated adult patients prior to and/or during diagnostic or surgical procedures requiring sedation, i.e. procedural/awake sedation.

### 4.2 Posology and method of administration

**For sedation of adult ICU (Intensive Care Unit) patients requiring a sedation level not deeper than arousal in response to verbal stimulation (corresponding to Richmond Agitation-Sedation Scale (RASS) 0 to -3).**

For hospital use only. Dexdor should be administered by healthcare professionals skilled in the management of patients requiring intensive care.

#### Posology

Patients already intubated and sedated may switch to dexmedetomidine with an initial infusion rate of 0.7 micrograms/kg/h which may then be adjusted stepwise within the dose range 0.2 to 1.4 micrograms/kg/h in order to achieve the desired level of sedation, depending on the patient's response. A lower starting infusion rate should be considered for frail patients. Dexmedetomidine is very potent and the infusion rate is given per **hour**. After dose adjustment, a new steady state sedation level may not be reached for up to one hour.

### *Maximum dose*

The maximum dose of 1.4 micrograms/kg/h should not be exceeded. Patients failing to achieve an adequate level of sedation with the maximum dose of dexmedetomidine should be switched to an alternative sedative agent.

Use of a loading dose of Dexdor in ICU sedation is not recommended and is associated with increased adverse reactions. Propofol or midazolam may be administered if needed until clinical effects of dexmedetomidine are established.

### *Duration*

There is no experience in the use of Dexdor for more than 14 days. The use of Dexdor for longer than this period should be regularly reassessed.

### **For sedation of non-intubated adult patients prior to and/or during diagnostic or surgical procedures requiring sedation, i.e. procedural/awake sedation.**

Dexdor should be administered only by health care professionals skilled in the anaesthetic management of patients in the operating room or during diagnostic procedures. When Dexdor is administered for conscious sedation, patients should be continuously monitored by persons not involved in the conduct of the diagnostic or surgical procedure. Patients should be monitored continuously for early signs of hypotension, hypertension, bradycardia, respiratory depression, airway obstruction, apnoea, dyspnoea and/or oxygen desaturation (see section 4.8).

Supplemental oxygen should be immediately available and provided when indicated. The oxygen saturation should be monitored by pulse oximetry.

Dexdor is given as a loading infusion followed by maintenance infusion. Depending on the procedure concomitant local anaesthesia or analgesia may be needed in order to achieve the desired clinical effect. Additional analgesia or sedatives (e.g. opioids, midazolam, or propofol) are recommended in case of painful procedures or if increased depth of sedation is necessary. The pharmacokinetic distribution half-life of Dexdor has been estimated to be around 6 min, which can be taken into consideration, together with the effects of other administered medications, when assessing the appropriate time needed for titration to desired clinical effect of Dexdor.

### *Initiation of Procedural Sedation:*

- A loading infusion of 1.0 microgram/kg over 10 minutes. For less invasive procedures such as ophthalmic surgery, a loading infusion of 0.5 micrograms/kg given over 10 minutes may be suitable.

### *Maintenance of Procedural Sedation:*

- The maintenance infusion is generally initiated at 0.6–0.7 microgram/kg/hour and titrated to achieve desired clinical effect with doses ranging from 0.2 to 1 microgram/kg/hour. The rate of the maintenance infusion should be adjusted to achieve the targeted level of sedation.

### Special populations

#### *Elderly*

No dose adjustment is normally required for elderly patients (see section 5.2). Elderly patients appear to have an increased risk for hypotension (see section 4.4) but the limited data available from procedural sedation do not suggest a clear dose dependency.

### *Renal impairment*

No dose adjustment is required for patients with renal impairment.

### *Hepatic impairment*

Dexmedetomidine is metabolised in the liver and should be used with caution in patients with hepatic impairment. A reduced maintenance dose may be considered (see sections 4.4 and 5.2).

### *Paediatric population*

The safety and efficacy of Dexdor in children aged 0 to 18 years have not been established. Currently available data are described in sections 4.8, 5.1 and 5.2 but no recommendation on a posology can be made.

### Method of administration

Dexdor must be administered only as a diluted intravenous infusion using a controlled infusion device. For instructions on dilution of the medicinal product before administration, see section 6.6.

## **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Advanced heart block (grade 2 or 3) unless paced.

Uncontrolled hypotension.

Acute cerebrovascular conditions.

## **4.4 Special warnings and precautions for use**

### Monitoring

Dexdor is intended for use in an intensive care setting, operating room and during diagnostic procedures. The use in other environments is not recommended. All patients should have continuous cardiac monitoring during Dexdor infusion. Respiration should be monitored in non-intubated patients due to the risk of respiratory depression and in some case apnoea (see section 4.8).

The time to recovery after the use of dexmedetomidine was reported to be approximately one hour. When used in an outpatient setting close monitoring should continue for at least one hour (or longer based on the patient condition), with medical supervision continued for at least one further hour to ensure the safety of the patient.

### General precautions

Dexdor should not be given as a bolus dose and in the ICU a loading dose is not recommended. Users should therefore be ready to use an alternative sedative for acute control of agitation or during procedures, especially during the first few hours of treatment. During procedural sedation a small bolus of another sedative may be used if a rapid increase in sedation level is required. Some patients receiving Dexdor have been observed to be arousable and alert when stimulated. This alone should not be considered as evidence of lack of efficacy in the absence of other clinical signs and symptoms.

Dexmedetomidine normally does not cause deep sedation and patients may be easily roused. Dexmedetomidine is therefore not suitable in patients who will not tolerate this profile of effects, for example those requiring continuous deep sedation.

Dexdor should not be used as a general anaesthetic induction agent for intubation or to provide sedation during muscle relaxant use.

Dexmedetomidine lacks the anticonvulsant action of some other sedatives and so will not suppress underlying seizure activity.

Care should be taken if combining dexmedetomidine with other substances with sedative or cardiovascular actions as additive effects may occur.

Dexdor is not recommended for patient controlled sedation. Adequate data is not available.

When Dexdor is used in an outpatient setting patients should normally be discharged into the care of a suitable third party. Patients should be advised to refrain from driving or other hazardous tasks and where possible to avoid the use of other agents that may sedate (e.g. benzodiazepines, opioids, alcohol) for a suitable period of time based on observed effects of dexmedetomidine, the procedure, concomitant medications, the age and the condition of the patient.

Caution should be exercised when administering dexmedetomidine to elderly patients. Elderly patients over 65 years of age may be more prone to hypotension with the administration of dexmedetomidine, including a loading dose, for procedures. A dose reduction should be considered. Please refer to section 4.2.

#### Mortality in ICU patients $\leq$ 65 years old

In the SPICE III pragmatic randomised controlled trial of 3 904 critically ill adult ICU patients dexmedetomidine was used as primary sedative and compared with usual care. There was no overall difference in 90-day mortality between the dexmedetomidine and usual care group (mortality 29.1% in both groups), but a heterogeneity of effect from age on mortality was observed. Dexmedetomidine was associated with an increased mortality in the age-group  $\leq$  65 years (odds ratio 1.26; 95% credibility interval 1.02 to 1.56) compared to alternative sedatives. While the mechanism is unclear, this heterogeneity of effect on mortality from age was most prominent in patients admitted for reasons other than post-operative care, and increased with increasing APACHE II scores and with decreasing age. These findings should be weighed against the expected clinical benefit of dexmedetomidine compared to alternative sedatives in younger patients.

#### Cardio-vascular effects and precautions

Dexmedetomidine reduces heart rate and blood pressure through central sympatholysis but at higher concentrations causes peripheral vasoconstriction leading to hypertension (see section 5.1). Dexmedetomidine is therefore not suitable in patients with severe cardiovascular instability.

Caution should be exercised when administering dexmedetomidine to patients with pre-existing bradycardia. Data on the effects of Dexdor in patients with heart rate  $<$  60 are very limited and particular care should be taken with such patients. Bradycardia does not normally require treatment, but has commonly responded to anti-cholinergic medicine or dose reduction where needed. Patients with high physical fitness and slow resting heart rate may be particularly sensitive to bradycardic effects of alpha-2 receptor agonists and cases of transient sinus arrest have been reported. Also cases of cardiac arrest, often preceded by bradycardia or atrioventricular block, have been reported (see section 4.8).

The hypotensive effects of dexmedetomidine may be of greater significance in those patients with pre-existing hypotension (especially if not responsive to vasopressors), hypovolaemia, chronic hypotension or reduced functional reserve such as patients with severe ventricular dysfunction and the

elderly and special care is warranted in these cases (see section 4.3). Hypotension does not normally require specific treatment but, where needed, users should be ready to intervene with dose reduction, fluids and/or vasoconstrictors .

Patients with impaired peripheral autonomic activity (e.g. due to spinal cord injury) may have more pronounced haemodynamic changes after starting dexmedetomidine and so should be treated with care.

Transient hypertension has been observed primarily during the loading dose in association with the peripheral vasoconstrictive effects of dexmedetomidine and a loading dose is not recommended in ICU sedation. Treatment of hypertension has generally not been necessary but decreasing the continuous infusion rate may be advisable.

Local vasoconstriction at higher concentration may be of greater significance in patients with ischaemic heart disease or severe cerebrovascular disease who should be monitored closely. Dose reduction or discontinuation should be considered in a patient developing signs of myocardial or cerebral ischaemia.

Caution is advised when administering dexmedetomidine together with spinal or epidural anaesthesia due to possible increased risk of hypotension or bradycardia.

#### Patients with hepatic impairment

Care should be taken in severe hepatic impairment as excessive dosing may increase the risk of adverse reactions, over-sedation or prolonged effect as a result of reduced dexmedetomidine clearance.

#### Patients with neurological disorders

Experience of dexmedetomidine in severe neurological disorders such as head injury and after neurosurgery is limited and it should be used with caution here, especially if deep sedation is required. Dexmedetomidine may reduce cerebral blood flow and intracranial pressure and this should be considered when selecting therapy.

#### Other

Alpha-2 agonists have rarely been associated with withdrawal reactions when stopped abruptly after prolonged use. This possibility should be considered if the patient develops agitation and hypertension shortly after stopping dexmedetomidine.

Dexmedetomidine may induce hyperthermia that may be resistant to traditional cooling methods. Dexmedetomidine treatment should be discontinued in the event of a sustained unexplained fever and is not recommended for use in malignant hyperthermia-sensitive patients.

Diabetes insipidus has been reported in association with dexmedetomidine treatment. If polyuria occurs, it is recommended to stop dexmedetomidine and check serum sodium level and urine osmolality.

Dexdor contains less than 1 mmol sodium (23 mg) per ml.

### **4.5 Interaction with other medicinal products and other forms of interaction**

Interaction studies have only been performed in adults.

Co-administration of dexmedetomidine with anaesthetics, sedatives, hypnotics, and opioids is likely to lead to an enhancement of effects, including sedative, anaesthetic and cardiorespiratory effects. Specific studies have confirmed enhanced effects with isoflurane, propofol, alfentanil, and midazolam.

No pharmacokinetic interactions between dexmedetomidine and isoflurane, propofol, alfentanil and midazolam have been demonstrated. However, due to possible pharmacodynamic interactions, when co-administered with dexmedetomidine, a reduction in dosage of dexmedetomidine or the concomitant anaesthetic, sedative, hypnotic or opioid may be required.

Inhibition of CYP enzymes including CYP2B6 by dexmedetomidine has been studied in human liver microsome incubations. *In vitro* study suggests that interaction potential *in vivo* exists between dexmedetomidine and substrates with dominant CYP2B6 metabolism.

Induction of dexmedetomidine *in vitro* was observed on CYP1A2, CYP2B6, CYP2C8, CYP2C9 and CYP3A4, and induction *in vivo* cannot be excluded. The clinical significance is unknown.

The possibility of enhanced hypotensive and bradycardic effects should be considered in patients receiving other medicinal products causing these effects, for example beta blockers, although additional effects in an interaction study with esmolol were modest.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

There are no or limited amount of data from the use of dexmedetomidine in pregnant women.

Studies in animals have shown reproductive toxicity (see section 5.3). Dexdor should not be used during pregnancy unless the clinical condition of the woman requires treatment with dexmedetomidine.

##### Breastfeeding

Dexmedetomidine is excreted in human milk, however levels will be below the limit of detection by 24 hours following treatment discontinuation. A risk to infants cannot be excluded. A decision must be made whether to discontinue breastfeeding or to discontinue dexmedetomidine therapy taking into account the benefit of breastfeeding for the child and the benefit of therapy for the woman.

##### Fertility

In the rat fertility study, dexmedetomidine had no effect on male or female fertility. No human data on fertility are available.

#### **4.7 Effects on ability to drive and use machines**

Patients should be advised to refrain from driving or other hazardous tasks for a suitable period of time after receiving Dexdor for procedural sedation.

#### **4.8 Undesirable effects**

##### Summary of the safety profile

##### Sedation of adult ICU (Intensive Care Unit) patients

The most frequently reported adverse reactions with dexmedetomidine in ICU setting are hypotension, hypertension and bradycardia, occurring in approximately 25%, 15% and 13% of patients respectively. Hypotension and bradycardia were also the most frequent dexmedetomidine-related serious adverse reactions occurring in 1.7% and 0.9% of randomised Intensive Care Unit (ICU) patients respectively.

##### Procedural/awake sedation

The most frequently reported adverse reactions with dexmedetomidine in procedural sedation are listed below (the protocols of phase III studies contained pre-defined thresholds for reporting changes in blood pressure, respiratory rate and heart rate as AEs).

- Hypotension (55% in dexmedetomidine-group vs. 30% in placebo-group receiving rescue midazolam and fentanyl)
- Respiratory depression (38% in dexmedetomidine-group vs. 35% in placebo-group receiving rescue midazolam and fentanyl)
- Bradycardia (14% in dexmedetomidine-group vs. 4% in placebo-group receiving rescue midazolam and fentanyl)

#### Tabulated list of adverse reactions

The adverse reactions listed in Table 1 have been accumulated from pooled data of clinical trials in intensive care.

Adverse reactions are ranked under headings of frequency, the most frequent first, using the following convention: Very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1\ 000$  to  $< 1/100$ ); rare ( $\geq 1/10\ 000$  to  $< 1/1\ 000$ ), very rare ( $< 1/10\ 000$ ), not known (cannot be estimated from the available data).

#### **Table 1.** Adverse reactions

##### ***Endocrine disorders***

Not known: Diabetes insipidus

##### ***Metabolism and nutrition disorders***

Common: Hyperglycaemia, hypoglycaemia

Uncommon: Metabolic acidosis, hypoalbuminaemia

##### ***Psychiatric disorders***

Common: Agitation

Uncommon: Hallucination

##### ***Cardiac disorders***

Very common: Bradycardia<sup>1,2</sup>

Common: Myocardial ischaemia or infarction, tachycardia

Uncommon: Atrioventricular block<sup>1</sup>, cardiac output decreased, cardiac arrest<sup>1</sup>

##### ***Vascular disorders:***

Very common: Hypotension<sup>1,2</sup>, hypertension<sup>1,2</sup>

##### ***Respiratory, thoracic and mediastinal disorders***

Very common: Respiratory depression<sup>2,3</sup>

Uncommon: Dyspnoea, apnoea

##### ***Gastrointestinal disorders***

Common: Nausea<sup>2</sup>, vomiting, dry mouth<sup>2</sup>

Uncommon: Abdominal distension

##### ***General disorders and administration site conditions***

Common: Withdrawal syndrome, hyperthermia

Uncommon: Drug ineffective, thirst

<sup>1</sup> See section on Description of selected adverse reactions

<sup>2</sup> Adverse reaction observed also in procedural sedation studies

<sup>3</sup> Incidence ‘common’ in ICU sedation studies

#### Description of selected adverse reactions

Clinically significant hypotension or bradycardia should be treated as described in section 4.4.

In relatively healthy non-ICU subjects treated with dexmedetomidine, bradycardia has occasionally led to sinus arrest or pause. The symptoms responded to leg raising and anticholinergics such as atropine or glycopyrrolate. In isolated cases bradycardia has progressed to periods of asystole in patients with pre-existing bradycardia. Also cases of cardiac arrest, often preceded by bradycardia or atrioventricular block, have been reported.

Hypertension has been associated with the use of a loading dose and this reaction can be reduced by avoiding such a loading dose or reducing the infusion rate or size of the loading dose.

#### Paediatric population

Children > 1 month post-natal, predominantly post-operative, have been evaluated for treatment up to 24 hours in the ICU and demonstrated a similar safety profile as in adults. Data in new-born infants (28–44 weeks gestation) is very limited and restricted to maintenance doses  $\leq 0.2$  mcg/kg/h. A single case of hypothermic bradycardia in a neonate has been reported in the literature.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via **the national reporting system** listed in [Appendix V](#).

## **4.9 Overdose**

### Symptoms

Several cases of dexmedetomidine overdose have been reported both in the clinical trial and the post-marketing data. The reported highest infusion rates of dexmedetomidine in these cases have reached up to 60  $\mu\text{g}/\text{kg}/\text{h}$  for 36 minutes and 30  $\mu\text{g}/\text{kg}/\text{h}$  for 15 minutes in a 20 month–old child and in an adult, respectively. The most common adverse reactions reported in conjunction with overdose include bradycardia, hypotension, hypertension, oversedation, respiratory depression and cardiac arrest.

### Management

In cases of overdose with clinical symptoms, dexmedetomidine infusion should be reduced or stopped. Expected effects are primarily cardiovascular and should be treated as clinically indicated (see section 4.4). At high concentration hypertension may be more prominent than hypotension. In clinical studies, cases of sinus arrest reversed spontaneously or responded to treatment with atropine and glycopyrrolate. Resuscitation was required in isolated cases of severe overdose resulting in cardiac arrest.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Psycholeptics, other hypnotics and sedatives, ATC code: N05CM18

Dexmedetomidine is a selective alpha-2 receptor agonist with a broad range of pharmacological properties. It has a sympatholytic effect through decrease of the release of noradrenaline in sympathetic nerve endings. The sedative effects are mediated through decreased firing of locus coeruleus, the predominant noradrenergic nucleus, situated in the brainstem. Dexmedetomidine has analgesic and anaesthetic/analgesic-sparing effects. The cardiovascular effects depend on the dose; with lower infusion rates the central effects dominate leading to decrease in heart rate and blood pressure. With higher doses, peripheral vasoconstricting effects prevail leading to an increase in systemic vascular resistance and blood pressure, while the bradycardic effect is further emphasised. Dexmedetomidine is relatively free from respiratory depressive effects when given as monotherapy to healthy subjects.

#### Sedation of adult ICU (Intensive Care Unit) patients

In placebo controlled trials in a post-operative ICU population previously intubated and sedated with midazolam or propofol, Dexdor significantly reduced the requirement for both rescue sedative (midazolam or propofol) and opioids during sedation for up to 24 hours. Most dexmedetomidine patients required no additional sedative treatment. Patients could be successfully extubated without stopping the Dexdor infusion. Studies from outside the ICU have confirmed that Dexdor can be administered safely to patients without endotracheal intubation provided adequate monitoring is in place.

Dexmedetomidine was similar to midazolam (Ratio 1.07; 95% CI 0.971, 1.176) and propofol (Ratio 1.00; 95% CI 0.922, 1.075) on the time in target sedation range in a predominantly medical population requiring prolonged light to moderate sedation (RASS 0 to -3) in the ICU for up to 14 days, reduced the duration of mechanical ventilation compared to midazolam and reduced the time to extubation compared to midazolam and propofol. Compared to both propofol and midazolam, patients were more easily roused, more cooperative and better able to communicate whether or not they had pain. Dexmedetomidine treated patients had more frequent hypotension and bradycardia but less tachycardia than those receiving midazolam and more frequent tachycardia but similar hypotension to propofol-treated patients. Delirium measured by the CAM-ICU scale was reduced in a study compared to midazolam and delirium-related adverse events were lower on dexmedetomidine compared to propofol. Those patients who withdrew due to insufficient sedation were switched to either propofol or midazolam. The risk of insufficient sedation was increased in patients who were difficult to sedate with standard care immediately prior to switching.

Evidence of paediatric efficacy was seen in a dose-controlled ICU study in a largely post-operative population aged 1 month to  $\leq 17$  years. Approximately 50% of patients treated with dexmedetomidine did not require rescue addition of midazolam during a median treatment period of 20.3 hours, not exceeding 24 hours. Data on treatment for  $> 24$  hours is not available. Data in new-born infants (28–44 weeks gestation) is very limited and restricted to low doses ( $\leq 0.2$  mcg/kg/h) (see sections 5.2 and 4.4). New-born infants may be particularly sensitive to the bradycardic effects of Dexdor in the presence of hypothermia and in conditions of heart rate-dependent cardiac output.

In double blind comparator controlled ICU studies the incidence of cortisol suppression in patients treated with dexmedetomidine ( $n = 778$ ) was 0.5% compared with 0% in patients treated with either midazolam ( $n = 338$ ) or propofol ( $n = 275$ ). The event was reported as mild in 1 and moderate in 3 cases.

#### Procedural/awake sedation

The safety and efficacy of dexmedetomidine for sedation of non-intubated patients prior to and/or during surgical and diagnostic procedures was evaluated in two randomised, double-blind, placebo-controlled multicentre clinical trials.

- Study 1 randomised patients undergoing elective surgeries/procedures under monitored anaesthesia care and local/regional anaesthesia to receive a loading infusion of dexmedetomidine

either 1 µg/kg (n = 129) or 0.5 µg/kg (n = 134), or placebo (normal saline; n = 63) given over 10 minutes and followed by a maintenance infusion started at 0.6 µg/kg/h. The maintenance infusion of study drug could be titrated from 0.2 µg/kg/h to 1 µg/kg/h. The proportion of patients that achieved the targeted sedation level (Observer's Assessment of Alertness/Sedation Scale ≤ 4) without need for rescue midazolam was 54% of the patients receiving dexmedetomidine 1 µg/kg and 40% of the patients receiving dexmedetomidine 0.5 µg/kg compared to 3% of patients receiving the placebo. The risk difference in proportion of subjects randomised to dexmedetomidine 1 µg/kg group and dexmedetomidine 0.5 µg/kg group not requiring rescue midazolam was 48% (95% CI: 37%–57%) and 40% (95% CI: 28%–48%), respectively compared placebo. The median (range) midazolam rescue dose was 1.5 (0.5–7.0) mg in the dexmedetomidine 1.0 µg/kg group, 2.0 (0.5–8.0) mg in the dexmedetomidine 0.5 µg/kg group, and 4.0 (0.5–14.0) mg in the placebo group. The difference in means in dose of rescue midazolam in dexmedetomidine 1 µg/kg and dexmedetomidine 0.5 µg/kg group compared to placebo was -3.1 mg (95% CI: -3.8 – -2.5) and -2.7 mg (95% CI: -3.3 – -2.1), respectively favouring dexmedetomidine. The median time to first rescue dose was 114 minutes in the dexmedetomidine 1.0 µg/kg group, 40 minutes in the dexmedetomidine 0.5 µg/kg group, and 20 minutes in the placebo group.

- Study 2 randomised patients undergoing awake fiberoptic intubation under topical anaesthesia to receive a loading infusion of dexmedetomidine 1 µg/kg (n = 55) or placebo (normal saline) (n = 50) given over 10 minutes and followed by a fixed maintenance infusion of 0.7 µg/kg/h. To maintain a Ramsay Sedation Scale ≥ 2 53% of the patients receiving dexmedetomidine did not require midazolam rescue vs. 14% of patients receiving placebo. The risk difference in proportion of subjects randomised to dexmedetomidine not requiring rescue midazolam was 43% (95% CI: 23 %–57%) compared placebo. The mean midazolam rescue dose was 1.1 mg in the dexmedetomidine group, and 2.8 mg in the placebo group. The difference in means in dose of rescue midazolam was -1.8 mg (95% CI: -2.7 – -0.86) favouring dexmedetomidine.

## 5.2 Pharmacokinetic properties

The pharmacokinetics of dexmedetomidine has been assessed following short term IV administration in healthy volunteers and long term infusion in ICU population.

### Distribution

Dexmedetomidine exhibits a two-compartment disposition model. In healthy volunteers it exhibits a rapid distribution phase with a central estimate of the distribution half-life ( $t_{1/2\alpha}$ ) of about 6 minutes. The mean estimate of the terminal elimination half-life ( $t_{1/2}$ ) is approximately 1.9 to 2.5 h (min 1.35, max 3.68 h) and the mean estimate of the steady-state volume of distribution ( $V_{ss}$ ) is approximately 1.16 to 2.16 l/kg (90 to 151 litres). Plasma clearance (Cl) has a mean estimated value of 0.46 to 0.73 l/h/kg (35.7 to 51.1 l/h). The mean body weight associated with these  $V_{ss}$  and Cl estimates was 69 kg. Plasma pharmacokinetics of dexmedetomidine is similar in the ICU population following infusion >24 h. The estimated pharmacokinetic parameters are:  $t_{1/2}$  approximately 1.5 hours,  $V_{ss}$  approximately 93 litres and Cl approximately 43 l/h. The pharmacokinetics of dexmedetomidine is linear in the dosing range from 0.2 to 1.4 µg/kg/h and it does not accumulate in treatments lasting up to 14 days. Dexmedetomidine is 94% bound to plasma proteins. Plasma protein binding is constant over the concentration range of 0.85 to 85 ng/ml. Dexmedetomidine binds to both human serum albumin and Alpha-1-acid glycoprotein with serum albumin as the major binding protein of dexmedetomidine in plasma.

### Biotransformation and Elimination

Dexmedetomidine is eliminated by extensive metabolism in the liver. There are three types of initial metabolic reactions; direct N-glucuronidation, direct N-methylation and cytochrome P450 catalysed oxidation. The most abundant circulating dexmedetomidine metabolites are two isomeric N-glucuronides. Metabolite H-1, N-methyl 3-hydroxymethyl dexmedetomidine O-glucuronide, is also a major circulating product of dexmedetomidine biotransformation. Cytochrome P-450 catalyses the

formation of two minor circulating metabolites, 3-hydroxymethyl dexmedetomidine produced by hydroxylation at the 3-methyl group of dexmedetomidine and H-3 produced by oxidation in the imidazole ring. Available data suggest that the formation of the oxidised metabolites is mediated by several CYP forms (CYP2A6, CYP1A2, CYP2E1, CYP2D6 and CYP2C19). These metabolites have negligible pharmacological activity.

Following IV administration of radiolabeled dexmedetomidine an average 95% of radioactivity was recovered in the urine and 4% in the faeces after nine days. The major urinary metabolites are the two isomeric N-glucuronides, which together accounted for approximately 34% of the dose and N-methyl 3-hydroxymethyl dexmedetomidine O-glucuronide that accounted for 14.51% of the dose. The minor metabolites dexmedetomidine carboxylic acid, 3-hydroxymethyl dexmedetomidine and its O-glucuronide individually comprised 1.11 to 7.66% of the dose. Less than 1% of unchanged parent drug was recovered in the urine. Approximately 28% of the urinary metabolites are unidentified minor metabolites.

### Special Populations

No major pharmacokinetic differences have been observed based on gender or age.

Dexmedetomidine plasma protein binding is decreased in subjects with hepatic impairment compared with healthy subjects. The mean percentage of unbound dexmedetomidine in plasma ranged from 8.5% in healthy subjects to 17.9% in subjects with severe hepatic impairment. Subjects with varying degrees of hepatic impairment (Child-Pugh Class A, B, or C) had decreased hepatic clearance of dexmedetomidine and prolonged plasma elimination  $t_{1/2}$ . The mean plasma clearance values of unbound dexmedetomidine for subjects with mild, moderate, and severe hepatic impairment were 59%, 51% and 32% of those observed in the normal healthy subjects, respectively. The mean  $t_{1/2}$  for the subjects with mild, moderate or severe hepatic impairment was prolonged to 3.9, 5.4, and 7.4 hours, respectively. Although dexmedetomidine is administered to effect, it may be necessary to consider initial/maintenance dose reduction in patients with hepatic impairment depending on the degree of impairment and the response.

The pharmacokinetics of dexmedetomidine in subjects with severe renal impairment (creatinine clearance <30 ml/min) is not altered relative to healthy subjects.

Data in new-born infants (28–44 weeks gestation) to children 17 years of age are limited. Dexmedetomidine half life in children (1 months to 17 years) appears similar to that seen in adults, but in new-born infants (under 1 month) it appears higher. In the age groups 1 months to 6 years, body weight-adjusted plasma clearance appeared higher but decreased in older children. Body weight-adjusted plasma clearance in new-born infants (under 1 month) appeared lower (0.9 l/h/kg) than in the older groups due to immaturity. The available data is summarised in the following table;

Age	N	Mean (95% CI)	
		Cl (l/h/kg)	$t_{1/2}$ (h)
Under 1 month	28	0.93 (0.76, 1.14)	4.47 (3.81, 5.25)
1 to < 6 months	14	1.21 (0.99, 1.48)	2.05 (1.59, 2.65)
6 to < 12 months	15	1.11 (0.94, 1.31)	2.01 (1.81, 2.22)
12 to < 24 months	13	1.06 (0.87, 1.29)	1.97 (1.62, 2.39)
2 to < 6 years	26	1.11 (1.00, 1.23)	1.75 (1.57, 1.96)
6 to < 17 years	28	0.80 (0.69, 0.92)	2.03 (1.78, 2.31)

### **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, single and repeated dose toxicity and genotoxicity.

In the reproductive toxicity studies, dexmedetomidine had no effect on male or female fertility in the rat, and no teratogenic effects were observed in the rat or rabbit. In the rabbit study intravenous administration of the maximum dose, 96 µg/kg/day, produced exposures that are similar to those observed clinically. In the rat, subcutaneous administration at the maximum dose, 200 µg/kg/day, caused an increase in embryofetal death and reduced the fetal body weight. These effects were associated with clear maternal toxicity. Reduced fetal body weight was noted also in the rat fertility study at dose 18 µg/kg/day and was accompanied with delayed ossification at dose 54 µg/kg/day. The observed exposure levels in the rat are below the clinical exposure range.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Sodium chloride  
Water for injections

### **6.2 Incompatibilities**

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

Compatibility studies have shown potential for adsorption of dexmedetomidine to some types of natural rubber. Although dexmedetomidine is dosed to effect, it is advisable to use components with synthetic or coated natural rubber gaskets.

### **6.3 Shelf life**

3 years

#### *After dilution*

Chemical and physical in-use stability has been demonstrated for 24 hours at 25°C.

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to the use are the responsibility of the user and would not normally be longer than 24 hours at 2° to 8°C, unless dilution has taken place in controlled and validated aseptic conditions.

### **6.4 Special precautions for storage**

This medicinal product does not require any special temperature storage conditions. Keep the ampoules or vials in the outer carton in order to protect from light.

For storage conditions after dilution of the medicinal product, see section 6.3

### **6.5 Nature and contents of container**

2 ml Type I glass ampoules  
2, 5 or 10 ml Type I glass vials (with filling volumes of 2, 4 and 10 ml), grey bromobutyl rubber closure with fluoropolymer coating

*Pack sizes*

5 x 2 ml ampoules  
25 x 2 ml ampoules  
5 x 2 ml vials  
4 x 4 ml vials  
4 x 10 ml vials

Not all pack sizes may be marketed.

## 6.6 Special precautions for disposal and other handling

Ampoules and vials are intended for single patient use only.

*Preparation of solution*

Dexdor can be diluted in glucose 50 mg/ml (5%), Ringers, mannitol or sodium chloride 9 mg/ml (0.9%) solution for injection to achieve the required concentration of either 4 micrograms/ml or 8 micrograms/ml prior to administration. Please see below in tabulated form the volumes needed to prepare the infusion.

### **In case the required concentration is 4 micrograms/ml:**

<b>Volume of Dexdor 100 micrograms/ml concentrate for solution for infusion</b>	<b>Volume of diluent</b>	<b>Total volume of infusion</b>
2 ml	48 ml	50 ml
4 ml	96 ml	100 ml
10 ml	240 ml	250 ml
20 ml	480 ml	500 ml

### **In case the required concentration is 8 micrograms/ml:**

<b>Volume of Dexdor 100 micrograms/ml concentrate for solution for infusion</b>	<b>Volume of diluent</b>	<b>Total volume of infusion</b>
4 ml	46 ml	50 ml
8 ml	92 ml	100 ml
20 ml	230 ml	250 ml
40 ml	460 ml	500 ml

The solution should be shaken gently to mix well.

Dexdor should be inspected visually for particulate matter and discoloration prior to administration.

Dexdor has been shown to be compatible when administered with the following intravenous fluids and medicinal products:

Lactated Ringers, 5% glucose solution, sodium chloride 9 mg/ml (0.9%) solution for injection, mannitol 200 mg/ml (20%), thiopental sodium, etomidate, vecuronium bromide, pancuronium bromide, succinylcholine, atracurium besylate, mivacurium chloride, rocuronium bromide, glycopyrrolate bromide, phenylephrine HCl, atropine sulfate, dopamine, noradrenaline, dobutamine, midazolam, morphine sulfate, fentanyl citrate, and a plasma-substitute.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

#### **7. MARKETING AUTHORISATION HOLDER**

Orion Corporation  
Orionintie 1  
FI-02200 Espoo  
Finland

#### **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/11/718/001-002, EU/1/11/718/004, EU/1/11/718/006-007

#### **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 16 September 2011  
Date of latest renewal: 26 May 2016

#### **10. DATE OF REVISION OF THE TEXT**

Detailed information on this product is available on the website of the European Medicines Agency  
<http://www.ema.europa.eu>

## **ANNEX II**

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer(s) responsible for batch release

Orion Corporation  
Orionintie 1  
FI-02200 Espoo  
Finland

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Dexdor 100 micrograms/ml concentrate for solution for infusion  
dexmedetomidine

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each ml of concentrate contains dexmedetomidine hydrochloride equivalent to 100 micrograms dexmedetomidine.

**3. LIST OF EXCIPIENTS**

Also contains sodium chloride, water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Concentrate for solution for infusion

5 x 2 ml ampoules  
25 x 2 ml ampoules  
5 x 2 ml vials  
4 x 4 ml vials  
4 x 10 ml vials

200 micrograms/2 ml  
400 micrograms/4 ml  
1000 micrograms/10 ml

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.  
Intravenous use  
Dexdor should be used immediately after dilution.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Keep the ampoules/vials in the outer carton in order to protect from light.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Orion Corporation  
Orionintie 1  
FI-02200 Espoo  
Finland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/11/718/001  
EU/1/11/718/002  
EU/1/11/718/004  
EU/1/11/718/006  
EU/1/11/718/007

**13. BATCH NUMBER**

Batch

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC {number}  
SN {number}  
NN {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS  
AMPOULES OR VIALS**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE OF ADMINISTRATION**

Dexdor 100 micrograms/ml sterile concentrate  
dexmedetomidine  
IV

**2. METHOD OF ADMINISTRATION**

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Batch

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

200 mcg/ 2 ml  
400 mcg/ 4 ml  
1000 mcg/ 10 ml

**6. OTHER**

**B. PACKAGE LEAFLET**

## Package leaflet: Information for the user

### Dexdor 100 micrograms/ml concentrate for solution for infusion dexmedetomidine

**Read all of this leaflet carefully before you are given this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or nurse.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

1. What Dexdor is and what it is used for
2. What you need to know before you are given Dexdor
3. How to use Dexdor
4. Possible side effects
5. How to store Dexdor
6. Contents of the pack and other information

#### **1. What Dexdor is and what it is used for**

Dexdor contains an active substance called dexmedetomidine which belongs to a medicine group called sedatives. It is used to provide sedation (a state of calm, drowsiness or sleep) for adult patients in hospital intensive care settings or awake sedation during different diagnostic or surgical procedures.

#### **2. What you need to know before you are given Dexdor**

##### **You must not be given Dexdor**

- if you are allergic to dexmedetomidine or any of the other ingredients of this medicine (listed in section 6)
- if you have some disorders of heart rhythm (heart block grade 2 or 3)
- if you have very low blood pressure which does not respond to treatment
- if you have recently had a stroke or other serious condition affecting blood supply to the brain.

##### **Warnings and precautions**

Before you have this medicine, tell your doctor or nurse if any of the following apply as Dexdor should be used cautiously:

- if you have an abnormally slow heart rate (either due to illness or high levels of physical fitness) as it may increase the risk for cardiac arrest
- if you have low blood pressure
- if you have low blood volume, for example after bleeding
- if you have certain heart disorders
- if you are elderly
- if you have a neurological disorder (for instance head or spinal cord injury or stroke)
- if you have severe liver problems
- if you have ever developed a serious fever after some medicines, especially anaesthetics.

This medicine may cause large amount of urine and excessive thirst, contact a doctor if these side effects occur. See section 4 for more information.

An increased mortality risk has been seen for patients 65 years of age and under when using this medicine, especially for patients admitted to the intensive care unit for other reasons than after surgery with a more severe disease condition on admission to the intensive care unit and with a younger age. The doctor will decide if this medicine is still suitable for you. The doctor will take into account the benefit and risks of this medicine for you, compared to treatment with other sedatives.

### **Other medicines and Dexdor**

Tell your doctor or nurse if you are taking, have recently taken or might take any other medicines.

The following medicines may enhance the effect of Dexdor:

- medicines that help you sleep or cause sedation (e.g. midazolam, propofol)
- strong pain medicines (e.g. opioids such as morphine, codeine)
- anaesthetic medicines (e.g. sevoflurane, isoflurane).

If you are using medicines which lower your blood pressure and heart rate, co-administration with Dexdor may enhance this effect. Dexdor should not be used with medicines that cause temporary paralysis.

### **Pregnancy and breast-feeding**

Dexdor should not be used during pregnancy or breast-feeding unless clearly necessary. Ask your doctor for advice before having this medicine

### **Driving and using machines**

Dexdor has major impact on the ability to drive and use machines. After you have been given Dexdor you must not drive, operate machinery, or work in dangerous situations until the effects are completely gone. Ask your doctor when you can start doing these activities again and when you can go back to this kind of work.

### **Excipients**

Dexdor contains less than 1 mmol sodium (23 mg) per ml, that is to say essentially 'sodium-free'.

## **3. How to use Dexdor**

### **Hospital intensive care**

Dexdor is administered to you by a doctor or nurse in hospital intensive care.

### **Procedural sedation/awake sedation**

Dexdor is administered to you by a doctor or a nurse prior to and/or during diagnostic or surgical procedures requiring sedation, i.e. procedural/awake sedation.

Your doctor will decide on a suitable dose for you. The amount of Dexdor depends on your age, size, general condition of health, the level of sedation needed and how you respond to the medicine. Your doctor may change your dose if needed and will monitor your heart and blood pressure during the treatment.

Dexdor is diluted and it is given to you as an infusion (drip) into your veins.

### **After sedation/wake-up**

- The doctor will keep you under medical supervision for some hours after the sedation to make sure that you feel well.
- You should not go home unaccompanied.
- Medicines to help you sleep, cause sedation or strong painkillers may not be appropriate for some time after you have been given Dexdor. Talk to your doctor about the use of these medicines and about the use of alcohol.

### **If you have been given more Dexdor than you should**

If you are given too much Dexdor, your blood pressure may go up or down, your heartbeat may slow down, you may breathe more slowly and you may feel more drowsy. Your doctor will know how to treat you based on your condition.

If you have any further questions on the use of this medicine, ask your doctor.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Very common (*may affect more than 1 in 10 people*):

- slow heart rate
- low or high blood pressure
- change in breathing pattern or stopping breathing.

Common (*may affect up to 1 in 10 people*):

- chest pain or heart attack
- fast heart rate
- low or high blood sugar
- nausea, vomiting or dry mouth
- restlessness
- high temperature
- symptoms after stopping the medicine.

Uncommon (*may affect up to 1 in 100 people*):

- reduced heart function, cardiac arrest
- swelling of the stomach
- thirst
- a condition where there is too much acid in the body
- low albumin level in blood
- shortness of breath
- hallucinations
- the medicine is not effective enough.

Not known (*frequency cannot be estimated from the available data*):

- large amount of urine and excessive thirst – may be symptoms of a hormonal disorder called diabetes insipidus. Contact a doctor if these occur.

### **Reporting of side effects**

If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Dexdor**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label and carton after EXP.

This medicine does not require any special temperature storage conditions. Keep the ampoules or vials in the outer carton in order to protect from light.

## **6. Contents of the pack and other information**

### **What Dexdor contains**

- The active substance is dexmedetomidine. Each ml of concentrate contains dexmedetomidine hydrochloride equivalent to 100 micrograms dexmedetomidine.
- The other ingredients are sodium chloride and water for injections.

Each 2 ml ampoule contains 200 micrograms of dexmedetomidine (as hydrochloride).

Each 2 ml vial contains 200 micrograms of dexmedetomidine (as hydrochloride).

Each 4 ml vial contains 400 micrograms of dexmedetomidine (as hydrochloride).

Each 10 ml vial contains 1000 micrograms of dexmedetomidine (as hydrochloride).

The concentration of the final solution after dilution should be either 4 micrograms/ml or 8 micrograms/ml.

### **What Dexdor looks like and contents of the pack**

Concentrate for solution for infusion (sterile concentrate).

The concentrate is a clear, colourless solution.

#### Containers

2 ml glass ampoules

2, 5 or 10 ml glass vials

#### Pack sizes

5 x 2 ml ampoules

25 x 2 ml ampoules

5 x 2 ml vials

4 x 4 ml vials

4 x 10 ml vials

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder**

Orion Corporation

Orionintie 1

FI-02200 Espoo

Finland

### **Manufacturer**

Orion Corporation Orion Pharma

Orionintie 1

FI-02200 Espoo

Finland

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

**Luxembourg/Luxemburg**

Orion Pharma BVBA/SPRL

Tél/Tel: +32 (0)15 64 10 20

**Nederland**

Orion Pharma BVBA/SPRL

Tel: +32 (0)15 64 10 20

**Česká republika**

Orion Pharma s.r.o.

Tel: +420 234 703 305

**Deutschland**

**Österreich**

Orion Pharma GmbH

Tel: + 49 40 899 6890

**Ελλάδα**

Orion Pharma Hellas M.E.Π.E

Τηλ: + 30 210 980 3355

**France**

Orion Pharma

Tél: +33 (0) 1 85 18 00 00

**Ísland**

Vistor hf.

Simi: +354 535 7000

**Κύπρος**

Lifepharm (ZAM) Ltd

Τηλ.: +357 22056300

**Lietuva**

UAB Orion Pharma

Tel: +370 5 276 9499

**Norge**

Orion Pharma AS

Tlf: + 47 4000 4210

**Portugal**

Orionfin Unipessoal Lda

Tel: + 351 21 154 68 20

**България**

Orion Pharma Poland Sp. z.o.o.

**Danmark**

Orion Pharma A/S

Tlf: +45 8614 00 00

**Eesti**

Orion Pharma Eesti Oü

Tel: +372 6 644 550

**España**

Orion Pharma S.L.

Tel: + 34 91 599 86 01

**Ireland**

Orion Pharma (Ireland) Ltd.

Tel: + 353 1 428 7777

**Italia**

Orion Pharma S.r.l.

Tel: + 39 02 67876111

**Latvija**

Orion Corporation

Orion Pharma pārstāvniecība

Tel: +371 20028332

**Magyarország**

Orion Pharma Kft.

Tel.: +36 1 239 9095

**Polska**

Orion Pharma Poland Sp. z.o.o.

Tel.: + 48 22 8 333 177

**Slovenija**

Orion Pharma d.o.o.

Tel: +386 (0) 1 600 8015

**Suomi/Finland**

Orion Corporation

Puh/Tel: + 358 10 4261

**Hrvatska**

PHOENIX Farmacija d.o.o.

Tel.: + 48 22 8 333 177

Tel: +385 1 6370450

**Malta**

Orion Corporation  
Tel: + 358 10 4261

**România**

Orion Corporation  
Tel: + 358 10 4261

**Slovenská republika**

Orion Pharma s.r.o.  
Tel: +420 234 703 305

**United Kingdom (Northern Ireland):**

Orion Pharma (Ireland) Ltd.  
Tel: +353 1 428 7777

**Sverige**

Orion Pharma AB  
Tel: + 46 8 623 6440

**This leaflet was last revised in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>

-----  
The following information is intended for healthcare professionals only:

**Dexdor 100 micrograms/ml concentrate for solution for infusion**

Method of administration

Dexdor should be administered by healthcare professionals skilled in the management of patients requiring intensive care or in the anaesthetic management of patients in the operating room. It must be administered only as a diluted intravenous infusion using a controlled infusion device

*Preparation of solution*

Dexdor can be diluted in glucose 50 mg/ml (5%), Ringers, mannitol or sodium chloride 9 mg/ml (0.9%) solution for injection to achieve the required concentration of either 4 micrograms/ml or 8 micrograms/ml prior to administration. Please see below in tabulated form the volumes needed to prepare the infusion.

**In the case the required concentration is 4 micrograms/ml:**

<b>Volume of Dexdor 100 micrograms/ml concentrate for solution for infusion</b>	<b>Volume of diluent</b>	<b>Total volume of infusion</b>
2 ml	48 ml	50 ml
4 ml	96 ml	100 ml
10 ml	240 ml	250 ml
20 ml	480 ml	500 ml

**In the case the required concentration is 8 micrograms/ml:**

<b>Volume of Dexdor 100 micrograms/ml concentrate for solution for infusion</b>	<b>Volume of diluent</b>	<b>Total volume of infusion</b>
4 ml	46 ml	50 ml
8 ml	92 ml	100 ml
20 ml	230 ml	250 ml
40 ml	460 ml	500 ml

The solution should be shaken gently to mix well.

Dexdor should be inspected visually for particulate matter and discoloration prior to administration.

Dexdor has been shown to be compatible when administered with the following intravenous fluids and medicinal products:

Lactated Ringers, 5% glucose solution, sodium chloride 9 mg/ml (0.9%) solution for injection, mannitol 200 mg/ml (20%), thiopental sodium, etomidate, vecuronium bromide, pancuronium bromide, succinylcholine, atracurium besylate, mivacurium chloride, rocuronium bromide, glycopyrrolate bromide, phenylephrine HCl, atropine sulfate, dopamine, noradrenaline, dobutamine, midazolam, morphine sulfate, fentanyl citrate, and a plasma-substitute.

Compatibility studies have shown potential for adsorption of dexmedetomidine to some types of natural rubber. Although dexmedetomidine is dosed to effect, it is advisable to use components with synthetic or coated natural rubber gaskets.

### **Shelf life**

Chemical and physical in-use stability has been demonstrated for 24 hours at 25°C.

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to the use are the responsibility of the user and would not normally be longer than 24 hours at 2° to 8°C, unless dilution has taken place in controlled and validated aseptic conditions.