

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

1. NAME OF THE MEDICINAL PRODUCT

Empliciti 300 mg powder for concentrate for solution for infusion.
Empliciti 400 mg powder for concentrate for solution for infusion.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Empliciti 300 mg powder for concentrate for solution for infusion
Each vial contains 300 mg elotuzumab*.

Empliciti 400 mg powder for concentrate for solution for infusion
Each vial contains 400 mg elotuzumab.

After reconstitution, each mL of concentrate contains 25 mg elotuzumab.

* Elotuzumab is produced in NS0 cells by recombinant DNA technology.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Powder for concentrate for solution for infusion (powder for concentrate).

The powder is white to off white whole or fragmented cake.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Empliciti is indicated in combination with lenalidomide and dexamethasone for the treatment of multiple myeloma in adult patients who have received at least one prior therapy (see sections 4.2 and 5.1).

4.2 Posology and method of administration

Elotuzumab therapy should be initiated and supervised by physicians experienced in the treatment of multiple myeloma.

Premedication for prevention of infusion reaction

Patients must be administered with the following premedications 45-90 minutes prior to Empliciti infusion (see section 4.4):

Dexamethasone 8 mg intravenous

H1 blocker: diphenhydramine (25-50 mg orally or intravenous) or equivalent H1 blocker.

H2 blocker: ranitidine (50 mg intravenous or 150 mg orally) or equivalent H2 blocker.

Paracetamol (650-1000 mg orally).

Management of infusion reaction

If a \geq Grade 2 infusion reaction occurs during Empliciti administration, the infusion must be interrupted. Upon resolution to \leq Grade 1, Empliciti should be restarted at 0.5 mL/min and may be gradually increased at a rate of 0.5 mL/min every 30 minutes as tolerated to the rate at which the infusion reaction occurred. If there is no recurrence of the infusion reaction, the escalation can be resumed (see Table 2).

In patients who experience an infusion reaction, vital signs should be monitored every 30 minutes for 2 hours after the end of the Empliciti infusion. If the infusion reaction recurs, the Empliciti infusion must be stopped and not restarted on that day (see section 4.4). Very severe infusion reactions (\geq Grade 3) may require permanent discontinuation of Empliciti therapy and emergency treatment.

Posology for administration with lenalidomide and dexamethasone

The dosing schedule is presented in Table 1.
Treatment should continue until disease progression or unacceptable toxicity.

The recommended dose of Empliciti is 10 mg/kg administered intravenously every week (28-day cycle), on days 1, 8, 15, and 22 for the first two cycles and every 2 weeks thereafter on days 1 and 15.

The recommended dose of lenalidomide is 25 mg orally once daily on days 1-21 of repeated 28-day cycles, and at least 2 hours after Empliciti infusion when administered on the same day.

The administration of dexamethasone is as follows:

- On days that Empliciti is administered, dexamethasone should be given as 28 mg orally once daily between 3 and 24 hours before Empliciti plus 8 mg intravenously between 45 and 90 minutes before Empliciti on days 1, 8, 15, and 22 of repeated 28-day cycles.
- On days that Empliciti is not administered but a dose of dexamethasone is scheduled (Days 8 and 22 of cycle 3 and all subsequent cycles), dexamethasone should be given 40 mg orally.

Table 1: Recommended dosing schedule of Empliciti in combination with lenalidomide and dexamethasone

Cycle	28-Day Cycles 1 & 2				28-Day Cycles 3+			
	1	8	15	22	1	8	15	22
Day of Cycle	1	8	15	22	1	8	15	22
Premedication	✓	✓	✓	✓	✓		✓	
Empliciti (mg/kg) Intravenous	10	10	10	10	10		10	
Lenalidomide (25 mg) Oral	Days 1-21				Days 1-21			
Dexamethasone (mg) Oral	28	28	28	28	28	40	28	40
Day of Cycle	1	8	15	22	1	8	15	22

For additional information concerning lenalidomide and dexamethasone, see the corresponding Summary of Product Characteristics.

See Method of administration below for instruction on infusion rates.

Dose delay, interruption, or discontinuation

If the dose of one medicine in the regimen is delayed, interrupted, or discontinued, the treatment with the other medicinal products may continue as scheduled. However, if oral or intravenous

dexamethasone is delayed or discontinued, the administration of Empliciti should be based on clinical judgment (e.g. risk of hypersensitivity) (see section 4.4).

Special populations

Paediatric population

There is no relevant use of Empliciti in the paediatric population for the indication of multiple myeloma.

Elderly

No dose adjustment is required for elotuzumab in patients over 65 years of age (see section 5.2). Data on the efficacy and safety of elotuzumab in patients ≥ 85 years of age are very limited.

Renal impairment

No dose adjustment of Empliciti is required for patients with mild ($\text{CrCl} = 60 - 89 \text{ mL/min}$), moderate ($\text{CrCl} = 30 - 59 \text{ mL/min}$), severe ($\text{CrCl} < 30 \text{ mL/min}$) renal impairment or end stage renal disease requiring dialysis (see section 5.2).

Hepatic impairment

No dose adjustment for Empliciti is required for patients with mild hepatic impairment (total bilirubin [TB] \leq to the upper limit of normal [ULN] and $\text{AST} > \text{ULN}$ or $\text{TB} < 1$ to $1.5 \times \text{ULN}$ and any AST). Empliciti has not been studied in patients with moderate ($\text{TB} > 1.5$ to $3 \times \text{ULN}$ and any AST) or severe ($\text{TB} > 3 \times \text{ULN}$ and any AST) hepatic impairment (see section 5.2).

Method of administration

Empliciti is for intravenous use only.

The administration of the reconstituted and diluted solution must be initiated at an infusion rate of 0.5 mL/min. If the infusion is well tolerated the infusion rate may be increased in a stepwise fashion as described in Table 2. The maximum infusion rate should not exceed 5 mL/min.

Table 2: Infusion rate for Empliciti

Cycle 1, Dose 1		Cycle 1, Dose 2		Cycle 1, Dose 3 and 4 and all subsequent Cycles
Time interval	Rate	Time interval	Rate	
0 - 30 min	0.5 mL/min	0 - 30 min	3 mL/min	5 mL/min*
30 - 60 min	1 mL/min	≥ 30 min	4 mL/min*	
≥ 60 min	2 mL/min*	-	-	

* Continue this rate until infusion is completed, approximately 1 hour based on patient weight.

For instructions on reconstitution and dilution of Empliciti before administration, see section 6.6.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

The Summary of Product Characteristics for all medicinal products used in combination with Empliciti must be consulted before starting therapy.

4.4 Special warnings and precautions for use

Infusion reaction

Infusion reactions have been reported in patients receiving elotuzumab (see section 4.8).

Premedication consisting of dexamethasone, H1 blocker, H2 blocker, and paracetamol must be administered prior to Empliciti infusion (see section 4.2 Premedication). The rate of infusion reactions was much higher in patients who were not premedicated.

If any of the symptoms of infusion reaction reach Grade ≥ 2 , Empliciti infusion must be interrupted and appropriate medical and supportive measures instituted. Vital signs should be monitored every 30 minutes for 2 hours after the end of the Empliciti infusion. Once the reaction has resolved (symptoms \leq Grade 1), Empliciti can be restarted at the initial infusion rate of 0.5 mL/min. If symptoms do not recur, the infusion rate may be gradually escalated every 30 minutes to a maximum of 5 mL/min (see section 4.2 Method of administration).

Very severe infusion reactions may require permanent discontinuation of Empliciti therapy and emergency treatment. Patients with mild or moderate infusion reactions may receive Empliciti with a reduced infusion rate and close monitoring (see section 4.2 Method of administration).

Conditions for use of medicinal products used with Empliciti

Empliciti is used in combination with other medicinal products; therefore, the conditions for use applicable to those medicinal products also apply to the combination therapy. The Summary of Product Characteristics for all medicinal products used in combination with Empliciti must be consulted before starting therapy.

Infections

In clinical trials of patients with multiple myeloma, the incidence of all infections, including pneumonia, were higher in patients treated with Empliciti (see section 4.8). Patients should be monitored and infections should be managed with standard treatment.

Second primary malignancies (SPMs)

In a clinical trial of patients with multiple myeloma that compared Empliciti combined with lenalidomide and dexamethasone treatment to lenalidomide and dexamethasone treatment (Study 1), the incidence of SPMs, and specifically of solid tumours and non-melanoma skin cancer, was higher in patient treated with Empliciti (see section 4.8). SPMs are known to be associated with lenalidomide exposure, which was extended in patients treated with Empliciti combined with lenalidomide and dexamethasone vs. lenalidomide and dexamethasone. The rate of haematologic malignancies was the same between the two treatment arms. Patients should be monitored for the development of SPMs.

4.5 Interaction with other medicinal products and other forms of interaction

Pharmacokinetic interaction studies have not been conducted. Empliciti, as a humanised monoclonal antibody, is not expected to be metabolised by cytochrome P450 (CYP) enzymes or other drug metabolising enzymes, inhibition or induction of these enzymes by co-administered medicinal products is not anticipated to affect the pharmacokinetics of Empliciti.

Empliciti may be detected in the serum protein electrophoresis (SPEP) and serum immunofixation assays of myeloma patients and could interfere with correct response classification. The presence of elotuzumab in patient's serum may cause a small peak in the early gamma region on SPEP that is IgG κ on serum immunofixation. This interference can impact the determination of complete response and possibly relapse from complete response in patients with IgG kappa myeloma protein.

In case of detection of additional peaks on serum immunofixation, the possibility of a biconal gammopathy should be excluded.

The Summary of Product Characteristics for all medicinal products used in combination with Empliciti must be consulted before starting therapy.

4.6 Fertility, pregnancy and lactation

Woman of childbearing potential/Contraception in the males and females

Empliciti should not be used in women of childbearing potential, unless the clinical condition of the woman requires treatment with elotuzumab. Women of childbearing potential should use effective contraception.

Male patients must use effective contraception measures during and for 180 days following treatment if their partner is pregnant or of childbearing potential and not using effective contraception.

Pregnancy

There is no human experience with elotuzumab during pregnancy. Elotuzumab will be given in combination with lenalidomide, which is contraindicated during pregnancy. No animal data are present regarding the effect on reproductive toxicity because of the lack of an adequate animal model. Empliciti should not be used during pregnancy unless the clinical condition of the woman requires treatment with elotuzumab.

The Summary of Product Characteristics for all medicinal products used in combination with Empliciti must be consulted before starting therapy. When Empliciti is used with lenalidomide there is a risk of foetal harm, including severe life-threatening human birth defects associated with these agents and the need to follow requirements regarding pregnancy avoidance, including testing and contraception. Lenalidomide is present in the blood and sperm of patients receiving the medicine. Refer to the Summary of Product Characteristics for requirements regarding contraception due to presence and transmission in sperm and for additional detail. Patients receiving Empliciti in combination with lenalidomide should adhere to the pregnancy prevention programme of lenalidomide.

Breast-feeding

Elotuzumab is not expected to be excreted into human milk. Elotuzumab will be given in combination with lenalidomide and breast-feeding should be stopped because of the use of lenalidomide.

Fertility

Studies to evaluate the effect of elotuzumab on fertility have not been performed. Thus, the effect of elotuzumab on male and female fertility is unknown.

4.7 Effects on ability to drive and use machines

On the basis of reported adverse reactions, Empliciti is not expected to influence the ability to drive or use machines. Patients experiencing infusion reactions should be advised not to drive and use machines until symptoms abate.

4.8 Undesirable effects

Summary of safety profile

The safety data of elotuzumab have been assessed from a total of 554 patients with multiple myeloma treated with elotuzumab in combination with lenalidomide and dexamethasone (451 patients) or bortezomib and dexamethasone (103 patients) pooled across 6 clinical trials. The majority of adverse reactions were mild to moderate (Grade 1 or 2).

The most serious adverse reaction that may occur during elotuzumab treatment is pneumonia.

The most common adverse reactions (occurring in > 10% of patients) with elotuzumab treatment were infusion related reactions, diarrhoea, herpes zoster, nasopharyngitis, cough, pneumonia, upper respiratory tract infection, lymphopenia and weight decreased.

Tabulated list of adverse reactions

Adverse reactions reported in 554 patients with multiple myeloma who were treated with elotuzumab in 6 clinical trials are presented in Table 3.

These reactions are presented by system organ class and by frequency. Frequencies are defined as: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$); and not known (cannot be estimated from available data).

Table 3: Adverse reactions in patients with multiple myeloma treated with Empliciti

System Organ Class	Adverse reactions	Frequency overall	Grade 3/4 frequency
<i>Infections and infestations</i>	Herpes zoster ^a	Very common	Common
	Nasopharyngitis	Very common	None reported
	Pneumonia ^b	Very common	Very common
	Upper respiratory tract infection	Very common	Common
<i>Blood and lymphatic system disorders</i>	Lymphopenia ^c	Very common	Very common
<i>Immune system disorders</i>	Anaphylactic reaction	Uncommon	Uncommon
	Hypersensitivity	Common	Uncommon
<i>Psychiatric disorders</i>	Mood altered	Common	None reported
<i>Nervous system disorders</i>	Headache	Very common	Uncommon
	Hypoaesthesia	Common	Uncommon
<i>Vascular disorders</i>	Deep vein thrombosis	Common	Common
<i>Respiratory, thoracic and mediastinal disorders</i>	Cough ^d	Very common	Uncommon
	Oropharyngeal pain	Common	None reported
<i>Gastrointestinal disorders</i>	Diarrhoea	Very common	Common
<i>Skin and subcutaneous tissue disorders</i>	Night sweats	Common	None reported
<i>General disorders and administration site conditions</i>	Chest pain	Common	Common
	Fatigue	Very common	Common
	Pyrexia	Very common	Common
<i>Investigations</i>	Weight decreased	Very common	Uncommon
<i>Injury, poisoning and procedural complications</i>	Infusion related reaction	Common	Common

^a The term herpes zoster is a grouping of the following terms: herpes zoster, oral herpes, and herpes virus infection.

^b The term pneumonia is a grouping of the following terms: pneumonia, atypical pneumonia, bronchopneumonia, lobar pneumonia, bacterial pneumonia, fungal pneumonia, pneumonia influenza, and pneumococcal pneumonia.

^c The term lymphopenia includes the following terms: lymphopenia and lymphocyte count decreased.

^d The term cough includes the following terms: cough, productive cough, and upper airway cough syndrome.

Exposure-adjusted rates for adverse reactions (all Grades and Grade 3/4) in Study 1, a clinical trial in patients with multiple myeloma comparing Empliciti combined with lenalidomide and dexamethasone treatment (N = 318) to lenalidomide and dexamethasone treatment (N = 317), is shown in Table 4.

Table 4: Exposure-adjusted rates for adverse reactions for Empliciti-treated patients versus lenalidomide and dexamethasone-treated patients [includes multiple occurrences of all treated patients]

Adverse reaction	Empliciti + Lenalidomide and Dexamethasone N = 318				Lenalidomide and Dexamethasone N = 317			
	All grades		Grade 3/4		All grades		Grade 3/4	
	Event count	Rate (incidence rate/100 patient years)	Event count	Rate (incidence rate/100 patient years)	Event count	Rate (incidence rate/100 patient years)	Event count	Rate (incidence rate/100 patient years)
Diarrhoea	303	59.2	19	3.7	206	49.3	13	3.1
Pyrexia	220	43.0	8	1.6	116	27.7	10	2.4
Fatigue	205	40.0	33	6.4	145	34.7	26	6.2
Cough ^a	170	33.2	1	0.2	85	20.3	-	-
Nasopharyngitis	151	29.5	-	-	116	27.7	-	-
Upper respiratory tract infection	129	25.2	2	0.4	95	22.7	4	1.0
Lymphopenia ^b	90	17.6	65	12.7	57	13.6	31	7.4
Headache	88	17.2	1	0.2	40	9.6	1	0.2
Pneumonia ^c	80	15.6	54	10.5	54	12.9	34	8.1
Herpes zoster ^d	51	10.0	5	1.0	24	5.7	3	0.7
Oropharyngeal pain	45	8.8	-	-	17	4.1	-	-
Weight decreased	44	8.6	4	0.8	20	4.8	-	-
Night sweats	31	6.1	-	-	12	2.9	-	-
Chest pain	29	5.7	2	0.4	12	2.9	1	0.2
Deep vein thrombosis	26	5.1	18	3.5	12	2.9	7	1.7
Hypoaesthesia	25	4.9	1	0.2	12	2.9	-	-
Mood altered	23	4.5	-	-	8	1.9	-	-
Hypersensitivity	10	2.0	-	-	4	1.0	1	0.2

^a The term cough includes the following terms: cough, productive cough, and upper airway cough syndrome.

^b The term lymphopenia includes the following terms: lymphopenia and lymphocyte count decreased.

^c The term pneumonia is a grouping of the following terms: pneumonia, atypical pneumonia, bronchopneumonia, lobar pneumonia, bacterial pneumonia, fungal pneumonia, pneumonia influenza, and pneumococcal pneumonia.

^d The term herpes zoster is a grouping of the following terms: herpes zoster, oral herpes, and herpes virus infection.

Description of selected adverse reactions

Infusion reactions

In a clinical trial of patients with multiple myeloma (Study 1), infusion reactions were reported in approximately 10% of premedicated patients treated with Empliciti combined with lenalidomide and

dexamethasone (N = 318) (see section 4.4). The rate of mild to moderate infusion reactions was > 50% in patients who were not premedicated. All reports of infusion reaction were ≤ Grade 3. Grade 3 infusion reactions occurred in 1% of patients. The most common symptoms of an infusion reaction included fever, chills, and hypertension. Five percent (5%) of patients required interruption of the administration of Empliciti for a median of 25 minutes due to infusion reaction, and 1% of patients discontinued due to infusion reactions. Of the patients who experienced an infusion reaction, 70% (23/33) had the reaction during the first dose.

Infections

The incidence of infections, including pneumonia, was higher with Empliciti treatment than with control (see section 4.4). In a clinical trial of patients with multiple myeloma (Study 1), infections were reported in 81.4% of patients in the Empliciti combined with lenalidomide and dexamethasone arm (N = 318) and 74.4% in lenalidomide and dexamethasone arm (N = 317). Grade 3-4 infections were noted in 28% and 24.3% of Empliciti combined with lenalidomide and dexamethasone and lenalidomide and dexamethasone treated patients, respectively. Fatal infections were infrequent and were reported in 2.5% of Empliciti combined with lenalidomide and dexamethasone and 2.2% of lenalidomide and dexamethasone treated patients. The incidence of pneumonia was higher in the Empliciti combined with lenalidomide and dexamethasone arm compared to lenalidomide and dexamethasone arm reported at 15.1% vs. 11.7% with a fatal outcome at 0.6% vs. 0%, respectively.

Second Primary Malignancies

The incidence of SPMs was higher with Empliciti treatment than with control (see section 4.4). In a clinical trial of patients with multiple myeloma (Study 1), invasive SPMs have been observed in 6.9% of patients treated with Empliciti combined with lenalidomide and dexamethasone (N = 318) and 4.1% of patients treated with lenalidomide and dexamethasone (N = 317). Second Primary Malignancies are known to be associated with lenalidomide exposure which was extended in patients treated with Empliciti combined with lenalidomide and dexamethasone vs. lenalidomide and dexamethasone. The rate of haematologic malignancies were the same between the two treatment arms (1.6%). Solid tumours were reported in 2.5% and 1.9% of Empliciti combined with lenalidomide and dexamethasone and lenalidomide and dexamethasone treated patients, respectively. Non-melanoma skin cancer was reported in 3.1% and 1.6% of patients treated with Empliciti combined with lenalidomide and dexamethasone and lenalidomide and dexamethasone, respectively.

Deep vein thrombosis

In a clinical trial of patients with multiple myeloma (Study 1), deep vein thromboses were reported in 7.2% of patients treated with Empliciti combined with lenalidomide and dexamethasone (N = 318) and 3.8% of patients treated with lenalidomide and dexamethasone (N = 317). Among, patients treated with aspirin, deep vein thromboses were reported in 4.1% of patients treated with Empliciti combined with lenalidomide and dexamethasone (E-Ld) and 1.4% of patients treated with lenalidomide and dexamethasone (Ld). The rates of deep vein thromboses observed between treatment arms were similar for patients given prophylaxis with low molecular weight heparin (2.2% in both treatment arms), and for patients given vitamin K antagonists the rates were 0% for patients treated with E-Ld and 6.7% for patients treated with Ld.

Immunogenicity

As with all therapeutic proteins, there is a potential for immunogenicity to Empliciti. Of 390 patients across four clinical studies who were treated with Empliciti and evaluable for the presence of anti-product antibodies, 72 patients (18.5%) tested positive for treatment-emergent anti-product antibodies by an electrochemiluminescent (ECL) assay. Neutralizing antibodies were detected in 19 of 299 patients in Study 1. In the majority of patients, immunogenicity occurred early in treatment and was transient resolving by 2 to 4 months. There was no clear causal evidence of altered pharmacokinetic, efficacy, or toxicity profiles with anti-product antibody development based on the population pharmacokinetic and exposure-response analyses.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare

professionals are asked to report any suspected adverse reactions via [the national reporting system](#) listed in [Appendix V](#).

4.9 Overdose

One patient was reported to be overdosed with 23.3 mg/kg of elotuzumab in combination with lenalidomide and dexamethasone. The patient had no symptoms, did not require any treatment for the overdose, and was able to continue on elotuzumab therapy.

In clinical studies, approximately 78 patients were evaluated with elotuzumab at 20 mg/kg without apparent toxic effects.

In case of overdose, patients should be closely monitored for signs or symptoms of adverse reactions, and appropriate symptomatic treatment instituted.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antineoplastic agents, monoclonal antibodies. ATC code: L01XC23.

Mechanism of action

Elotuzumab is an immunostimulatory humanised, IgG1 monoclonal antibody that specifically targets the SLAMF7 (signaling lymphocyte activation molecule family member 7) protein. SLAMF7 is highly expressed on multiple myeloma cells independent of cytogenetic abnormalities. SLAMF7 is also expressed on natural killer cells, normal plasma cells, and other immune cells including some T cell subsets, monocytes, B cells, and pDCs (plasmacytoid dendritic cells), but is not detected on normal solid tissues or haematopoietic stem cells.

Elotuzumab directly activates natural killer cells through both the SLAMF7 pathway and Fc receptors enhancing anti-myeloma activity *in vitro*. Elotuzumab also targets SLAMF7 on myeloma cells and facilitates the interaction with natural killer cells to mediate the killing of myeloma cells through antibody-dependent cellular cytotoxicity (ADCC). In nonclinical models, elotuzumab has demonstrated synergistic activity when combined with lenalidomide or bortezomib.

Clinical efficacy and safety

Two randomised, open-label studies were conducted to evaluate the efficacy and safety of Empliciti (elotuzumab) in adult patients with multiple myeloma who have received one or more prior therapies. Study 1 provided the pivotal data for the indication for Empliciti in combination with lenalidomide and dexamethasone.

Study 1

A randomised, open-label study was conducted to evaluate the efficacy and safety of Empliciti in combination with lenalidomide and dexamethasone in patients with multiple myeloma who have received one to three prior therapies. All patients had documented progression following their most recent therapy. Patients who were refractory to lenalidomide were excluded and 6% of patients had prior lenalidomide treatment. Patients had to recover after transplant for a minimum of 12 weeks from autologous stem cell transplant (SCT), and 16 weeks from allogeneic SCT. Patients with cardiac amyloidosis or plasma cell leukemia were excluded from this study.

Eligible patients were randomised in a 1:1 ratio to receive either Empliciti in combination with lenalidomide and dexamethasone or lenalidomide and dexamethasone. Treatment was administered in 4-week cycles until disease progression or unacceptable toxicity. Elotuzumab 10 mg/kg was administered intravenously each week for the first 2 cycles and every 2 weeks thereafter. Prior to Empliciti infusion, dexamethasone was administered as a divided dose: an oral dose of 28 mg and an

intravenous dose of 8 mg. In the control group and on weeks without Empliciti, dexamethasone 40 mg was administered as a single oral dose weekly. Lenalidomide 25 mg was taken orally once daily for the first 3 weeks of each cycle. Assessment of tumour response was conducted every 4 weeks.

A total of 646 patients were randomised to receive treatment: 321 to Empliciti in combination with lenalidomide and dexamethasone and 325 to lenalidomide and dexamethasone.

Demographics and baseline characteristics were well balanced between treatment arms. The median age was 66 years (range 37 to 91); 57% of patients were older than 65 years; 60% of patients were male; Whites comprised 84% of the study population, Asians 10%, and blacks 4%. The International Staging System (ISS) Stage was I in 43%, II in 32% and III in 21% of patients. The high risk cytogenetic categories of del17p and t(4;14) were present in 32% and 9% of patients, respectively. The median number of prior therapies was 2. Thirty-five percent (35%) of patients were refractory (progression during or within 60 days of last therapy) and 65% were relapsed (progression after 60 days of last therapy). Prior therapies included: stem cell transplant (55%), bortezomib (70%) melphalan (65%), thalidomide (48%), and lenalidomide (6%).

The primary endpoints of this study, progression-free survival (PFS), as assessed by hazard ratio, and overall response rate (ORR) were determined based on assessments made by a blinded Independent Review Committee. Efficacy results are presented in Table 5 and Figure 1. The median number of treatment cycles was 19 for the Empliciti arm and 14 for the comparator arm.

Table 5: Efficacy results for Study 1


	Empliciti + Lenalidomide/ Dexamethasone N = 321	Lenalidomide/ Dexamethasone N = 325
PFS (ITT)		
Hazard Ratio [97.61% CI]	0.68 [0.55, 0.85]	
Stratified log-rank test p-value ^a	0.0001	
1-Year PFS rate (%) [95% CI]	68 [63, 73]	56 [50, 61]
2-Year PFS rate (%) [95% CI]	39 [34, 45]	26 [21, 31]
3-Year PFS rate ^b (%) [95% CI]	23 [18, 28]	15 [10, 20]
Median PFS in months [95% CI]	18.5 [16.5, 21.4]	14.3 [12.0, 16.0]
Response		
Overall Response (ORR) ^c n (%) [95% CI]	252 (78.5) [73.6, 82.9]	213 (65.5) [60.1, 70.7]
p-value ^d	0.0002	
Complete Response (CR + sCR) ^c n (%)	14 (4.4) ^f	24 (7.4)
Very Good Partial Response (VGPR) n (%)	91 (28.3)	67 (20.6)
Partial Response (RR/PR) n (%)	147 (45.8)	122 (37.5)
Combined Responses (CR+sCR+VGPR) n (%)	105 (32.7)	91 (28.0)
Overall Survival^g		
Hazard Ratio [95% CI]	0.77 [0.61, 0.97]	
Stratified log-rank test p-value	0.0257 ^h	
Median OS in months [95% CI]	43.7 [40.34, NE]	39.6 [33.25, NE]

^a p-value based on the log-rank test stratified by B2 microglobulins (<3.5 mg/L versus ≥ 3.5 mg/L), number of prior lines of therapy (1 versus 2 or 3), and prior immunomodulatory therapy (no versus prior thalidomide only versus other).

^b A pre-specified analysis for 3-year PFS rate was performed based on a minimum follow-up time of 33 months.

- c European Group for Blood and Marrow Transplantation (EBMT) criteria.
- d p-value based on the Cochran-Mantel-Haenszel chi-square test stratified by B2 microglobulins (<3.5 mg/L versus ≥ 3.5 mg/L), number of prior lines of therapy (1 versus 2 or 3), and prior immunomodulatory therapy (no versus prior thalidomide only versus other).
- e Complete response (CR) + stringent complete response (sCR).
- f Complete response rates in Emluciti group may be underestimated due to interference of elotuzumab monoclonal antibody with immunofixation assay and serum protein electrophoresis assay.
- g A pre-specified interim analysis for OS was performed based on a minimum follow-up time of 35.4 months.
- h The interim OS analysis did not meet the protocol-specified early stopping boundary for OS ($p \leq 0.014$).

Figure 1: Progression free survival



Number of subjects at risk

E-Ld	321	282	240	206	164	133	87	43	12	1
Ld	325	262	204	168	130	97	53	24	7	


Improvements observed in PFS were consistent across subsets regardless of age (< 65 versus ≥ 65), risk status, presence or absence of cytogenetic categories del17p or t(4;14), ISS stage, number of prior therapies, prior immunomodulatory exposure, prior bortezomib exposure, relapsed or refractory status or renal function as shown in Table 6.

Table 6: Efficacy results for subsets

Subset description	E-Ld N = 321 Median PFS (months) [95% CI]	Ld N = 325 Median PFS (months) [95% CI]	HR [95% CI]
Age			
< 65 years	19.4 [15.9, 23.1]	15.7 [11.2, 18.5]	0.74 [0.55, 1.00]
≥ 65 years	18.5 [15.7, 22.2]	12.9 [10.9, 14.9]	0.64 [0.50, 0.82]
Risk factors			
High risk	14.8 [9.1, 19.6]	7.2 [5.6, 11.2]	0.63 [0.41, 0.95]
Standard risk	19.4 [16.5, 22.7]	16.4 [13.9, 18.5]	0.75 [0.59, 0.94]
Cytogenetic category			
Presence of del17p	19.6 [15.8, NE]	14.9 [10.6, 17.5]	0.65 [0.45, 0.93]
Absence of del17p	18.5 [15.8, 22.1]	13.9 [11.1, 16.4]	0.68 [0.54, 0.86]
Presence of t(4;14)	15.8 [8.4, 18.4]	5.5 [3.1, 10.3]	0.55 [0.32, 0.98]
Absence of t(4;14)	19.6 [17.0, 23.0]	14.9 [12.4, 17.1]	0.68 [0.55, 0.84]
ISS Stage			
I	22.2 [17.8, 31.3]	16.4 [14.5, 18.6]	0.61 [0.45, 0.83]
II	15.9 [9.5, 23.1]	12.9 [11.1, 18.5]	0.83 [0.60, 1.16]
III	14.0 [9.3, 17.3]	7.4 [5.6, 11.7]	0.70 [0.48, 1.04]
Prior therapies			
Lines of prior therapy = 1	18.5 [15.8, 20.7]	14.5 [10.9, 17.5]	0.71 [0.54, 0.94]
Lines of prior therapy = 2 or 3	18.5 [15.9, 23.9]	14.0 [11.1, 15.7]	0.65 [0.50, 0.85]
Prior thalidomide exposure	18.4 [14.1, 23.1]	12.3 [9.3, 14.9]	0.61 [0.46, 0.80]
No prior immunomodulatory exposure	18.9 [15.8, 22.2]	17.5 [13.0, 20.0]	0.78 [0.59, 1.04]
Prior bortezomib exposure	17.8 [15.8, 20.3]	12.3 [10.2, 14.9]	0.67 [0.53, 0.84]
No prior bortezomib exposure	21.4 [16.6, NE]	17.5 [13.1, 21.3]	0.70 [0.48, 1.00]
Response to therapy			
Relapsed	19.4 [16.6, 22.2]	16.6 [13.0, 18.9]	0.75 [0.59, 0.96]
Refractory	16.6 [14.5, 23.3]	10.4 [6.6, 13.3]	0.55 [0.40, 0.76]
Renal function			
Baseline CrCl < 60 mL/min	18.5 [14.8, 23.3]	11.7 [7.5, 17.4]	0.56 [0.39, 0.80]
Baseline CrCl ≥ 60 mL/min	18.5 [15.9, 22.2]	14.9 [12.1, 16.7]	0.72 [0.57, 0.90]

The 1-, 2- and 3-year rates of overall survival for Empliciti in combination with lenalidomide and dexamethasone treatment were 91%, 73%, and 60% respectively, compared with 83%, 69%, and 53% respectively, for lenalidomide and dexamethasone treatment (See Figure 2).

Figure 2: Overall survival



Number of subjects at risk		0	4	8	12	16	20	24	28	32	36	40	44	48
E-Ld		321	308	296	283	264	242	224	210	191	152	84	23	5
Ld		325	298	278	255	237	222	208	193	174	134	69	22	3

Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies in all subsets of the paediatric population in treatment of multiple myeloma (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

The pharmacokinetics (PK) of elotuzumab was studied in patients with multiple myeloma. Elotuzumab exhibits nonlinear PK with decrease in clearance with increase in dose from 0.5-20 mg/kg.

Absorption

Elotuzumab is dosed via intravenous route and therefore is immediately and completely bioavailable.

Distribution

The geometric mean volume of distribution of elotuzumab at 10 mg/kg (in combination with lenalidomide and dexamethasone) at steady state is 6.02 L (CV: 22.1%).

Biotransformation

The metabolic pathway of elotuzumab has not been characterized. As an IgG monoclonal antibody, elotuzumab is expected to be degraded into small peptides and amino acids via catabolic pathways.

Elimination

The geometric mean total clearance of elotuzumab at 10 mg/kg (in combination with lenalidomide and dexamethasone) at steady state is 0.194 L/day (CV: 62.9%). Upon discontinuation of elotuzumab in combination with lenalidomide and dexamethasone, concentrations of elotuzumab will decrease to approximately 3% (approximately 97% washout as estimated by 5 half-lives) of the population predicted steady-state maximal serum concentration by 3 months.

Special populations

Based on a population PK analysis using data from 375 patients, the clearance of elotuzumab increased with increasing body weight supporting a weight-based dose. The population PK analysis suggested that the following factors had no clinically important effect on the clearance of elotuzumab: age (37 to 88 years), gender, race, baseline LDH, albumin, renal impairment, and mild hepatic impairment.

Renal impairment

An open-label study evaluated the pharmacokinetics of elotuzumab in combination with lenalidomide and dexamethasone in patients with multiple myeloma with varying degrees of renal impairment (classified using the CrCl values). The effect of renal impairment on the pharmacokinetics of elotuzumab was evaluated in patients with normal renal function (CrCl > 90 mL/min; N = 8), severe renal impairment not requiring dialysis (CrCl <30 mL/min; N = 9), or end-stage renal disease requiring dialysis (CrCl < 30 mL/min; N = 9). No clinically important differences in the pharmacokinetics of elotuzumab were found between patients with severe renal impairment (with and without dialysis) and patients with normal renal function (see section 4.2).

Hepatic impairment

Empliciti is an IgG1 monoclonal antibody, which is principally cleared by catabolism. Thus, hepatic functional impairment is not likely to alter its clearance. The effect of hepatic impairment on the clearance of Empliciti was evaluated by population PK analyses in patients with mild hepatic impairment (total bilirubin [TB] ≤ the upper limit of normal [ULN] and AST > ULN or TB < 1 to 1.5 × ULN and any AST; N = 33). No clinically important differences in the clearance of Empliciti were found between patients with mild hepatic impairment and patients with normal hepatic function. Elotuzumab has not been studied in patients with moderate (TB > 1.5 to 3 × ULN and any AST) or severe hepatic impairment (TB > 3 × ULN and any AST) (see section 4.2).

5.3 Preclinical safety data

Elotuzumab only recognizes human SLAMF7 protein. Because elotuzumab does not recognize non-human forms of SLAMF7 protein, *in vivo* safety data from animal studies are irrelevant. In the same line, no carcinogenicity data are available for elotuzumab in animals, nor were fertility and embryo-foetal toxicity studies performed. Non-clinical safety information primarily consists of limited *in vitro* human cell/tissue studies where no safety findings were identified.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sucrose
Sodium citrate
Citric acid monohydrate
Polysorbate 80

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

Unopened vial

3 years.

After reconstitution and dilution

The reconstituted solution should be transferred from the vial into the infusion bag immediately.

Chemical and physical in use stability of the reconstituted and diluted solution has been demonstrated for 24 hours at 2°C - 8°C and protected from light.

From a microbiological point of view, the solution for infusion should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C protected from light. Do not freeze the reconstituted or diluted solution. The solution for infusion may be stored for a maximum of 8 hours of the total 24 hours at 20°C – 25°C and room light. This 8-hour period should be inclusive of the product administration period.

6.4 Special precautions for storage

Store in a refrigerator (2°C - 8°C).

Do not freeze.

Store in the original package in order to protect from light.

For storage conditions after reconstitution or dilution of the medicinal product, see section 6.3.

6.5 Nature and contents of container

20 ml Type I glass vial, closed with a grey butyl stopper and sealed with aluminium crimp seal with a polypropylene flip off button, containing either 300 mg or 400 mg elotuzumab. The flip-off seal button colour is ivory for the 300 mg presentation and blue for the 400 mg presentation.

Pack size of 1 vial.

6.6 Special precautions for disposal and other handling

Calculating the dose

Calculate the dose (mg) and determine the number of vials needed for the 10 mg/kg dose based on patient weight. More than one vial of Empliciti may be needed to give the total dose for the patient.

- The total elotuzumab dose in mg = the patient's weight in kg x 10.

Preparing the infusion

Aseptically reconstitute each Empliciti vial with a syringe of adequate size and an 18 gauge or smaller needle as shown in Table 7. A slight back pressure may be experienced during administration of the water for injections, which is considered normal.

Table 7: Reconstitution instructions

Strength	Amount of water for injections, required for reconstitution	Final volume of reconstituted Empliciti in the vial (including volume displaced by the solid cake)	Post-reconstitution concentration
300 mg vial	13.0 mL	13.6 mL	25 mg/mL
400 mg vial	17.0 mL	17.6 mL	25 mg/mL

- Hold the vial upright and swirl the solution by rotating the vial to dissolve the lyophilised cake. Then invert the vial a few times in order to dissolve any powder that may be present on top of the vial or the stopper. Avoid vigorous agitation, DO NOT SHAKE. The lyophilised powder should dissolve in less than 10 minutes.
- After the remaining solids are completely dissolved, allow the reconstituted solution to stand for 5 to 10 minutes. The reconstituted solution is colourless to slightly yellow, and clear to very

opalescent. Empliciti should be inspected visually for particulate matter and discolouration prior to administration. Discard the solution if any particulate matter or discolouration is observed.

- Once the reconstitution is completed, withdraw the necessary volume for the calculated dose from each vial, up to a maximum of 16 mL from 400 mg vial and 12 mL from 300 mg vial. Dilute the reconstituted solution with 230 mL of either sodium chloride 9 mg/mL (0.9%) solution for injection or 5% glucose injection, into an infusion bag made of polyvinyl chloride or polyolefin. The volume of sodium chloride 9 mg/mL (0.9%) solution for injection or 5% glucose injection should be adjusted so as not to exceed 5 mL/kg of patient weight at any given dose of Empliciti.

Administration

The entire Empliciti infusion should be administered with an infusion set and a sterile, non-pyrogenic, low-protein-binding filter (with a pore size of 0.2-1.2 µm) using an automated infusion pump.

Empliciti infusion is compatible with:

- PVC and polyolefin containers
- PVC infusion sets
- polyethersulfone and nylon in-line filters with pore sizes of 0.2 µm to 1.2 µm.

Empliciti should be initiated at an infusion rate of 0.5 mL/min. If well tolerated, the infusion rate may be increased stepwise as described in Table 2 (see section 4.2 Method of administration). The maximum infusion rate should not exceed 5 mL/min.

The Empliciti infusion solution should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C – 8°C protected from light. Do not freeze the reconstituted or diluted solution. The solution for infusion may be stored for a maximum of 8 hours of the total 24 hours at 20°C – 25°C and room light. This 8-hour period should be inclusive of the product administration period.

Disposal

Do not store any unused portion of the infusion solution for reuse. Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Bristol-Myers Squibb Pharma EEIG
Plaza 254
Blanchardstown Corporate Park 2
Dublin 15, D15 T867
Ireland

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/16/1088/001-002

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 11 May 2016

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

ANNEX II

- A. MANUFACTURERS OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURERS OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturers of the biological active substance

Bristol-Myers Squibb Company
6000 Thompson Road, East Syracuse
New York 13057
United States

Bristol-Myers Squibb Company
38 Jackson Road, Devens
MA 01434
United States

Name and address of the manufacturers responsible for batch release

Bristol-Myers Squibb S.r.l.
Loc. Fontana del Ceraso
03012 Anagni (FR)
Italy

Swords Laboratories t/a Bristol-Myers Squibb Cruiserath Biologics
Cruiserath Road, Mulhuddart
Dublin 15
Ireland

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

• **Periodic safety update reports**

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

The marketing authorisation holder shall submit the first periodic safety update report for this product within 6 months following authorisation.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

- **Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON

1. NAME OF THE MEDICINAL PRODUCT

Empliciti 300 mg powder for concentrate for solution for infusion
elotuzumab

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each vial contains 300 mg elotuzumab.
After reconstitution, each mL of concentrate contains 25 mg elotuzumab.

3. LIST OF EXCIPIENTS

Excipients: sucrose, sodium citrate, citric acid monohydrate, and polysorbate 80. See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Powder for concentrate for solution for infusion
1 vial

5. METHOD AND ROUTE(S) OF ADMINISTRATION

For single use only.
Read the package leaflet before use.
Intravenous use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store in a refrigerator.
Do not freeze.
Store in the original package in order to protect from light.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Bristol-Myers Squibb Pharma EEIG
Plaza 254
Blanchardstown Corporate Park 2
Dublin 15, D15 T867
Ireland

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/16/1088/001

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC:
SN:
NN:

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING

VIAL LABEL

1. NAME OF THE MEDICINAL PRODUCT

Empliciti 300 mg powder for concentrate
elotuzumab

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each vial contains 300 mg elotuzumab.
After reconstitution, each mL of concentrate contains 25 mg elotuzumab.

3. LIST OF EXCIPIENTS

Excipients: sucrose, sodium citrate, citric acid monohydrate, and polysorbate 80. See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

300 mg powder for concentrate

5. METHOD AND ROUTE(S) OF ADMINISTRATION

For single use only.
Read the package leaflet before use.
Intravenous use.
IV use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store in a refrigerator.
Do not freeze.
Store in the original package in order to protect from light.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Bristol-Myers Squibb Pharma EEIG
Plaza 254
Blanchardstown Corporate Park 2
Dublin 15, D15 T867
Ireland

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/16/1088/001

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON

1. NAME OF THE MEDICINAL PRODUCT

Empliciti 400 mg powder for concentrate for solution for infusion
elotuzumab

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each vial contains 400 mg elotuzumab.
After reconstitution, each mL of concentrate contains 25 mg elotuzumab.

3. LIST OF EXCIPIENTS

Excipients: sucrose, sodium citrate, citric acid monohydrate, and polysorbate 80. See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Powder for concentrate for solution for infusion
1 vial

5. METHOD AND ROUTE(S) OF ADMINISTRATION

For single use only.
Read the package leaflet before use.
Intravenous use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store in a refrigerator.
Do not freeze.
Store in the original package in order to protect from light.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Bristol-Myers Squibb Pharma EEIG
Plaza 254
Blanchardstown Corporate Park 2
Dublin 15, D15 T867
Ireland

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/16/1088/002

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC:
SN:
NN:

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING

VIAL LABEL

1. NAME OF THE MEDICINAL PRODUCT

Empliciti 400 mg powder for concentrate
elotuzumab

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each vial contains 400 mg elotuzumab.
After reconstitution, each mL of concentrate contains 25 mg elotuzumab.

3. LIST OF EXCIPIENTS

Excipients: sucrose, sodium citrate, citric acid monohydrate, and polysorbate 80. See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

400 mg powder for concentrate

5. METHOD AND ROUTE(S) OF ADMINISTRATION

For single use only.
Read the package leaflet before use.
Intravenous use.
IV use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store in a refrigerator.
Do not freeze.
Store in the original package in order to protect from light.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Bristol-Myers Squibb Pharma EEIG
Plaza 254
Blanchardstown Corporate Park 2
Dublin 15, D15 T867
Ireland

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/16/1088/002

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

B. PACKAGE LEAFLET

Package leaflet: Information for the user

Empliciti 300 mg powder for concentrate for solution for infusion Empliciti 400 mg powder for concentrate for solution for infusion

elotuzumab

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, or pharmacist or nurse.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Empliciti is and what it is used for
2. What you need to know before you use Empliciti
3. How to use Empliciti
4. Possible side effects
5. How to store Empliciti
6. Contents of the pack and other information

1. What Empliciti is and what it is used for

Empliciti contains the active substance elotuzumab, which is a monoclonal antibody, a type of protein designed to recognise and attach to a specific target substance in the body. Elotuzumab attaches to a target protein called SLAMF7. SLAMF7 is found in large amounts on the surface of multiple myeloma cells and on certain cells of your immune system (natural killer cells). When elotuzumab binds to SLAMF7 on the multiple myeloma or natural killer cells, it stimulates your immune system to attack and destroy the multiple myeloma cells.

Empliciti is used to treat multiple myeloma (a cancer of the bone marrow) in adults. It is used in combination with lenalidomide and dexamethasone. Multiple myeloma is a cancer of a type of white blood cell called plasma cells. These cells divide out of control and collect in the bone marrow. This results in damage to the bones and kidneys.

Empliciti is used when you have already had at least one multiple myeloma treatment before. One of these treatments may include bone marrow transplantation.

2. What you need to know before you use Empliciti

You should not be given Empliciti:

- if you are allergic to elotuzumab or any of the other ingredients of this medicine (listed in section 6 “Contents of the pack and other information”). Talk to your doctor if you are not sure.

Warnings and precautions

Infusion reaction

Tell your doctor or nurse straight away if you get any of the infusion reactions listed at the top of section 4. These side effects mostly occur during or after the infusion of the first dose. You will be monitored for signs of such effects during and after the infusion.

Depending on the seriousness of the infusion reactions, you may require additional treatment to prevent complications and reduce your symptoms, or your infusion of Empliciti may be interrupted. When the symptoms go away or improve, the infusion can be continued more slowly and speeded up gradually if the symptoms do not recur. Your doctor may decide not to continue Empliciti treatment if you have a strong infusion reaction.

Before each infusion of Empliciti, you will be given medicines to reduce infusion reaction (see section 3 “How to use Empliciti, Medicines given before each infusion”).

Before starting treatment with Empliciti, you must also read the package leaflet warnings and precautions of all medicines to be taken in combination with Empliciti for information related to these medicines. When lenalidomide is used, particular attention to pregnancy testing and prevention requirements is needed (see “Pregnancy and breast-feeding” in this section).

Children and adolescents

Empliciti is not recommended for use in children and adolescents aged under 18 years.

Other medicines and Empliciti

Tell your doctor if you are taking, have recently taken, or might take any other medicines.

Pregnancy and breast-feeding

For women taking Empliciti

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

You should not use Empliciti if you are pregnant, unless your doctor specifically recommends it. The effects of Empliciti in pregnant women or its possible harm to an unborn baby are unknown.

- You must use effective contraception while you are being treated with Empliciti, if there is any chance you could become pregnant.
- If you become pregnant while using Empliciti, tell your doctor.

When Empliciti is given in combination with lenalidomide, you must follow the pregnancy prevention programme for lenalidomide (see package leaflet for lenalidomide). **Lenalidomide is expected to be harmful for an unborn baby.**

It is not known, whether elotuzumab passes into breast milk or if there is any risk to the breast-fed infant. Elotuzumab will be given in combination with lenalidomide and breast-feeding should be stopped because of the use of lenalidomide.

For men taking Empliciti

You should use a condom while taking Empliciti and for 180 days after stopping treatment to ensure your partner does not become pregnant.

Driving and using machines

Empliciti is unlikely to affect your ability to drive or use machines. However, if you get an infusion reaction (fever, chills, high blood pressure see section 4 “Possible side effects”), do not drive, cycle or use machines until the reaction stops.

Empliciti contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per dose, i.e. it is essentially ‘sodium free’.

3. How to use Empliciti

How much Empliciti is given

The amount of Empliciti you will be given will be calculated based on your body weight. The recommended dose is 10 mg of elotuzumab per kilogram of your body weight.

How Empliciti is given

You will receive Empliciti under the supervision of an experienced healthcare professional. It will be given into a vein (intravenously) as a drip (infusion) over several hours.

Empliciti is taken in treatment cycles that are 28 days (4 weeks) long in combination with lenalidomide and dexamethasone.

- In cycles 1 and 2, Empliciti is administered once weekly on days 1, 8, 15, and 22.
- In cycles 3 and beyond, Empliciti is administered once every 2 weeks on days 1 and 15.

Your doctor will continue to treat you with Empliciti for as long as the disease improves or remains stable and side effects are tolerable.

You will also be given lenalidomide and dexamethasone together with Empliciti.

Medicines given before each infusion

You must receive the following medicines before each infusion of Empliciti to help reduce possible infusion reactions:

- medicine to reduce an allergic reaction (an anti-histamine)
- medicine to reduce inflammation (dexamethasone)
- medicine to reduce pain and fever (paracetamol)

If you miss a dose of Empliciti

Empliciti is used in combination with other medicines for multiple myeloma. If any medicine in the treatment is delayed, interrupted, or discontinued, your doctor will decide how your treatment should be continued.

If you are given too much Empliciti

As Empliciti will be given to you by a healthcare professional, it is unlikely you will be given too much. In the unlikely case of an overdose, your doctor will monitor you for side effects.

If you stop using Empliciti

Stopping your treatment with Empliciti may stop the effect of the medicine. Do not stop treatment unless you have discussed this with your doctor.

If you have any further questions on the use of this medicine, ask your doctor.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Your doctor will discuss these with you and will explain the risks and benefits of your treatment.

The following side effects have been reported in clinical trials with elotuzumab:

Infusion reactions

Empliciti has been associated with infusion reactions (see section 2 “Warnings and precautions”). **Tell your doctor or nurse straight away if you feel unwell during infusion.** Below is a list of typical symptoms associated with infusion reactions:

- Fever
- Chills
- High blood pressure

Other symptoms may occur as well. Your doctor may consider slowing the Empliciti infusion or interrupting it to manage these symptoms.

Other side effects

Very common (may affect more than 1 in 10 people)

- Fever
- Sore throat
- Pneumonia
- Painful skin rash with blisters (shingles, zona)
- Weight decrease
- Low white blood cell count
- Cough
- Common cold
- Headache
- Diarrhoea
- Feeling tired or weak

Common (may affect up to 1 in 10 people)

- Chest pain
- Blood clots in the veins (thrombosis)
- Night sweats
- Mood changes
- Decreased sensitivity, especially in the skin
- Allergic reactions (hypersensitivity)
- Pain in the mouth/throat region/sore throat

Uncommon (may affect up to 1 in 100 people)

- Sudden life-threatening allergic reaction (anaphylactic reaction)

Tell your doctor immediately if you get any of the side effects listed above. Do not try to treat your symptoms with other medicines.

Reporting of side effects

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Empliciti

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the vial label and carton after EXP. The expiry date refers to the last day of that month.

Store in a refrigerator (2°C - 8°C).

Do not freeze.

Store in the original package in order to protect from light.

After reconstitution, the reconstituted solution should be transferred from the vial to the infusion bag immediately.

After dilution, the infusion must be completed within 24 hours of preparation. The product should be used immediately. If not used immediately, the solution for infusion may be stored in the refrigerator (2 °C - 8 °C) for up to 24 hours.

Any unused medicine or waste material should be disposed of in accordance with local requirements.

6. Contents of the pack and other information

What Empliciti contains

- The active substance is elotuzumab.
Each vial contains either 300 mg or 400 mg of elotuzumab.
After reconstitution, each mL of concentrate contains 25 mg of elotuzumab.
- The other ingredients (excipients) are sucrose, sodium citrate (see section 2 “Empliciti contains sodium”), citric acid monohydrate, and polysorbate 80.

What Empliciti looks like and contents of the pack

Empliciti powder for concentrate for solution for infusion (powder for concentrate) is a white to off white whole or fragmented cake provided in a glass vial.

Empliciti is available in packs containing 1 vial.

Marketing Authorisation Holder

Bristol-Myers Squibb Pharma EEIG
Plaza 254
Blanchardstown Corporate Park 2
Dublin 15, D15 T867
Ireland

Manufacturer

Bristol-Myers Squibb S.r.l.
Loc. Fontana del Ceraso
03012 Anagni (FR)
Italy

Swords Laboratories t/a Bristol-Myers Squibb Cruiserath Biologics
Cruiserath Road, Mulhuddart
Dublin 15
Ireland

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

Belgique/België/Belgien

N.V. Bristol-Myers Squibb Belgium S.A.
Tél/Tel: + 32 2 352 76 11

Lietuva

Bristol-Myers Squibb Gyógyszerkereskedelmi Kft.
Tel: +370 52 369140

България

Bristol-Myers Squibb Kft.
Тел.: + 359 2 4942 480

Luxembourg/Luxemburg

N.V. Bristol-Myers Squibb Belgium S.A.
Tél/Tel: + 32 2 352 76 11

Česká republika

Bristol-Myers Squibb spol. s r.o.
Tel: + 420 221 016 111

Magyarország

Bristol-Myers Squibb Gyógyszerkereskedelmi Kft.
Tel.: + 36 1 301 9700

Danmark

Bristol-Myers Squibb
Tlf: + 45 45 93 05 06

Deutschland

Bristol-Myers Squibb GmbH & Co. KGaA
Tel: + 49 89 121 42-0

Eesti

Bristol-Myers Squibb Gyógyszerkereskedelmi Kft. Bristol-Myers Squibb Norway Ltd
Tel: +372 640 1030 Tlf: + 47 67 55 53 50

Ελλάδα

BRISTOL-MYERS SQUIBB A.E.
Τηλ: + 30 210 6074300

España

BRISTOL-MYERS SQUIBB, S.A.
Tel: + 34 91 456 53 00

France

Bristol-Myers Squibb SARL
Tél: + 33 (0)1 58 83 84 96

Hrvatska

Bristol-Myers Squibb spol. s r.o.
TEL: +385 1 2078 508

Ireland

Bristol-Myers Squibb Pharmaceuticals Ltd
Tel: + 353 (1 800) 749 749

Ísland

Vistor hf.
Sími: + 354 535 7000

Italia

BRISTOL-MYERS SQUIBB S.R.L.
Tel: + 39 06 50 39 61

Κύπρος

BRISTOL-MYERS SQUIBB A.E.
Τηλ: + 357 800 92666

Latvija

Bristol-Myers Squibb Gyógyszerkereskedelmi Kft. Bristol-Myers Squibb Pharmaceuticals Ltd
Tel: +371 67708347 Tel: + 44 (0800) 731 1736

Malta

BRISTOL-MYERS SQUIBB S.R.L.
Tel: + 39 06 50 39 61

Nederland

Bristol-Myers Squibb B.V.
Tel: + 31 (0)30 300 2222

Norge

Bristol-Myers Squibb Norway Ltd
Tlf: + 47 67 55 53 50

Österreich

Bristol-Myers Squibb GesmbH
Tel: + 43 1 60 14 30

Polska

Bristol-Myers Squibb Polska Sp. z o.o.
Tel.: + 48 22 2606400

Portugal

Bristol-Myers Squibb Farmacêutica Portuguesa,
S.A.
Tel: + 351 21 440 70 00

România

Bristol-Myers Squibb Gyógyszerkereskedelmi Kft.
Tel: + 40 (0)21 272 16 00

Slovenija

Bristol-Myers Squibb spol. s r.o.
Tel: +386 1 2355 100

Slovenská republika

Bristol-Myers Squibb spol. s r.o.
Tel: + 421 2 20833 600

Suomi/Finland

Oy Bristol-Myers Squibb (Finland) Ab
Puh/Tel: + 358 9 251 21 230

Sverige

Bristol-Myers Squibb AB
Tel: + 46 8 704 71 00

United Kingdom

Bristol-Myers Squibb Pharmaceuticals Ltd
Tel: + 44 (0800) 731 1736

This leaflet was last revised in

Detailed information on this medicine is available on the European Medicines Agency web site:
<http://www.ema.europa.eu>. There are also links to other websites about rare diseases and treatments.

This leaflet is available in all EU/EEA languages on the European Medicines Agency website.

The following information is intended for healthcare professionals only:

Preparation and administration of Empliciti

Calculating the dose

Calculate the dose (mg) and determine the number of vials needed for the 10 mg/kg dose based on patient weight. More than one vial of Empliciti may be needed to give the total dose for the patient.

- The total elotuzumab dose in mg = the patient’s weight in kg x 10.

Preparing the infusion

Aseptically reconstitute each Empliciti vial with a syringe of adequate size and an 18 gauge or smaller needle as shown in Table 1. A slight back pressure may be experienced during administration of the water for injections, which is considered normal.

Table 1: Reconstitution instructions

Strength	Amount of water for injections, required for reconstitution	Final volume of reconstituted Empliciti in the vial	Post-reconstitution concentration
300 mg vial	13.0 mL	13.6 mL	25 mg/mL
400 mg vial	17.0 mL	17.6 mL	25 mg/mL

- Hold the vial upright and swirl the solution by rotating the vial to dissolve the lyophilised cake. Then invert the vial a few times in order to dissolve any powder that may be present on top of the vial or the stopper. Avoid vigorous agitation, DO NOT SHAKE. The lyophilised powder should dissolve in less than 10 minutes.
- After the remaining solids are completely dissolved, allow the reconstituted solution to stand for 5 to 10 minutes. The reconstituted solution is colourless to slightly yellow and clear to very opalescent. Empliciti should be inspected visually for particulate matter and discolouration prior to administration. Discard the solution if any particulate matter or discolouration is observed.
- Once the reconstitution is completed, withdraw the necessary volume for the calculated dose from each vial, up to a maximum of 16 mL from 400 mg vial and 12 mL from 300 mg vial. Dilute the reconstituted solution with 230 mL of either sodium chloride 9 mg/mL (0.9%) solution for injection or 5% glucose injection into an infusion bag made of polyvinyl chloride or polyolefin. The volume of sodium chloride 9 mg/mL (0.9%) solution for injection or 5% glucose injection should be adjusted so as not to exceed 5 mL/kg of patient weight at any given dose of Empliciti.

Administration

The entire Empliciti infusion should be administered with an infusion set and a sterile, non-pyrogenic, low-protein-binding filter (with a pore size of 0.2-1.2 µm) using an automated infusion pump.

Empliciti infusion is compatible with:

- PVC and polyolefin containers
- PVC infusion sets
- polyethersulfone and nylon in-line filters with pore sizes of 0.2 µm to 1.2 µm.

Empliciti should be initiated at an infusion rate of 0.5 mL/min. If well tolerated, the infusion rate may be increased stepwise as described in Table 2. The maximum infusion rate should not exceed 5 mL/min.

Table 2: Infusion rate for Empliciti

Cycle 1, Dose 1		Cycle 1, Dose 2		Cycle 1, Dose 3 and 4 and all subsequent Cycles
Time interval	Rate	Time interval	Rate	Rate
0 - 30 min	0.5 mL/min	0 - 30 min	3 mL/min	5 mL/min*
30 - 60 min	1 mL/min	≥ 30 min	4 mL/min*	
≥ 60 min	2 mL/min*	-	-	

* Continue this rate until infusion is completed, approximately 1 hour based on patient weight.

The Empliciti infusion should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C – 8°C protected from light. Do not freeze the reconstituted or diluted solution. The solution for infusion may be stored for a maximum of 8 hours of the total 24 hours at 20°C – 25°C and room light. This 8-hour period should be inclusive of the product administration period.

Disposal

Do not store any unused portion of the infusion solution for reuse. Any unused medicine or waste material should be disposed of in accordance with local requirements.