

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Evalon suspension and solvent for oral spray for chickens.

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

### Evalon:

Each dose (0.007 ml) of undiluted vaccine contains:

### **Active substances:**

<i>Eimeria acervulina</i> , strain 003 .....	332 – 450*
<i>Eimeria brunetti</i> , strain 034.....	213 – 288*
<i>Eimeria maxima</i> , strain 013.....	196 – 265*
<i>Eimeria necatrix</i> , strain 033 .....	340 – 460*
<i>Eimeria tenella</i> , strain 004 .....	276 – 374*

\* Number of sporulated oocysts derived from precocious attenuated lines of coccidia, according to *in vitro* procedures of the manufacturer at the time of blending.

### HIPRAMUNE T (solvent):

### **Adjuvant:**

Montanide IMS

### **Excipients:**

Brilliant Blue (E133)

Red AC (E129)

Vanillin

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Suspension and solvent for oral spray.

Suspension: White turbid suspension.

Solvent: Dark brownish solution.

## 4. CLINICAL PARTICULARS

### 4.1 Target species

Chickens.

### 4.2 Indications for use, specifying the target species

For active immunisation of chicks from 1 day of age to reduce clinical signs (diarrhoea), intestinal lesions and oocysts output associated with coccidiosis caused by *Eimeria acervulina*, *Eimeria brunetti*, *Eimeria maxima*, *Eimeria necatrix* and *Eimeria tenella*.

Onset of immunity: 3 weeks post-vaccination.

Duration of immunity: 60 weeks post-vaccination in an environment that permits oocysts recycling.

### **4.3 Contraindications**

None.

### **4.4 Special warnings for each target species**

Vaccinate healthy animals only.

The vaccine will not protect species other than chickens against coccidiosis and is only effective against the *Eimeria* species indicated.

It is normal to find vaccinal oocysts in the intestine or litter of vaccinated flocks. Generally the number is higher the first weeks post-vaccination and lower once the flock has achieved a proper protection.

### **4.5 Special precautions for use**

#### Special precautions for use in animals

Chickens must be strictly floor-reared in the first 3 weeks after vaccination.

It is recommended that litter should be removed and facilities and material cleaned between production cycles to reduce field infections.

#### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash and disinfect hands and equipment after use.

### **4.6 Adverse reactions (frequency and seriousness)**

None.

### **4.7 Use during pregnancy, lactation or lay**

The safety of the veterinary medicinal product has not been established during lay. Do not use in birds in lay and within 2 weeks before the onset of the laying period.

### **4.8 Interaction with other medicinal products and other forms of interaction**

No information is available on the safety and efficacy of this immunological veterinary medicinal product when used with any other veterinary medicinal product. A decision to use this immunological veterinary medicinal product before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

No anticoccidial substances or other agents having anticoccidial activity via feed or water should be used for at least 3 weeks following vaccination of the chickens. The correct replication of the vaccine oocysts and consequently, the development of a solid immunity could be hindered. Additionally, the enhancement of protection produced by oocyst re-infections would also be limited.

#### 4.9 Amounts to be administered and administration route

Oral use.

The method of administration is by coarse spray.

##### Vaccination schedule:

One dose of vaccine (0.007 ml) from 1 day of age.

##### Administration route:

The method of administration is by coarse spray by using a suitable device (volume delivered: 28 ml/100 chicks, droplet size: 200–250 µm and working pressure: 2 to 3 bars). Before starting the preparation, make certain to have a clean container available with sufficient capacity for preparing the diluted vaccine suspension. Dilute the vaccine with the corresponding volumes:

Doses	Water	Vaccine	Solvent	Total
<b>1,000</b>	223 ml	7 ml	50 ml	280 ml
<b>5,000</b>	1,115 ml	35 ml	250 ml	1,400 ml
<b>10,000</b>	2,230 ml	70 ml	500 ml	2,800 ml

Shake the solvent vial. Dilute the content of the vial with clean room temperature water into an appropriate container.

Shake the vaccine vial and dilute the content into the previous solution.

Fill the reservoir of the spraying device with all the vaccine suspension prepared.

Maintain the diluted vaccine suspension in continuous homogenisation by using a magnetic stirrer while the vaccine is being administered via coarse spray to the chicks.

To improve the uniformity of the vaccination maintain the chicks inside the transportation box for at least 1 hour in order to let them ingest all the vaccine droplets.

After this time, place the chicks carefully in the litter and continue with regular management practices.

The device should be cleaned after each use. See the manufacturer's instructions to ensure proper disinfection and maintenance of the device.

#### 4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Severe overdose (10-fold) may result in a temporary reduction in daily live weight gain within the first week without any consequences on the final performances.

#### 4.11 Withdrawal period(s)

Zero days.

### 5. IMMUNOLOGICAL PROPERTIES

Pharmacotherapeutic group: Immunological for Aves, live parasitic vaccines for domestic fowl.  
ATC vet code: QI01AN01.

To stimulate active immunity against coccidiosis caused by *Eimeria acervulina*, *Eimeria brunetti*, *Eimeria maxima*, *Eimeria necatrix* and *Eimeria tenella*.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Evalon (vaccine)

Phosphate buffered solution (PBS):

- Potassium chloride
- Disodium phosphate dodecahydrate
- Potassium dihydrogen phosphate
- Sodium chloride

#### HIPRAMUNE T (solvent)

- Brilliant blue (E 133)
- Red AC (E 129)
- Vanillin
- Montanide IMS

### **6.2 Major incompatibilities**

Do not mix with any other veterinary medicinal product, except the solvent supplied for use with the veterinary medicinal product.

### **6.3 Shelf life**

#### Evalon (vaccine):

Shelf life of the veterinary medicinal product as packaged for sale: 10 months.

Shelf life after first opening the immediate packaging: use immediately.

Shelf life after dilution according to directions: 10 hours.

#### HIPRAMUNE T (solvent):

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

### **6.4. Special precautions for storage**

Store and transport refrigerated (2 °C–8 °C).

Do not freeze.

### **6.5 Nature and composition of immediate packaging**

#### Evalon (vaccine)

10 ml, 50 ml or 100 ml type I colourless glass vials containing 7 ml, 35 ml or 70 ml of suspension (1,000, 5,000 and 10,000 doses) closed with type I polymeric elastomer closures and aluminium caps.

#### HIPRAMUNE T (solvent)

Polypropylene (PP) vials containing 50 ml, 250 ml and 500 ml of solvent closed with type I polymeric elastomer closures and aluminium caps.

#### Pack sizes

Cardboard box with one vial of 1,000 doses (7 ml) and one vial with 50 ml of solvent.

Cardboard box with one vial of 5,000 doses (35 ml) and one vial with 250 ml of solvent.

Cardboard box with one vial of 10,000 doses (70 ml) and one vial with 500 ml of solvent.

Not all pack sizes may be marketed.

**6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

LABORATORIOS HIPRA, S.A.

Avda. la Selva, 135

17170 Amer (Girona)

SPAIN

Tel.: +34 972 430660

Fax: +34 972 430661

E-mail: [hipra@hipra.com](mailto:hipra@hipra.com)

**8. MARKETING AUTHORISATION NUMBER(S)**

EU/2/16/194/001–003

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 18/04/2016

Date of last renewal: {DD/MM/YYYY}

**10. DATE OF REVISION OF THE TEXT**

{DD/MM/YYYY}

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency (<http://www.ema.europa.eu/>).

**PROHIBITION OF SALE, SUPPLY AND/OR USE**

Not applicable.

## **ANNEX II**

- A. MANUFACTURERS OF THE BIOLOGICAL ACTIVE SUBSTANCES AND  
MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. STATEMENT OF THE MRLs**

**A. MANUFACTURERS OF THE BIOLOGICAL ACTIVE SUBSTANCES AND  
MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer of the biological active substances

Laboratorios Hipra, S.A.  
Avda. La Selva 135, Amer, 17170 Gerona, Spain

Laboratorios Hipra, S.A.  
Crt. de Susqueda, Amer, 17170 Gerona, Spain

Laboratorios Hipra, S.A.  
Carretera C-63, km 48.300,  
Polígono Industrial El Rieral  
Amer, 17170 Gerona, Spain

Name and address of the manufacturer responsible for batch release

Laboratorios Hipra, S.A.  
Avda. La Selva 135, Amer, 17170 Gerona, Spain

**B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Veterinary medicinal product subject to prescription.

**C. STATEMENT OF THE MRLs**

The active substances being principles of biological origin intended to produce active immunity are not within the scope of Regulation (EC) No 470/2009.

The excipients (including adjuvants) listed in section 6.1 of the SPC are either allowed substances for which table 1 of the annex to Commission Regulation (EU) No 37/2010 indicates that no MRLs are required or considered as not falling within the scope of Regulation (EC) No 470/2009 when used as in this veterinary medicinal product.



**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGE**

Cardboard box

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Evalon suspension and solvent for oral spray for chickens.

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each dose (0.007 ml) of undiluted vaccine contains:

*Eimeria acervulina*, strain 003 .....332–450  
*Eimeria brunetti*, strain 034.....213–288  
*Eimeria maxima*, strain 013.....196–265  
*Eimeria necatrix*, strain 033 .....340–460  
*Eimeria tenella*, strain 004 .....276–374

**3. PHARMACEUTICAL FORM**

Suspension and solvent for oral spray.

**4. PACKAGE SIZE**

One vial of 1,000 doses and one vial with 50 ml of HIPRAMUNE T (solvent).  
One vial of 5,000 doses and one vial with 250 ml of HIPRAMUNE T (solvent).  
One vial of 10,000 doses and one vial with 500 ml of HIPRAMUNE T (solvent).

**5. TARGET SPECIES**

Chickens.

**6. INDICATION(S)****7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use.  
Coarse spray.  
Read the package leaflet before use.

**8. WITHDRAWAL PERIOD(S)**

Withdrawal period: Zero days.

**9. SPECIAL WARNING(S), IF NECESSARY**

Read package leaflet before use.

**10. EXPIRY DATE**

EXP {month/year}

Once diluted use within 10 hours.

**11. SPECIAL STORAGE CONDITIONS**

Store and transport refrigerated. Do not freeze.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

Disposal: read package leaflet.

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

Keep out of the sight and reach of children.

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Laboratorios Hipra, S.A.  
Avda. la Selva, 135  
17170 Amer (Girona)  
SPAIN

**16. MARKETING AUTHORISATION NUMBER(S)**

EU/2/16/194/001

EU/2/16/194/002

EU/2/16/194/003

**17. MANUFACTURER’S BATCH NUMBER**

Batch {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS****Vaccine vial of 1,000 or 5,000 doses****1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Evalon suspension for oral spray for chickens.

**2. QUANTITY OF THE ACTIVE SUBSTANCE(S)**Each dose (0.007 ml) of undiluted vaccine contains:

*Eimeria acervulina*, strain 003 .....332–450  
*Eimeria brunetti*, strain 034.....213–288  
*Eimeria maxima*, strain 013.....196–265  
*Eimeria necatrix*, strain 033 .....340–460  
*Eimeria tenella*, strain 004 .....276–374

**3. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

1,000 doses

5,000 doses

**4. ROUTE(S) OF ADMINISTRATION**

Oral use.

To be mixed with the solvent. Read the package leaflet before use.

**5. WITHDRAWAL PERIOD(S)**

Withdrawal period: Zero days.

**6. BATCH NUMBER**

Batch {number}

**7. EXPIRY DATE**

EXP {month/year}

Once diluted use within 10 hours.

**8. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGE**

Vaccine vial of 10,000 doses

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Evalon suspension for oral spray for chickens.

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each dose (0.007 ml) of undiluted vaccine contains:

*Eimeria acervulina*, strain 003 .....332–450  
*Eimeria brunetti*, strain 034.....213–288  
*Eimeria maxima*, strain 013.....196–265  
*Eimeria necatrix*, strain 033 .....340–460  
*Eimeria tenella*, strain 004 .....276–374

**3. PHARMACEUTICAL FORM**

Suspension for oral spray.

**4. PACKAGE SIZE**

10,000 doses

**5. TARGET SPECIES**

Chickens.

**6. INDICATION(S)****7. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use.

Coarse spray.

To be mixed with the solvent. Read the package leaflet before use.

**8. WITHDRAWAL PERIOD(S)**

Withdrawal period: Zero days.

**9. SPECIAL WARNING(S), IF NECESSARY**

Read package leaflet before use.

**10. EXPIRY DATE**

EXP {month/year}

Once diluted use within 10 hours.

**11. SPECIAL STORAGE CONDITIONS**

Store and transport refrigerated. Do not freeze.

**12. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS, IF ANY**

**13. THE WORDS “FOR ANIMAL TREATMENT ONLY” AND CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE, IF APPLICABLE**

For animal treatment only. To be supplied only on veterinary prescription.

**14. THE WORDS “KEEP OUT OF THE SIGHT AND REACH OF CHILDREN”**

**15. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Laboratorios Hipra, S.A.  
Avda. la Selva, 135  
17170 Amer (Girona)  
SPAIN

**16. MARKETING AUTHORISATION NUMBER(S)**

EU/2/16/194/003

**17. MANUFACTURER’S BATCH NUMBER**

Batch {number}

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

Solvent vial of 50 ml, 250 ml or 500 ml

**1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

HIPRAMUNE T, solvent for Evalon

**2. CONTENTS BY WEIGHT, BY VOLUME OR BY NUMBER OF DOSES**

50 ml

250 ml

500 ml

**3. ROUTE(S) OF ADMINISTRATION****4. STORAGE CONDITIONS**

Store and transport refrigerated. Do not freeze.

**5. BATCH NUMBER**

Batch {number}

**6. EXPIRY DATE**

EXP {month/year}

**7. THE WORDS “FOR ANIMAL TREATMENT ONLY”**

For animal treatment only.



## **B. PACKAGE LEAFLET**

**PACKAGE LEAFLET:**  
**Evalon suspension and solvent for oral spray for chickens**

**1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder <and manufacturer responsible for batch release:

Laboratorios Hipra, S.A.  
Avda. la Selva 135  
17170 Amer (Girona)  
SPAIN

**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Evalon suspension and solvent for oral spray for chickens.

**3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)**

Evalon

Active substances:

Each dose (0.007 ml) of undiluted vaccine contains

<i>Eimeria acervulina</i> , strain 003 .....	332 – 450 *
<i>Eimeria brunetti</i> , strain 034.....	213 – 288 *
<i>Eimeria maxima</i> , strain 013.....	196 – 265 *
<i>Eimeria necatrix</i> , strain 033 .....	340 – 460 *
<i>Eimeria tenella</i> , strain 004 .....	276 – 374 *

\* Number of sporulated oocysts derived from precocious attenuated lines of coccidia, according to *in vitro* procedures of the manufacturer at the time of blending.

HIPRAMUNE T (solvent)

**Adjuvant:**

Montanide IMS

**Excipients:**

Brilliant Blue (E133)

Red AC (E129)

Vanillin

**4. INDICATION(S)**

For active immunisation of chicks from 1 day of age to reduce clinical signs (diarrhoea), intestinal lesions and oocysts output associated with coccidiosis caused by *Eimeria acervulina*, *Eimeria brunetti*, *Eimeria maxima*, *Eimeria necatrix* and *Eimeria tenella*.

Onset of immunity: 3 weeks post-vaccination.

Duration of immunity: 60 weeks post-vaccination in an environment that permits oocysts recycling.

## **5. CONTRAINDICATIONS**

None.

## **6. ADVERSE REACTIONS**

None.

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

## **7. TARGET SPECIES**

Chickens.

## **8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

One dose of vaccine (0.007 ml) from 1 day of age.

Oral use.

The method of administration is by coarse spray.

## **9. ADVICE ON CORRECT ADMINISTRATION**

The method of administration is by coarse spray by using a suitable device (volume delivered: 28 ml/100 chicks, droplet size: 200–250 µm and working pressure: 2 to 3 bars). Before starting the preparation, make certain to have a clean container available with sufficient capacity for preparing the diluted vaccine suspension. Dilute the vaccine with the corresponding volumes:

<b>DOSES</b>	<b>WATER</b>	<b>VACCINE</b>	<b>SOLVENT</b>	<b>TOTAL</b>
<b>1,000</b>	223 ml	7 ml	50 ml	280 ml
<b>5,000</b>	1115 ml	35 ml	250 ml	1400 ml
<b>10,000</b>	2230 ml	70 ml	500 ml	2800 ml

Shake the solvent vial. Dilute the contents of the vial with clean room temperature water into an appropriate container.

Shake the vaccine vial and dilute the content into the previous solution.

Fill the reservoir of the spraying device with all the vaccine suspension prepared.

Maintain the diluted vaccine suspension in continuous homogenisation by using a magnetic stirrer while the vaccine is being administered via coarse spray to the chicks.

To improve the uniformity of the vaccination maintain the chicks inside the transportation box for at least 1 hour in order to let them ingest all the vaccine droplets.

After this time, place the chicks carefully in the litter and continue with regular management practices.

The device should be cleaned after each use. See the manufacturer's instructions to ensure proper disinfection and maintenance of the device.

#### **10. WITHDRAWAL PERIOD(S)**

Zero days.

#### **11. SPECIAL STORAGE PRECAUTIONS**

Keep out of the sight and reach of children.

Store and transport refrigerated (2 °C - 8 °C). Do not freeze.

Shelf life of Evalon as packaged for sale: 10 months.

Shelf life after first opening the immediate packaging: use immediately.

Shelf life after dilution according to directions: 10 hours.

Shelf life of HIPRAMUNE T as packaged for sale: 2 years.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and the label.

#### **12. SPECIAL WARNING(S)**

Special warnings for each target species:

Vaccinate healthy animals only.

The vaccine will not protect species other than chickens against coccidiosis and is only effective against the *Eimeria* species indicated.

It is normal to find vaccinal oocysts in the intestine or litter of vaccinated flocks. Generally the number is higher the first weeks post-vaccination and lower once the flock has achieved a proper protection.

Special precautions for use in animals:

Chickens must be strictly floor-reared in the first 3 weeks after vaccination.

It is recommended that litter should be removed and facilities and material cleaned between production cycles to reduce field infections.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Wash and disinfect hands and equipment after use.

Lay:

The safety of the veterinary medicinal product has not been established during lay. Do not use in birds in lay and within 2 weeks before the onset of the laying period

Interaction with other medicinal products and other forms of interaction:

No information is available on the safety and efficacy of this immunological veterinary medicinal product when used with any other veterinary medicinal product. A decision to use this immunological veterinary medicinal product before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

No anticoccidial substances or other agents having anticoccidial activity via feed or water should be used for at least 3 weeks following vaccination of the chickens. The correct replication of the vaccine oocysts and consequently, the development of a solid immunity could be hindered. Additionally, the enhancement of protection produced by oocyst re-infections would also be limited.

Overdose (symptoms, emergency procedures, antidotes):

Severe overdose (10-fold) may result in a temporary reduction in daily live weight gain within the first week without any consequences on the final performances.

Incompatibilities:

Do not mix with any other veterinary medicinal product, except the solvent supplied for use with the veterinary medicinal product.

**13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY**

Medicines should not be disposed of via wastewater or household waste.

Ask your veterinary surgeon or pharmacist how to dispose of medicines no longer required. These measures should help to protect the environment.

**14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED**

Detailed information on this veterinary medicinal product is available on the website of the European Medicines Agency (<http://www.ema.europa.eu/>).

**15. OTHER INFORMATION>**

Pack sizes:

Cardboard box with one vial of 1,000 doses (7 ml) and one vial with 50 ml of solvent.

Cardboard box with one vial of 5,000 doses (35 ml) and one vial with 250 ml of solvent.

Cardboard box with one vial of 10,000 doses (70 ml) and one vial with 500 ml of solvent.

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorization holder.

<b>België/Belgique/Belgien</b> HIPRA BENELUX NV Tél/Tel: +32 09 2964464	<b>Lietuva</b> LABORATORIOS HIPRA, S.A. Tel: +34 972 43 06 60
<b>Република България</b> LABORATORIOS HIPRA, S.A. Тел: +34 972 43 06 60	<b>Luxembourg/Luxemburg</b> HIPRA BENELUX NV Tél/Tel: +32 09 2964464
<b>Česká republika</b> HIPRA SLOVENSKO, s.r.o. Tel: +421 02 32 335 223	<b>Magyarország</b> LABORATORIOS HIPRA, S.A. Tel: +34 972 43 06 60

<b>Danmark</b> LABORATORIOS HIPRA, S.A. Tel: +34 972 43 06 60	<b>Malta</b> LABORATORIOS HIPRA, S.A. Tel: +34 972 43 06 60
<b>Deutschland</b> HIPRA DEUTSCHLAND GmbH Tel: +49 211 698236 – 0	<b>Nederland</b> HIPRA BENELUX NV Tel: +32 09 2964464
<b>Eesti</b> LABORATORIOS HIPRA, S.A. Tel: +34 972 43 06 60	<b>Norge</b> LABORATORIOS HIPRA, S.A. Tlf: +34 972 43 06 60
<b>Ελλάδα</b> HIPRA ΕΛΛΑΣ Α.Ε. Τηλ: +30 210 4978660	<b>Österreich</b> HIPRA DEUTSCHLAND GmbH Tel: +49 211 698236 – 0
<b>España</b> LABORATORIOS HIPRA, S.A. Tel: +34 972 43 06 60	<b>Polska</b> HIPRA POLSKA Sp.z.o.o. Tel: +48 22 642 33 06
<b>France</b> HIPRA FRANCE Tél: +33 02 51 80 77 91	<b>Portugal</b> ARBUSET, Produtos Farmacêuticos e Sanitários De Uso Animal, Lda Tel:+351 219 663 450
<b>Hrvatska</b> LABORATORIOS HIPRA, S.A. Tel: +34 972 43 06 60	<b>România</b> LABORATORIOS HIPRA, S.A. Tel: +34 972 43 06 60
<b>Ireland</b> LABORATORIOS HIPRA, S.A. Puh/Tel: +34 972 43 06 60	<b>Slovenija</b> LABORATORIOS HIPRA, S.A. Tel: +34 972 43 06 60
<b>Ísland</b> LABORATORIOS HIPRA, S.A. Sími: +34 972 43 06 60	<b>Slovenská republika</b> HIPRA SLOVENSKO, s.r.o. Tel: +421 02 32 335 223
<b>Italia</b> Hipra Italia S.r.l. Tel: +39 030 7241821	<b>Suomi/Finland</b> LABORATORIOS HIPRA, S.A. Puh/Tel: +34 972 43 06 60
<b>Κύπρος</b> LABORATORIOS HIPRA, S.A. Τηλ: +34 972 43 06 60	<b>Sverige</b> LABORATORIOS HIPRA, S.A. Tel. +34 972 43 06 60
<b>Latvija</b> LABORATORIOS HIPRA, S.A. Tel. +34 972 43 06 60	<b>United Kingdom (Northern Ireland)</b> LABORATORIOS HIPRA, S.A. Puh/Tel: +34 972 43 06 60