

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Fingolimod Mylan 0.5 mg hard capsules

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each capsule contains 0.5 mg fingolimod (as hydrochloride).

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Hard capsule (capsule)

Brown-orange opaque cap and white opaque body, printed with 'MYLAN' over 'FD 0.5' in black ink on both the cap and body. Dimension: approximately 16 mm in length.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Fingolimod Mylan is indicated as single disease modifying therapy in highly active relapsing remitting multiple sclerosis for the following groups of adult patients and paediatric patients aged 10 years and older:

- Patients with highly active disease despite a full and adequate course of treatment with at least one disease modifying therapy (for exceptions and information about washout periods see sections 4.4 and 5.1).

or

- Patients with rapidly evolving severe relapsing remitting multiple sclerosis defined by 2 or more disabling relapses in one year and with 1 or more Gadolinium enhancing lesions on brain magnetic resonance imaging (MRI) or a significant increase in T2 lesion load as compared to a previous recent MRI.

4.2 Posology and method of administration

The treatment should be initiated and supervised by a physician experienced in multiple sclerosis.

Posology

In adults, the recommended dose of fingolimod is one 0.5 mg capsule taken orally once daily.

In paediatric patients (10 years of age and above), the recommended dose is dependent on body weight:

- Paediatric patients with body weight $\leq$ 40 kg: one 0.25 mg capsule taken orally once daily.
- Paediatric patients with body weight $>$ 40 kg: one 0.5 mg capsule taken orally once daily.

Paediatric patients who start on 0.25 mg capsules and subsequently reach a stable body weight above 40 kg should be switched to 0.5 mg capsules.

When switching from a 0.25 mg to a 0.5 mg daily dose, it is recommended to repeat the same first dose monitoring as for treatment initiation.

Fingolimod Mylan is not available in 0.25 mg strength. For this dosage, other medicinal products containing fingolimod, available on the market, should be used.

The same first dose monitoring as for treatment initiation is recommended when treatment is interrupted for:

- One day or more during the first 2 weeks of treatment.
- More than 7 days during weeks 3 and 4 of treatment.
- More than 2 weeks after one month of treatment.

If the treatment interruption is of shorter duration than the above, the treatment should be continued with the next dose as planned (see section 4.4).

Special populations

Elderly

Fingolimod Mylan should be used with caution in patients aged 65 years and over due to insufficient data on safety and efficacy (see section 5.2).

Renal impairment

Fingolimod was not studied in patients with renal impairment in the multiple sclerosis pivotal studies. Based on clinical pharmacology studies, no dose adjustments are needed in patients with mild to severe renal impairment.

Hepatic impairment

Fingolimod Mylan must not be used in patients with severe hepatic impairment (Child-Pugh class C) (see section 4.3). Although no dose adjustments are needed in patients with mild or moderate hepatic impairment, caution should be exercised when initiating treatment in these patients (see sections 4.4 and 5.2).

Paediatric population

There are very limited data available in children between 10–12 years old (see sections 4.4, 4.8 and 5.1).

The safety and efficacy of fingolimod in children aged below 10 years have not yet been established. No data are available.

Method of administration

This medicinal product is for oral use.

Fingolimod Mylan can be taken with or without food (see section 5.2).

The capsules should always be swallowed intact, without opening them.

4.3 Contraindications

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Immunodeficiency syndrome.
- Patients with increased risk for opportunistic infections, including immunocompromised patients (including those currently receiving immunosuppressive therapies or those immunocompromised by prior therapies).
- Suspected or confirmed progressive multifocal leukoencephalopathy (PML) (see section 4.4).
- Severe active infections, active chronic infections (hepatitis, tuberculosis).
- Active malignancies.
- Severe liver impairment (Child-Pugh class C).

- Patients who in the previous 6 months had myocardial infarction (MI), unstable angina pectoris, stroke/transient ischaemic attack (TIA), decompensated heart failure (requiring inpatient treatment), or New York Heart Association (NYHA) class III/IV heart failure (see section 4.4).
- Patients with severe cardiac arrhythmias requiring anti-arrhythmic treatment with class Ia or class III anti-arrhythmic medicinal products (see section 4.4).
- Patients with second-degree Mobitz type II atrioventricular (AV) block or third-degree AV block, or sick-sinus syndrome, if they do not wear a pacemaker (see section 4.4).
- Patients with a baseline QTc interval ≥ 500 msec (see section 4.4).
- During pregnancy and in women of childbearing potential not using effective contraception (see sections 4.4 and 4.6).

4.4 Special warnings and precautions for use

Bradyarrhythmia

Initiation of treatment results in a transient decrease in heart rate and may also be associated with atrioventricular conduction delays, including the occurrence of isolated reports of transient, spontaneously resolving complete AV block (see sections 4.8 and 5.1).

After the first dose, the decline in heart rate starts within one hour and is maximal within 6 hours. This post-dose effect persists over the following days, although usually to a milder extent and usually abates over the next weeks. With continued administration, the average heart rate returns towards baseline within one month. However individual patients may not return to baseline heart rate by the end of the first month. Conduction abnormalities were typically transient and asymptomatic. They usually did not require treatment and resolved within the first 24 hours on treatment. If necessary, the decrease in heart rate induced by fingolimod can be reversed by parenteral doses of atropine or isoprenaline.

All patients should have an electrocardiogram (ECG) and blood pressure measurement performed prior to and 6 hours after the first dose of Fingolimod Mylan. All patients should be monitored for a period of 6 hours for signs and symptoms of bradycardia with hourly heart rate and blood pressure measurement. Continuous (real time) ECG monitoring during this 6-hour period is recommended.

The same precautions as for the first dose are recommended when patients are switched from the 0.25 mg to the 0.5 mg daily dose.

Should post-dose bradyarrhythmia-related symptoms occur, appropriate clinical management should be initiated and monitoring should be continued until the symptoms have resolved. Should a patient require pharmacological intervention during the first-dose monitoring, overnight monitoring in a medical facility should be instituted and the first-dose monitoring should be repeated after the second dose of Fingolimod Mylan.

If the heart rate at 6 hours is the lowest since the first dose was administered (suggesting that the maximum pharmacodynamic effect on the heart may not yet be manifest), monitoring should be extended by at least 2 hours and until heart rate increases again. Additionally, if after 6 hours, the heart rate is < 45 bpm in adults, < 55 bpm in paediatric patients aged 12 years and above, or < 60 bpm in paediatric patients aged 10 to below 12 years, or the ECG shows new onset second degree or higher grade AV block or a QTc interval ≥ 500 msec, extended monitoring (at least overnight monitoring), should be performed and until the findings have resolved. The occurrence at any time of third degree AV block should also lead to extended monitoring (at least overnight monitoring).

The effects on heart rate and atrioventricular conduction may recur on re-introduction of fingolimod treatment depending on duration of the interruption and time since start of treatment. The same first dose monitoring as for treatment initiation is recommended when treatment is interrupted (see section 4.2).

Very rare cases of T-wave inversion have been reported in adult patients treated with fingolimod. In

case of T-wave inversion, the prescriber should ensure that there are no associated myocardial ischaemia signs or symptoms. If myocardial ischaemia is suspected, it is recommended to seek advice from a cardiologist.

Due to the risk of serious rhythm disturbances or significant bradycardia, Fingolimod Mylan should not be used in patients with sino-atrial heart block, a history of symptomatic bradycardia, recurrent syncope or cardiac arrest, or in patients with significant QT prolongation ($QT_c > 470$ msec [adult female], $QT_c > 460$ msec [paediatric female] or > 450 msec [adult and paediatric male]), uncontrolled hypertension or severe sleep apnoea (see also section 4.3). In such patients, treatment with this medicinal product should be considered only if the anticipated benefits outweigh the potential risks and advice from a cardiologist sought prior to initiation of treatment in order to determine the most appropriate monitoring. At least overnight extended monitoring is recommended for treatment initiation (see also section 4.5).

Fingolimod has not been studied in patients with arrhythmias requiring treatment with class Ia (e.g. quinidine, disopyramide) or class III (e.g. amiodarone, sotalol) antiarrhythmic medicinal products. Class Ia and class III antiarrhythmic medicinal products have been associated with cases of torsades de pointes in patients with bradycardia (see section 4.3).

Experience with fingolimod is limited in patients receiving concurrent therapy with beta blockers, heart-rate-lowering calcium channel blockers (such as verapamil or diltiazem), or other substances which may decrease heart rate (e.g. ivabradine, digoxin, anticholinesteratic agents or pilocarpine). Since the initiation of fingolimod treatment is also associated with slowing of the heart rate (see also section 4.8, Bradyarrhythmia), concomitant use of these substances during treatment initiation may be associated with severe bradycardia and heart block. Because of the potential additive effect on heart rate treatment with Fingolimod Mylan should not be initiated in patients who are concurrently treated with these substances (see also section 4.5). In such patients, treatment with fingolimod should be considered only if the anticipated benefits outweigh the potential risks. If treatment is considered, advice from a cardiologist should be sought regarding the switch to non heart-rate lowering medicinal products prior to initiation of treatment. If the heart-rate-lowering treatment cannot be stopped, cardiologist's advice should be sought to determine appropriate first dose monitoring, at least overnight extended monitoring is recommended (see also section 4.5).

QT interval

In a thorough QT interval study of doses of 1.25 or 2.5 mg fingolimod at steady-state, when a negative chronotropic effect of fingolimod was still present, treatment resulted in a prolongation of QT_{cI} , with the upper limit of the 90% CI ≤ 13.0 ms. There is no dose- or exposure-response relationship of fingolimod and QT_{cI} prolongation. There is no consistent signal of increased incidence of QT_{cI} outliers, either absolute or change from baseline, associated with treatment.

The clinical relevance of this finding is unknown. In the multiple sclerosis studies, clinically relevant effects on prolongation of the QT_c -interval have not been observed but patients at risk for QT prolongation were not included in clinical studies.

Medicinal products that may prolong QT_c interval are best avoided in patients with relevant risk factors, for example, hypokalaemia or congenital QT prolongation.

Immunosuppressive effects

Fingolimod has an immunosuppressive effect that predisposes patients to an infection risk, including opportunistic infections that can be fatal and increases the risk of developing lymphomas and other malignancies, particularly those of the skin. Physicians should carefully monitor patients, especially those with concurrent conditions or known factors, such as previous immunosuppressive therapy. If this risk is suspected, discontinuation of treatment should be considered by the physician on a case-by-case basis (see also section 4.4 'Infections' and 'Cutaneous malignancies' and section 4.8 'Lymphomas').

Infections

A core pharmacodynamic effect of fingolimod is a dose-dependent reduction of the peripheral lymphocyte count to 20-30% of baseline values. This is due to the reversible sequestration of lymphocytes in lymphoid tissues (see section 5.1).

Before initiating treatment with Fingolimod Mylan, a recent complete blood count (CBC) (i.e. within 6 months or after discontinuation of prior therapy) should be available. Assessments of CBC are also recommended periodically during treatment, at month 3 and at least yearly thereafter, and in case of signs of infection. Absolute lymphocyte count $< 0.2 \times 10^9/l$, if confirmed, should lead to treatment interruption until recovery, because in clinical studies, fingolimod treatment was interrupted in patients with absolute lymphocyte count $< 0.2 \times 10^9/l$.

Initiation of treatment with Fingolimod Mylan should be delayed in patients with severe active infection until resolution.

The immune system effects of Fingolimod Mylan may increase the risk of infections, including opportunistic infections (see section 4.8). Effective diagnostic and therapeutic strategies should be employed in patients with symptoms of infection while on therapy. When evaluating a patient with a suspected infection that could be serious, referral to a physician experienced in treating infections should be considered. During treatment, patients should be instructed to report promptly symptoms of infection to their physician.

Suspension of Fingolimod Mylan should be considered if a patient develops a serious infection and consideration of benefit-risk should be undertaken prior to re-initiation of therapy.

Elimination of fingolimod following discontinuation of therapy may take up to two months and vigilance for infection should therefore be continued throughout this period. Patients should be instructed to report symptoms of infection up to 2 months after discontinuation of treatment.

Herpes viral infection

Serious, life-threatening, and sometimes fatal cases of encephalitis, meningitis or meningoencephalitis caused by herpes simplex and varicella zoster viruses have occurred with fingolimod at any time during treatment. If herpes encephalitis, meningitis or meningoencephalitis occur, treatment should be discontinued and appropriate treatment for the respective infection should be administered.

Patients need to be assessed for their immunity to varicella (chickenpox) prior to Fingolimod Mylan treatment. It is recommended that patients without a health care professional confirmed history of chickenpox or documentation of a full course of vaccination with varicella vaccine undergo antibody testing to varicella zoster virus (VZV) before initiating fingolimod therapy. A full course of vaccination for antibody-negative patients with varicella vaccine is recommended prior to commencing treatment with this medicinal product (see section 4.8). Initiation of treatment with fingolimod should be postponed for one month to allow full effect of vaccination to occur.

Cryptococcal meningitis

Cases of cryptococcal meningitis (a fungal infection), sometimes fatal, have been reported in the post-marketing setting after approximately 2-3 years of treatment, although an exact relationship with the duration of treatment is unknown (see section 4.8). Patients with symptoms and signs consistent with cryptococcal meningitis (e.g. headache accompanied by mental changes such as confusion, hallucinations and/or personality changes) should undergo prompt diagnostic evaluation. If cryptococcal meningitis is diagnosed, fingolimod should be suspended and appropriate treatment should be initiated. A multidisciplinary consultation (i.e. infectious disease specialist) should be undertaken if re-initiation of fingolimod is warranted.

Progressive multifocal leukoencephalopathy (PML)

PML has been reported under fingolimod treatment since marketing authorisation (see section 4.8). It

is an opportunistic infection caused by John Cunningham virus (JCV), which may be fatal or result in severe disability. The majority of PML cases have occurred after 2 or more years of fingolimod treatment. In addition to duration of fingolimod exposure, other potential risk factors for PML include prior therapy with immunosuppressants or immunomodulators, and/or severe lymphopenia ($<0.5 \times 10^9/l$). Patients at increased risk should be closely monitored for any signs or symptoms of PML. PML can only occur in the presence of a JCV infection. If JCV testing is undertaken, it should be considered that the influence of lymphopenia on the accuracy of anti-JCV antibody testing has not been studied in fingolimod-treated patients. A negative anti-JCV antibody test does not preclude the possibility of subsequent JCV infection. Before initiating treatment with fingolimod, a baseline MRI should be available (usually within 3 months) as a reference. During routine MRI (in accordance with national and local recommendations), physicians should pay attention to PML suggestive lesions. MRI findings may be apparent before clinical signs or symptoms. Annual MRIs may be considered as part of increased vigilance especially in patients at increased risk of PML. Cases of asymptomatic PML based on MRI findings and positive JCV DNA in the cerebrospinal fluid have been reported in patients treated with fingolimod. If PML is suspected, MRI should be performed immediately for diagnostic purposes and treatment should be suspended until PML has been excluded. If PML is confirmed, treatment with fingolimod must be permanently discontinued (see also section 4.3).

Immune reconstitution inflammatory syndrome (IRIS) has been reported in patients treated with sphingosine 1-phosphate (S1P) receptor modulators, including fingolimod, who developed PML and subsequently discontinued treatment. IRIS presents as a clinical decline in the patient's condition that may be rapid, can lead to serious neurological complications or death, and is often associated with characteristic changes on MRI. The time to onset of IRIS in patients with PML was usually from weeks to months after S1P receptor modulator discontinuation. Monitoring for development of IRIS and appropriate treatment of the associated inflammation should be undertaken.

Human papilloma virus (HPV) infection

HPV infection, including papilloma, dysplasia, warts and HPV-related cancer, has been reported under treatment with fingolimod in the post-marketing setting (see section 4.8). Due to the immunosuppressive properties of fingolimod, vaccination against HPV should be considered prior to treatment initiation taking into account vaccination recommendations. Cancer screening, including Pap test, is recommended as per standard of care.

Macular oedema

Macular oedema with or without visual symptoms has been reported in 0.5% of patients treated with fingolimod 0.5 mg, occurring predominantly in the first 3-4 months of therapy (see section 4.8). An ophthalmological evaluation is therefore recommended at 3-4 months after treatment initiation. If patients report visual disturbances at any time while on therapy, evaluation of the fundus, including the macula, should be carried out.

Patients with history of uveitis and patients with diabetes mellitus are at increased risk of macular oedema (see section 4.8). Fingolimod has not been studied in multiple sclerosis patients with concomitant diabetes mellitus. It is recommended that multiple sclerosis patients with diabetes mellitus or a history of uveitis undergo an ophthalmological evaluation prior to initiating therapy and have follow-up evaluations while receiving therapy.

Continuation of treatment in patients with macular oedema has not been evaluated. It is recommended that Fingolimod Mylan be discontinued if a patient develops macular oedema. A decision on whether or not therapy should be re-initiated after resolution of macular oedema needs to take into account the potential benefits and risks for the individual patient.

Liver injury

Increased hepatic enzymes, in particular alanine aminotransaminase (ALT) but also gamma glutamyltransferase (GGT) and aspartate transaminase (AST) have been reported in multiple sclerosis patients treated with fingolimod. Some cases of acute liver failure requiring liver transplant and

clinically significant liver injury have also been reported. Signs of liver injury, including markedly elevated serum hepatic enzymes and elevated total bilirubin, have occurred as early as ten days after the first dose and have also been reported after prolonged use. In clinical trials, elevations 3-fold the upper limit of normal (ULN) or greater in ALT occurred in 8.0% of adult patients treated with fingolimod 0.5 mg compared to 1.9% of placebo patients. Elevations 5-fold the ULN occurred in 1.8% of patients on fingolimod and 0.9% of patients on placebo. In clinical trials, treatment was discontinued if the elevation exceeded 5 times the ULN. Recurrence of liver transaminase elevations occurred with rechallenge in some patients, supporting a relationship to fingolimod. In clinical studies, transaminase elevations occurred at any time during treatment although the majority occurred within the first 12 months. Serum transaminase levels returned to normal within approximately 2 months after discontinuation of treatment.

Fingolimod has not been studied in patients with severe pre-existing hepatic injury (Child-Pugh class C) and should not be used in these patients (see section 4.3).

Due to the immunosuppressive properties of fingolimod, initiation of treatment should be delayed in patients with active viral hepatitis until resolution.

Recent (i.e. within last 6 months) transaminase and bilirubin levels should be available before initiation of treatment. In the absence of clinical symptoms, liver transaminases and serum bilirubin should be monitored at months 1, 3, 6, 9 and 12 on therapy and periodically thereafter until 2 months after fingolimod discontinuation. In the absence of clinical symptoms, if liver transaminases are greater than 3 but less than 5 times the ULN without increase in serum bilirubin, more frequent monitoring including serum bilirubin and alkaline phosphatase (ALP) measurement should be instituted to determine if further increases occur and in order to discern if an alternative aetiology of hepatic dysfunction is present. If liver transaminases are at least 5 times the ULN or at least 3 times the ULN associated with any increase in serum bilirubin, treatment should be discontinued. Hepatic monitoring should be continued. If serum levels return to normal (including if an alternative cause of the hepatic dysfunction is discovered), fingolimod may be restarted based on a careful benefit-risk assessment of the patient.

Patients who develop symptoms suggestive of hepatic dysfunction, such as unexplained nausea, vomiting, abdominal pain, fatigue, anorexia, or jaundice and/or dark urine, should have liver enzymes and bilirubin checked promptly and treatment should be discontinued if significant liver injury is confirmed.

Treatment should not be resumed unless a plausible alternative aetiology for the signs and symptoms of liver injury can be established.

Although there are no data to establish that patients with pre-existing liver disease are at increased risk of developing elevated liver function tests when taking fingolimod, caution should be exercised in patients with a history of significant liver disease.

Blood pressure effects

Patients with hypertension uncontrolled by medication were excluded from participation in premarketing clinical trials and special care is indicated if patients with uncontrolled hypertension are treated with Fingolimod Mylan.

In multiple sclerosis (MS) clinical trials, patients treated with fingolimod 0.5 mg had an average increase of approximately 3 mmHg in systolic pressure and approximately 1 mmHg in diastolic pressure, first detected approximately one month after treatment initiation and persisting with continued treatment. In the two-year placebo-controlled study, hypertension was reported as an adverse event in 6.5% of patients on fingolimod 0.5 mg and in 3.3% of patients on placebo. Therefore, blood pressure should be regularly monitored during treatment.

Respiratory effects

Minor dose-dependent reductions in values for forced expiratory volume (FEV₁) and diffusion capacity for carbon monoxide (DLCO) were observed with fingolimod treatment starting at month-1 and remaining stable thereafter. It should be used with caution in patients with severe respiratory disease, pulmonary fibrosis and chronic obstructive pulmonary disease (see section 4.8).

Posterior reversible encephalopathy syndrome (PRES)

Rare cases of PRES have been reported at the 0.5 mg dose in clinical trials and in the post-marketing setting (see section 4.8). Symptoms reported included sudden onset of severe headache, nausea, vomiting, altered mental status, visual disturbances and seizure. Symptoms of PRES are usually reversible but may evolve into ischaemic stroke or cerebral haemorrhage. Delay in diagnosis and treatment may lead to permanent neurological sequelae. If PRES is suspected, Fingolimod Mylan should be discontinued.

Prior treatment with immunosuppressive or immunomodulatory therapies

There have been no studies performed to evaluate the efficacy and safety of fingolimod when switching patients from teriflunomide, dimethyl fumarate or alemtuzumab treatment to fingolimod. When switching patients from another disease modifying therapy to fingolimod, the elimination half-life and mode of action of the other therapy must be considered in order to avoid an additive immune effect whilst at the same time minimising the risk of disease reactivation. A CBC is recommended prior to initiating Fingolimod Mylan to ensure that immune effects of the previous therapy (i.e. cytopenia) have resolved.

Fingolimod Mylan can generally be started immediately after discontinuation of interferon or glatiramer acetate.

For dimethyl fumarate, the washout period should be sufficient for CBC to recover before treatment is started.

Due to the long elimination half-life of natalizumab, elimination usually takes up to 2-3 months following discontinuation. Teriflunomide is also eliminated slowly from the plasma. Without an accelerated elimination procedure, clearance of teriflunomide from plasma can take from several months up to 2 years. An accelerated elimination procedure as defined in the teriflunomide summary of product characteristics is recommended or alternatively washout period should not be shorter than 3.5 months. Caution regarding potential concomitant immune effects is required when switching patients from natalizumab or teriflunomide to fingolimod.

Alemtuzumab has profound and prolonged immunosuppressive effects. As the actual duration of these effects is unknown, initiating treatment with fingolimod after alemtuzumab is not recommended unless the benefits of such treatment clearly outweigh the risks for the individual patient.

A decision to use prolonged concomitant treatment with corticosteroids should be taken after careful consideration.

Co-administration with potent CYP450 inducers

The combination of fingolimod with potent CYP450 inducers should be used with caution. Concomitant administration with St John's Wort (*Hypericum perforatum*) is not recommended (see section 4.5).

Malignancies

Cutaneous malignancies

Basal cell carcinoma (BCC) and other cutaneous neoplasms, including malignant melanoma, squamous cell carcinoma, Kaposi's sarcoma and Merkel cell carcinoma, have been reported in patients receiving fingolimod (see section 4.8). Vigilance for skin lesions is warranted and a medical

evaluation of the skin is recommended at initiation and then every 6 to 12 months taking into consideration clinical judgement. The patient should be referred to a dermatologist in case suspicious lesions are detected.

Since there is a potential risk of malignant skin growths, patients treated with fingolimod should be cautioned against exposure to sunlight without protection. These patients should not receive concomitant phototherapy with UV-B-radiation or PUVA-photochemotherapy.

Lymphomas

There have been cases of lymphoma in clinical studies and the post-marketing setting (see section 4.8). The cases reported were heterogeneous in nature, mainly non-Hodgkin's lymphoma, including B-cell and T-cell lymphomas. Cases of cutaneous T-cell lymphoma (mycosis fungoides) have been observed. A fatal case of Epstein-Barr virus (EBV) positive B-cell lymphoma has also been observed. If lymphoma is suspected, treatment should be discontinued.

Women of childbearing potential

Due to risk to the foetus, fingolimod is contraindicated during pregnancy and in women of childbearing potential not using effective contraception. Before initiation of treatment, women of childbearing potential must be informed of this risk to the foetus, must have a negative pregnancy test and must use effective contraception during treatment and for 2 months after treatment discontinuation (see sections 4.3 and 4.6 and the information contained in the educational materials).

Tumefactive lesions

Rare cases of tumefactive lesions associated with MS relapse were reported in the post-marketing setting. In case of severe relapses, MRI should be performed to exclude tumefactive lesions. Discontinuation of treatment should be considered by the physician on a case-by-case basis taking into account individual benefits and risks.

Return of disease activity (rebound) after fingolimod discontinuation

In the post-marketing setting, severe exacerbation of disease has been observed rarely in some patients stopping fingolimod. This has generally been observed within 12 weeks after stopping treatment, but has also been reported up to 24 weeks after fingolimod discontinuation. Caution is therefore indicated when stopping treatment. If discontinuation of fingolimod is deemed necessary, the possibility of recurrence of exceptionally high disease activity should be considered and patients should be monitored for relevant signs and symptoms and appropriate treatment initiated as required (see 'Stopping therapy' below).

Stopping therapy

If a decision is made to stop treatment with Fingolimod Mylan a 6 week interval without therapy is needed, based on half-life, to clear it from the circulation (see section 5.2). Lymphocyte counts progressively return to normal range within 1-2 months of stopping therapy in most patients (see section 5.1) although full recovery can take significantly longer in some patients. Starting other therapies during this interval will result in concomitant exposure to fingolimod. Use of immunosuppressants soon after the discontinuation of Fingolimod Mylan may lead to an additive effect on the immune system and caution is therefore indicated.

After stopping fingolimod in the setting of PML, it is recommended to monitor patients for development of immune reconstitution inflammatory syndrome (PML-IRIS) (see "Progressive multifocal leukoencephalopathy" above).

Caution is also indicated when stopping therapy due to the risk of a rebound (see 'Return of disease activity (rebound) after fingolimod discontinuation' above). If discontinuation of Fingolimod Mylan is deemed necessary, patients should be monitored during this time for relevant signs of a possible

rebound.

Interference with serological testing

Since fingolimod reduces blood lymphocyte counts via re-distribution in secondary lymphoid organs, peripheral blood lymphocyte counts cannot be utilised to evaluate the lymphocyte subset status of a patient treated with Fingolimod Mylan. Laboratory tests involving the use of circulating mononuclear cells require larger blood volumes due to reduction in the number of circulating lymphocytes.

Paediatric population

The safety profile in paediatric patients is similar to that in adults and the warnings and precautions for adults therefore also apply to paediatric patients.

In particular, the following should be noted when prescribing Fingolimod Mylan to paediatric patients:

- Precautions should be followed at the time of the first dose (see 'Bradyarrhythmia' above). The same precautions as for the first dose are recommended when patients are switched from the 0.25 mg to the 0.5 mg daily dose.
- In the controlled paediatric trial D2311, cases of seizures, anxiety, depressed mood and depression have been reported with a higher incidence in patients treated with fingolimod compared to patients treated with interferon beta-1a. Caution is required in this subgroup population (see 'Paediatric population' in section 4.8).
- Mild isolated bilirubin increases have been noted in paediatric patients on fingolimod.
- It is recommended that paediatric patients complete all immunisations in accordance with current immunisation guidelines before starting Fingolimod Mylan therapy (see 'Infections' above).
- There are very limited data available in children between 10–12 years old, less than 40 kg or at Tanner stage < 2 (see sections 4.8 and 5.1). Caution is required in these subgroups due to very limited knowledge available from the clinical study.
- Long-term safety data in the paediatric population are not available.

4.5 Interaction with other medicinal products and other forms of interaction

Anti-neoplastic, immunomodulatory or immunosuppressive therapies

Anti-neoplastic, immunomodulatory or immunosuppressive therapies should not be co-administered due to the risk of additive immune system effects (see sections 4.3 and 4.4).

Caution should also be exercised when switching patients from long-acting therapies with immune effects such as natalizumab, teriflunomide or mitoxantrone (see section 4.4). In multiple sclerosis clinical studies the concomitant treatment of relapses with a short course of corticosteroids was not associated with an increased rate of infection.

Vaccination

During and for up to two months after treatment with Fingolimod Mylan, vaccination may be less effective. The use of live attenuated vaccines may carry a risk of infections and should therefore be avoided (see sections 4.4 and 4.8).

Bradycardia-inducing substances

Fingolimod has been studied in combination with atenolol and diltiazem. When it was used with atenolol in an interaction study in healthy volunteers, there was an additional 15% reduction of heart rate at fingolimod treatment initiation, an effect not seen with diltiazem. Treatment with Fingolimod Mylan should not be initiated in patients receiving beta blockers, or other substances which may decrease heart rate, such as class Ia and III antiarrhythmics, calcium channel blockers (such as verapamil or diltiazem), ivabradine, digoxin, anticholinesteratic agents or pilocarpine because of the

potential additive effects on heart rate (see sections 4.4 and 4.8). If treatment with this medicinal product is considered in such patients, advice from a cardiologist should be sought regarding the switch to non-heart-rate lowering medicinal products or appropriate monitoring for treatment initiation, at least overnight monitoring is recommended, if the heart-rate-lowering medication cannot be stopped.

Pharmacokinetic interactions of other substances on fingolimod

Fingolimod is metabolised mainly by CYP4F2. Other enzymes like CYP3A4 may also contribute to its metabolism, notably in the case of strong induction of CYP3A4. Potent inhibitors of transporter proteins are not expected to influence fingolimod disposition. Co-administration with ketoconazole resulted in a 1.7-fold increase in fingolimod and fingolimod phosphate exposure (AUC) by inhibition of CYP4F2. Caution should be exercised with substances that may inhibit CYP3A4 (protease inhibitors, azole antifungals, some macrolides such as clarithromycin or telithromycin).

Co-administration of carbamazepine 600 mg twice daily at steady-state and a single dose of fingolimod 2 mg reduced the AUC of fingolimod and its metabolite by approximately 40%. Other strong CYP3A4 enzyme inducers, for example rifampicin, phenobarbital, phenytoin, efavirenz and St. John's Wort, may reduce the AUC of fingolimod and its metabolite at least to this extent. As this could potentially impair the efficacy, their co-administration should be used with caution. Concomitant administration with St. John's Wort is however not recommended (see section 4.4).

Pharmacokinetic interactions of fingolimod on other substances

Fingolimod is unlikely to interact with substances mainly cleared by the CYP450 enzymes or by substrates of the main transporter proteins.

Co-administration of fingolimod with ciclosporin did not elicit any change in the ciclosporin or fingolimod exposure. Therefore, fingolimod is not expected to alter the pharmacokinetics of medicinal products that are CYP3A4 substrates.

Co-administration of fingolimod with oral contraceptives (ethinylestradiol and levonorgestrel) did not elicit any change in oral contraceptive exposure. No interaction studies have been performed with oral contraceptives containing other progestagens, however an effect of fingolimod on their exposure is not expected.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential/Contraception in females

Fingolimod is contraindicated in women of childbearing potential not using effective contraception (see section 4.3). Therefore, before initiation of treatment in women of childbearing potential, a negative pregnancy test result must be available and counselling should be provided regarding the serious risk to the foetus. Women of childbearing potential must use effective contraception during treatment and for 2 months after discontinuation of fingolimod, since it takes approximately 2 months to eliminate from the body after treatment discontinuation (see section 4.4).

Specific measures are also included in the educational materials. These measures must be implemented before fingolimod is prescribed to female patients and during treatment.

When stopping fingolimod therapy for planning a pregnancy the possible return of disease activity should be considered (see section 4.4).

Pregnancy

Based on human experience, post-marketing data suggest that use of fingolimod is associated with a 2-fold increased risk of major congenital malformations when administered during pregnancy compared

with the rate observed in the general population (2-3%; EUROCAT).

The following major malformations were most frequently reported:

- Congenital heart disease such as atrial and ventricular septal defects, tetralogy of Fallot
- Renal abnormalities
- Musculoskeletal abnormalities

There are no data on the effects of fingolimod on labour and delivery.

Animal studies have shown reproductive toxicity including foetal loss and organ defects, notably persistent truncus arteriosus and ventricular septal defect (see section 5.3). Furthermore, the receptor affected by fingolimod (sphingosine 1-phosphate receptor) is known to be involved in vascular formation during embryogenesis.

Consequently, fingolimod is contraindicated during pregnancy (see section 4.3). It should be stopped 2 months before planning a pregnancy (see section 4.4). If a woman becomes pregnant during treatment, fingolimod must be discontinued. Medical advice should be given regarding the risk of harmful effects to the foetus associated with treatment and ultrasonography examinations should be performed.

Breast-feeding

Fingolimod is excreted in milk of treated animals during lactation (see section 5.3). Due to the potential for serious adverse reactions in nursing infants, women receiving Fingolimod Mylan should not breastfeed.

Fertility

Data from preclinical studies do not suggest that fingolimod would be associated with an increased risk of reduced fertility (see section 5.3).

4.7 Effects on ability to drive and use machines

Fingolimod has no or negligible influence on the ability to drive and use machines.

However, dizziness or drowsiness may occasionally occur when initiating treatment. On initiation of Fingolimod Mylan treatment it is recommended that patients be observed for a period of 6 hours (see section 4.4, 'Bradyarrhythmia').

4.8 Undesirable effects

Summary of the safety profile

The most frequent adverse reactions (incidence $\geq 10\%$) at the 0.5 mg dose were headache (24.5%), hepatic enzyme increased (15.2%), diarrhoea (12.6%), cough (12.3%), influenza (11.4%), sinusitis (10.9%) and back pain (10.0%).

Tabulated list of adverse reactions

Adverse reactions reported in clinical trials and derived from post-marketing experience via spontaneous case reports or literature cases are shown below. Frequencies were defined using the following convention: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1\ 000$ to $< 1/100$); rare ($\geq 1/10\ 000$ to $< 1/1\ 000$); very rare ($< 1/10\ 000$); not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in the order of decreasing seriousness.

Infections and infestations	
Very common	Influenza
	Sinusitis
Common	Herpes viral infections
	Bronchitis
	<i>Tinea versicolor</i>
Uncommon	Pneumonia
Not known	Progressive multifocal leukoencephalopathy (PML)**
	Cryptococcal infections**
Neoplasms benign, malignant and unspecified (incl cysts and polyps)	
Common	Basal cell carcinoma
Uncommon	Malignant melanoma****
Rare	Lymphoma***
	Squamous cell carcinoma****
Very rare	Kaposi's sarcoma****
Not known	Merkel cell carcinoma***
Blood and lymphatic system disorders	
Common	Lymphopenia
	Leucopenia
Uncommon	Thrombocytopenia
Not known	Autoimmune haemolytic anaemia***
	Peripheral oedema***
Immune system disorders	
Not known	Hypersensitivity reactions, including rash, urticaria and angioedema upon treatment initiation***
	Immune reconstitution inflammatory syndrome (IRIS)**
Psychiatric disorders	
Common	Depression
Uncommon	Depressed mood
Nervous system disorders	
Very common	Headache
Common	Dizziness
	Migraine
Uncommon	Seizure
Rare	Posterior reversible encephalopathy syndrome (PRES)*
Not known	Severe exacerbation of disease after fingolimod discontinuation***
Eye disorders	
Common	Vision blurred
Uncommon	Macular oedema
Cardiac disorders	
Common	Bradycardia
	Atrioventricular block
Very rare	T-wave inversion***
Vascular disorders	
Common	Hypertension
Respiratory, thoracic and mediastinal disorders	
Very common	Cough
Common	Dyspnoea

Gastrointestinal disorders	
Very common	Diarrhoea
Uncommon	Nausea***
Hepatobiliary disorders	
Not known:	Acute hepatic failure***
Skin and subcutaneous tissue disorders	
Common	Eczema
	Alopecia
	Pruritus
Musculoskeletal and connective tissue disorders	
Very common	Back pain
Common	Myalgia
	Arthralgia
General disorders and administration site conditions	
Common	Asthenia
Investigations	
Very common	Hepatic enzyme increased (increased alanine transaminase, gamma glutamyltransferase, aspartate transaminase)
Common	Weight decreased***
	Blood triglycerides increased
Uncommon	Neutrophil count decreased
<p>* The frequency category was based on an estimated exposure of approximately 10 000 patients to fingolimod in all clinical trials.</p> <p>** PML, IRIS and cryptococcal infections (including cases of cryptococcal meningitis) have been reported in the post-marketing setting (see section 4.4).</p> <p>*** Adverse reactions from spontaneous reports and literature.</p> <p>**** The frequency category and risk assessment were based on an estimated exposure of more than 24 000 patients to fingolimod 0.5 mg in all clinical trials.</p>	

Description of selected adverse reactions

Infections

In multiple sclerosis clinical studies the overall rate of infections (65.1%) at the 0.5 mg dose was similar to placebo. However, lower respiratory tract infections, primarily bronchitis and to a lesser extent herpes infection and pneumonia were more common in fingolimod-treated patients.

Some cases of disseminated herpes infection, including fatal cases, have been reported even at the 0.5 mg dose.

In the post-marketing setting, cases of infections with opportunistic pathogens, such as viral (e.g. varicella zoster virus [VZV], JCV causing PML, herpes simplex virus [HSV]), fungal (e.g. cryptococci including cryptococcal meningitis) or bacterial (e.g. atypical mycobacterium), have been reported, some of which have been fatal (see section 4.4).

HPV infection, including papilloma, dysplasia, warts and HPV-related cancer, has been reported under treatment with fingolimod in the post-marketing setting (see section 4.4). Due to the immunosuppressive properties of fingolimod, vaccination against HPV should be considered prior to treatment initiation taking into account vaccination recommendations. Cancer screening, including Pap test, is recommended as per standard of care.

Macular oedema

In multiple sclerosis clinical studies macular oedema occurred in 0.5% of patients treated with the recommended dose of 0.5 mg and 1.1% of patients treated with the higher dose of 1.25 mg. The majority of cases occurred within the first 3-4 months of therapy. Some patients presented with

blurred vision or decreased visual acuity, but others were asymptomatic and diagnosed on routine ophthalmological examination. The macular oedema generally improved or resolved spontaneously after discontinuation of treatment. The risk of recurrence after re-challenge has not been evaluated.

Macular oedema incidence is increased in multiple sclerosis patients with a history of uveitis (17% with a history of uveitis vs. 0.6% without a history of uveitis). Fingolimod has not been studied in multiple sclerosis patients with diabetes mellitus, a disease which is associated with an increased risk for macular oedema (see section 4.4). In renal transplant clinical studies in which patients with diabetes mellitus were included, therapy with fingolimod 2.5 mg and 5 mg resulted in a 2-fold increase in the incidence of macular oedema.

Bradycardia

Initiation of treatment results in a transient decrease in heart rate and may also be associated with atrioventricular conduction delays. In multiple sclerosis clinical studies the maximal decline in heart rate was seen within 6 hours after treatment initiation, with declines in mean heart rate of 12-13 beats per minute for fingolimod 0.5 mg. Heart rate below 40 beats per minute in adults and below 50 beats per minute in paediatric patients, was rarely observed in patients on fingolimod 0.5 mg. The average heart rate returned towards baseline within one month of chronic treatment. Bradycardia was generally asymptomatic but some patients experienced mild to moderate symptoms, including hypotension, dizziness, fatigue and/or palpitations, which resolved within the first 24 hours after treatment initiation (see also sections 4.4 and 5.1).

In multiple sclerosis clinical studies first-degree atrioventricular block (prolonged PR interval on ECG) was detected after treatment initiation in adult and paediatric patients. In adult clinical trials it occurred in 4.7% of patients on fingolimod 0.5 mg, in 2.8% of patients on intramuscular interferon beta-1a and in 1.6% of patients on placebo. Second-degree atrioventricular block was detected in less than 0.2% adult patients on fingolimod 0.5 mg. In the post-marketing setting, isolated reports of transient, spontaneously resolving complete AV block have been observed during the 6 hour monitoring period following the first dose of fingolimod. The patients recovered spontaneously. The conduction abnormalities observed both in clinical trials and post-marketing were typically transient, asymptomatic and resolved within the first 24 hours after treatment initiation. Although most patients did not require medical intervention, one patient on fingolimod 0.5 mg received isoprenaline for asymptomatic second-degree Mobitz I atrioventricular block.

In the post-marketing setting, isolated delayed onset events, including transient asystole and unexplained death, have occurred within 24 hours of the first dose. These cases have been confounded by concomitant medicinal products and/or pre-existing disease. The relationship of such events to fingolimod is uncertain.

Blood pressure

In multiple sclerosis clinical studies fingolimod 0.5 mg was associated with an average increase of approximately 3 mmHg in systolic pressure and approximately 1 mmHg in diastolic pressure, manifesting approximately one month after treatment initiation. This increase persisted with continued treatment. Hypertension was reported in 6.5% of patients on fingolimod 0.5 mg and in 3.3% of patients on placebo. In the post-marketing setting, cases of hypertension have been reported within the first month of treatment initiation and on the first day of treatment that may require treatment with antihypertensive agents or discontinuation of fingolimod (see also section 4.4, Blood pressure effects).

Liver function

Increased hepatic enzymes have been reported in adult and paediatric multiple sclerosis patients treated with fingolimod. In clinical studies 8.0% and 1.8% of adult patients treated with fingolimod 0.5 mg experienced an asymptomatic elevation in serum levels of ALT of ≥ 3 x ULN (upper limit of normal) and ≥ 5 x ULN, respectively. Recurrence of liver transaminase elevations has occurred upon re-challenge in some patients, supporting a relationship to the medicinal product. In clinical studies, transaminase elevations occurred at any time during treatment although the majority occurred within the first 12 months. ALT levels returned to normal within approximately 2 months after discontinuation of treatment. In a small number of patients (N=10 on 1.25 mg, N=2 on 0.5 mg) who

experienced ALT elevations $\geq 5 \times$ ULN and who continued on fingolimod therapy, the ALT levels returned to normal within approximately 5 months (see also section 4.4, Liver function).

Nervous system disorders

In clinical studies, rare events involving the nervous system occurred in patients treated with fingolimod at higher doses (1.25 or 5.0 mg) including ischaemic and haemorrhagic strokes and neurological atypical disorders, such as acute disseminated encephalomyelitis (ADEM)-like events.

Cases of seizures, including status epilepticus, have been reported with the use of fingolimod in clinical studies and in the post-marketing setting.

Vascular disorders

Rare cases of peripheral arterial occlusive disease occurred in patients treated with fingolimod at higher doses (1.25 mg).

Respiratory system

Minor dose-dependent reductions in values for forced expiratory volume (FEV₁) and diffusion capacity for carbon monoxide (DLCO) were observed with fingolimod treatment starting at month one and remaining stable thereafter. At month 24, the reduction from baseline values in percentage of predicted FEV₁ was 2.7% for fingolimod 0.5 mg and 1.2% for placebo, a difference that resolved after treatment discontinuation. For DLCO the reductions at month 24 were 3.3% for fingolimod 0.5 mg and 2.7% for placebo (see also section 4.4, Respiratory effects).

Lymphomas

There have been cases of lymphoma of different varieties, in both clinical studies and the post-marketing setting, including a fatal case of Epstein-Barr virus (EBV) positive B-cell lymphoma. The incidence of non-Hodgkin's lymphoma (B-cell and T-cell) cases was higher in clinical trials than expected in the general population. Some T-cell lymphoma cases were also reported in the post-marketing setting, including cases of cutaneous T-cell lymphoma (mycosis fungoides) (see also section 4.4, Malignancies).

Haemophagocytic syndrome (HPS)

Very rare cases of HPS with fatal outcome have been reported in patients treated with fingolimod in the context of an infection. HPS is a rare condition that has been described in association with infections, immunosuppression and a variety of autoimmune diseases.

Paediatric population

In the controlled paediatric trial D2311 (see section 5.1), the safety profile in paediatric patients (10 to below 18 years of age) receiving fingolimod 0.25 mg or 0.5 mg daily was overall similar to that seen in adult patients. There were, nevertheless, more neurological and psychiatric disorders observed in the study. Caution is needed in this subgroup due to very limited knowledge available from the clinical study.

In the paediatric study, cases of seizures were reported in 5.6% of fingolimod-treated patients and 0.9% of interferon beta-1a-treated patients.

Depression and anxiety are known to occur with increased frequency in the multiple sclerosis population. Depression and anxiety have also been reported in paediatric patients treated with fingolimod.

Mild isolated bilirubin increases have been noted in paediatric patients on fingolimod.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare

professionals are asked to report any suspected adverse reactions the national reporting system listed in [Appendix V](#).

4.9 Overdose

Single doses up to 80 times the recommended dose (0.5 mg) were well tolerated in healthy adult volunteers. At 40 mg, 5 of 6 subjects reported mild chest tightness or discomfort which was clinically consistent with small airway reactivity.

Fingolimod can induce bradycardia upon treatment initiation. The decline in heart rate usually starts within one hour of the first dose and is steepest within 6 hours. The negative chronotropic effect of fingolimod persists beyond 6 hours and progressively attenuates over subsequent days of treatment (see section 4.4 for details). There have been reports of slow atrioventricular conduction, with isolated reports of transient, spontaneously resolving complete AV block (see sections 4.4 and 4.8).

If the overdose constitutes first exposure to Fingolimod Mylan, it is important to monitor patients with a continuous (real time) ECG and hourly measurement of heart rate and blood pressure, at least during the first 6 hours (see section 4.4).

Additionally, if after 6 hours the heart rate is < 45 bpm in adults, < 55 bpm in paediatric patients aged 12 years and above, or < 60 bpm in paediatric patients aged 10 years to below 12 years, or if the ECG at 6 hours after the first dose shows second degree or higher AV block, or if it shows a QTc interval ≥ 500 msec, monitoring should be extended at least for overnight and until the findings have resolved. The occurrence at any time of third degree AV block should also lead to extended monitoring including overnight monitoring.

Neither dialysis nor plasma exchange results in removal of fingolimod from the body.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Immunosuppressants, selective immunosuppressants,
ATC code: L04AE01

Mechanism of action

Fingolimod is a sphingosine 1-phosphate receptor modulator. It is metabolised by sphingosine kinase to the active metabolite fingolimod phosphate. Fingolimod phosphate binds at low nanomolar concentrations to sphingosine 1-phosphate (S1P) receptor 1 located on lymphocytes and readily crosses the blood-brain barrier to bind to S1P receptor 1 located on neural cells in the central nervous system (CNS). By acting as a functional antagonist of S1P receptors on lymphocytes, fingolimod phosphate blocks the capacity of lymphocytes to egress from lymph nodes, causing a redistribution, rather than depletion, of lymphocytes. Animal studies have shown that this redistribution reduces the infiltration of pathogenic lymphocytes, including pro-inflammatory Th17 cells, into the CNS, where they would be involved in nerve inflammation and nervous tissue damage. Animal studies and *in vitro* experiments indicate that fingolimod may also act via interaction with S1P receptors on neural cells.

Pharmacodynamic effects

Within 4-6 hours after the first dose of fingolimod 0.5 mg, the lymphocyte count decreases to approximately 75% of baseline in peripheral blood. With continued daily dosing, the lymphocyte count continues to decrease over a two-week period, reaching a minimal count of approximately 500 cells/microlitre or approximately 30% of baseline. Eighteen percent of patients reached a minimal count below 200 cells/microlitre on at least one occasion. Low lymphocyte counts are maintained with chronic daily dosing. The majority of T and B lymphocytes regularly traffic through lymphoid organs

and these are the cells mainly affected by fingolimod. Approximately 15-20% of T lymphocytes have an effector memory phenotype, cells that are important for peripheral immune surveillance. Since this lymphocyte subset typically does not traffic to lymphoid organs it is not affected by fingolimod. Peripheral lymphocyte count increases are evident within days of stopping treatment and typically normal counts are reached within one to two months. Chronic fingolimod dosing leads to a mild decrease in the neutrophil count to approximately 80% of baseline. Monocytes are unaffected by fingolimod.

Fingolimod causes a transient reduction in heart rate and decrease in atrioventricular conduction at treatment initiation (see sections 4.4 and 4.8). The maximal decline in heart rate is seen within 6 hours post dose, with 70% of the negative chronotropic effect achieved on the first day. With continued administration heart rate returns to baseline within one month. The decrease in heart rate induced by fingolimod can be reversed by parenteral doses of atropine or isoprenaline. Inhaled salmeterol has also been shown to have a modest positive chronotropic effect. With initiation of fingolimod treatment there is an increase in atrial premature contractions, but there is no increased rate of atrial fibrillation/flutter or ventricular arrhythmias or ectopy. Fingolimod treatment is not associated with a decrease in cardiac output. Autonomic responses of the heart, including diurnal variation of heart rate and response to exercise are not affected by treatment.

S1P4 could partially contribute to the effect but was not the main receptor responsible for the lymphoid depletion. The mechanism of action of bradycardia and vasoconstriction were also studied *in vitro* in guinea pigs and isolated rabbit aorta and coronary artery. It was concluded that bradycardia could be mediated primarily by activation of inward-rectifying potassium channel or G-protein activated inwardly rectifying K⁺ channel (IKACH/GIRK) and that vasoconstriction seems to be mediated by a Rho kinase and calcium dependent mechanism.

Fingolimod treatment with single or multiple doses of 0.5 and 1.25 mg for two weeks is not associated with a detectable increase in airway resistance as measured by FEV₁ and forced expiratory flow rate (FEF) 25-75. However, single fingolimod doses $\geq$ 5 mg (10-fold the recommended dose) are associated with a dose-dependent increase in airway resistance. Treatment with multiple doses of 0.5, 1.25, or 5 mg is not associated with impaired oxygenation or oxygen desaturation with exercise or an increase in airway responsiveness to methacholine. Subjects on fingolimod treatment have a normal bronchodilator response to inhaled beta-agonists.

Clinical efficacy and safety

The efficacy of fingolimod has been demonstrated in two studies which evaluated once-daily doses of 0.5 mg and 1.25 mg in adult patients with relapsing-remitting multiple sclerosis (RRMS). Both studies included adult patients who had experienced $\geq$ 2 relapses in the prior 2 years or $\geq$ 1 relapse during the prior year. Expanded Disability Status Score (EDSS) was between 0 and 5.5. A third study targeting the same adult patient population was completed after registration of fingolimod.

Study D2301 (FREEDOMS) was a 2-year randomised, double-blind, placebo-controlled Phase III study of 1 272 patients (n=425 on 0.5 mg, 429 on 1.25 mg, 418 on placebo). Median values for baseline characteristics were: age 37 years, disease duration 6.7 years and EDSS score 2.0. Outcome results are shown in Table 1. There were no significant differences between the 0.5 mg and the 1.25 mg doses as regards either endpoint.

Table 1 Study D2301 (FREEDOMS): main results

	Fingolimod 0.5 mg	Placebo
Clinical endpoints		
Annualised relapse rate (primary endpoint)	0.18**	0.4
Percentage of patients remaining relapse-free at 24 months	70%**	46%

Proportion with 3-month Confirmed Disability Progression† Hazard ratio (95% CI)	17% 0.70 (0.52, 0.96)*	24%
MRI endpoints		
Median (mean) number of new or enlarging T2 lesions over 24 months	0.0 (2.5)**	5.0 (9.8)
Median (mean) number of Gd-enhancing lesions at month 24	0.0 (0.2)**	0.0 (1.1)
Median (mean) % change in brain volume over 24 months	-0.7 (-0.8)**	-1.0 (-1.3)
† Disability progression defined as 1-point increase in EDSS confirmed 3 months later ** p < 0.001, *p < 0.05 compared to placebo All analyses of clinical endpoints were intent-to-treat. MRI analyses used evaluable dataset.		

Patients who completed the 24-month core FREEDOMS study could enter a dose-blinded extension study (D2301E1) and receive fingolimod. In total, 920 patients entered (n=331 continued on 0.5 mg, 289 continued on 1.25 mg, 155 switched from placebo to 0.5 mg and 145 switched from placebo to 1.25 mg). After 12 months (month 36), 856 patients (93%) were still enrolled. Between months 24 and 36, the annualised relapse rate (ARR) for patients on fingolimod 0.5 mg in the core study who remained on 0.5 mg was 0.17 (0.21 in the core study). The ARR for patients who switched from placebo to fingolimod 0.5 mg was 0.22 (0.42 in the core study).

Comparable results were shown in a replicate 2-year randomised, double-blind, placebo-controlled Phase III study on fingolimod in 1 083 patients (n=358 on 0.5 mg, 370 on 1.25 mg, 355 on placebo) with RRMS (D2309; FREEDOMS 2). Median values for baseline characteristics were: age 41 years, disease duration 8.9 years, EDSS score 2.5.

Table 2 Study D2309 (FREEDOMS 2): main results

	Fingolimod 0.5 mg	Placebo
Clinical endpoints		
Annualised relapse rate (primary endpoint)	0.21**	0.4
Percentage of patients remaining relapse-free at 24 months	71.5%**	52.7%
Proportion with 3-month Confirmed Disability Progression† Hazard ratio (95% CI)	25% 0.83 (0.61, 1.12)	29%
MRI endpoints		
Median (mean) number of new or enlarging T2 lesions over 24 months	0.0 (2.3)**	4.0 (8.9)
Median (mean) number of Gd-enhancing lesions at month 24	0.0 (0.4)**	0.0 (1.2)
Median (mean) % change in brain volume over 24 months	-0.71 (-0.86)**	-1.02 (-1.28)
† Disability progression defined as 1-point increase in EDSS confirmed 3 months later ** p < 0.001 compared to placebo All analyses of clinical endpoints were intent-to-treat. MRI analyses used evaluable dataset.		

Study D2302 (TRANSFORMS) was a 1-year randomised, double-blind, double-dummy, active (interferon beta-1a)-controlled Phase III study of 1 280 patients (n=429 on 0.5 mg, 420 on 1.25 mg, 431 on interferon beta-1a, 30 µg by intramuscular injection once weekly). Median values for baseline characteristics were: age 36 years, disease duration 5.9 years and EDSS score 2.0. Outcome results are shown in Table 3. There were no significant differences between the 0.5 mg and the 1.25 mg doses as regards study endpoints.

Table 3 Study D2302 (TRANSFORMS): main results

	Fingolimod 0.5 mg	Interferon beta-1a, 30 µg
Clinical endpoints		
Annualised relapse rate (primary endpoint)	0.16**	0.33
Percentage of patients remaining relapse-free at 12 months	83%**	71%
Proportion with 3-month Confirmed Disability Progression† Hazard ratio (95% CI)	6% 0.71 (0.42, 1.21)	8%
MRI endpoints		
Median (mean) number of new or enlarging T2 lesions over 12 months	0.0 (1.7)*	1.0 (2.6)
Median (mean) number of Gd-enhancing lesions at 12 months	0.0 (0.2)**	0.0 (0.5)
Median (mean) % change in brain volume over 12 months	-0.2 (-0.3)**	-0.4 (-0.5)
† Disability progression defined as 1-point increase in EDSS confirmed 3 months later. * p < 0.01, ** p < 0.001, compared to interferon beta-1a All analyses of clinical endpoints were intent-to-treat. MRI analyses used evaluable dataset.		

Patients who completed the 12-month core TRANSFORMS study could enter a dose-blinded extension (D2302E1) and receive fingolimod. In total, 1,030 patients entered, however, 3 of these patients did not receive treatment (n=356 continued on 0.5 mg, 330 continued on 1.25 mg, 167 switched from interferon beta-1a to 0.5 mg and 174 from interferon beta-1a to 1.25 mg). After 12 months (month 24), 882 patients (86%) were still enrolled. Between months 12 and 24, the ARR for patients on fingolimod 0.5 mg in the core study who remained on 0.5 mg was 0.20 (0.19 in the core study). The ARR for patients who switched from interferon beta-1a to fingolimod 0.5 mg was 0.33 (0.48 in the core study).

Pooled results of Studies D2301 and D2302 showed a consistent and statistically significant reduction in annualised relapse rate compared to comparator in subgroups defined by gender, age, prior multiple sclerosis therapy, disease activity or disability levels at baseline.

Further analyses of clinical trial data demonstrate consistent treatment effects in highly active subgroups of relapsing remitting multiple sclerosis patients.

Paediatric population

The efficacy and safety of once-daily doses of fingolimod 0.25 mg or 0.5 mg (dose selected based on body weight and exposure measurements) have been established in paediatric patients aged 10 to < 18 years with relapsing-remitting multiple sclerosis.

Study D2311 (PARADIGMS) was a double-blind, double-dummy, active-controlled study with flexible duration up to 24 months, with 215 patients 10 to < 18 years old (n=107 on fingolimod, 108 on interferon beta-1a 30 µg by intramuscular injection once weekly).

Median values for baseline characteristics were: age 16 years, median disease duration 1.5 years and EDSS score 1.5. The majority of patients were Tanner stage 2 or higher (94.4%) and were > 40 kg (95.3%). Overall, 180 (84%) of patients completed the core phase on study drug (n=99 [92.5%] on fingolimod, 81 [75%] on interferon beta-1a). Outcome results are shown in Table 4.

Table 4 Study D2311 (PARADIGMS): main results

	Fingolimod 0.25 mg or 0.5 mg	Interferon beta-1a 30 µg
Clinical endpoints	N=107	N=107 [#]
Annualised relapse rate (primary endpoint)	0.122**	0.675
Percentage of patients remaining relapse-free at 24 months	85.7**	38.8
MRI endpoints		
Annualised rate of the number of new or newly enlarging T2 lesions	n=106	n=102
Adjusted mean	4.393**	9.269
Number of Gd-enhancing T1 lesions per scan up to month 24	n=106	n=101
Adjusted mean	0.436**	1.282
Annualised rate of brain atrophy from baseline up to month 24	n=96	n=89
Least Square Mean	-0.48*	-0.80
# One patient randomised to receive interferon beta-1a by intramuscular injection was unable to swallow the double-dummy medication and discontinued from study. The patient was excluded from the full analysis and safety set.		
* p < 0.05, ** p < 0.001, compared to interferon beta-1a. All analyses of clinical endpoints were on the full analysis set.		

5.2 Pharmacokinetic properties

Pharmacokinetic data were obtained in healthy adult volunteers, in renal transplant adult patients and in multiple sclerosis adult patients.

The pharmacologically active metabolite responsible for efficacy is fingolimod phosphate.

Absorption

Fingolimod absorption is slow (t_{max} of 12-16 hours) and extensive (≥ 85%). The apparent absolute oral bioavailability is 93% (95% confidence interval: 79-111%). Steady-state-blood concentrations are reached within 1 to 2 months following once-daily administration and steady-state levels are approximately 10-fold greater than with the initial dose.

Food intake does not alter C_{max} or exposure (AUC) of fingolimod. Fingolimod phosphate C_{max} was slightly decreased by 34% but AUC was unchanged. Therefore, Fingolimod Mylan may be taken

without regard to meals (see section 4.2).

Distribution

Fingolimod highly distributes in red blood cells, with the fraction in blood cells of 86%. Fingolimod phosphate has a smaller uptake in blood cells of < 17%. Fingolimod and fingolimod phosphate are highly protein bound (> 99%).

Fingolimod is extensively distributed to body tissues with a volume of distribution of about $1\,200 \pm 260$ litres. A study in four healthy subjects who received a single intravenous dose of a radioiodolabelled analogue of fingolimod demonstrated that it penetrates into the brain. In a study in 13 male multiple sclerosis patients who received fingolimod 0.5 mg/day, the mean amount of fingolimod (and fingolimod phosphate) in seminal ejaculate, at steady-state, was approximately 10 000 times lower than the oral dose administered (0.5 mg).

Biotransformation

Fingolimod is transformed in humans by reversible stereoselective phosphorylation to the pharmacologically active (S)-enantiomer of fingolimod phosphate. Fingolimod is eliminated by oxidative biotransformation catalysed mainly via CYP4F2 and possibly other isoenzymes and subsequent fatty acid-like degradation to inactive metabolites. Formation of pharmacologically inactive non-polar ceramide analogues of fingolimod was also observed. The main enzyme involved in the metabolism of fingolimod is partially identified and may be either CYP4F2 or CYP3A4.

Following single oral administration of [^{14}C] fingolimod, the major fingolimod-related components in blood, as judged from their contribution to the AUC up to 34 days post dose of total radiolabelled components, are fingolimod itself (23%), fingolimod phosphate (10%) and inactive metabolites (M3 carboxylic acid metabolite (8%), M29 ceramide metabolite (9%) and M30 ceramide metabolite (7%)).

Elimination

Fingolimod blood clearance is 6.3 ± 2.3 l/h and the average apparent terminal elimination half-life ($t_{1/2}$) is 6-9 days. Blood levels of fingolimod and fingolimod phosphate decline in parallel in the terminal phase, leading to similar half-lives for both.

After oral administration, about 81% of the dose is slowly excreted in the urine as inactive metabolites. Fingolimod and fingolimod phosphate are not excreted intact in urine but are the major components in the faeces, with amounts representing less than 2.5% of the dose each. After 34 days, the recovery of the administered dose is 89%.

Linearity

Fingolimod and fingolimod phosphate concentrations increase in an apparently dose proportional manner after multiple once-daily doses of 0.5 mg or 1.25 mg.

Characteristics in specific groups of patients

Gender, ethnicity and renal impairment

The pharmacokinetics of fingolimod and fingolimod phosphate do not differ in males and females, in patients of different ethnic origin, or in patients with mild to severe renal impairment.

Hepatic impairment

In subjects with mild, moderate, or severe hepatic impairment (Child-Pugh class A, B and C), no change in fingolimod $C_{\max}$ was observed, but fingolimod AUC was increased respectively by 12%, 44% and 103%. In patients with severe hepatic impairment (Child-Pugh class C), fingolimod-phosphate $C_{\max}$ was decreased by 22% and AUC was not substantially changed. The pharmacokinetics of fingolimod-phosphate were not evaluated in patients with mild or moderate hepatic impairment.

The apparent elimination half-life of fingolimod is unchanged in subjects with mild hepatic impairment but is prolonged by about 50% in patients with moderate or severe hepatic impairment.

Fingolimod should not be used in patients with severe hepatic impairment (Child-Pugh class C) (see section 4.3). It should be introduced cautiously in mild and moderate hepatic impaired patients (see section 4.2).

Elderly population

Clinical experience and pharmacokinetic information in patients aged above 65 years are limited. Fingolimod Mylan should be used with caution in patients aged 65 years and over (see section 4.2).

Paediatric population

In paediatric patients (10 years of age and above), fingolimod-phosphate concentrations increase in an apparent dose proportional manner between 0.25 mg and 0.5 mg.

Fingolimod-phosphate concentration at steady state is approximately 25% lower in paediatric patients (10 years of age and above) following daily administration of 0.25 mg or 0.5 mg fingolimod compared to the concentration in adult patients treated with fingolimod 0.5 mg once daily.

There are no data available for paediatric patients below 10 years old.

5.3 Preclinical safety data

The preclinical safety profile of fingolimod was assessed in mice, rats, dogs and monkeys. The major target organs were the lymphoid system (lymphopenia and lymphoid atrophy), lungs (increased weight, smooth muscle hypertrophy at the bronchio-alveolar junction) and heart (negative chronotropic effect, increase in blood pressure, perivascular changes and myocardial degeneration) in several species; blood vessels (vasculopathy) in rats only at doses of 0.15 mg/kg and higher in a 2 year study, representing an approximate 4-fold margin based on the human systemic exposure (AUC) at a daily dose of 0.5 mg.

No evidence of carcinogenicity was observed in a 2-year bioassay in rats at oral doses of fingolimod up to the maximally tolerated dose of 2.5 mg/kg, representing an approximate 50-fold margin based on human systemic exposure (AUC) at the 0.5 mg dose. However, in a 2-year mouse study, an increased incidence of malignant lymphoma was seen at doses of 0.25 mg/kg and higher, representing an approximate 6-fold margin based on the human systemic exposure (AUC) at a daily dose of 0.5 mg.

Fingolimod was neither mutagenic nor clastogenic in animal studies.

It had no effect on sperm count/motility or on fertility in male and female rats up to the highest dose tested (10 mg/kg), representing an approximate 150-fold margin based on human systemic exposure (AUC) at a daily dose of 0.5 mg.

Fingolimod was teratogenic in the rat when given at doses of 0.1 mg/kg or higher. Drug exposure in rats at this dose was similar to that in patients at the therapeutic dose (0.5 mg). The most common foetal visceral malformations included persistent truncus arteriosus and ventricular septum defect. The teratogenic potential in rabbits could not be fully assessed, however an increased embryo-foetal mortality was seen at doses of 1.5 mg/kg and higher and a decrease in viable foetuses as well as foetal growth retardation was seen at 5 mg/kg. Drug exposure in rabbits at these doses was similar to that in patients.

In rats, F1 generation pup survival was decreased in the early postpartum period at doses that did not cause maternal toxicity. However, F1 body weights, development, behaviour and fertility were not affected by treatment with fingolimod. It was excreted in milk of treated animals during lactation at concentrations 2-fold to 3-fold higher than that found in maternal plasma. Fingolimod and its metabolites crossed the placental barrier in pregnant rabbits.

Juvenile animal studies

Results from two toxicity studies in juvenile rats showed slight effects on neurobehavioural response, delayed sexual maturation and a decreased immune response to repeated stimulations with keyhole limpet haemocyanin (KLH), which were not considered adverse. Overall, the treatment-related effects of fingolimod in juvenile animals were comparable to those seen in adult rats at similar dose levels, with the exception of changes in bone mineral density and neurobehavioural impairment (reduced auditory startle response) observed at doses of 1.5 mg/kg and higher in juvenile animals and the absence of smooth muscle hypertrophy in the lungs of the juvenile rats.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule content

Calcium hydrogen phosphate dihydrate
Glycine
Silica, colloidal anhydrous
Magnesium stearate

Capsule shell

Gelatin
Titanium dioxide (E171)
Yellow iron oxide (E172)
Red iron oxide (E172)

Printing ink

Shellac (E904)
Propylene glycol (E1520)
Black iron oxide (E172)
Potassium hydroxide

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

Do not store above 25°C.
Store in the original package in order to protect from moisture.

6.5 Nature and contents of container

PVC/PCTFE-alu blister

Pack size:
28, 30, 84 or 98 hard capsules
Multipacks containing 84 (3 packs of 28) hard capsules

Calendar packs containing 28 or 84 hard capsules
Unit dose blister packs containing 7 x 1, 28 x 1, 90 x 1 or 98 x 1 hard capsules

PVC/PE/PVdC-alu blister

Pack size:

28, 30, 84 or 98 hard capsules

Multipacks containing 84 (3 packs of 28) hard capsules

Calendar packs containing 28 or 84 hard capsules

Unit dose blister packs containing 7 x 1, 28 x 1, 90 x 1 or 98 x 1 hard capsules

White round HDPE bottle with white opaque child-resistant PP closure with wad containing aluminium induction sealing liner

Pack size: 90 or 100 hard capsules.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Mylan Pharmaceuticals Limited
Damastown Industrial Park,
Mulhuddart, Dublin 15,
DUBLIN
Ireland

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/21/1573/001
EU/1/21/1573/002
EU/1/21/1573/003
EU/1/21/1573/004
EU/1/21/1573/005
EU/1/21/1573/006
EU/1/21/1573/007
EU/1/21/1573/008
EU/1/21/1573/009
EU/1/21/1573/010
EU/1/21/1573/011
EU/1/21/1573/012
EU/1/21/1573/013
EU/1/21/1573/014
EU/1/21/1573/015
EU/1/21/1573/016
EU/1/21/1573/017
EU/1/21/1573/018
EU/1/21/1573/019
EU/1/21/1573/020
EU/1/21/1573/021
EU/1/21/1573/022

EU/1/21/1573/023
EU/1/21/1573/024

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 18 August 2021

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <https://www.ema.europa.eu/en>

ANNEX II

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer(s) responsible for batch release

Mylan Hungary Kft
Mylan utca 1
Komarom
H-2900
Hungary

Viatrix Germany GmbH,
Zweigniederlassung Bad Homburg v. d. Hoehe,
Benzstrasse 1,
Bad Homburg v. d. Hoehe,
Hessen, 61352,
Germany.

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

- **Additional risk minimisation measures**

Prior to launch of Fingolimod Mylan in each Member State the Marketing Authorisation Holder (MAH) must agree about the content and format of the educational programme, including

communication media, distribution modalities, and any other aspects of the programme, with the National Competent Authority (NCA).

The MAH shall ensure that in each Member State (MS) where Fingolimod Mylan is marketed, all physicians who intend to prescribe it are provided with the following educational materials:

1. Summary of Product Characteristics (SmPC)
2. Physician's checklist for adult and paediatric patients, to consider prior to prescribing Fingolimod Mylan.
3. The Patient / Parent / Caregiver's guide, to be provided to all patients, their parents (or legal representatives), and caregivers.
4. The pregnancy-specific patient reminder card, to be provided to all patients, their parents (or legal representatives), and caregivers, as applicable.

Physician's checklist

The physician's checklist shall contain the following key messages:

Safety topics	Key safety messages
Bradyarrhythmia (including conduction defects and bradycardia complicated by hypotension) occurring post-first dose	<ul style="list-style-type: none">• Do not initiate Fingolimod Mylan in patients with a cardiac condition or who are taking medicinal products for which Fingolimod Mylan is contraindicated.• Prior to initiating Fingolimod Mylan in patients with underlying medical conditions or who are concomitantly taking medicinal products which carry an increased risk of serious rhythm disturbance or bradycardia, ensure the anticipated benefits outweigh the potential risks and seek advice from a cardiologist regarding appropriate monitoring (at least overnight extended monitoring for treatment initiation) and/or adjustment of the concomitant medicinal product.• Monitor all patients for signs and symptoms of bradycardia for a period of at least 6 hours after the first dose of Fingolimod Mylan, including performing an electrocardiogram (ECG) and blood pressure measurement prior to and 6 hours after first dose.• If post-dose signs and symptoms of bradyarrhythmia occur, extend first dose monitoring per guidelines until resolution; be familiar with criteria (i.e. need for pharmacological intervention, age-specific heart rate limits, new ECG findings) that would warrant overnight monitoring.• Follow first-dose monitoring recommendations after treatment interruption or increase in daily dose.

Liver transaminase elevation	<ul style="list-style-type: none"> • Do not initiate Fingolimod Mylan in patients with severe liver impairment (Child-Pugh class C) • Transaminase and bilirubin levels should be obtained prior to initiation of Fingolimod Mylan, monitored every 3 months for the first year on therapy, and periodically thereafter, up to 2 months after Fingolimod Mylan discontinuation. • For asymptomatic elevations in liver function tests (LFTs), perform LFTs more frequently if increases in transaminase are greater than 3 times to less than 5 times upper limit of normal (ULN) without increase in serum bilirubin. Discontinue Fingolimod Mylan for transaminase increase at least 5 times the ULN or at least 3 times the ULN associated with any increase in serum bilirubin. Restart Fingolimod Mylan only after careful consideration of benefit-risk. • For patients with clinical symptoms of liver dysfunction, evaluate promptly and discontinue Fingolimod Mylan if significant liver injury is confirmed. If serum levels return to normal (including if an alternative cause of the liver dysfunction is discovered), Fingolimod Mylan may be restarted based on a careful benefit-risk assessment of the patient.
Macular oedema	<ul style="list-style-type: none"> • Obtain an ophthalmological assessment before starting Fingolimod Mylan in patients with diabetes or a history of uveitis. • Obtain an ophthalmological assessment in all patients 3 to 4 months after initiating Fingolimod Mylan. • It is recommended to discontinue Fingolimod Mylan in patients who develop macular degeneration. Restart Fingolimod Mylan only after careful consideration of benefit-risk.

<p>Opportunistic infections including varicella zoster virus (VZV), herpes viral infections other than VZV, fungal infections</p>	<ul style="list-style-type: none"> • Do not initiate Fingolimod Mylan in patients with immunodeficiency syndrome, increased risk for opportunistic infections including immunocompromised patients or severe active or active chronic infections (i.e. hepatitis or tuberculosis). • Fingolimod Mylan may be initiated in patients who have had severe active infection that has resolved. • Anti-neoplastic, immunomodulatory or immunosuppressive therapies should not be co-administered due to the risk of additive immune system effects. Carefully consider any decision regarding prolonged concomitant corticosteroid use. • Monitor peripheral blood lymphocyte counts prior to and during treatment with Fingolimod Mylan. Interrupt treatment for lymphocyte count $<0.2 \times 10^9/l$ until recovery. • Instruct patients to report signs and symptoms of infections during, and for up to two months after, treatment with Fingolimod Mylan. • For potentially serious infections, evaluate the patient promptly and consider an infectious disease referral. Consider suspending Fingolimod Mylan and the benefit-risk of any subsequent reinitiation. • Be aware that serious, life-threatening and sometimes fatal cases of opportunistic infections of the central nervous system (CNS) have occurred on Fingolimod Mylan treatment, including herpes viral infection (encephalitis, meningitis and meningo-encephalitis; observed at any time) and cryptococcal meningitis, (observed after approximately 2-3 years). • Fingolimod Mylan should be discontinued in patients with CNS herpes infections. Fingolimod Mylan should be suspended in patients with cryptococcal meningitis with careful consideration with a specialist before reinitiating. • Inform patients that during Fingolimod Mylan treatment, they should not receive live attenuated vaccines and that other vaccines may be less effective. • Prior to initiation of Fingolimod Mylan, check varicella status and recommend a full course of vaccination for VZV in anti-body negative patients. Postpone initiation of treatment for 1 month to allow the full effect of vaccination. • Recommend vaccination against human papilloma virus (HPV) prior to initiation of treatment.
---	--

Progressive multifocal leukoencephalopathy (PML)	<ul style="list-style-type: none"> Do not treat with Fingolimod Mylan in patients with suspected or confirmed PML. Be aware that PML has been predominantly observed after 2 or more years of fingolimod treatment. Ensure patients have a baseline magnetic resonance imaging (MRI) usually within 3 months before initiating Fingolimod Mylan. Annual MRIs may be considered especially in patients with multiple risk factors generally associated with PML. If there is suspicion of PML, perform a diagnostic MRI immediately and suspend Fingolimod Mylan until PML has been excluded. If PML is confirmed, treatment with Fingolimod Mylan must be permanently discontinued. Immune reconstitution inflammatory syndrome (IRIS) has been reported in patients treated with S1P receptor modulators, including fingolimod, who developed PML and subsequently discontinued treatment. The time to onset of IRIS in patients with PML was usually from weeks to months after S1P receptor modulator discontinuation. Monitoring for development of IRIS and appropriate treatment of the associated inflammation should be undertaken.
Reproductive toxicity	<ul style="list-style-type: none"> Fingolimod Mylan is teratogenic and contraindicated in women of childbearing potential not using effective contraception or who are pregnant. Women of childbearing potential must use effective contraception during treatment and for two months following treatment discontinuation. Before initiating treatment and regularly thereafter, counsel women of childbearing potential, including adolescent females, their parents or legal representatives, about the risks to a foetus and to use effective contraception during treatment and for two months after discontinuation. Confirm a negative pregnancy test prior to starting treatment and repeat at suitable intervals. Discontinue Fingolimod Mylan if a woman becomes pregnant and consider the possible return of disease activity. Instruct patient to stop Fingolimod Mylan two months before attempting to become pregnant.
Skin cancer (basal cell carcinoma, Kaposi's sarcoma, malignant melanoma, Merkel cell carcinoma, squamous cell carcinoma)	<ul style="list-style-type: none"> Perform skin examination prior to treatment initiation and every 6 to 12 months. Refer patients to a dermatologist if suspicious lesions are detected. Caution against exposure to sunlight without protection. Instruct patient to avoid concomitant phototherapy with UV-B-radiation or PUVA-photochemotherapy.
Use in paediatric patients, including impact on growth and development	<ul style="list-style-type: none"> All warnings and precautions and monitoring in adults also apply to paediatric patients. Assess Tanner staging, height, and weight according to standard of care. Ensure that vaccination status is up to date before initiation of Fingolimod Mylan. Monitor for signs and symptoms of depression and anxiety.

Patient / Parent / Caregiver guide

The patient/parents/caregiver guide shall contain the following key messages:

Safety topics	Key safety messages
Bradyarrhythmia (including conduction defects and bradycardia complicated by hypotension) occurring post-first dose	<ul style="list-style-type: none"> • Inform your doctor if you have underlying cardiac conditions or are taking medicines known to decrease your heart rate. • Your doctor will perform an ECG and blood pressure measurement before the first dose of Fingolimod Mylan. • Your doctor will monitor your heart rate after the first dose. Prolonged and overnight monitoring may be required. Follow-up monitoring may be required at treatment re-initiation. • Immediately inform your doctor of symptoms indicating low heart rate (such as dizziness, vertigo, nausea or palpitations), that develop after the first dose of Fingolimod Mylan. • Call your doctor in case of missed doses as the first dose monitoring may need to be repeated.
Liver transaminase elevation	<ul style="list-style-type: none"> • Inform your doctor if you have liver problems. • Your doctor will perform liver function tests before starting treatment, at specified intervals during treatment and for up to 2 months after discontinuation. • Inform your doctor if you notice any of the signs of liver injury (such as yellowing of your skin or the whites of your eyes, abnormally dark urine, pain on the right side of the stomach area, unexplained nausea and vomiting).
Macular oedema	<ul style="list-style-type: none"> • Your doctor may arrange an eye assessment before starting Fingolimod Mylan and as required during treatment. A follow-up eye assessment may be done 3-4 months after starting Fingolimod Mylan. • Immediately inform your doctor of any symptoms of visual changes during treatment and for up to two months after the end of treatment with Fingolimod Mylan.
Opportunistic infections including varicella zoster virus (VZV), herpes viral infections other than VZV, fungal infections	<ul style="list-style-type: none"> • Your doctor will monitor blood lymphocyte counts prior to and during treatment with Fingolimod Mylan. Treatment with Fingolimod Mylan may be interrupted if blood lymphocyte count is too low. • Immediately inform your doctor of signs and symptoms of infection, during and up to two months after Fingolimod Mylan treatment (such as fever, flu-like symptoms, headache accompanied by stiff neck, sensitivity to light, nausea, shingles and/or confusion or seizures [may be symptoms of meningitis and/or encephalitis]).
Progressive multifocal leukoencephalopathy (PML)	<ul style="list-style-type: none"> • PML is a rare brain disorder caused by an infection that may lead to severe disability or death. • Your doctor will arrange magnetic resonance imaging (MRI) scans before you start treatment and during treatment to monitor the risk of PML. • Immediately inform your physician if you believe your MS is getting worse or if you notice any new symptoms during and after treatment with Fingolimod Mylan, for example changes in mood or behaviour, new or worsening weakness on one side of the body, changes in vision, confusion, memory lapses or speech and communication difficulties. These may be symptoms of PML or of an inflammatory reaction (known as immune reconstitution inflammatory syndrome or IRIS) that can occur in patients with PML as Fingolimod Mylan is removed from their body after they stop taking it. • Speak with your partner or caregivers and inform them about your treatment. Symptoms might arise that you might not become aware of by yourself.

Skin cancer (basal cell carcinoma, Kaposi's sarcoma, malignant melanoma, Merkel cell carcinoma, squamous cell carcinoma)	<ul style="list-style-type: none"> Immediately inform your doctor if any skin nodules (e.g. shiny, pearly nodules), patches or open sores that do not heal within weeks are noted. Skin cancers have been reported in multiple sclerosis patients treated with Fingolimod Mylan. Symptoms of skin cancer may include abnormal growth or changes of skin tissue (e.g. unusual moles) with a change in colour, shape or size over time.
Reproductive toxicity	<ul style="list-style-type: none"> Fingolimod Mylan must not be used in women who could become pregnant and are not using effective contraception or who are pregnant. If you are a woman of childbearing potential, you must use effective contraception during treatment and for two months following treatment discontinuation. Immediately report to your doctor any (intended or unintended) pregnancy during treatment and up to two months following discontinuation of Fingolimod Mylan treatment.
Specifically for paediatric patients	<p>All warnings and precautions and monitoring in adults also apply to paediatric patients. In addition:</p> <ul style="list-style-type: none"> Your doctor will assess height, weight, and puberty status according to standard of care. Your doctor will ensure that your vaccination status is up to date before you start treatment with Fingolimod Mylan. Monitor for signs and symptoms of depression and anxiety.

Pregnancy-specific patient reminder card

The pregnancy-specific patient reminder card shall contain the following key messages:

<ul style="list-style-type: none"> IF USED DURING PREGNANCY, FINGOLIMOD MYLAN CAN HARM YOUR UNBORN BABY. Fingolimod Mylan is contraindicated during pregnancy and in women of childbearing potential not using effective contraception. It is important that you use effective contraception while taking Fingolimod Mylan and for 2 months after you stop taking it to avoid becoming pregnant. Your doctor will provide counselling regarding effective contraception. Your doctor will provide counselling before treatment initiation and regularly thereafter regarding the risk of Fingolimod Mylan harming an unborn baby and required actions to minimise this risk. A pregnancy test must be carried out and negative results verified by your doctor before starting treatment. A pregnancy test must be repeated at suitable intervals. Women must NOT become pregnant while on treatment. If you become pregnant or want to become pregnant, Fingolimod Mylan must be discontinued. Inform your doctor immediately if you think you are pregnant. Your doctor will provide counselling in the event of pregnancy and evaluation of the outcome of any pregnancy. Immediately inform your doctor if there is worsening of multiple sclerosis after stopping treatment with Fingolimod Mylan.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON

1. NAME OF THE MEDICINAL PRODUCT

Fingolimod Mylan 0.5 mg hard capsules
fingolimod

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each capsule contains 0.5 mg of fingolimod (as hydrochloride).

3. LIST OF EXCIPIENTS

4. PHARMACEUTICAL FORM AND CONTENTS

Hard capsule

28 hard capsules

30 hard capsules

84 hard capsules

98 hard capsules

Calendar pack: 28 hard capsules

Calendar pack: 84 hard capsules

Unit dose blister pack: 7 x 1 hard capsule

Unit dose blister pack: 28 x 1 hard capsule

Unit dose blister pack: 90 x 1 hard capsule

Unit dose blister pack: 98 x 1 hard capsule

Bottle pack: 90 hard capsules

Bottle pack: 100 hard capsules

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.

Oral use.

Swallow each capsule whole.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.

Store in the original package in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Mylan Pharmaceuticals Limited, Damastown Industrial Park, Mulhuddart, Dublin 15, DUBLIN, Ireland

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/21/1573/001
EU/1/21/1573/002
EU/1/21/1573/003
EU/1/21/1573/004
EU/1/21/1573/005
EU/1/21/1573/006
EU/1/21/1573/007
EU/1/21/1573/008
EU/1/21/1573/010
EU/1/21/1573/011
EU/1/21/1573/012
EU/1/21/1573/013
EU/1/21/1573/014
EU/1/21/1573/015
EU/1/21/1573/016
EU/1/21/1573/017
EU/1/21/1573/018
EU/1/21/1573/019
EU/1/21/1573/020
EU/1/21/1573/021
EU/1/21/1573/023
EU/1/21/1573/024

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Fingolimod Mylan 0.5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA
--

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**OUTER CARTON FOR MULTIPACK (WITH BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Fingolimod Mylan 0.5 mg hard capsules
fingolimod

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each capsule contains 0.5 mg of fingolimod (as hydrochloride).

3. LIST OF EXCIPIENTS**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule

Multipack: 84 (3 packs of 28) hard capsules

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.
Swallow each capsule whole.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.
Store in the original package in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Mylan Pharmaceuticals Limited, Damastown Industrial Park, Mulhuddart, Dublin 15, DUBLIN, Ireland

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/21/1573/009

EU/1/21/1573/022

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Fingolimod Mylan 0.5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC
SN
NN

PARTICULARS TO APPEAR ON THE OUTER PACKAGING**INTERMEDIATE CARTON FOR MULTIPACK (WITHOUT BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Fingolimod Mylan 0.5 mg hard capsules
fingolimod

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each capsule contains 0.5 mg of fingolimod (as hydrochloride).

3. LIST OF EXCIPIENTS**4. PHARMACEUTICAL FORM AND CONTENTS**

28 hard capsules. Component of a multipack can't be sold separately.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.
Swallow each capsule whole.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.
Store in the original package in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Mylan Pharmaceuticals Limited, Damastown Industrial Park, Mulhuddart, Dublin 15, DUBLIN, Ireland

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/21/1573/009

EU/1/21/1573/022

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Fingolimod Mylan 0.5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

MINIMUM PARTICULARS TO APPEAR ON BLISTERS

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Fingolimod Mylan 0.5 mg hard capsules
fingolimod

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Mylan Pharmaceuticals Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

[For calendar packs]
SUN→MON→TUE→WED→THU→FRI→SAT

MINIMUM PARTICULARS TO APPEAR ON BLISTERS

UNIT DOSE BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Fingolimod Mylan 0.5 mg hard capsules
fingolimod

2. NAME OF THE MARKETING AUTHORISATION HOLDER

Mylan Pharmaceuticals Limited

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

Oral use

PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**BOTTLE****1. NAME OF THE MEDICINAL PRODUCT**

Fingolimod Mylan 0.5 mg hard capsules
fingolimod

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each capsule contains 0.5 mg of fingolimod (as hydrochloride).

3. LIST OF EXCIPIENTS**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule

90 hard capsules

100 hard capsules

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.
Swallow each capsule whole.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY**8. EXPIRY DATE**

EXP

9. SPECIAL STORAGE CONDITIONS

Do not store above 25°C.
Store in the original package in order to protect from moisture.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Mylan Pharmaceuticals Limited, Damastown Industrial Park, Mulhuddart, Dublin 15, DUBLIN, Ireland

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/21/1573/012

EU/1/21/1573/013

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Fingolimod Mylan 0.5 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

B. PACKAGE LEAFLET

Package leaflet: Information for the patient

Fingolimod Mylan 0.5 mg hard capsules fingolimod

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Fingolimod Mylan is and what it is used for
2. What you need to know before you take Fingolimod Mylan
3. How to take Fingolimod Mylan
4. Possible side effects
5. How to store Fingolimod Mylan
6. Contents of the pack and other information

1. What Fingolimod Mylan is and what it is used for

What Fingolimod Mylan is

Fingolimod Mylan contains the active substance fingolimod.

What Fingolimod Mylan is used for

Fingolimod Mylan is used in adults and in children and adolescents (10 years of age and above) to treat relapsing-remitting multiple sclerosis (MS), more specifically in:

- Patients who have failed to respond despite treatment with an MS treatment.
- or
- Patients who have rapidly evolving severe MS.

Fingolimod Mylan does not cure MS, but it helps to reduce the number of relapses and to slow down the progression of physical disabilities due to MS.

What is multiple sclerosis

MS is a long-term condition that affects the central nervous system (CNS), comprised of the brain and spinal cord. In MS inflammation destroys the protective sheath (called myelin) around the nerves in the CNS and stops the nerves from working properly. This is called demyelination.

Relapsing-remitting MS is characterised by repeated attacks (relapses) of nervous system symptoms that reflect inflammation within the CNS. Symptoms vary from patient to patient but typically involve walking difficulties, numbness, vision problems or disturbed balance. Symptoms of a relapse may disappear completely when the relapse is over, but some problems may remain.

How Fingolimod Mylan works

Fingolimod Mylan helps to protect against attacks on the CNS by the immune system by reducing the ability of some white blood cells (lymphocytes) to move freely within the body and by stopping them from reaching the brain and spinal cord. This limits nerve damage caused by MS. This medicine also reduces some of the immune reactions of your body.

2. What you need to know before you take Fingolimod Mylan

Do not take Fingolimod Mylan

- **if you are allergic** to fingolimod or any of the other ingredients of this medicine (listed in section 6).
- if you have a **lowered immune response** (due to an immunodeficiency syndrome, a disease or to medicines that suppress the immune system).
- if your doctor suspects you may have a **rare brain infection called progressive multifocal leukoencephalopathy (PML) or if PML has been confirmed**.
- if you have a **severe active infection or active chronic infection** such as hepatitis or tuberculosis.
- if you have an **active cancer**.
- if you have **severe liver problems**.
- **if, in the last 6 months, you have had heart attack, angina, stroke or warning of a stroke or certain types of heart failure**.
- if you have certain types of **irregular or abnormal heartbeat** (arrhythmia), including patients in whom the electrocardiogram (ECG) shows prolonged QT interval.
- **if you are taking or have recently taken medicine for irregular heartbeat** such as quinidine, disopyramide, amiodarone or sotalol.
- if you are **pregnant or a woman of childbearing potential not using effective contraception**.

If this applies to you or you are unsure, **talk to your doctor before taking Fingolimod Mylan**.

Warnings and precautions

Talk to your doctor before taking Fingolimod Mylan:

- **if you have severe breathing problems during sleep (severe sleep apnoea)**.
- **if you have been told you have an abnormal electrocardiogram**.
- **if you suffer from symptoms of slow heart rate (e.g. dizziness, nausea, or palpitations)**.
- **if you are taking or have recently taken medicines that slow your heart rate** (such as beta blockers, verapamil, diltiazem or ivabradine, digoxin, anticholinesteratic agents or pilocarpine).
- **if you have a history of sudden loss of consciousness or fainting (syncope)**.
- **if you plan to get vaccinated**.
- **if you have never had chickenpox**.
- **if you have or have had visual disturbances** or other signs of swelling in the central vision area (macula) at the back of the eye (a condition known as macular oedema, see below), inflammation or infection of the eye (uveitis), **or if you have diabetes** (which can cause eye problems).
- **if you have liver problems**.
- if you have **high blood pressure that cannot be controlled by medicines**.
- if you have **severe lung problems** or smoker's cough.

If any of these applies to you or you are unsure, **talk to your doctor before taking Fingolimod Mylan**.

Slow heart rate (bradycardia) and irregular heartbeat

At the beginning of treatment or after taking the first dose of 0.5 mg when you switch from a 0.25 mg daily dose, Fingolimod Mylan causes the heart rate to slow down. As a result, you may feel dizzy or tired, or be consciously aware of your heartbeat, or your blood pressure may drop. **If these effects are severe, tell your doctor, because you may need treatment right away**. This medicine can also cause an irregular heartbeat, especially after the first dose. Irregular heartbeat usually returns to normal in less than one day. Slow heart rate usually returns to normal within one month. During this period, no clinically significant heart rate effects are usually expected.

Your doctor will ask you to stay at the surgery or clinic for at least 6 hours, with hourly pulse and blood pressure measurements, after taking the first dose of Fingolimod Mylan or after taking the first dose of 0.5 mg when you switch from a 0.25 mg daily dose, so that appropriate measures can be taken in the event of side effects that occur at the start of treatment. You should have an electrocardiogram

performed prior to the first dose of this medicine and after the 6-hour monitoring period. Your doctor may monitor your electrocardiogram continuously during that time. If after the 6-hour period you have a very slow or decreasing heart rate, or if your electrocardiogram shows abnormalities, you may need to be monitored for a longer period (at least 2 more hours and possibly overnight) until these have resolved. The same may apply if you are resuming Fingolimod Mylan after a break in treatment, depending on both how long the break was and how long you had been taking it before the break.

If you have, or if you are at risk for, an irregular or abnormal heartbeat, if your electrocardiogram is abnormal, or if you have heart disease or heart failure, Fingolimod Mylan may not be appropriate for you.

If you have a history of sudden loss of consciousness or decreased heart rate, Fingolimod Mylan may not be appropriate for you. You will be evaluated by a cardiologist (heart specialist) to advise how you should start treatment, including overnight monitoring.

If you are taking medicines that can cause your heart rate to decrease, Fingolimod Mylan may not be appropriate for you. You will need to be evaluated by a cardiologist, who will check whether you can be switched to alternative medicine that does not decrease your heart rate in order to allow treatment with Fingolimod Mylan. If such a switch is impossible, the cardiologist will advise how you should start treatment with Fingolimod Mylan, including overnight monitoring.

If you have never had chickenpox

If you have never had chickenpox, your doctor will check your immunity against the virus that causes it (varicella zoster virus). If you are not protected against the virus, you may need a vaccination before you start treatment with Fingolimod Mylan. If this is the case, your doctor will delay the start of treatment until one month after the full course of vaccination is completed.

Infections

Fingolimod Mylan lowers the white blood cell count (particularly the lymphocyte count). White blood cells fight infection. While you are taking this medicine (and for up to 2 months after you stop taking it), you may get infections more easily. Any infection that you already have may get worse. Infections could be serious and life-threatening. If you think you have an infection, have fever, feel like you have the flu, have shingles or have a headache accompanied by stiff neck, sensitivity to light, nausea, rash and/or confusion or seizures (fits) (these may be symptoms of meningitis and/or encephalitis caused by a fungal or herpes viral infection), contact your doctor straight away, because it could be serious and life-threatening.

Human papilloma virus (HPV) infection, including papilloma, dysplasia, warts and HPV-related cancer, has been reported in patients treated with Fingolimod Mylan. Your doctor will consider whether you need to have a vaccination against HPV before starting treatment. If you are a woman, your doctor will also recommend HPV screening.

PML

PML is a rare brain disorder caused by an infection that may lead to severe disability or death. Your doctor will arrange magnetic resonance imaging (MRI) scans before you start treatment and during treatment to monitor the risk of PML.

If you believe your MS is getting worse or if you notice any new symptoms, for example changes in mood or behaviour, new or worsening weakness on one side of the body, changes in vision, confusion, memory lapses or speech and communication difficulties, talk to your doctor as soon as possible. These may be symptoms of PML. Also speak with your partner or caregivers and inform them about your treatment. Symptoms might arise that you might not become aware of by yourself.

If you get PML it can be treated and your treatment with Fingolimod Mylan will be stopped. Some people get an inflammatory reaction as Fingolimod Mylan is removed from the body. This reaction (known as immune reconstitution inflammatory syndrome or IRIS) may lead to your condition getting worse, including worsening of brain function.

Macular oedema

Before you start Fingolimod Mylan, if you have or have had visual disturbances or other signs of swelling in the central vision area (macula) at the back of the eye, inflammation or infection of the eye (uveitis) or diabetes, your doctor may want you to undergo an eye examination.

Your doctor may want you to undergo an eye examination 3 to 4 months after starting Fingolimod Mylan treatment.

The macula is a small area of the retina at the back of the eye which enables you to see shapes, colours and details clearly and sharply. Fingolimod Mylan may cause swelling in the macula, a condition that is known as macular oedema. The swelling usually happens in the first 4 months of treatment.

Your chance of developing macular oedema is higher if you have **diabetes** or have had an inflammation of the eye called uveitis. In these cases, your doctor will want you to undergo regular eye examinations in order to detect macular oedema.

If you have had macular oedema, talk to your doctor before you resume treatment with Fingolimod Mylan.

Macular oedema can cause some of the same vision symptoms as an MS attack (optic neuritis). Early on, there may not be any symptoms. Be sure to tell your doctor about any changes in your vision. Your doctor may want you to undergo an eye examination, especially if:

- the centre of your vision gets blurry or has shadows.
- you develop a blind spot in the centre of your vision.
- you have problems seeing colours or fine detail.

Liver function tests

If you have severe liver problems, you should not take Fingolimod Mylan. It may affect your liver function. You will probably not notice any symptoms but if you notice yellowing of your skin or the whites of your eyes, abnormally dark urine (brown coloured), pain on the right side of your stomach area (abdomen), tiredness, feeling less hungry than usual or unexplained nausea and vomiting, **tell your doctor straight away**.

If you get any of these symptoms after starting Fingolimod Mylan, **tell your doctor straight away**.

Before, during and after the treatment, your doctor will request blood tests to monitor your liver function. If your test results indicate a problem with your liver, you may have to interrupt treatment with Fingolimod Mylan.

High blood pressure

As Fingolimod Mylan causes a slight elevation of blood pressure, your doctor may want to check your blood pressure regularly.

Lung problems

Fingolimod Mylan has a slight effect on the lung function. Patients with severe lung problems or with smoker's cough may have a higher chance of developing side effects.

Blood count

The desired effect of Fingolimod Mylan treatment is to reduce the amount of white blood cells in your blood. This will usually go back to normal within 2 months of stopping treatment. If you need to have any blood tests, tell the doctor that you are taking this medicine. Otherwise, it may not be possible for the doctor to understand the results of the test and for certain types of blood test your doctor may need to take more blood than usual.

Before you start Fingolimod Mylan, your doctor will confirm whether you have enough white blood cells in your blood and may want to repeat a check regularly. In case you do not have enough white

blood cells, you may have to interrupt treatment.

Posterior reversible encephalopathy syndrome (PRES)

A condition called PRES has been rarely reported in MS patients treated with fingolimod. Symptoms may include sudden onset of severe headache, confusion, seizures and vision changes. Tell your doctor straight away if you experience any of these symptoms during your treatment, because it could be serious.

Cancer

Skin cancers have been reported in MS patients treated with fingolimod. Talk to your doctor straight away if you notice any skin nodules (e.g. shiny pearly nodules), patches or open sores that do not heal within weeks. Symptoms of skin cancer may include abnormal growth or changes of skin tissue (e.g. unusual moles) with a change in colour, shape or size over time. Before you start Fingolimod Mylan, a skin examination is required to check whether you have any skin nodules. Your doctor will also carry out regular skin examinations during your treatment. If you develop problems with your skin, your doctor may refer you to a dermatologist, who after consultation may decide that it is important for you to be seen on a regular basis.

A type of cancer of the lymphatic system (lymphoma) has been reported in MS patients treated with fingolimod.

Exposure to the sun and protection against the sun

Fingolimod weakens your immune system. This increases your risk of developing cancers, in particular skin cancers. You should limit your exposure to the sun and UV rays by:

- wearing appropriate protective clothing.
- regularly applying sunscreen with a high degree of UV protection.

Unusual brain lesions associated with MS relapse

Rare cases of unusually large brain lesions associated with MS relapse have been reported in patients treated with fingolimod. In case of severe relapse, your doctor will consider performing MRI to evaluate this condition and will decide whether you need to stop treatment.

Switch from other treatments to Fingolimod Mylan

Your doctor may switch you directly from beta interferon, glatiramer acetate or dimethyl fumarate to Fingolimod Mylan if there are no signs of abnormalities caused by your previous treatment. Your doctor may have to do a blood test in order to exclude such abnormalities. After stopping natalizumab you may have to wait for 2-3 months before starting treatment with Fingolimod Mylan. To switch from teriflunomide, your doctor may advise you to wait for a certain time or to go through an accelerated elimination procedure. If you have been treated with alemtuzumab, a thorough evaluation and discussion with your doctor is required to decide if Fingolimod Mylan is appropriate for you.

Women of childbearing potential

If used during pregnancy, Fingolimod Mylan can harm the unborn baby. Before you start treatment your doctor will explain the risk to you and ask you to do a pregnancy test in order to ensure that you are not pregnant. Your doctor will give you a card which explains why you should not become pregnant while taking Fingolimod Mylan. It also explains what you should do to avoid becoming pregnant while you are taking this medicine. You must use effective contraception during treatment and for 2 months after stopping treatment (see section “Pregnancy and breastfeeding”).

Worsening of MS after stopping Fingolimod Mylan treatment

Do not stop taking this medicine or change your dose without talking to your doctor first.

Tell your doctor straight away if you think your MS is getting worse after you have stopped treatment with Fingolimod Mylan. This could be serious (see “If you stop taking Fingolimod Mylan” in section 3, and also section 4, “Possible side effects”).

Elderly

Experience with fingolimod in elderly patients (over 65 years) is limited. Talk to your doctor if you have any concerns.

Children and adolescents

Fingolimod Mylan is not intended for use in children below 10 years old as it has not been studied in MS patients in this age group.

The warnings and precautions listed above also apply to children and adolescents. The following information is particularly important for children and adolescents and their caregivers:

- Before you start Fingolimod Mylan, your doctor will check your vaccination status. If you have not had certain vaccinations, it may be necessary for you to be given them before this medicine can be started.
- The first time you take Fingolimod Mylan, or when you switch from a 0.25 mg daily dose to a 0.5 mg daily dose, your doctor will monitor your heart rate and heartbeat (see ‘Slow heart rate (bradycardia) and irregular heartbeat’ above).
- If you experience convulsions or fits before or whilst taking Fingolimod Mylan, let your doctor know.
- If you suffer from depression or anxiety or if you become depressed or anxious while you are taking Fingolimod Mylan, let your doctor know. You may need to be monitored more closely.

Other medicines and Fingolimod Mylan

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. Tell your doctor if you are taking any of the following medicines:

- **Medicines that suppress or modulate the immune system**, including **other medicines used to treat MS**, such as beta interferon, glatiramer acetate, natalizumab, mitoxantrone, teriflunomide, dimethyl fumarate or alemtuzumab. You must not use Fingolimod Mylan together with such medicines as this could intensify the effect on the immune system (see also ‘Do not take Fingolimod Mylan’).
- **Corticosteroids**, due to a possible added effect on the immune system.
- **Vaccines**. If you need to receive a vaccine, seek your doctor’s advice first. During and for up to 2 months after treatment with Fingolimod Mylan, you should not receive certain types of vaccine (live attenuated vaccines) as they could trigger the infection that they were supposed to prevent. Other vaccines may not work as well as usual if given during this period.
- **Medicines that slow the heartbeat** (for example beta blockers, such as atenolol). Use of Fingolimod Mylan together with such medicines could intensify the effect on heartbeat in the first days after starting treatment.
- **Medicines for irregular heartbeat**, such as quinidine, disopyramide, amiodarone or sotalol. You must not use Fingolimod Mylan if you are taking such a medicine because it could intensify the effect on irregular heartbeat (see also ‘Do not take Fingolimod Mylan’).
- **Other medicines:**
 - protease inhibitors, anti-infectives such as ketoconazole,azole antifungals, clarithromycin or telithromycin.
 - carbamazepine, rifampicin, phenobarbital, phenytoin, efavirenz or St. John’s Wort (*Hypericum perforatum*) (potential risk of reduced efficacy of Fingolimod Mylan).

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

Pregnancy

Do not use Fingolimod Mylan during pregnancy, if you are trying to become pregnant or if you are a woman who could become pregnant and you are not using effective contraception. If this medicine is used during pregnancy, there is a risk of harm to the unborn baby. The rate of congenital malformations observed in babies exposed to fingolimod during pregnancy is about 2 times the rate observed in the general population (in whom the rate of congenital malformations is about 2-3%). The most frequently reported malformations included cardiac, renal and musculoskeletal malformations.

Therefore, if you are a woman of childbearing potential:

- before you start treatment with Fingolimod Mylan your doctor will inform you about the risk to an unborn baby and ask you to do a pregnancy test in order to ensure that you are not pregnant, and,
- you must use effective contraception while taking this medicine and for two months after you stop taking it to avoid becoming pregnant. Talk to your doctor about reliable methods of contraception.

Your doctor will give you a card which explains why you should not become pregnant while taking Fingolimod Mylan.

If you do become pregnant while taking Fingolimod Mylan, tell your doctor straight away. Your doctor will decide to stop treatment (see “If you stop taking Fingolimod Mylan” in section 3, and also section 4, “Possible side effects”). Specialised pre-natal monitoring will be performed.

Breast-feeding

You should not breast-feed while you are taking Fingolimod Mylan. It can pass into breast milk and there is a risk of serious side effects for the baby.

Driving and using machines

Your doctor will tell you whether your illness allows you to drive vehicles, including a bicycle and use machines safely. Fingolimod Mylan is not expected to have an influence on your ability to drive and use machines.

However, at initiation of treatment you will have to stay at the doctor’s surgery or clinic for 6 hours after taking the first dose of this medicine. Your ability to drive and use machines may be impaired during and potentially after this time period.

3. How to take Fingolimod Mylan

Treatment with Fingolimod Mylan will be overseen by a doctor who is experienced in the treatment of multiple sclerosis.

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

The recommended dose is:

Adults

The dose is one 0.5 mg capsule per day.

Children and adolescents (10 years of age and above)

The dose depends on body weight:

- *Children and adolescents with body weight equal to or below 40 kg:* one 0.25 mg capsule per day.
- *Children and adolescents with a body weight above 40 kg:* one 0.5 mg capsule per day.

Children and adolescents who start on one 0.25 mg capsule per day and later reach a stable body weight above 40 kg will be instructed by their doctor to switch to one 0.5 mg capsule per day. In this case, it is recommended to repeat the first-dose observation period.

Fingolimod Mylan is only available as 0.5 mg hard capsules which are not suitable for children and adolescents with a body weight equal to or below 40 kg.

Other fingolimod-containing medicines are available in 0.25 mg strength.

Ask your doctor or pharmacist.

Do not exceed the recommended dose.

Fingolimod Mylan is for oral use.

Take Fingolimod Mylan once a day with a glass of water. The capsules should always be swallowed intact, without opening them. This medicine can be taken with or without food.

Taking Fingolimod Mylan at the same time each day will help you remember when to take your medicine.

If you have questions about how long to take this medicine, talk to your doctor or your pharmacist.

If you take more Fingolimod Mylan than you should

If you have taken too much call your doctor straight away.

If you forget to take Fingolimod Mylan

If you have been taking this medicine for less than one month and you forget to take one dose for a whole day, call your doctor before you take the next dose. Your doctor may decide to keep you under observation at the time you take the next dose.

If you have been taking Fingolimod Mylan for at least one month and have forgotten to take your treatment for more than 2 weeks, call your doctor before you take the next dose. Your doctor may decide to keep you under observation at the time you take the next dose. However, if you have forgotten to take your treatment for up to 2 weeks, you can take the next dose as planned.

Never take a double dose to make up for a forgotten dose.

If you stop taking Fingolimod Mylan

Do not stop taking this medicine or change your dose without talking to your doctor first.

Fingolimod Mylan will stay in your body for up to 2 months after you stop taking it. Your white blood cell count (lymphocyte count) may also remain low during this time and the side effects described in this leaflet may still occur. After stopping this medicine you may have to wait for 6-8 weeks before starting a new MS treatment.

If you have to restart Fingolimod Mylan more than 2 weeks after you stop taking it, the effect on heart rate normally seen when treatment is first started may re-occur and you will need to be monitored at the doctor's surgery or clinic for re-initiation of treatment. Do not restart this medicine after stopping it for more than two weeks without seeking advice from your doctor.

Your doctor will decide whether and how you need to be monitored after stopping Fingolimod Mylan. Tell your doctor straight away if you think your MS is getting worse after you have stopped treatment. This could be serious.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Some side effects could be or could become serious:

Common (may affect up to 1 in 10 people)

- Coughing with phlegm, chest discomfort, fever (signs of lung disorders)
- Herpes virus infection (shingles or herpes zoster) with symptoms such as blisters, burning, itching or pain of the skin, typically on the upper body or the face. Other symptoms may be

fever and weakness in the early stages of infection, followed by numbness, itching or red patches with severe pain

- Slow heartbeat (bradycardia), irregular heart rhythm
- A type of skin cancer called basal cell carcinoma (BCC) which often appears as a pearly nodule, although it can also take other forms
- Depression and anxiety are known to occur with increased frequency in the MS population and have also been reported in paediatric patients treated with fingolimod
- Weight loss

Uncommon (may affect up to 1 in 100 people)

- Pneumonia with symptoms such as fever, cough, difficulty breathing
- Macular oedema (swelling in the central vision area of the retina at the back of the eye) with symptoms such as shadows or blind spot in the centre of the vision, blurred vision, problems seeing colours or details
- Reduction in blood platelets which increases risk of bleeding or bruising
- Malignant melanoma (a type of skin cancer which usually develops from an unusual mole). Possible signs of melanoma include moles which may change size, shape, elevation or colour over time, or new moles. The moles may itch, bleed or ulcerate
- Convulsion, fits (more frequent in children and adolescents than in adults)

Rare (may affect up to 1 in 1 000 people)

- A condition called posterior reversible encephalopathy syndrome (PRES). Symptoms may include sudden onset of severe headache, confusion, seizures and/or vision disturbances
- Lymphoma (a type of cancer that affects the lymph system)
- Squamous cell carcinoma: a type of skin cancer which may present as a firm red nodule, a sore with crust, or a new sore on an existing scar

Very rare (may affect up to 1 in 10 000 people)

- Electrocardiogram anomaly (T-wave inversion)
- Tumour related to infection with human herpes virus 8 (Kaposi's sarcoma)

Not known (frequency cannot be estimated from the available data)

- Allergic reactions, including symptoms of rash or itchy hives, swelling of lips, tongue or face, which are more likely to occur on the day you start Fingolimod Mylan treatment
- Signs of liver disease (including liver failure), such as yellowing of your skin or the whites of your eyes (jaundice), nausea or vomiting, pain on the right side of your stomach area (abdomen), dark urine (brown coloured), feeling less hungry than usual, tiredness and abnormal liver function tests. In a very small number of cases, liver failure could lead to liver transplantation
- Risk of a rare brain infection called progressive multifocal leukoencephalopathy (PML). The symptoms of PML may be similar to an MS relapse. Symptoms might also arise that you might not become aware of by yourself, such as changes in mood or behaviour, memory lapses, speech and communication difficulties, which your doctor may need to investigate further to rule out PML. Therefore, if you believe your MS is getting worse or if you or those close to you notice any new or unusual symptoms, it is very important that you speak to your doctor as soon as possible
- Inflammatory disorder after stopping Fingolimod Mylan treatment (known as immune reconstitution inflammatory syndrome or IRIS)
- Cryptococcal infections (a type of fungal infection), including cryptococcal meningitis with symptoms such as headache accompanied by stiff neck, sensitivity to light, nausea and/or confusion
- Merkel cell carcinoma (a type of skin cancer). Possible signs of Merkel cell carcinoma include flesh-coloured or bluish-red, painless nodule, often on the face, head or neck. Merkel cell carcinoma can also present as a firm painless nodule or mass. Long-term exposure to the sun and a weak immune system can affect the risk of developing Merkel cell carcinoma

- After Fingolimod Mylan treatment is stopped, symptoms of MS can return and may become worse than they were before or during treatment.
- Autoimmune form of anaemia (decreased amount of red blood cells) where red blood cells are destroyed (autoimmune haemolytic anaemia).

If you experience any of these, **tell your doctor straight away**.

Other side effects

Very common (may affect more than 1 in 10 people)

- Infection from flu virus with symptoms such as tiredness, chills, sore throat, aching in the joints or muscles, fever
- Feeling of pressure or pain in the cheeks and forehead (sinusitis)
- Headache
- Diarrhoea
- Back pain
- Blood testing showing higher levels of liver enzymes
- Cough

Common (may affect up to 1 in 10 people)

- Ringworm, a fungal infection of the skin (tinea versicolor)
- Dizziness
- Severe headache often accompanied by nausea, vomiting and sensitivity to light (migraine)
- Low level of white blood cells (lymphocytes, leucocytes)
- Weakness
- Itchy, red, burning rash (eczema)
- Itching
- Blood fat (triglycerides) level increased
- Hair loss
- Breathlessness
- Depression
- Blurred vision (see also the section on macular oedema under ‘Some side effects could be or could become serious’)
- Hypertension (Fingolimod Mylan may cause a mild increase in blood pressure)
- Muscle pain
- Joint pain

Uncommon (may affect up to 1 in 100 people)

- Low level of certain white blood cells (neutrophils)
- Depressed mood
- Nausea

Rare (may affect up to 1 in 1 000 people)

- Cancer of the lymphatic system (lymphoma)

Not known (frequency cannot be estimated from the available data)

- Peripheral swelling

If any of these affects you severely, **tell your doctor**

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via **the national reporting system** listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Fingolimod Mylan

Keep this medicine out of the sight and reach of children.

- Do not use this medicine after the expiry date which is stated on the carton and blister foil/bottle after 'EXP'. The expiry date refers to the last day of that month.
- Do not store above 25°C.
- Store in the original package in order to protect from moisture.
- Do not use this medicine if you notice that the pack is damaged or shows signs of tampering.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Fingolimod Mylan contains

- The active substance is fingolimod. Each capsule contains 0.5 mg fingolimod (as hydrochloride).
- The other ingredients are:
 - Capsule content: calcium hydrogen phosphate dihydrate, glycine, colloidal anhydrous silica and magnesium stearate.
 - Capsule shell: gelatin, titanium dioxide (E171), yellow iron oxide (E172) and red iron oxide (E172).
 - Printing ink: shellac (E904), propylene glycol (E1520), black iron oxide (E172) and potassium hydroxide.

What Fingolimod Mylan looks like and contents of the pack

A hard capsule (capsule) with brown-orange opaque cap and white opaque body, printed with 'MYLAN' over 'FD 0.5' in black ink on both the cap and body.

Fingolimod Mylan 0.5 mg capsules are available in:

Blister packs containing 28, 30, 84 or 98 hard capsules

Multipacks comprising 3 cartons, each containing 28 hard capsules

Calendar packs containing 28 or 84 hard capsules

Unit dose blister packs containing 7 x 1, 28 x 1, 90 x 1 or 98 x 1 hard capsules

Bottle packs containing 90 or 100 hard capsules

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Mylan Pharmaceuticals Limited, Damastown Industrial Park, Mulhuddart, Dublin 15, DUBLIN, Ireland

Manufacturer(s)

Mylan Hungary Kft, Mylan utca 1, Komarom, H-2900, Hungary

Viartis Germany GmbH, Zweigniederlassung Bad Homburg v. d. Hoehe, Benzstrasse 1, Bad Homburg v. d. Hoehe, Hessen, 61352, Germany.

For any other information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

Viатris

Tél/Tel: + 32 (0)2 658 61 00

България

Виатрис ЕООД

Тел: +359 2 44 55 400

Česká republika

Viатris CZ s.r.o.

Tel: + 420 222 004 400

Danmark

Viатris ApS

Tlf: +45 28 11 69 32

Deutschland

Viатris Healthcare GmbH

Tel: +49 800 0700 800

Eesti

Viатris OÜ

Tel: + 372 6363 052

Ελλάδα

Viатris Hellas Ltd

Τηλ: +30 2100 100 002

España

Viатris Pharmaceuticals, S.L.

Tel: + 34 900 102 712

France

Viатris Santé

Tél: +33 4 37 25 75 00

Hrvatska

Viатris Hrvatska d.o.o.

Tel: +385 1 23 50 599

Ireland

Viатris Limited

Tel: +353 1 8711600

Ísland

Icepharma hf.

Sími: +354 540 8000

Italia

Viатris Italia S.r.l.

Tel: + 39 (0) 2 612 46921

Κύπρος

CPO Pharmaceuticals Limited

Τηλ: +357 22863100

Lietuva

Viатris UAB

Tel: +370 5 205 1288

Luxembourg/Luxemburg

Viатris

Tél/Tel: + 32 (0)2 658 61 00

(Belgique/Belgien)

Magyarország

Viатris Healthcare Kft.

Tel.: + 36 1 465 2100

Malta

V.J. Salomone Pharma Ltd

Tel: + 356 21 22 01 74

Nederland

Mylan BV

Tel: +31 (0)20 426 3300

Norge

Viатris AS

Tlf: + 47 66 75 33 00

Österreich

Viатris Austria GmbH

Tel: +43 1 86390

Polska

Viатris Healthcare Sp. z o.o.

Tel: + 48 22 546 64 00

Portugal

Mylan, Lda.

Tel: + 351 214 127 200

România

BGP Products SRL

Tel: +40 372 579 000

Slovenija

Viатris d.o.o.

Tel: + 386 1 23 63 180

Slovenská republika

Viатris Slovakia s.r.o.

Tel: +421 2 32 199 100

Suomi/Finland

Viатris Oy

Puh/Tel: +358 20 720 9555

Sverige

Viатris AB

Tel: + 46 (0)8 630 19 00

Latvija
Viatis SIA
Tel: +371 676 055 80

This leaflet was last revised in

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency website:
<http://www.ema.europa.eu>.