

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Glyxambi 10 mg/5 mg film-coated tablets  
Glyxambi 25 mg/5 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Glyxambi 10 mg/5 mg film-coated tablets

Each film-coated tablet contains 10 mg empagliflozin and 5 mg linagliptin.

Glyxambi 25 mg/5 mg film-coated tablets

Each film-coated tablet contains 25 mg empagliflozin and 5 mg linagliptin.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet (tablet).

Glyxambi 10 mg/5 mg film-coated tablets

Pale yellow, arc triangular, flat faced, bevel-edged, film-coated tablets. One side is debossed with the Boehringer Ingelheim company symbol, the other side is debossed with “10/5” (tablet dimensions: 8 mm each side).

Glyxambi 25 mg/5 mg film-coated tablets

Pale pink, arc triangular, flat faced, bevel-edged, film-coated tablets. One side is debossed with the Boehringer Ingelheim company symbol, the other side is debossed with “25/5” (tablet dimensions: 8 mm each side).

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Glyxambi, fixed dose combination of empagliflozin and linagliptin, is indicated in adults aged 18 years and older with type 2 diabetes mellitus:

- to improve glycaemic control when metformin and/or sulphonylurea (SU) and one of the monocomponents of Glyxambi do not provide adequate glycaemic control
- when already being treated with the free combination of empagliflozin and linagliptin

(See sections 4.2, 4.4, 4.5 and 5.1 for available data on combinations studied)

### 4.2 Posology and method of administration

#### Posology

The recommended starting dose is one film-coated tablet of Glyxambi 10 mg/5 mg (10 mg empagliflozin plus 5 mg linagliptin) once daily.

In patients who tolerate this starting dose and require additional glycaemic control, the dose can be increased to one film-coated tablet of Glyxambi 25 mg/5 mg (25 mg empagliflozin plus 5 mg

linagliptin) once daily.

When Glyxambi is used in combination with metformin, the metformin dose should be continued.

When Glyxambi is used in combination with a sulphonylurea or with insulin, a lower dose of the sulphonylurea or insulin may be considered to reduce the risk of hypoglycaemia (see sections 4.4, 4.5 and 4.8).

Patients switching from empagliflozin (either 10 mg or 25 mg daily dose) and linagliptin (5 mg daily dose) to Glyxambi should receive the same daily dose of empagliflozin and linagliptin in the fixed dose combination as in separate tablets.

#### Missed doses

If a dose is missed, and it is 12 hours or more until the next dose, the dose should be taken as soon as the patient remembers. The next dose should be taken at the usual time. If a dose is missed, and it is less than 12 hours until the next dose, the dose should be skipped and the next dose should be taken at the usual time. A double dose should not be taken to compensate for a forgotten dose.

#### Special populations

##### *Renal impairment*

The glycaemic efficacy of empagliflozin is dependent on renal function. For cardiovascular risk reduction as add on to standard of care, a dose of 10 mg empagliflozin once daily should be used in patients with an eGFR below 60 ml/min/1.73 m<sup>2</sup> (see Table 1). Because the glycaemic lowering efficacy of empagliflozin is reduced in patients with moderate renal impairment and likely absent in patients with severe renal impairment, if further glycaemic control is needed, the addition of other anti-hyperglycaemic agents should be considered.

For dose adjustment recommendations according to eGFR or CrCL refer to Table 1.

Table 1: Dose adjustment recommendations<sup>a</sup>

eGFR [ml/min/1.73 m <sup>2</sup> ] or CrCL [ml/min]	Empagliflozin	Linagliptin
≥60	Initiate with 10 mg.  In patients tolerating 10 mg and requiring additional glycaemic control, the dose can be increased to 25 mg.	5 mg  No dose adjustment for linagliptin is required.
45 to <60	Initiate with 10 mg. <sup>b</sup>  Continue with 10 mg in patients already taking empagliflozin.	
30 to <45	Initiate with 10 mg. <sup>b</sup>  Continue with 10 mg in patients already taking empagliflozin. <sup>b</sup>	
<30	Empagliflozin is not recommended.	

<sup>a</sup> See sections 4.4, 4.8, 5.1 and 5.2

<sup>b</sup> patients with type 2 diabetes mellitus and established cardiovascular disease

Glyxambi should not be used in patients with end stage renal disease (ESRD) or in patients on dialysis, as there are insufficient data on empagliflozin to support use in these patients (see sections 4.4, 5.1 and 5.2).

### *Hepatic impairment*

No dose adjustment is required in patients with mild to moderate hepatic impairment.

Empagliflozin exposure is increased in patients with severe hepatic impairment and therapeutic experience in such patients is limited (see section 5.2). Therefore, Glyxambi is not recommended for use in this population.

### *Elderly*

No dose adjustment based on age is required. However, renal function and risk of volume depletion should be taken into account in elderly patients (see sections 4.4 and 4.8). Based on very limited experience in patients 75 years and older, initiation of Glyxambi therapy is not recommended in this population (see sections 4.4 and 5.2).

### *Paediatric population*

Safety and efficacy of Glyxambi in paediatric patients below 18 years of age have not been established. No data are available.

### Method of administration

Glyxambi tablets are for oral use and can be taken with or without a meal at any time of the day at regular intervals. The tablets should be swallowed whole with water.

## **4.3 Contraindications**

Hypersensitivity to the active substances, to any other Sodium-Glucose-Co-Transporter-2 (SGLT2) inhibitor, to any other Dipeptidyl-Peptidase-4 (DPP-4) inhibitor, or to any of the excipients listed in section 6.1.

## **4.4 Special warnings and precautions for use**

### Diabetic ketoacidosis

Rare cases of diabetic ketoacidosis (DKA), including life-threatening and fatal cases, have been reported in patients treated with SGLT2 inhibitors, including empagliflozin. In a number of cases, the presentation of the condition was atypical with only moderately increased blood glucose values, below 14 mmol/L (250 mg/dL). It is not known if DKA is more likely to occur with higher doses of empagliflozin.

The risk of DKA must be considered in the event of non-specific symptoms such as nausea, vomiting, anorexia, abdominal pain, excessive thirst, difficulty breathing, confusion, unusual fatigue or sleepiness. Patients should be assessed for ketoacidosis immediately if these symptoms occur, regardless of blood glucose level.

In patients where DKA is suspected or diagnosed, treatment with empagliflozin should be discontinued immediately.

Treatment should be interrupted in patients who are hospitalised for major surgical procedures or acute serious medical illnesses. Monitoring of ketones is recommended in these patients. Measurement of blood ketone levels is preferred to urine. Treatment with empagliflozin may be restarted when the ketone values are normal and the patient's condition has stabilised.

Before initiating empagliflozin, factors in the patient history that may predispose to ketoacidosis should be considered.

Patients who may be at higher risk of DKA include patients with a low beta-cell function reserve (e.g. type 2 diabetes patients with low C-peptide or latent autoimmune diabetes in adults (LADA) or patients with a history of pancreatitis), patients with conditions that lead to restricted food intake or

severe dehydration, patients for whom insulin doses are reduced and patients with increased insulin requirements due to acute medical illness, surgery or alcohol abuse. SGLT2 inhibitors should be used with caution in these patients.

Restarting SGLT2 inhibitor treatment in patients with previous DKA while on SGLT2 inhibitor treatment is not recommended, unless another clear precipitating factor is identified and resolved.

Glyxambi should not be used for treatment of patients with type 1 diabetes. Data from a clinical trial program in patients with type 1 diabetes showed increased DKA occurrence with common frequency in patients treated with empagliflozin 10 mg and 25 mg as an adjunct to insulin compared to placebo.

### Renal impairment

In patients with an eGFR below 60 mL/min/1.73 m<sup>2</sup> or CrCl <60 mL/min, the daily dose of empagliflozin/linagliptin is limited to 10 mg/5 mg (see section 4.2). Empagliflozin/linagliptin is not recommended when eGFR is below 30 mL/min/1.73 m<sup>2</sup> or CrCl is below 30 mL/min. Empagliflozin/linagliptin should not be used in patients with ESRD or in patients on dialysis. There are insufficient data to support use in these patients (see sections 4.2, 5.1 and 5.2).

### Monitoring of renal function

Assessment of renal function is recommended as follows:

- prior to empagliflozin/linagliptin initiation and periodically during treatment, i.e. at least yearly (see sections 4.2, 5.1 and 5.2).
- prior to initiation of any concomitant medicinal product that may have a negative impact on renal function.

### Hepatic injury

Cases of hepatic injury have been reported with empagliflozin in clinical trials. A causal relationship between empagliflozin and hepatic injury has not been established.

### Elevated haematocrit

Haematocrit increase was observed with empagliflozin treatment (see section 4.8).

### Chronic kidney disease

There is experience with empagliflozin for the treatment of diabetes in patients with chronic kidney disease (eGFR  $\geq$ 30 mL/min/1.73 m<sup>2</sup>) both with and without albuminuria. Patients with albuminuria may benefit more from treatment with empagliflozin.

### Risk for volume depletion

Based on the mode of action of SGLT2 inhibitors, osmotic diuresis accompanying therapeutic glucosuria may lead to a modest decrease in blood pressure (see section 5.1). Therefore, caution should be exercised in patients for whom an empagliflozin-induced drop in blood pressure could pose a risk, such as patients with known cardiovascular disease, patients on anti-hypertensive therapy (e.g. thiazide and loop diuretics, see also section 4.5) with a history of hypotension or patients aged 75 years and older.

In case of conditions that may lead to fluid loss (e.g. gastrointestinal illness), careful monitoring of volume status (e.g. physical examination, blood pressure measurements, laboratory tests including haematocrit) and electrolytes is recommended for patients receiving empagliflozin. Temporary interruption of treatment with Glyxambi should be considered until the fluid loss is corrected.

## Elderly

A higher risk of volume depletion adverse reactions were reported in patients aged 75 years and older, treated with empagliflozin, especially at 25 mg/day (see section 4.8). Therefore, special attention should be given to their volume intake in case of co-administered medicinal products which may lead to volume depletion (e.g. diuretics, ACE inhibitors). Therapeutic experience is limited with Glyxambi in patients > 75 years of age, and, no experience is available in patients aged 85 years and older. Initiation of therapy with Glyxambi in this population is not recommended (see section 4.2).

## Urinary tract infections

In Glyxambi clinical trials, the incidence of urinary tract infections was overall similar between the patients treated with Glyxambi and the patients treated with empagliflozin or linagliptin. The frequencies were comparable to the incidence of urinary tract infections in empagliflozin clinical trials (see section 4.8).

In a pool of placebo-controlled double-blind trials of 18 to 24 weeks duration, the overall frequency of urinary tract infection reported as adverse event was similar in patients treated with empagliflozin 25 mg and placebo and higher in patients treated with empagliflozin 10 mg (see section 4.8). Post-marketing cases of complicated urinary tract infections including pyelonephritis and urosepsis have been reported in patients treated with empagliflozin. Pyelonephritis and urosepsis were not reported from the clinical trials in patients treated with Glyxambi. However, temporary interruption of Glyxambi should be considered in patients with complicated urinary tract infections.

## Necrotising fasciitis of the perineum (Fournier's gangrene)

Post-marketing cases of necrotising fasciitis of the perineum, (also known as Fournier's gangrene), have been reported in female and male patients taking SGLT2 inhibitors. This is a rare but serious and potentially life-threatening event that requires urgent surgical intervention and antibiotic treatment.

Patients should be advised to seek medical attention if they experience a combination of symptoms of pain, tenderness, erythema, or swelling in the genital or perineal area, with fever or malaise. Be aware that either uro-genital infection or perineal abscess may precede necrotising fasciitis. If Fournier's gangrene is suspected, Glyxambi should be discontinued and prompt treatment (including antibiotics and surgical debridement) should be instituted.

## Lower limb amputations

An increase in cases of lower limb amputation (primarily of the toe) has been observed in long-term clinical trials with another SGLT2 inhibitor. It is unknown whether this constitutes a class effect. Like for all diabetic patients it is important to counsel patients on routine preventative foot-care.

## Cardiac failure

Experience with empagliflozin in New York Heart Association (NYHA) class I-II is limited, and there is no experience in clinical trials with empagliflozin in NYHA class III-IV. In the EMPA-REG OUTCOME trial, 10.1% of the patients were reported with cardiac failure at baseline. The reduction of cardiovascular death in these patients was consistent with the overall trial population.

## Urine laboratory assessments

Due to the mechanism of action of empagliflozin, patients taking Glyxambi will test positive for glucose in their urine.

### Interference with 1,5-anhydroglucitol (1,5-AG) assay

Monitoring glycaemic control with 1,5-AG assay is not recommended as measurements of 1,5-AG are unreliable in assessing glycaemic control in patients taking SGLT2 inhibitors. Use of alternative methods to monitor glycaemic control is advised.

### Acute pancreatitis

Use of dipeptidyl peptidase-4 (DPP-4) inhibitors has been associated with a risk of developing acute pancreatitis. Acute pancreatitis has been observed in patients taking linagliptin. In a cardiovascular and renal safety trial (CARMELINA) with median observation period of 2.2 years, adjudicated acute pancreatitis was reported in 0.3% of patients treated with linagliptin and in 0.1% of patients treated with placebo. Patients should be informed of the characteristic symptoms of acute pancreatitis.

If pancreatitis is suspected, Glyxambi should be discontinued; if acute pancreatitis is confirmed, Glyxambi should not be restarted. Caution should be exercised in patients with a history of pancreatitis.

### Bullous pemphigoid

Bullous pemphigoid has been observed in patients taking linagliptin. In the CARMELINA trial, bullous pemphigoid was reported in 0.2% of patients on treatment with linagliptin and in no patient on placebo. If bullous pemphigoid is suspected, Glyxambi should be discontinued.

### Use with medicinal products known to cause hypoglycaemia

Empagliflozin and linagliptin as single agents showed an incidence of hypoglycaemia comparable to placebo when used alone or in combination with other antidiabetics not known to cause hypoglycaemia (e.g. metformin, thiazolidinediones). When used in combination with antidiabetics known to cause hypoglycaemia (e.g. sulphonylureas and/or insulin), the incidence of hypoglycaemia of both agents was increased (see section 4.8).

There are no data about the hypoglycaemic risk of Glyxambi when used with insulin and/or sulphonylurea. However, caution is advised when Glyxambi is used in combination with antidiabetics. A dose reduction of the sulphonylurea or insulin may be considered (see section 4.2 and 4.5).

## **4.5 Interaction with other medicinal products and other forms of interaction**

No drug interaction studies have been performed with Glyxambi and other medicinal products; however, such studies have been conducted with the individual active substances. Based on results of pharmacokinetic studies, no dose adjustment of Glyxambi is recommended when co-administered with commonly prescribed medicinal products, except those mentioned below.

### Pharmacodynamic interactions

#### *Insulin and sulphonylureas*

Insulin and sulphonylureas may increase the risk of hypoglycaemia. Therefore, a lower dose of insulin or sulphonylureas may be required to reduce the risk of hypoglycaemia when used in combination with Glyxambi (see sections 4.2, 4.4 and 4.8).

#### *Diuretics*

Empagliflozin may add to the diuretic effect of thiazide and loop diuretics and may increase the risk of dehydration and hypotension (see section 4.4).

## Pharmacokinetic interactions

### Effects of other medicinal products on empagliflozin

Empagliflozin is mainly excreted unchanged. A minor fraction is metabolised via uridine 5'-diphosphoglucuronosyltransferases (UGT); therefore, a clinically relevant effect of UGT inhibitors on empagliflozin is not expected (see section 5.2). The effect of UGT induction on empagliflozin (e.g. induction by rifampicin or phenytoin) has not been studied. Co-treatment with known inducers of UGT enzymes is not recommended due to a potential risk of decreased efficacy of empagliflozin. If an inducer of these UGT enzymes must be co-administered, monitoring of glycaemic control to assess response to Glyxambi is appropriate.

Co-administration of empagliflozin with probenecid, an inhibitor of UGT enzymes and OAT3, resulted in a 26% increase in peak empagliflozin plasma concentrations ( $C_{max}$ ) and a 53% increase in area under the concentration-time curve (AUC). These changes were not considered to be clinically meaningful.

An interaction study with gemfibrozil, an *in vitro* inhibitor of OAT3 and OATP1B1/1B3 transporters, showed that empagliflozin  $C_{max}$  increased by 15% and AUC increased by 59% following co-administration. These changes were not considered to be clinically meaningful.

Inhibition of OATP1B1/1B3 transporters by co-administration with rifampicin resulted in a 75% increase in  $C_{max}$  and a 35% increase in AUC of empagliflozin. These changes were not considered to be clinically meaningful.

Interaction studies suggest that the pharmacokinetics of empagliflozin were not influenced by co-administration with metformin, glimepiride, pioglitazone, sitagliptin, linagliptin, warfarin, verapamil, ramipril, simvastatin, torasemide and hydrochlorothiazide.

### Effects of empagliflozin on other medicinal products

Empagliflozin may increase renal lithium excretion and the blood lithium levels may be decreased. Serum concentration of lithium should be monitored more frequently after empagliflozin initiation and dose changes. Please refer the patient to the lithium prescribing doctor in order to monitor serum concentration of lithium.

Interaction studies conducted in healthy volunteers suggest that empagliflozin had no clinically relevant effect on the pharmacokinetics of metformin, glimepiride, pioglitazone, sitagliptin, linagliptin, simvastatin, warfarin, ramipril, digoxin, diuretics and oral contraceptives.

### Effects of other medicinal products on linagliptin

Co-administration of rifampicin decreased linagliptin exposure by 40%, suggesting that the efficacy of linagliptin may be reduced when administered in combination with a strong P-glycoprotein (P-gp) or cytochrome P450 (CYP) isozyme CYP3A4 inducer, particularly if these are administered long-term (see section 5.2). Co-administration with other potent inducers of P-gp and CYP3A4, such as carbamazepine, phenobarbital and phenytoin, has not been studied.

Co-administration of a single 5 mg oral dose of linagliptin and multiple 200 mg oral doses of ritonavir, a potent inhibitor of P-glycoprotein and CYP3A4, increased the AUC and  $C_{max}$  of linagliptin approximately twofold and threefold, respectively. The unbound concentrations, which are usually less than 1% at the therapeutic dose of linagliptin, were increased 4 to 5-fold after co-administration with ritonavir. Simulations of steady-state plasma concentrations of linagliptin with and without ritonavir indicated that the increase in exposure will be not associated with an increased accumulation. These changes in linagliptin pharmacokinetics were not considered to be clinically relevant. Therefore, clinically relevant interactions would not be expected with other P-glycoprotein/CYP3A4 inhibitors.



Interaction studies conducted in healthy volunteers suggest that the pharmacokinetics of linagliptin were not influenced by co-administration with metformin and glibenclamide.

#### Effects of linagliptin on other medicinal products

Linagliptin is a weak competitive and a weak to moderate mechanism-based inhibitor of CYP isozyme CYP3A4, but does not inhibit other CYP isozymes. It is not an inducer of CYP isozymes. Linagliptin is a P-glycoprotein substrate, and inhibits P-glycoprotein mediated transport of digoxin with low potency.

Linagliptin had no clinically relevant effect on the pharmacokinetics of metformin, glibenclamide, simvastatin, pioglitazone, warfarin, digoxin, empagliflozin or oral contraceptives providing *in vivo* evidence of a low propensity for causing drug interactions with substrates of CYP3A4, CYP2C9, CYP2C8, P-gp and organic cationic transporter (OCT).

### **4.6 Fertility, pregnancy and lactation**

#### Pregnancy

There are no data from the use of empagliflozin and linagliptin in pregnant women. Animal studies have shown that empagliflozin and linagliptin cross the placenta during late gestation, but do not indicate direct or indirect harmful effects with respect to early embryonic development with either empagliflozin or linagliptin (see section 5.3). Animal studies with empagliflozin have shown adverse effects on postnatal development (see section 5.3). As a precautionary measure it is preferable to avoid the use of Glyxambi during pregnancy.

#### Breast-feeding

No data in humans are available on excretion of empagliflozin and linagliptin into milk. Available non-clinical data in animals have shown excretion of empagliflozin and linagliptin in milk. A risk to newborns or infants cannot be excluded. Glyxambi should not be used during breast-feeding.

#### Fertility

No trials on the effect on human fertility have been conducted with Glyxambi or with the individual active substances. Non-clinical studies with empagliflozin and linagliptin as single agents do not indicate direct or indirect harmful effects with respect to fertility (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

Glyxambi has minor influence on the ability to drive and use machines. Patients should be advised to take precautions to avoid hypoglycaemia while driving and using machines, in particular when Glyxambi is used in combination with other antidiabetic medicinal products known to cause hypoglycaemia (e.g. insulin and analogues, sulphonylureas).

### **4.8 Undesirable effects**

#### Summary of the safety profile

The most frequent adverse reaction was urinary tract infection (7.5% with Glyxambi 10 mg empagliflozin/5 mg linagliptin and 8.5% with Glyxambi 25 mg empagliflozin/5 mg linagliptin) (see Description of selected adverse reactions). The most serious adverse reactions were ketoacidosis (< 0.1%), pancreatitis (0.2 %), hypersensitivity (0.6 %), and hypoglycaemia (2.4 %) (see section 4.4).

Overall, the safety profile of Glyxambi was in line with the safety profiles of the individual active substances (empagliflozin and linagliptin). No additional adverse reactions were identified with Glyxambi.

### Tabulated list of adverse reactions

The adverse reactions shown in the table below (see Table 2) are listed by system organ class and are based on the safety profiles of empagliflozin and linagliptin monotherapy. Frequency categories are defined as very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1\ 000$  to  $< 1/100$ ), rare ( $\geq 1/10\ 000$  to  $< 1/1\ 000$ ), very rare ( $< 1/10\ 000$ ) and not known (cannot be estimated from the available data).

Table 2 Tabulated list of adverse reactions (MedDRA) from reported placebo-controlled trials and from post-marketing experience

System organ class	Frequency	Adverse reaction
Infections and infestations	Common	Urinary tract infection <sup>1,*</sup> (including pyelonephritis and urosepsis) <sup>4</sup>
	Common	Vaginal moniliasis, vulvovaginitis, balanitis and other genital infections <sup>1,*</sup>
	Common	Nasopharyngitis <sup>2</sup>
	Rare	Necrotising fasciitis of the perineum (Fournier's gangrene) <sup>#</sup>
Immune system disorders	Uncommon	Hypersensitivity <sup>2</sup>
	Uncommon	Angioedema <sup>3,4</sup> , urticaria <sup>3,4</sup>
Metabolism and nutrition disorders	Common	Hypoglycaemia (when used with sulphonylurea or insulin) <sup>*</sup>
	Common	Thirst
	Rare	Diabetic ketoacidosis <sup>4,#</sup>
Vascular disorders	Uncommon	Volume depletion <sup>1,*b</sup>
Respiratory, thoracic and mediastinal disorders	Common	Cough <sup>2</sup>
Gastrointestinal disorders	Common	Constipation
	Uncommon	Pancreatitis <sup>2</sup>
	Rare	Mouth ulceration <sup>3</sup>
Skin and subcutaneous tissue disorders	Common	Pruritus <sup>1</sup>
	Common	Rash <sup>3,4</sup>
	Not known	Bullous pemphigoid <sup>2,a</sup>
Renal and urinary disorders	Common	Increased urination <sup>1,*</sup>
	Uncommon	Dysuria <sup>1</sup>
	Very rare	Tubulointerstitial nephritis <sup>4</sup>
Investigations	Common	Amylase increased <sup>2</sup>
	Common	Lipase increased <sup>2</sup>
	Uncommon	Haematocrit increased <sup>1,5</sup>
	Uncommon	Serum lipids increased <sup>1,6</sup>
	Uncommon	Blood creatinine increased/Glomerular filtration rate decreased <sup>1,*</sup>

<sup>1</sup> derived from empagliflozin experiences

<sup>2</sup> derived from linagliptin experiences

<sup>3</sup> derived from linagliptin postmarketing experience

<sup>4</sup> derived from empagliflozin postmarketing experience

<sup>5</sup> Mean changes from baseline in haematocrit were 3.3% and 4.2% for Glyxambi 10 mg/5 mg and 25 mg/5 mg, respectively, compared to 0.2% for placebo. In a clinical trial with empagliflozin, haematocrit values returned towards baseline values after a follow-up period of 30 days after treatment stop.

<sup>6</sup> Mean percent increases from baseline for Glyxambi 10 mg/5 mg and 25 mg/5 mg versus placebo, respectively, were total cholesterol 3.2% and 4.6% versus 0.5%; HDL-cholesterol 8.5% and 6.2% versus 0.4%; LDL-cholesterol 5.8% and 11.0% versus 3.3%; triglycerides -0.5% and 3.3% versus 6.4%.

<sup>a</sup> In the CARMELINA trial (see section 5.1), bullous pemphigoid was reported in 0.2% patients treated with linagliptin and in no patients treated with placebo.

<sup>b</sup> Pooled data of empagliflozin trials in patients with heart failure (where half of the patients had type 2 diabetes mellitus) showed a higher frequency of volume depletion ("very common": 11.4% for empagliflozin versus 9.7% for placebo).

<sup>#</sup> see section 4.4

\* see subsection below for additional information

## Description of selected adverse reactions

### Hypoglycaemia

In pooled clinical trials of Glyxambi in patients with type 2 diabetes and inadequate glycaemic control on background metformin, the frequency of the reported hypoglycaemic events was 2.4%. The incidence of confirmed hypoglycaemic events was low (< 1.5%). There was no notable difference of the incidence in patients treated with different dose strengths of Glyxambi compared to the treatment with empagliflozin or linagliptin.

One patient administered Glyxambi experienced a confirmed (investigator-defined), major hypoglycaemic event (defined as an event requiring assistance) in the active- or placebo-controlled trials (overall frequency 0.1%).

Based on the experience with empagliflozin and linagliptin, an increase of the risk of hypoglycaemia is expected with the concomitant treatment of insulin and/or sulphonylurea (see section 4.4 and information below)

#### *Hypoglycaemia with empagliflozin*

The frequency of hypoglycaemia depended on the background therapy in the respective trials and was similar for empagliflozin and placebo as monotherapy, as add-on to metformin, and as add-on to pioglitazone +/- metformin. The frequency of patients with hypoglycaemia was increased in patients treated with empagliflozin compared to placebo when given as add-on to metformin plus sulphonylurea (empagliflozin 10 mg: 16.1%, empagliflozin 25 mg: 11.5%, placebo: 8.4%), add-on to basal insulin +/- metformin and +/-sulphonylurea (empagliflozin 10 mg: 19.5%, empagliflozin 25 mg: 28.4%, placebo: 20.6% during initial 18 weeks treatment when insulin could not be adjusted; empagliflozin 10 mg and 25 mg: 36.1%, placebo 35.3% over the 78 week trial), and add-on to MDI insulin with or without metformin (empagliflozin 10 mg: 39.8%, empagliflozin 25 mg: 41.3%, placebo: 37.2% during initial 18 weeks treatment when insulin could not be adjusted; empagliflozin 10 mg: 51.1%, empagliflozin 25 mg: 57.7%, placebo: 58% over the 52-week trial).

#### *Major hypoglycaemia with empagliflozin (events requiring assistance)*

The frequency of patients with major hypoglycaemic events was low (< 1%) and similar for empagliflozin and placebo as monotherapy, as add-on to metformin +/- sulphonylurea, and as add-on to pioglitazone +/- metformin.

The frequency of patients with major hypoglycaemic events was increased in patients treated with empagliflozin compared to placebo when given as add-on to basal insulin +/- metformin and +/-sulphonylurea (empagliflozin 10 mg: 0%, empagliflozin 25 mg: 1.3%, placebo: 0% during initial 18 weeks treatment when insulin could not be adjusted; empagliflozin 10 mg: 0%, empagliflozin 25 mg: 1.3%, placebo 0% over the 78-week trial), and add-on to MDI insulin with or without metformin (empagliflozin 10 mg: 1.6%, empagliflozin 25 mg: 0.5%, placebo: 1.6% during initial 18 weeks treatment when insulin could not be adjusted and over the 52-week trial).

#### *Hypoglycaemia with linagliptin*

The most frequently reported adverse event in clinical trials with linagliptin was hypoglycaemia observed under the triple combination, linagliptin plus metformin plus sulphonylurea (22.9% vs 14.8% in placebo).

Hypoglycaemias in the placebo-controlled trials (10.9%; N=471) were mild (80%; N=384), moderate (16.6%; N=78) or severe (1.9%; N=9) in intensity.

### Urinary tract infection

In clinical trials with Glyxambi, there was no notable difference of the frequency of urinary tract infections in patients treated with Glyxambi (Glyxambi 25 mg/5 mg: 8.5%; Glyxambi 10 mg/5 mg: 7.5%) compared to the patients treated with empagliflozin and linagliptin. The frequencies have been comparable to those reported from the empagliflozin clinical trials (see also section 4.4).

In empagliflozin trials, the overall frequency of urinary tract infection was similar in patients treated with empagliflozin 25 mg and placebo (7.0% and 7.2%), and higher in patients treated with empagliflozin 10 mg (8.8%). Similar to placebo, urinary tract infection was reported more frequently for empagliflozin in patients with a history of chronic or recurrent urinary tract infections. The intensity of urinary tract infections was similar to placebo for mild, moderate and severe intensity reports. Urinary tract infection was reported more frequently in female patients treated with empagliflozin compared to placebo, but not in male patients.

### Vaginal moniliasis, vulvovaginitis, balanitis and other genital infection

In clinical trials with Glyxambi, genital infections in patients treated with Glyxambi (Glyxambi 25 mg/5 mg: 3.0%; Glyxambi 10 mg/5 mg: 2.5%) were reported more frequently than for linagliptin but less frequently than for empagliflozin. Overall, the frequencies for Glyxambi have been comparable to those reported from the empagliflozin clinical trials.

In empagliflozin trials, vaginal moniliasis, vulvovaginitis, balanitis and other genital infections were reported more frequently for empagliflozin 10 mg (4.0%) and empagliflozin 25 mg (3.9%) compared to placebo (1.0%). These infections were reported more frequently for empagliflozin compared to placebo in female patients, and the difference in frequency was less pronounced in male patients. The genital tract infections were mild and moderate in intensity, none was severe in intensity.

### Increased urination

In clinical trials with Glyxambi, increased urination in patients treated with Glyxambi (Glyxambi 25 mg/5 mg: 2.6%; Glyxambi 10 mg/5 mg: 1.4%) was reported more frequently than for linagliptin and with similar frequency than for empagliflozin. Overall, the frequencies for Glyxambi have been comparable to those reported from the empagliflozin clinical trials.

In clinical trials with empagliflozin, increased urination (including the predefined terms pollakiuria, polyuria, nocturia) was observed at higher frequencies in patients treated with empagliflozin (empagliflozin 10 mg: 3.5%, empagliflozin 25 mg: 3.3%) compared to placebo (1.4%). Increased urination was mostly mild or moderate in intensity. The frequency of reported nocturia was comparable between placebo and empagliflozin (< 1%).

### Volume depletion

In clinical trials with Glyxambi, there was no notable difference in the frequency of volume depletion in patients treated with Glyxambi (Glyxambi 25 mg/5 mg: 0.4%; Glyxambi 10 mg/5 mg: 0.8%) compared to the patients treated with empagliflozin and linagliptin. The frequencies have been comparable to those reported from the empagliflozin clinical trials.

In clinical trials with empagliflozin, the overall frequency of volume depletion (including the predefined terms blood pressure (ambulatory) decreased, blood pressure systolic decreased, dehydration, hypotension, hypovolaemia, orthostatic hypotension, and syncope) was similar in patients treated with empagliflozin (empagliflozin 10 mg: 0.6%, empagliflozin 25 mg: 0.4%) and placebo (0.3%). The frequency of volume depletion events was increased in patients 75 years and older treated with empagliflozin 10 mg (2.3%) or empagliflozin 25 mg (4.3%) compared to placebo (2.1%).

### Blood creatinine increased/Glomerular filtration rate decreased

In clinical trials with Glyxambi, the frequency of patients with increased blood creatinine (Glyxambi 25 mg/5 mg: 0.4%; Glyxambi 10 mg/5 mg: 0%) and decreased glomerular filtration rate (Glyxambi 25 mg/5 mg: 0.4%; Glyxambi 10 mg/5 mg: 0.6%) has been comparable to those reported from the empagliflozin clinical trials.

In clinical trials with empagliflozin, the overall frequency of patients with increased blood creatinine and decreased glomerular filtration rate were similar between empagliflozin and placebo (blood creatinine increased: empagliflozin 10 mg 0.6%, empagliflozin 25 mg 0.1%, placebo 0.5%; glomerular filtration rate decreased: empagliflozin 10 mg 0.1%, empagliflozin 25 mg 0%, placebo 0.3%).

### Elderly

In clinical trials, nineteen patients 75 years or older were treated with Glyxambi. No patient was older than 85 years. The safety profile of Glyxambi did not differ in the elderly. Based on empagliflozin experiences, elderly patients may be at increased risk of volume depletion (see sections 4.2, 4.4 and 5.2)

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

## **4.9 Overdose**

### Symptoms

In controlled clinical trials single doses of up to 800 mg empagliflozin (equivalent to 32 times the highest recommended daily dose) in healthy volunteers and multiple daily doses of up to 100 mg empagliflozin (equivalent to 4 times the highest recommended daily dose) in patients with type 2 diabetes did not show any toxicity. Empagliflozin increased urine glucose excretion leading to an increase in urine volume. The observed increase in urine volume was not dose-dependent. There is no experience with doses above 800 mg in humans.

During controlled clinical trials in healthy subjects, single doses of up to 600 mg linagliptin (equivalent to 120 times the recommended dose) were generally well tolerated. There is no experience with doses above 600 mg in humans.

### Treatment

In the event of an overdose, it is reasonable to employ the usual supportive measures, e.g., remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring and institute clinical measures as required.

The removal of empagliflozin by haemodialysis has not been studied. Linagliptin is not expected to be eliminated to a therapeutically significant degree by haemodialysis or peritoneal dialysis.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Drugs used in diabetes, combinations of oral blood glucose lowering drugs, ATC code: A10BD19

#### Mechanism of action

Glyxambi combines two antihyperglycaemic medicinal products with complementary mechanisms of action to improve glycaemic control in patients with type 2 diabetes: empagliflozin, a sodium-glucose co-transporter (SGLT2) inhibitor, and linagliptin, DPP-4 inhibitor.

#### Empagliflozin

Empagliflozin is a reversible, highly potent (IC<sub>50</sub> of 1.3 nmol) and selective competitive inhibitor of SGLT2. Empagliflozin does not inhibit other glucose transporters important for glucose transport into peripheral tissues and is 5 000 times more selective for SGLT2 versus SGLT1, the major transporter responsible for glucose absorption in the gut.

SGLT2 is highly expressed in the kidney, whereas expression in other tissues is absent or very low. It is responsible, as the predominant transporter, for the reabsorption of glucose from the glomerular filtrate back into the circulation. In patients with type 2 diabetes and hyperglycaemia a higher amount of glucose is filtered and reabsorbed.

Empagliflozin improves glycaemic control in patients with type 2 diabetes mellitus by reducing renal glucose re-absorption. The amount of glucose removed by the kidney through this glucuretic mechanism is dependent upon the blood glucose concentration and GFR. Inhibition of SGLT2 in patients with type 2 diabetes mellitus and hyperglycaemia leads to excess glucose excretion in the urine. In addition, initiation of empagliflozin increases excretion of sodium resulting in osmotic diuresis and reduced intravascular volume.

In patients with type 2 diabetes, urinary glucose excretion increased immediately following the first dose of empagliflozin and was continuous over the 24-hour dosing interval. Increased urinary glucose excretion was maintained at the end of the 4-week treatment period, averaging approximately 78 g/day. Increased urinary glucose excretion resulted in an immediate reduction in plasma glucose levels in patients with type 2 diabetes.

Empagliflozin improves both fasting and post prandial plasma glucose levels. The mechanism of action of empagliflozin is independent of beta cell function and insulin pathway and this contributes to a low risk of hypoglycaemia. Improvement of surrogate markers of beta cell function including Homeostasis Model Assessment  $\beta$  (HOMA  $\beta$ ) was noted. In addition, urinary glucose excretion triggers calorie loss, associated with body fat loss and body weight reduction. The glucosuria observed with empagliflozin is accompanied by diuresis which may contribute to sustained and moderate reduction of blood pressure. The glucosuria, natriuresis and osmotic diuresis observed with empagliflozin may contribute to the improvement in cardiovascular outcomes.

#### Linagliptin

Linagliptin is an inhibitor of DPP-4 an enzyme which is involved in the inactivation of the incretin hormones GLP-1 and GIP (glucagon-like peptide-1, glucose-dependent insulinotropic polypeptide). These hormones are rapidly degraded by the enzyme DPP-4. Both incretin hormones are involved in the physiological regulation of glucose homeostasis. Incretins are secreted at a low basal level throughout the day and levels rise immediately after meal intake. GLP-1 and GIP increase insulin biosynthesis and secretion from pancreatic beta cells in the presence of normal and elevated blood glucose levels. Furthermore GLP-1 also reduces glucagon secretion from pancreatic alpha cells, resulting in a reduction in hepatic glucose output. Linagliptin binds very effectively to DPP-4 in a

reversible manner and thus leads to a sustained increase and a prolongation of active incretin levels. Linagliptin glucose-dependently increases insulin secretion and lowers glucagon secretion thus resulting in an overall improvement in the glucose homeostasis. Linagliptin binds selectively to DPP-4 and exhibits a > 10,000-fold selectivity versus DPP-8 or DPP-9 activity *in vitro*.

### Clinical efficacy and safety

A total of 2 173 patients with type 2 diabetes mellitus and inadequate glycaemic control were treated in clinical trials to evaluate the safety and efficacy of Glyxambi; 1 005 patients were treated with Glyxambi 10 mg empagliflozin/5 mg linagliptin or 25 mg empagliflozin/5 mg linagliptin. In clinical trials, patients were treated for up to 24 or 52 weeks.

### Glyxambi added to metformin

In a factorial design trial patients inadequately controlled on metformin were treated for 24-weeks with Glyxambi 10 mg/5 mg, Glyxambi 25 mg/5 mg, empagliflozin 10 mg, empagliflozin 25 mg or linagliptin 5 mg. The treatment with Glyxambi resulted in statistically significant improvements in HbA<sub>1c</sub> (see Table 3) and fasting plasma glucose (FPG) compared to linagliptin 5 mg and also compared to empagliflozin 10 mg or 25 mg. Glyxambi also provided statistically significant improvements in body weight compared to linagliptin 5 mg.

Table 3 Efficacy parameters in clinical trial comparing Glyxambi to individual active substances as add-on therapy in patients inadequately controlled on metformin

	<b>Glyxambi 25 mg/5 mg</b>	<b>Glyxambi 10 mg/5 mg</b>	<b>Empagliflo zin 25 mg</b>	<b>Empagliflo zin 10 mg</b>	<b>Linaglipti n 5 mg</b>
<b>Primary endpoint: HbA<sub>1c</sub> ( %) – 24 weeks</b>					
<b>Number of patients analysed</b>	<b>134</b>	<b>135</b>	<b>140</b>	<b>137</b>	<b>128</b>
Baseline mean (SE)	7.90 (0.07)	7.95 (0.07)	8.02 (0.07)	8.00 (0.08)	8.02 (0.08)
Change from baseline at week 24 <sup>1</sup> : - adjusted mean <sup>2</sup> (SE)	-1.19 (0.06)	-1.08 (0.06)	-0.62 (0.06)	-0.66 (0.06)	-0.70 (0.06)
Comparison vs. empagliflozin <sup>1</sup> : - adjusted mean <sup>2</sup> (SE) - 95.0 % CI - p-value	vs. 25 mg -0.58 (0.09) -0.75, -0.41 <0.0001	vs. 10 mg -0.42 (0.09) -0.59, -0.25 <0.0001	--	--	--
Comparison vs. linagliptin 5 mg <sup>1</sup> : - adjusted mean <sup>2</sup> (SE) - 95.0 % CI - p-value	-0.50 (0.09) -0.67, -0.32 <0.0001	-0.39 (0.09) -0.56, -0.21 <0.0001	--	--	--

<sup>1</sup> Last observation (prior to glycaemic rescue ) carried forward (LOCF)

<sup>2</sup> Mean adjusted for baseline value and stratification

In a pre-specified subgroup of patients with baseline HbA<sub>1c</sub> greater or equal than 8.5%, the reduction from baseline in HbA<sub>1c</sub> at 24 weeks with Glyxambi 25 mg/5 mg was -1.8% (p<0.0001 versus linagliptin 5 mg, p<0.001 versus empagliflozin 25 mg) and with Glyxambi 10 mg/5 mg -1.6% (p<0.01 versus linagliptin 5 mg, n.s. versus empagliflozin 10 mg).

Overall, the effects on HbA<sub>1c</sub> reduction observed at 24 weeks were sustained at week 52.



Empagliflozin in patients inadequately controlled on metformin and linagliptin

In patients inadequately controlled on maximally tolerated doses of metformin, open label linagliptin 5 mg was added for 16 weeks. In patients inadequately controlled after this 16 week period, patients received double-blind treatment with either empagliflozin 10 mg, empagliflozin 25 mg or placebo for 24-weeks. After this double-blind period, treatment with both empagliflozin 10 mg and empagliflozin 25 mg provided statistically significant improvements in HbA<sub>1c</sub>, FPG and body weight compared to placebo; all patients continued treatment with metformin and linagliptin 5 mg during the trial. A statistically significant greater number of patients with a baseline HbA<sub>1c</sub>  $\geq 7.0\%$  treated with both doses of empagliflozin achieved a target HbA<sub>1c</sub> of  $<7\%$  compared to placebo (see Table 4). After 24-weeks treatment with empagliflozin, both systolic and diastolic blood pressures were reduced, -2.6/-1.1 mmHg (n.s. versus placebo for SBP and DBP) for empagliflozin 25 mg and -1.3/-0.1 mmHg (n.s. versus placebo for SBP and DBP) for empagliflozin 10 mg.

After 24 weeks, rescue therapy was used in 4 (3.6%) patients treated with empagliflozin 25 mg and in 2 (1.8%) patients treated with empagliflozin 10 mg, compared to 13 (12.0%) patients treated with placebo (all patients on background metformin + linagliptin 5 mg).

Table 4 Efficacy parameters in the clinical trial comparing empagliflozin to placebo as add-on therapy in patients inadequately controlled on metformin and linagliptin 5 mg

	Metformin + linagliptin 5 mg		
	Empagliflozin 10 mg <sup>1</sup>	Empagliflozin 25 mg <sup>1</sup>	Placebo <sup>2</sup>
<b>HbA<sub>1c</sub> ( %) - 24 weeks<sup>3</sup></b>			
N	109	110	106
Baseline (mean)	7.97	7.97	7.96
Change from baseline (adjusted mean)	-0.65	-0.56	0.14
Comparison vs. placebo (adjusted mean) (95 % CI) <sup>2</sup>	-0.79 (-1.02, -0.55) p<0.0001	-0.70 (-0.93, -0.46) p<0.0001	
<b>Body Weight-24 weeks<sup>3</sup></b>			
N	109	110	106
Baseline (mean) in kg	88.4	84.4	82.3
Change from baseline (adjusted mean)	-3.1	-2.5	-0.3
Comparison vs. placebo (adjusted mean) (95 % CI) <sup>1</sup>	-2.8 (-3.5, -2.1) p<0.0001	-2.2 (-2.9, -1.5) p<0.0001	
<b>Patients ( %) achieving HbA<sub>1c</sub> &lt;7 % with baseline HbA<sub>1c</sub> <math>\geq 7\%</math> - 24 weeks<sup>4</sup></b>			
N	100	107	100
Patients ( %) achieving A1C <7 %	37.0	32.7	17.0
Comparison vs. placebo (odds ratio) (95 % CI) <sup>5</sup>	4.0 (1.9, 8.7)	2.9 (1.4, 6.1)	

<sup>1</sup> Patients randomised to the empagliflozin 10 mg or 25 mg groups were receiving Glyxambi 10 mg/5 mg or 25 mg/5 mg with background metformin

<sup>2</sup> Patients randomised to the placebo group were receiving the placebo plus linagliptin 5 mg with background metformin

<sup>3</sup> Mixed-effects models for repeated measurements (MMRM) on FAS (OC) includes baseline HbA<sub>1c</sub>, baseline eGFR (MDRD), geographical region, visit treatment, and treatment by visit interaction. For FPG, baseline FPG is also included. For weight, baseline weight is also included.

<sup>4</sup> Not evaluated for statistical significance; not part of sequential testing procedure for the secondary endpoints

<sup>5</sup> Logistic regression on FAS (NCF) includes baseline HbA<sub>1c</sub>, baseline eGFR (MDRD), geographical region, and treatment;

based on patients with HbA<sub>1c</sub> of 7 % and above at baseline

In a pre-specified subgroup of patients with baseline HbA<sub>1c</sub> greater or equal than 8.5% the reduction from baseline in HbA<sub>1c</sub> with empagliflozin 25 mg/linagliptin 5 mg was -1.3% at 24 weeks (p<0.0001 versus placebo and linagliptin 5 mg) and with empagliflozin 10 mg/linagliptin 5 mg -1.3% at 24 weeks (p<0.0001 versus placebo and linagliptin 5 mg).

Linagliptin 5 mg in patients inadequately controlled on metformin and empagliflozin 10 mg or empagliflozin 25 mg

In patients inadequately controlled on maximally tolerated doses of metformin, open label empagliflozin 10 mg or empagliflozin 25 mg was added for 16 weeks. In patients inadequately controlled after this 16 week period, patients received double-blind treatment with either linagliptin 5 mg or placebo for 24-weeks. After this double-blind period, treatment in both populations (metformin + empagliflozin 10 mg and metformin + empagliflozin 25 mg) linagliptin 5 mg provided statistically significant improvements in HbA<sub>1c</sub> compared to placebo; all patients continued treatment with metformin and empagliflozin during the trial. A statistically significant greater number of patients with a baseline HbA<sub>1c</sub> ≥7.0% and treated with linagliptin achieved a target HbA<sub>1c</sub> of <7% compared to placebo (see Table 5).

Table 5 Efficacy parameters in clinical trials comparing Glyxambi 10 mg/5 mg to empagliflozin 10 mg as well as Glyxambi 25 mg/5 mg to empagliflozin 25 mg as add-on therapy in patients inadequately controlled on empagliflozin 10 mg/25 mg and metformin

	<b>Metformin + empagliflozin 10 mg</b>		<b>Metformin + empagliflozin 25 mg</b>	
	<b>Linagliptin 5 mg</b>	<b>Placebo</b>	<b>Linagliptin 5 mg</b>	<b>Placebo</b>
<b>HbA<sub>1c</sub> (%) – 24 weeks<sup>1</sup></b>				
N	122	125	109	108
Baseline (mean)	8.04	8.03	7.82	7.88
Change from baseline (adjusted mean)	-0.53	-0.21	-0.58	-0.10
Comparison vs. placebo (adjusted mean) (95 % CI)	-0.32 (-0.52, -0.13) p=0.0013		-0.47 (-0.66, -0.28) p<0.0001	
<b>Patients (%) achieving HbA<sub>1c</sub> &lt;7 % with baseline HbA<sub>1c</sub> ≥7 % – 24 weeks<sup>2</sup></b>				
N	116	119	100	107
Patients (%) achieving HbA <sub>1c</sub> <7 %	25.9	10.9	36.0	15.0
Comparison vs. placebo (odds ratio) (95 % CI) <sup>3</sup>	3.965 (1.771, 8.876) p=0.0008		4.429 (2.097, 9.353) p<0.0001	

Patients randomised to the linagliptin 5 mg group were receiving either fixed dose combination tablets Glyxambi 10 mg/5 mg plus metformin or fixed dose combination tablets Glyxambi 25 mg/5 mg plus metformin; patients randomised to the placebo group were receiving placebo plus empagliflozin 10 mg plus metformin or placebo plus empagliflozin 25 mg plus metformin

<sup>1</sup> MMRM model on FAS (OC) includes baseline HbA<sub>1c</sub>, baseline eGFR (MDRD), geographical region, visit, treatment, and treatment by visit interaction. For FPG, baseline FPG is also included.

<sup>2</sup> Not evaluated for statistical significance; not part of sequential testing procedure for the secondary endpoints

<sup>3</sup> Logistic regression on FAS (NCF) includes baseline HbA<sub>1c</sub>, baseline eGFR (MDRD), geographical region, and treatment; based on patients with HbA<sub>1c</sub> of 7% and above at baseline

Cardiovascular safety

*Empagliflozin cardiovascular outcome (EMPA-REG OUTCOME) trial*

The double-blind, placebo-controlled EMPA-REG OUTCOME trial compared pooled doses of empagliflozin 10 mg and 25 mg with placebo as adjunct to standard care therapy in patients with

type 2 diabetes and established cardiovascular disease. A total of 7 020 patients were treated (empagliflozin 10 mg: 2 345, empagliflozin 25 mg: 2 342, placebo: 2 333) and followed for a median of 3.1 years. The mean age was 63 years, the mean HbA<sub>1c</sub> was 8.1%, and 71.5% were male. At baseline, 74% of patients were being treated with metformin, 48% with insulin, and 43% with a sulfonylurea. About half of the patients (52.2%) had an eGFR of 60-90 ml/min/1.73 m<sup>2</sup>, 17.8% of 45-60 ml/min/1.73 m<sup>2</sup> and 7.7% of 30-45 ml/min/1.73 m<sup>2</sup>.

At week 12, an adjusted mean (SE) improvement in HbA<sub>1c</sub> when compared to baseline of 0.11% (0.02) in the placebo group, 0.65% (0.02) and 0.71% (0.02) in the empagliflozin 10 and 25 mg groups was observed. After the first 12 weeks glycaemic control was optimized independent of investigative treatment. Therefore the effect was attenuated at week 94, with an adjusted mean (SE) improvement in HbA<sub>1c</sub> of 0.08% (0.02) in the placebo group, 0.50% (0.02) and 0.55% (0.02) in the empagliflozin 10 and 25 mg groups.

Empagliflozin was superior in preventing the primary combined endpoint of cardiovascular death, nonfatal myocardial infarction, or non-fatal stroke, as compared with placebo. The treatment effect was driven by a significant reduction in cardiovascular death with no significant change in non-fatal myocardial infarction, or non-fatal stroke. The reduction of cardiovascular death was comparable for empagliflozin 10 mg and 25 mg and confirmed by an improved overall survival (see Table 6). The effect of empagliflozin on the primary combined endpoint of CV death, non-fatal MI, or non-fatal stroke was largely independent of glycaemic control or renal function (eGFR) and generally consistent across eGFR categories down to an eGFR of 30 ml/min/1.73 m<sup>2</sup> in the EMPA-REG OUTCOME study.

Table 6 Treatment effect for the primary composite endpoint, its components and mortality<sup>a</sup>

	<b>Placebo</b>	<b>Empagliflozin<sup>b</sup></b>
<b>N</b>	2333	4687
<b>Time to first event of CV death, non-fatal MI, or non-fatal stroke N (%)</b>	282 (12.1)	490 (10.5)
Hazard ratio vs. placebo (95.02% CI)*		0.86 (0.74, 0.99)
p-value for superiority		0.0382
<b>CV Death N (%)</b>	137 (5.9)	172 (3.7)
Hazard ratio vs. placebo (95% CI)		0.62 (0.49, 0.77)
p-value		<0.0001
<b>Non-fatal MI N (%)</b>	121 (5.2)	213 (4.5)
Hazard ratio vs. placebo (95% CI)		0.87 (0.70, 1.09)
p-value		0.2189
<b>Non-fatal stroke N (%)</b>	60 (2.6)	150 (3.2)
Hazard ratio vs. placebo (95% CI)		1.24 (0.92, 1.67)
p-value		0.1638
<b>All-cause mortality N (%)</b>	194 (8.3)	269 (5.7)
Hazard ratio vs. placebo (95% CI)		0.68 (0.57, 0.82)
p-value		<0.0001
<b>Non-CV mortality N (%)</b>	57 (2.4)	97 (2.1)
Hazard ratio vs. placebo (95% CI)		0.84 (0.60, 1.16)

CV = cardiovascular, MI = myocardial infarction

<sup>a</sup> Treated set (TS), i.e. patients who had received at least one dose of trial drug

<sup>b</sup> Pooled doses of empagliflozin 10 mg and 25 mg

\* Since data from the trial were included in an interim analysis, a two-sided 95.02% confidence interval applied which corresponds to a p-value of less than 0.0498 for significance.

The efficacy for preventing cardiovascular mortality has not been conclusively established in patients using empagliflozin concomitantly with DPP-4 inhibitors or in Black patients because the representation of these groups in the EMPA-REG OUTCOME trial was limited.

### *Heart failure requiring hospitalization*

In the EMPA-REG OUTCOME trial, empagliflozin reduced the risk of heart failure requiring hospitalization compared with placebo (empagliflozin 2.7%; placebo 4.1%; HR 0.65, 95% CI 0.50, 0.85).

### *Nephropathy*

In the EMPA-REG OUTCOME trial, for time to first nephropathy event, the HR was 0.61 (95% CI 0.53, 0.70) for empagliflozin (12.7%) vs placebo (18.8%).

In addition, empagliflozin showed a higher (HR 1.82, 95% CI 1.40, 2.37) occurrence of sustained normo- or micro-albuminuria (49.7%) in patients with baseline macro-albuminuria compared with placebo (28.8%).

### *Linagliptin cardiovascular and renal safety (CARMELINA) trial*

The double-blind, placebo-controlled CARMELINA trial evaluated the cardiovascular and renal safety of linagliptin versus placebo as adjunct to standard care therapy in patients with type 2 diabetes and with increased CV risk evidenced by a history of established macrovascular or renal disease. A total of 6 979 patients were treated (linagliptin 5 mg: 3 494, placebo: 3 485) and followed for a median of 2.2 years. The trial population included 1 211 (17.4%) patients  $\geq$  75 years of age, the mean HbA<sub>1c</sub> was 8.0%, 63% were male. Approximately 19% of the population had an eGFR of 45-60 mL/min/1.73 m<sup>2</sup>, 28% of 30-45 mL/min/1.73 m<sup>2</sup> and 15% of <30 mL/min/1.73 m<sup>2</sup>.

Linagliptin did not increase the risk of the combined endpoint of CV death, non-fatal myocardial infarction or non-fatal stroke (MACE-3) [HR=1.02; (95% CI 0.89, 1.17); p=0.0002 for non-inferiority], or the risk of combined endpoint of renal death, ESRD, 40% or more sustained decrease in eGFR [HR=1.04; (95% CI 0.89, 1.22)]. In analyses for albuminuria progression (change from normoalbuminuria to micro- or macroalbuminuria, or from microalbuminuria to macroalbuminuria) the estimated hazard ratio was 0.86 (95% CI 0.78, 0.95) for linagliptin versus placebo. In addition, linagliptin did not increase the risk of hospitalization for heart failure [HR=0.90; (95% CI 0.74, 1.08)]. No increased risk of CV death or all-cause mortality was observed.

Safety data from this trial was in line with previous known safety profile of linagliptin.

### *Linagliptin cardiovascular safety (CAROLINA) trial*

The double-blind parallel group CAROLINA trial evaluated the cardiovascular safety of linagliptin versus glimepiride as adjunct to standard care therapy in patients with type 2 diabetes and with increased CV risk. A total of 6 033 patients were treated (linagliptin 5 mg: 3 023, glimepiride 1 mg to 4 mg: 3 010) and followed for a median of 6.25 years. The mean age was 64 years, the mean HbA<sub>1c</sub> was 7.15%, and 60% were male. Approximately 19% of the population had an eGFR <60 mL/min/1.73 m<sup>2</sup>.

The trial was designed to demonstrate non-inferiority for the primary cardiovascular endpoint which was a composite of the first occurrence of cardiovascular death or a non-fatal myocardial infarction (MI) or a non-fatal stroke (3P-MACE). Linagliptin did not increase the risk of the combined endpoint of CV death, non-fatal myocardial infarction or non-fatal stroke (MACE-3) [Hazard Ratio (HR)=0.98; (95% CI 0.84, 1.14); p<0.0001 for non-inferiority], when added to standard of care in adult patients with type 2 diabetes with increased CV risk compared to glimepiride (see Table 7).

Table 7 Major adverse cardiovascular events (MACE) and mortality by treatment group in the CAROLINA trial

	<b>Linagliptin 5mg</b>		<b>Glimepiride (1-4mg)</b>		<b>Hazard ratio (95% CI)</b>
	Number of Subjects (%)	Incidence Rate per 1000 PY*	Number of Subjects (%)	Incidence Rate per 1000 PY*	
Number of patients	3023		3010		
Primary CV composite (Cardiovascular death, non-fatal MI, non-fatal stroke)	356 (11.8)	20.7	362 (12.0)	21.2	0.98 (0.84, 1.14)**
All-cause mortality	308 (10.2)	16.8	336 (11.2)	18.4	0.91 (0.78,1.06)
CV death	169 (5.6)	9.2	168 (5.6)	9.2	1.00 (0.81, 1.24)
Hospitalization for heart failure (HHF)	112 (3.7)	6.4	92 (3.1)	5.3	1.21 (0.92, 1.59)

\* PY=patient years

\*\* Test on non-inferiority to demonstrate that the upper bound of the 95% CI for the hazard ratio is less than 1.3

### Paediatric population

The European Medicines Agency has waived the obligation to submit the results of trials with Glyxambi in all subsets of the paediatric population in type 2 diabetes mellitus (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

The rate and extent of absorption of empagliflozin and linagliptin in Glyxambi are equivalent to the bioavailability of empagliflozin and linagliptin when administered as individual tablets. The pharmacokinetics of empagliflozin and linagliptin as single agents have been extensively characterized in healthy subjects and patients with type 2 diabetes. Pharmacokinetics were generally similar in healthy subjects and in patients with type 2 diabetes.

Glyxambi showed a similar food effect as the individual active substances. Glyxambi can therefore be taken with or without food.

### Empagliflozin

#### Absorption

After oral administration, empagliflozin was rapidly absorbed with peak plasma concentrations occurring at a median  $t_{max}$  of 1.5 hours post dose. Thereafter, plasma concentrations declined in a biphasic manner with a rapid distribution phase and a relatively slow terminal phase. The steady state mean plasma area under the concentration-time curve (AUC) and  $C_{max}$  were 1,870 nmol.h and 259 nmol/L with empagliflozin 10 mg and 4,740 nmol.h and 687 nmol/L with empagliflozin 25 mg once daily. Systemic exposure of empagliflozin increased in a dose proportional manner. The single dose and steady state pharmacokinetic parameters of empagliflozin were similar suggesting linear pharmacokinetics with respect to time.

Administration of empagliflozin 25 mg after intake of a high-fat and high calorie meal resulted in slightly lower exposure; AUC decreased by approximately 16% and  $C_{max}$  by approximately 37% compared to fasted condition. The observed effect of food on empagliflozin pharmacokinetics was not considered clinically relevant and empagliflozin may be administered with or without food.

#### Distribution

The apparent steady-state volume of distribution was estimated to be 73.8 L based on the population pharmacokinetic analysis. Following administration of an oral [ $^{14}$ C]-empagliflozin solution to healthy

volunteers, the red blood cell partitioning was approximately 37% and plasma protein binding was 86%.

### Biotransformation

No major metabolites of empagliflozin were detected in human plasma and the most abundant metabolites were three glucuronide conjugates (2-, 3-, and 6-O-glucuronide). Systemic exposure of each metabolite was less than 10% of total drug-related material. *In vitro* studies suggest that the primary route of metabolism of empagliflozin in humans is glucuronidation by the uridine 5'-diphospho-glucuronosyltransferases UGT2B7, UGT1A3, UGT1A8 and UGT1A9.

### Elimination

Based on the population pharmacokinetic analysis, the apparent terminal elimination half life of empagliflozin was estimated to be 12.4 hours and apparent oral clearance was 10.6 L/hour. The inter subject and residual variabilities for empagliflozin oral clearance were 39.1% and 35.8%, respectively. With once daily dosing, steady state plasma concentrations of empagliflozin were reached by the fifth dose. Consistent with the half life, up to 22% accumulation, with respect to plasma AUC, was observed at steady state.

Following administration of an oral [<sup>14</sup>C]-empagliflozin solution to healthy volunteers, approximately 96% of the drug-related radioactivity was eliminated in faeces (41%) or urine (54%). The majority of drug-related radioactivity recovered in faeces was unchanged parent drug and approximately half of drug related radioactivity excreted in urine was unchanged parent drug.

### Linagliptin

#### Absorption

After oral administration of a 5 mg dose to healthy volunteers or patients, linagliptin was rapidly absorbed, with peak plasma concentrations (median  $T_{max}$ ) occurring 1.5 hours post-dose.

After once daily dosing of 5 mg linagliptin, steady-state plasma concentrations are reached by the third dose. Plasma AUC of linagliptin increased approximately 33% following 5 mg doses at steady-state compared to the first dose. The intra-subject and inter-subject coefficients of variation for linagliptin AUC were small (12.6% and 28.5%, respectively). Due to the concentration dependent binding of linagliptin to DPP-4, the pharmacokinetics of linagliptin based on total exposure is not linear; indeed total plasma AUC of linagliptin increased in a less than dose-proportional manner while unbound AUC increases in a roughly dose-proportional manner.

The absolute bioavailability of linagliptin is approximately 30%. Co-administration of a high-fat meal with linagliptin prolonged the time to reach  $C_{max}$  by 2 hours and lowered  $C_{max}$  by 15% but no influence on  $AUC_{0-72h}$  was observed. No clinically relevant effect of  $C_{max}$  and  $T_{max}$  changes is expected; therefore linagliptin may be administered with or without food.

The steady state plasma  $AUC_{\tau,ss}$  and  $C_{max,ss}$  concentrations of linagliptin were 153 nmol\*hr/L and 12.9 nmol/L for linagliptin 5 mg once daily for 7 days.

#### Distribution

As a result of tissue binding, the mean apparent volume of distribution at steady-state following a single 5 mg intravenous dose of linagliptin to healthy subjects is approximately 1,110 litres, indicating that linagliptin extensively distributes to the tissues. Plasma protein binding of linagliptin is concentration-dependent, decreasing from about 99% at 1 nmol/L to 75-89% at  $\geq 30$  nmol/L, reflecting saturation of binding to DPP-4 with increasing concentration of linagliptin. At high concentrations, where DPP-4 is fully saturated, 70-80% of linagliptin was bound to other plasma proteins than DPP-4, hence 30-20% were unbound in plasma.

### Biotransformation

Following a [<sup>14</sup>C] linagliptin oral 10 mg dose, approximately 5% of the radioactivity was excreted in urine. Metabolism plays a subordinate role in the elimination of linagliptin. One main metabolite with a relative exposure of 13.3% of linagliptin at steady-state was detected which was found to be pharmacologically inactive and thus to not contribute to the plasma DPP-4 inhibitory activity of linagliptin.

### Elimination

Plasma concentrations of linagliptin decline in a triphasic manner with a long terminal half-life (terminal half-life for linagliptin more than 100 hours) that is mostly related to the saturable, tight binding of linagliptin to DPP-4 and does not contribute to the accumulation of the medicinal product. The effective half-life for accumulation of linagliptin, as determined from oral administration of multiple doses of 5 mg linagliptin, is approximately 12 hours.

Following administration of an oral [<sup>14</sup>C] linagliptin dose to healthy subjects, approximately 85% of the administered radioactivity was eliminated in faeces (80%) or urine (5%) within 4 days of dosing. Renal clearance at steady-state was approximately 70 mL/min.

### Renal impairment

#### Empagliflozin

In patients with mild, moderate or severe renal impairment (eGFR <30 to <90 mL/min/1.73 m<sup>2</sup>) and patients with kidney failure or end stage renal disease (ESRD), AUC of empagliflozin increased by approximately 18%, 20%, 66%, and 48%, respectively compared to subjects with normal renal function. Peak plasma levels of empagliflozin were similar in subjects with moderate renal impairment and kidney failure/ESRD compared to patients with normal renal function. Peak plasma levels of empagliflozin were roughly 20% higher in subjects with mild and severe renal impairment as compared to subjects with normal renal function. The population pharmacokinetic analysis showed that the apparent oral clearance of empagliflozin decreased with a decrease in eGFR leading to an increase in drug exposure (see section 4.2).

#### Linagliptin

A multiple-dose, open-label trial was conducted to evaluate the pharmacokinetics of linagliptin (5 mg dose) in patients with varying degrees of chronic renal insufficiency compared to subjects with normal renal function. The trial included patients with renal insufficiency classified on the basis of creatinine clearance as mild (50 to <80 mL/min), moderate (30 to <50 mL/min), and severe (<30 mL/min), as well as patients with ESRD on haemodialysis. In addition patients with T2DM and severe renal impairment (<30 mL/min) were compared to T2DM patients with normal renal function. Under steady-state conditions, linagliptin exposure in patients with mild renal impairment was comparable to healthy subjects. In moderate renal impairment, a moderate increase in exposure of about 1.7-fold was observed compared with control. Exposure in T2DM patients with severe RI was increased by about 1.4-fold compared to T2DM patients with normal renal function. Steady-state predictions for AUC of linagliptin in patients with ESRD indicated comparable exposure to that of patients with moderate or severe renal impairment. In addition, linagliptin is not expected to be eliminated to a therapeutically significant degree by haemodialysis or peritoneal dialysis (see section 4.2).

### Hepatic impairment

#### Empagliflozin

In patients with mild, moderate and severe hepatic insufficiency (Child-Pugh classification), mean AUC and C<sub>max</sub> of empagliflozin increased (AUC by 23%, 47%, 75% and C<sub>max</sub> by 4%, 23%, 48%)

compared to subjects with normal hepatic function (see section 4.2).

### Linagliptin

In non-diabetic patients with mild, moderate and severe hepatic insufficiency (according to the Child-Pugh classification), mean AUC and C<sub>max</sub> of linagliptin were similar to healthy subjects following administration of multiple 5 mg doses of linagliptin.

### Body mass index

No dose adjustment is necessary for Glyxambi based on body mass index. Body mass index had no clinically relevant effect on the pharmacokinetics of empagliflozin or linagliptin based on population pharmacokinetic analysis.

### Gender

Gender had no clinically relevant effect on the pharmacokinetics of empagliflozin or linagliptin based on population pharmacokinetic analysis.

### Race

No clinically relevant difference in pharmacokinetics of empagliflozin and linagliptin were seen in population pharmacokinetic analysis and dedicated phase I trials.

### Elderly

Age did not have a clinically meaningful impact on the pharmacokinetics of empagliflozin or linagliptin based on population pharmacokinetic analysis. Elderly subjects (65 to 80 years) had comparable plasma concentrations of linagliptin compared to younger subjects.

### Paediatric patients

#### Empagliflozin

A paediatric Phase 1 trial examined the pharmacokinetics and pharmacodynamics of empagliflozin (5 mg, 10 mg and 25 mg) in children and adolescents  $\geq 10$  to  $< 18$  years of age with type 2 diabetes mellitus. The observed pharmacokinetic and pharmacodynamic responses were consistent with those found in adult subjects.

#### Linagliptin

A paediatric Phase 2 trial examined the pharmacokinetics and pharmacodynamics of 1 mg and 5 mg linagliptin in children and adolescents  $\geq 10$  to  $< 18$  years of age with type 2 diabetes mellitus. The observed pharmacokinetic and pharmacodynamic responses were consistent with those found in adult subjects. Linagliptin 5 mg showed superiority over 1 mg with regard to trough DPP-4 inhibition (72% vs 32%,  $p=0.0050$ ) and a numerically larger reduction with regard to adjusted mean change from baseline in HbA<sub>1c</sub> (-0.63% vs -0.48%, n.s.). Due to the limited nature of the data set the results should be interpreted cautiously.

### Drug interactions

No drug interaction trials have been performed with Glyxambi and other medicinal products; however, such trials have been conducted with the individual active substances.

#### In vitro assessment of empagliflozin

Based on *in vitro* studies, empagliflozin does not inhibit, inactivate, or induce CYP450 isoforms.



Empagliflozin does not inhibit UGT1A1, UGT1A3, UGT1A8, UGT1A9, or UGT2B7. Drug-drug interactions involving the major CYP450 and UGT isoforms with empagliflozin and concomitantly administered substrates of these enzymes are therefore considered unlikely.

*In vitro* data suggest that the primary route of metabolism of empagliflozin in humans is glucuronidation by uridine 5'-diphosphoglucuronosyltransferases UGT1A3, UGT1A8, UGT1A9, and UGT2B7.

Empagliflozin is a substrate of the human uptake transporters OAT3, OATP1B1, and OATP1B3, but not Organic Anion Transporter 1 (OAT1) and Organic Cation Transporter 2 (OCT2). Empagliflozin is a substrate of P-glycoprotein (P-gp) and breast cancer resistance protein (BCRP).

Empagliflozin does not inhibit P-gp at therapeutic doses. Based on *in vitro* studies, empagliflozin is considered unlikely to cause interactions with medicinal products that are P-gp substrates. Co-administration of digoxin, a P-gp substrate, with empagliflozin resulted in a 6% increase in AUC and 14% increase in  $C_{max}$  of digoxin. These changes were not considered to be clinically meaningful.

Empagliflozin does not inhibit human uptake transporters such as OAT3, OATP1B1, and OATP1B3 *in vitro* at clinically relevant plasma concentrations and, as such, drug-drug interactions with substrates of these uptake transporters are considered unlikely.

#### *In vitro* assessment of linagliptin

Linagliptin was a substrate for OATP8-, OCT2-, OAT4-, OCTN1- and OCTN2, suggesting a possible OATP8-mediated hepatic uptake, OCT2-mediated renal uptake and OAT4-, OCTN1- and OCTN2-mediated renal secretion and reabsorption of linagliptin *in vivo*. OATP2, OATP8, OCTN1, OCT1 and OATP2 activities were slightly to weakly inhibited by linagliptin.

### **5.3 Preclinical safety data**

General toxicity studies in rats up to 13 weeks were performed with the combination of empagliflozin and linagliptin.

Focal areas of hepatocellular necrosis were found in the combination groups at  $\geq 15$ : 30 mg/kg linagliptin: empagliflozin (3.8 times the clinical exposure for linagliptin and 7.8 times the clinical exposure for empagliflozin) as well as in the group treated with empagliflozin alone but not in the control group. The clinical relevance of this finding remains uncertain.

At exposures sufficiently in excess of exposure in humans after therapeutic doses, the combination of empagliflozin and linagliptin was not teratogenic and did not show maternal toxicity. Adverse effects on renal development were not observed after administration of empagliflozin alone, linagliptin alone or after administration of the combined products.

#### Empagliflozin

Non clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, genotoxicity, fertility and early embryonic development.

In long-term toxicity studies in rodents and dogs, signs of toxicity were observed at exposures greater than or equal to 10-times the clinical dose of empagliflozin. Most toxicity was consistent with secondary pharmacology related to urinary glucose loss and electrolyte imbalances including decreased body weight and body fat, increased food consumption, diarrhoea, dehydration, decreased serum glucose and increases in other serum parameters reflective of increased protein metabolism and gluconeogenesis, urinary changes such as polyuria and glucosuria, and microscopic changes including mineralisation in kidney and some soft and vascular tissues. Microscopic evidence of the effects of exaggerated pharmacology on the kidney observed in some species included tubular dilatation, and tubular and pelvic mineralisation at approximately 4-times the clinical AUC exposure of empagliflozin

associated with the 25 mg dose.

In a 2 year carcinogenicity study, empagliflozin did not increase the incidence of tumours in female rats up to the highest dose of 700 mg/kg/day, which corresponds to approximately 72 times the maximal clinical AUC exposure to empagliflozin. In male rats, treatment related benign vascular proliferative lesions (haemangiomas) of the mesenteric lymph node were observed at the highest dose, but not at 300 mg/kg/day, which corresponds to approximately 26 times the maximal clinical exposure to empagliflozin. Interstitial cell tumours in the testes were observed with a higher incidence in rats at 300 mg/kg/day and above, but not at 100 mg/kg/day which corresponds to approximately 18 times the maximal clinical exposure to empagliflozin. Both tumours are common in rats and are unlikely to be relevant to humans.

Empagliflozin did not increase the incidence of tumours in female mice at doses up to 1 000 mg/kg/day, which corresponds to approximately 62-times the maximal clinical exposure to empagliflozin. Empagliflozin induced renal tumours in male mice at 1 000 mg/kg/day, but not at 300 mg/kg/day, which corresponds to approximately 11-times the maximal clinical exposure to empagliflozin. The mode of action for these tumours is dependent on the natural predisposition of the male mouse to renal pathology and a metabolic pathway not reflective of humans. The male mouse renal tumours are considered not relevant to humans.

At exposures sufficiently in excess of exposure in humans after therapeutic doses, empagliflozin had no adverse effects on fertility or early embryonic development. Empagliflozin administered during the period of organogenesis was not teratogenic. Only at maternally toxic doses, empagliflozin also caused bent limb bones in the rat and increased embryofetal loss in the rabbit.

In pre- and postnatal toxicity studies with empagliflozin in rats, reduced weight gain in offspring was observed at maternal exposures approximately 4 times the maximal clinical exposure to empagliflozin. No such effect was seen at systemic exposure equal to the maximal clinical exposure to empagliflozin. The relevance of this finding to humans is unclear.

In a juvenile toxicity study in the rat, when empagliflozin was administered from postnatal day 21 until postnatal day 90, non-adverse, minimal to mild renal tubular and pelvic dilation in juvenile rats was seen only at 100 mg/kg/day, which approximates 11-times the maximum clinical dose of 25 mg. These findings were absent after a 13 weeks drug-free recovery period.

### Linagliptin

Non clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, genotoxicity, fertility and early embryonic development.

In long-term toxicity studies in rodents and Cynomolgus monkeys, signs of toxicity were observed at exposures greater than 300-times the clinical dose of linagliptin.

Liver, kidneys and gastrointestinal tract are the principal target organs of toxicity in mice and rats. At exposures greater than 1 500-times the clinical exposure, adverse reactions on reproductive organs, thyroid and the lymphoid organs were seen in rats. Strong pseudo-allergic reactions were observed in dogs at medium doses, secondarily causing cardiovascular changes, which were considered dog-specific. Liver, kidneys, stomach, reproductive organs, thymus, spleen, and lymph nodes were target organs of toxicity in Cynomolgus monkeys at more than 450-times the clinical exposure. At more than 100-times clinical exposure, irritation of the stomach was the major finding in monkeys.

Oral 2-year carcinogenicity studies in rats and mice revealed no evidence of carcinogenicity in rats or male mice. A significantly higher incidence of malignant lymphomas only in female mice at the highest dose (>200-times human exposure) is not considered relevant for humans. Based on these studies there is no concern for carcinogenicity in humans.

Linagliptin had no adverse effects on fertility or early embryonic development at exposures greater

than 900-times the clinical exposure. Linagliptin administered during the period of organogenesis was not teratogenic. Only at maternally toxic doses, linagliptin caused a slight retardation of skeletal ossification in the rat and increased embryofetal loss in the rabbit.

In the pre- and postnatal toxicity study with linagliptin in rats, reduced weight gain in offspring was observed at maternal exposures approximately 1 500-times the maximal clinical exposure to linagliptin. No such effect was seen at systemic exposure 49-times the maximal clinical exposure to linagliptin.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

#### Glyxambi 10 mg/5 mg film-coated tablets

##### Tablet core

Mannitol (E421)  
Pre-gelatinised starch (maize)  
Maize starch  
Copovidone (K-value nominally 28)  
Crospovidone (Type B)  
Talc  
Magnesium stearate

##### Film coating

Hypromellose 2910  
Mannitol (E421)  
Talc  
Titanium dioxide (E171)  
Macrogol 6000  
Iron oxide yellow (E172)

#### Glyxambi 25 mg/5 mg film-coated tablets

##### Tablet core

Mannitol (E421)  
Pre-gelatinised starch (maize)  
Maize starch  
Copovidone (K-value nominally 28)  
Crospovidone (Type B)  
Talc  
Magnesium stearate

##### Film coating

Hypromellose 2910  
Mannitol (E421)  
Talc  
Titanium dioxide (E171)  
Macrogol 6000  
Iron oxide red (E172)

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

3 years.

## **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

## **6.5 Nature and contents of container**

PVC/PVDC/aluminium perforated unit dose blisters.

Pack sizes of 7 x 1, 10 x 1, 14 x 1, 28 x 1, 30 x 1, 60 x 1, 70 x 1, 90 x 1 and 100 x 1 film-coated tablets.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

No special requirements for disposal.

## **7. MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim International GmbH  
Binger Str. 173  
55216 Ingelheim am Rhein  
Germany

## **8. MARKETING AUTHORISATION NUMBERS**

### Glyxambi 10 mg/5 mg film-coated tablets

EU/1/16/1146/001 (7 x 1 film-coated tablets)  
EU/1/16/1146/002 (10 x 1 film-coated tablets)  
EU/1/16/1146/003 (14 x 1 film-coated tablets)  
EU/1/16/1146/004 (28 x 1 film-coated tablets)  
EU/1/16/1146/005 (30 x 1 film-coated tablets)  
EU/1/16/1146/006 (60 x 1 film-coated tablets)  
EU/1/16/1146/007 (70 x 1 film-coated tablets)  
EU/1/16/1146/008 (90 x 1 film-coated tablets)  
EU/1/16/1146/009 (100 x 1 film-coated tablets)

### Glyxambi 25 mg/5 mg film-coated tablets

EU/1/16/1146/010 (7 x 1 film-coated tablets)  
EU/1/16/1146/011 (10 x 1 film-coated tablets)  
EU/1/16/1146/012 (14 x 1 film-coated tablets)  
EU/1/16/1146/013 (28 x 1 film-coated tablets)  
EU/1/16/1146/014 (30 x 1 film-coated tablets)  
EU/1/16/1146/015 (60 x 1 film-coated tablets)  
EU/1/16/1146/016 (70 x 1 film-coated tablets)

EU/1/16/1146/017 (90 x 1 film-coated tablets)  
EU/1/16/1146/018 (100 x 1 film-coated tablets)

#### **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 11 November 2016  
Date of latest renewal: 16 July 2021

#### **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## **ANNEX II**

- A. MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**

### Name and address of the manufacturers responsible for batch release

Boehringer Ingelheim Pharma GmbH & Co. KG  
Binger Strasse 173  
55216 Ingelheim am Rhein  
Germany

Rottendorf Pharma GmbH  
Ostenfelder Strasse 51 – 61  
59320 Ennigerloh  
Germany

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to medical prescription.

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

The marketing authorisation holder (MAH) shall submit the first PSUR for this product within 6 months following authorisation.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**



## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Glyxambi 10 mg/5 mg film-coated tablets  
empagliflozin/linagliptin

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each tablet contains 10 mg empagliflozin and 5 mg linagliptin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

film-coated tablet

7 x 1 film-coated tablets  
10 x 1 film-coated tablets  
14 x 1 film-coated tablets  
28 x 1 film-coated tablets  
30 x 1 film-coated tablets  
60 x 1 film-coated tablets  
70 x 1 film-coated tablets  
90 x 1 film-coated tablets  
100 x 1 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim International GmbH  
55216 Ingelheim am Rhein  
Germany

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1146/001 7 x 1 film-coated tablets  
EU/1/16/1146/002 10 x 1 film-coated tablets  
EU/1/16/1146/003 14 x 1 film-coated tablets  
EU/1/16/1146/004 28 x 1 film-coated tablets  
EU/1/16/1146/005 30 x 1 film-coated tablets  
EU/1/16/1146/006 60 x 1 film-coated tablets  
EU/1/16/1146/007 70 x 1 film-coated tablets  
EU/1/16/1146/008 90 x 1 film-coated tablets  
EU/1/16/1146/009 100 x 1 film-coated tablets

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Glyxambi 10 mg/5 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS**

**BLISTER (perforated)**

**1. NAME OF THE MEDICINAL PRODUCT**

Glyxambi 10 mg/5 mg tablets  
empagliflozin/linagliptin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Glyxambi 25 mg/5 mg film-coated tablets  
empagliflozin/linagliptin

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each tablet contains 25 mg empagliflozin and 5 mg linagliptin.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

film-coated tablet

7 x 1 film-coated tablets  
10 x 1 film-coated tablets  
14 x 1 film-coated tablets  
28 x 1 film-coated tablets  
30 x 1 film-coated tablets  
60 x 1 film-coated tablets  
70 x 1 film-coated tablets  
90 x 1 film-coated tablets  
100 x 1 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim International GmbH  
55216 Ingelheim am Rhein  
Germany

**12. MARKETING AUTHORISATION NUMBERS**

EU/1/16/1146/010 7 x 1 film-coated tablets  
EU/1/16/1146/011 10 x 1 film-coated tablets  
EU/1/16/1146/012 14 x 1 film-coated tablets  
EU/1/16/1146/013 28 x 1 film-coated tablets  
EU/1/16/1146/014 30 x 1 film-coated tablets  
EU/1/16/1146/015 60 x 1 film-coated tablets  
EU/1/16/1146/016 70 x 1 film-coated tablets  
EU/1/16/1146/017 90 x 1 film-coated tablets  
EU/1/16/1146/018 100 x 1 film-coated tablets

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Glyxambi 25 mg/5 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS**

**BLISTER (perforated)**

**1. NAME OF THE MEDICINAL PRODUCT**

Glyxambi 25 mg/5 mg tablets  
empagliflozin/linagliptin

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Boehringer Ingelheim

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**B. PACKAGE LEAFLET**



## Package leaflet: Information for the patient

### **Glyxambi 10 mg/5 mg film-coated tablets** **Glyxambi 25 mg/5 mg film-coated tablets** empagliflozin/linagliptin

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

### **What is in this leaflet**

1. What Glyxambi is and what it is used for
2. What you need to know before you take Glyxambi
3. How to take Glyxambi
4. Possible side effects
5. How to store Glyxambi
6. Contents of the pack and other information

#### **1. What Glyxambi is and what it is used for**

##### **What Glyxambi is**

Glyxambi contains two active substances called empagliflozin and linagliptin. Each belongs to a group of medicines called “oral anti-diabetics”. These are medicines taken by mouth to treat type 2 diabetes.

##### **What is type 2 diabetes?**

Type 2 diabetes is a condition that comes from both your genes and your lifestyle. If you have type 2 diabetes, your pancreas may not make enough insulin to control the level of glucose in your blood, and your body is unable to use its own insulin effectively. This results in high levels of sugar in your blood, which can lead to medical problems like heart disease, kidney disease, blindness, and poor circulation in your limbs.

##### **How Glyxambi works**

Empagliflozin belongs to a group of medicines called sodium glucose co-transporter-2 (SGLT2) inhibitors. It works by blocking the SGLT2 protein in your kidneys. This causes blood sugar (glucose) to be removed in your urine. Linagliptin works in a different way, namely by enabling the pancreas to produce more insulin to lower blood glucose levels. It does this by blocking a protein called DPP-4. Thereby Glyxambi lowers the amount of sugar in your blood.

##### **What Glyxambi is used for**

- Glyxambi is added to metformin and/or sulphonylurea (SU) to treat type 2 diabetes in adult patients aged 18 years and older whose diabetes cannot be controlled when treated with metformin and/or sulphonylurea in combination with empagliflozin, or when treated with metformin and/or sulphonylurea in combination with linagliptin.
- Glyxambi can also be used as an alternative to taking both empagliflozin and linagliptin as single tablets. To avoid overdose, do not continue taking empagliflozin and linagliptin tablets separately, if you are taking this medicine.

It is important that you continue with your diet and exercise plan as recommended by your doctor,

pharmacist or nurse.

## 2. What you need to know before you take Glyxambi

### Do not take Glyxambi

- if you are allergic to empagliflozin, linagliptin, any other SGLT2 inhibitor (e.g. dapagliflozin, canagliflozin), any other DPP-4 inhibitor (e.g. sitagliptin, vildagliptin), or any of the other ingredients of this medicine (listed in section 6).

### Warnings and precautions

Talk to your doctor, before taking this medicine, and during treatment:

- if you have type 1 diabetes (your body does not produce any insulin). Glyxambi should not be used to treat type 1 diabetes.
- if you experience rapid weight loss, feeling sick or being sick, stomach pain, excessive thirst, fast and deep breathing, confusion, unusual sleepiness or tiredness, a sweet smell to your breath, a sweet or metallic taste in your mouth, or a different odour to your urine or sweat. Contact a doctor or the nearest hospital straight away as these symptoms could be a sign of “diabetic ketoacidosis” – a rare, but serious, sometimes life-threatening problem you can get with diabetes because of increased levels of “ketone bodies” in your urine or blood, seen in tests. The risk of developing diabetic ketoacidosis may be increased with prolonged fasting, excessive alcohol consumption, dehydration or sudden reductions in insulin dose, or a higher need of insulin due to major surgery or serious illness.
- if you are taking other anti-diabetic medicines known as “sulphonylurea” (e.g. glimepiride, glipizide) and/or using insulin. Your doctor may want to reduce your dose of these medicines when you take them together with Glyxambi, in order to avoid too low blood sugar (hypoglycaemia).
- if you have or have had a disease of the pancreas.
- if you have serious kidney problems. Your doctor may limit your daily dose or ask you to take a different medicine (see also section 3, ‘How to take Glyxambi’).
- if you have serious liver problems. Your doctor may ask you to take a different medicine.
- if you might be at risk of dehydration, for example:
  - if you are being sick, have diarrhoea or fever, or if you are not able to eat or drink
  - if you are taking medicines that increase urine production [diuretics] or lower blood pressure
  - if you are over 75 years old

Possible signs are listed in section 4 under ‘dehydration’. Your doctor may ask you to stop taking Glyxambi until you recover to prevent loss of too much body fluid. Ask about ways to prevent dehydration.

- if you are 75 years old or older, you should not start taking Glyxambi.
- if you have an increase in the proportion of red blood cells in your blood (haematocrit), seen in laboratory blood tests (see also section 4, ‘Possible side effects’).

Contact your doctor if you experience any of the following during treatment with Glyxambi:

- if you develop symptoms of acute pancreatitis, like persistent, severe stomach ache (abdominal pain). Possible signs are listed in section 4, ‘Possible side effects’. Your doctor may need to change your treatment.
- if you have a serious infection of the kidney or the urinary tract with fever. Your doctor may ask you to stop taking Glyxambi until you have recovered.
- if you encounter blistering of the skin it may be a sign for a condition called bullous pemphigoid. Your doctor may ask you to stop Glyxambi.

Talk to your doctor immediately if you develop a combination of symptoms of pain, tenderness, redness, or swelling of the genitals or the area between the genitals and the anus with fever or feeling generally unwell. These symptoms could be a sign of a rare but serious or even life-threatening

infection, called necrotising fasciitis of the perineum or Fournier's gangrene which destroys the tissue under the skin. Fournier's gangrene has to be treated immediately.

#### Foot care

Like for all diabetic patients it is important to check your feet regularly and adhere to any other advice regarding foot care given by your health care professional.

#### Kidney function

Before you start treatment with Glyxambi and regularly during treatment, your doctor will check how well your kidneys are working.

#### Urine glucose

Because of how this medicine works, your urine will test positive for sugar while you are taking this medicine.

#### **Children and adolescents**

This medicine is not recommended for children and adolescents under 18 years, because it has not been studied in these patients.

#### **Other medicines and Glyxambi**

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines. In particular, you should tell your doctor if you are using the following medicines:

- other anti-diabetic medicines, such as insulin or a sulphonylurea. Your doctor may want to lower the dose of these other medicines, to prevent your blood sugar levels from getting too low.
- medicines used to remove water from your body (diuretics). Your doctor may ask you to stop taking Glyxambi.
- medicines that might have an effect on the break down of empagliflozin or linagliptin in your body such as rifampicin (an antibiotic used to treat tuberculosis) or certain medicines used to treat seizures (such as carbamazepine, phenobarbital or phenytoin). The effect of Glyxambi may be reduced.
- lithium because Glyxambi can lower the amount of lithium in your blood.

#### **Pregnancy, breast-feeding and fertility**

If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

It is not known whether Glyxambi is harmful to the unborn child. As a precautionary measure it is preferable to avoid the use of this medicine during pregnancy.

It is not known whether the active substances of Glyxambi pass into human breast milk. Do not use this medicine if you are breast-feeding.

It is not known whether Glyxambi has an effect on the fertility in humans.

#### **Driving and using machines**

Glyxambi has minor influence on the ability to drive and use machines.

Taking this medicine in combination with sulphonylureas or insulin, can cause your blood sugar levels to drop too low (hypoglycaemia), which may cause symptoms such as shaking, sweating and changes in vision, and may affect your ability to drive and use machines. Do not drive or use any tools or machines, if you experience any of these symptoms while taking Glyxambi.

### **3. How to take Glyxambi**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist

if you are not sure.

### **How much to take**

The usual starting dose is one film-coated tablet of Glyxambi 10 mg/5 mg (10 mg empagliflozin and 5 mg linagliptin) once a day.

Your doctor will decide whether you need to increase your dose to one film-coated tablet of Glyxambi 25 mg/5 mg (25 mg empagliflozin and 5 mg linagliptin) once a day. If you already take 25 mg empagliflozin and 5 mg linagliptin as separate tablets and you switch to Glyxambi, you can start directly with Glyxambi 25 mg/5 mg.

### Renal impairment

Talk to your doctor if you have kidney problems. Your doctor may limit your dose or decide to use an alternative medicine.

### Hepatic impairment

Talk to your doctor in case you suffer from severe hepatic impairment. Glyxambi is not recommended and your doctor may decide to use an alternative medicine.

### Elderly

There is very little experience in patients aged 75 years or older. Treatment with Glyxambi should not be started in patients above the age of 75.

### **Taking this medicine**

- Swallow the tablet whole with water.
- You can take Glyxambi with or without food.
- You can take the tablet at any time of the day. However, try to take it at the same time each day. This will help you to remember to take it.

Your doctor may prescribe Glyxambi together with another anti-diabetic medicine. Remember to take all medicines as directed by your doctor to achieve the best results for your health.

Appropriate diet and exercise help your body to use its blood sugar better. It is important to stay on the diet and exercise program recommended by your doctor while taking Glyxambi.

### **If you take more Glyxambi than you should**

If you take more of this medicine than you should, immediately talk to a doctor or go to a hospital. Take the medicine pack with you.

### **If you forget to take Glyxambi**

What to do if you forget to take a tablet depends on how long it is until your next dose:

- If it is 12 hours or more until your next dose, take Glyxambi as soon as you remember. Then take your next dose at the usual time.
- If it is less than 12 hours until your next dose, skip the missed dose. Then take your next dose at the usual time.
- Do not take a double dose of this medicine to make up for a forgotten dose.

### **If you stop taking Glyxambi**

Do not stop taking this medicine without first consulting your doctor. Your blood sugar levels may increase when you stop taking Glyxambi.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Contact a doctor or the nearest hospital straight away if you have any of the following side effects:**

**Diabetic ketoacidosis, seen rarely (may affect up to 1 in 1 000 people)**

These are the signs of diabetic ketoacidosis (see also section 2, 'Warnings and precautions'):

- increased levels of “ketone bodies” in your urine or blood
- rapid weight loss
- feeling sick or being sick
- stomach pain
- excessive thirst
- fast and deep breathing
- confusion
- unusual sleepiness or tiredness
- a sweet smell to your breath, a sweet or metallic taste in your mouth or a different odour to your urine or sweat.

This may occur regardless of blood glucose level. Your doctor may decide to temporarily or permanently stop your treatment with this medicine.

**Contact your doctor immediately if you notice any of the following side effects:**

Allergic reactions, seen uncommonly (may affect up to 1 in 100 people)

This medicine may cause allergic reactions, which may be serious, including hives (urticaria) and swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing (angioedema).

Inflammation of the pancreas (pancreatitis), seen uncommonly

This medicine may cause pancreatitis, which usually shows as persistent, severe abdominal (stomach) pain that might reach through to your back, often accompanied by feeling sick or being sick. Your doctor will need to change your treatment.

Low blood sugar (hypoglycaemia), seen commonly (may affect up to 1 in 10 people)

If you take Glyxambi with another medicine that can cause low blood sugar, such as a sulphonylurea or insulin, you are at risk of getting too low blood sugar (hypoglycaemia). The signs of too low blood sugar may include:

- shaking, sweating, feeling very anxious or confused, fast heart beat
- excessive hunger, headache

Your doctor will tell you how to treat low blood sugar levels and what to do if you get any of the signs above. If you have symptoms of low blood sugar, eat glucose tablets, a high sugar snack or drink fruit juice. Measure your blood sugar if possible and rest.

Urinary tract infection, seen commonly

The signs of urinary tract infection are:

- burning sensation when passing urine
- urine that appears cloudy
- pain in the pelvis, or mid-back pain (when kidneys are infected)

An urge to pass urine or more frequent urination may be due to the way this medicine works, but as they can also be signs of urinary tract infection, if you note an increase in such symptoms, you should also contact your doctor.

Loss of body fluid (dehydration), seen uncommonly

The signs of dehydration are not specific, but may include:

- unusual thirst
- lightheadedness or dizziness upon standing
- fainting or loss of consciousness

## **Other side effects while taking Glyxambi:**

### **Seen commonly**

- genital yeast infection like thrush
- inflamed nose or throat (nasopharyngitis)
- cough
- passing more urine than usual or needing to pass urine more often
- itching
- skin rash
- increased blood enzyme amylase
- increased pancreas enzyme lipase
- thirst
- constipation

### **Seen uncommonly**

- straining or pain when emptying the bladder
- laboratory blood tests may show changes in blood fat levels, an increase in the amount of red blood cells (increase in haematocrit), and changes related to kidney function (decrease in filtration rate and increase in blood creatinine)

### **Seen rarely**

- sore in the mouth
- necrotising fasciitis of the perineum or Fournier's gangrene, a serious soft tissue infection of the genitals or the area between the genitals and the anus

### **Seen very rarely**

- inflammation of the kidneys (tubulointerstitial nephritis)

### **Frequency not known (cannot be estimated from the available data)**

- blistering of skin (bullous pemphigoid)

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Glyxambi**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and the carton after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not use this medicine if you notice that the packaging is damaged or shows signs of tampering.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

What Glyxambi contains

### Glyxambi 10 mg/5 mg film-coated tablets

- The active substances are empagliflozin and linagliptin. Each film-coated tablet contains 10 mg empagliflozin and 5 mg linagliptin.
- The other ingredients are:  
Tablet core: mannitol (E421), pre-gelatinised starch (maize), maize starch, copovidone, crospovidone, talc and magnesium stearate.  
Film coating: hypromellose, mannitol (E421), talc, titanium dioxide (E171), macrogol 6000 and iron oxide yellow (E172).

### Glyxambi 25 mg/5 mg film-coated tablets

- The active substances are empagliflozin and linagliptin. Each film-coated tablet contains 25 mg empagliflozin and 5 mg linagliptin.
- The other ingredients are:  
Tablet core: mannitol (E421), pre-gelatinised starch (maize), maize starch, copovidone, crospovidone, talc and magnesium stearate.  
Film coating: hypromellose, mannitol (E421), talc, titanium dioxide (E171), macrogol 6000 and iron oxide red (E172).

### **What Glyxambi looks like and contents of the pack**

Glyxambi 10 mg/5 mg film-coated tablets (tablets) are pale yellow, arc triangular, flat faced and bevel-edged. They have “10/5” on one side and the Boehringer Ingelheim logo on the other side. Each side of the tablet is 8 mm long.

Glyxambi 25 mg/5 mg film-coated tablets (tablets) are pale pink, arc triangular, flat faced and bevel-edged. They have “25/5” on one side and the Boehringer Ingelheim logo on the other side. Each side of the tablet is 8 mm long.

Glyxambi is available in PVC/PVDC/aluminium perforated unit dose blisters.

The pack sizes are 7 x 1, 10 x 1, 14 x 1, 28 x 1, 30 x 1, 60 x 1, 70 x 1, 90 x 1 and 100 x 1 film-coated tablets.

Not all pack sizes may be marketed in your country.

**Marketing Authorisation Holder**

Boehringer Ingelheim International GmbH  
Binger Str. 173  
55216 Ingelheim am Rhein  
Germany

**Manufacturer**

Boehringer Ingelheim Pharma GmbH & Co. KG  
Binger Strasse 173  
55216 Ingelheim am Rhein  
Germany

Rottendorf Pharma GmbH  
Ostenfelder Strasse 51 – 61  
59320 Ennigerloh  
Germany



For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

SCS Boehringer Ingelheim Comm.V  
Tél/Tel: +32 2 773 33 11

**България**

Бьорингер Ингелхайм РЦВ ГмбХ и Ко. КГ -  
клон България  
Тел: +359 2 958 79 98

**Česká republika**

Boehringer Ingelheim spol. s r.o.  
Tel: +420 234 655 111

**Danmark**

Boehringer Ingelheim Danmark A/S  
Tlf: +45 39 15 88 88

**Deutschland**

Boehringer Ingelheim Pharma GmbH & Co. KG  
Tel: +49 (0) 800 77 90 900

Lilly Deutschland GmbH

Tel. +49 (0) 6172 273 2222

**Eesti**

Boehringer Ingelheim RCV GmbH & Co KG  
Eesti filiaal  
Tel: +372 612 8000

**Ελλάδα**

Boehringer Ingelheim Ελλάς Μονοπρόσωπη Α.Ε.  
Τηλ: +30 2 10 89 06 300

**España**

Boehringer Ingelheim España, S.A.  
Tel: +34 93 404 51 00

Lilly S.A.

Tel: +34 91 663 50 00

**France**

Boehringer Ingelheim France S.A.S.  
Tél: +33 3 26 50 45 33

Lilly France SAS

Tél: +33 1 55 49 34 34

**Hrvatska**

Boehringer Ingelheim Zagreb d.o.o.  
Tel: +385 1 2444 600

**Lietuva**

Boehringer Ingelheim RCV GmbH & Co KG  
Lietuvos filialas  
Tel: +370 5 2595942

**Luxembourg/Luxemburg**

SCS Boehringer Ingelheim Comm.V  
Belgique/Belgien  
Tél/Tel: +32 2 773 33 11

**Magyarország**

Boehringer Ingelheim RCV GmbH & Co KG  
Magyarországi Fióktelepe  
Tel: +36 1 299 89 00

**Malta**

Boehringer Ingelheim Ireland Ltd.  
Tel: +353 1 295 9620

**Nederland**

Boehringer Ingelheim b.v.  
Tel: +31 (0) 800 22 55 889

**Norge**

Boehringer Ingelheim Norway KS  
Tlf: +47 66 76 13 00

**Österreich**

Boehringer Ingelheim RCV GmbH & Co KG  
Tel: +43 1 80 105-7870

**Polska**

Boehringer Ingelheim Sp. z o.o.  
Tel: +48 22 699 0 699

**Portugal**

Boehringer Ingelheim Portugal, Lda.  
Tel: +351 21 313 53 00

Lilly Portugal Produtos Farmacêuticos, Lda

Tel: +351 21 412 66 00

**România**

Boehringer Ingelheim RCV GmbH & Co KG  
Viena - Sucursala București  
Tel: +40 21 302 28 00

**Ireland**

Boehringer Ingelheim Ireland Ltd.  
Tel: +353 1 295 9620

Eli Lilly and Company (Ireland) Limited  
Tel: +353 1 661 4377

**Ísland**

Vistor hf.  
Sími: +354 535 7000

**Italia**

Boehringer Ingelheim Italia S.p.A.  
Tel: +39 02 5355 1

Eli Lilly Italia S.p.A.  
Tel: +39 05 5425 71

**Κύπρος**

Boehringer Ingelheim Ελλάς Μονοπρόσωπη Α.Ε.  
Τηλ: +30 2 10 89 06 300

**Latvija**

Boehringer Ingelheim RCV GmbH & Co KG  
Latvijas filiāle  
Tel: +371 67 240 011

**Slovenija**

Boehringer Ingelheim RCV GmbH & Co KG  
Podružnica Ljubljana  
Tel: +386 1 586 40 00

**Slovenská republika**

Boehringer Ingelheim RCV GmbH & Co KG  
organizačná zložka  
Tel: +421 2 5810 1211

**Suomi/Finland**

Boehringer Ingelheim Finland Ky  
Puh/Tel: +358 10 3102 800

**Sverige**

Boehringer Ingelheim AB  
Tel: +46 8 721 21 00

**United Kingdom (Northern Ireland)**

Boehringer Ingelheim Ireland Ltd.  
Tel: +353 1 295 9620

Eli Lilly and Company (Ireland) Limited  
Tel: +353 1 661 4377

**This leaflet was last revised in {MM/YYYY}.**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.