

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

1. NAME OF THE MEDICINAL PRODUCT

Imlygic 10⁶ plaque forming units (PFU)/mL solution for injection
Imlygic 10⁸ plaque forming units (PFU)/mL solution for injection

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

2.1 General description

Talimogene laherparepvec is an attenuated herpes simplex virus type-1 (HSV-1) derived by functional deletion of 2 genes (ICP34.5 and ICP47) and insertion of coding sequence for human granulocyte macrophage colony-stimulating factor (GM-CSF) (see section 5.1).

Talimogene laherparepvec is produced in Vero cells by recombinant DNA technology.

2.2 Qualitative and quantitative composition

Imlygic 10⁶ plaque forming units (PFU)/mL solution for injection

Each vial contains 1 mL deliverable volume of Imlygic at a nominal concentration of 1 x 10⁶ (1 million) plaque forming units (PFU)/mL

Imlygic 10⁸ plaque forming units (PFU)/mL solution for injection

Each vial contains 1 mL deliverable volume of Imlygic at a nominal concentration of 1 x 10⁸ (100 million) plaque forming units (PFU)/mL.

Excipient with known effect

Each 4 mL dose contains approximately 30 mg (1.3 mmol) sodium and 80 mg sorbitol.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for injection.

Imlygic 10⁶ plaque forming units (PFU)/mL solution for injection

Clear to semi-translucent liquid following thaw from its frozen state.

It may contain white, visible, variously shaped, virus-containing particles.

Imlygic 10⁸ plaque forming units (PFU)/mL solution for injection

Semi-translucent to opaque liquid following thaw from its frozen state.

It may contain white, visible, variously shaped, virus-containing particles.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Imlygic is indicated for the treatment of adults with unresectable melanoma that is regionally or distantly metastatic (Stage IIIB, IIIC and IVM1a) with no bone, brain, lung or other visceral disease (see sections 4.4 and 5.1).

4.2 Posology and method of administration

Treatment with talimogene laherparepvec should be initiated and supervised by a qualified physician experienced in the treatment of cancer.

Patients treated with Imlygic must be given the patient alert card and be informed about the risks of Imlygic (see also package leaflet).

Posology

Imlygic is provided in single use vials of 1 mL each in two different concentrations:

- 10^6 (1 million) PFU/mL - For initial dose only.
- 10^8 (100 million) PFU/mL - For all subsequent doses.

The total injection volume for each treatment visit should be up to a maximum of 4 mL. The initial recommended dose is up to a maximum of 4 mL of Imlygic at a concentration of 10^6 (1 million) PFU/mL. Subsequent doses should be administered up to 4 mL of Imlygic at a concentration of 10^8 (100 million) PFU/mL.

The recommended dosing schedule for Imlygic is shown in table 1.

Table 1. Recommended dosing schedule for Imlygic

Treatment visit	Treatment interval	Maximum total injection volume	Dose concentrations	Prioritisation of lesions to be injected
Initial	-	Up to 4 mL	10^6 (1 million) PFU/mL	<ul style="list-style-type: none">• Inject largest lesion(s) first.• Prioritise injection of remaining lesions based on lesion size until maximum injection volume is reached.
Second	3 weeks after initial treatment	Up to 4 mL	10^8 (100 million) PFU/mL	<ul style="list-style-type: none">• First inject any new lesions (lesions that may have developed since initial treatment).• Prioritise injection of remaining lesions based on lesion size until maximum injection volume is reached.

Treatment visit	Treatment interval	Maximum total injection volume	Dose concentrations	Prioritisation of lesions to be injected
All subsequent treatment visits (including re-initiation)	2 weeks after previous treatment	Up to 4 mL	10 ⁸ (100 million) PFU/mL	<ul style="list-style-type: none"> • First inject any new lesions (lesions that may have developed since previous treatment). • Prioritise injection of remaining lesions based on lesion size until maximum injection volume is reached.

Determining Imlygic dose volume (per lesion)

The volume of Imlygic to be injected into each lesion is dependent on the size of the lesion and should be determined according to table 2. The total injection volume for each treatment session should be up to a maximum of 4 mL.

Table 2. Selection of Imlygic injection volume based on lesion size

Lesion size (longest dimension)	Imlygic injection volume
> 5 cm	up to 4 mL
> 2.5 cm to 5 cm	up to 2 mL
> 1.5 cm to 2.5 cm	up to 1 mL
> 0.5 cm to 1.5 cm	up to 0.5 mL
≤ 0.5 cm	up to 0.1 mL

Patients may experience increase in size of existing lesion(s) or the appearance of a new lesion prior to achieving a response. As long as there are injectable lesion(s) remaining, Imlygic should be continued for at least 6 months unless the physician considers that the patient is not benefitting from Imlygic treatment or that other treatment is required.

Imlygic treatment may be reinitiated if new lesions appear following a complete response and the physician considers that the patient will benefit from treatment.

Special populations

Paediatric population

The safety and efficacy of Imlygic in paediatric patients has not been established. No data are available.

Elderly population

No adjustment of the dose is required in patients ≥ 65 years old (see section 5.1).

Hepatic and renal impairment

No clinical studies have been conducted to evaluate the effect of hepatic or renal impairment on the pharmacokinetics of talimogene laherparepvec. However, no adjustment in dosage is necessary for patients with hepatic or renal impairment.

Method of administration

Imlygic is to be administered by intralesional injection into cutaneous, subcutaneous, and/or nodal lesions that are visible, palpable or detectable by ultrasound guidance.

If healthcare professionals are accidentally exposed to Imlygic, see sections 4.4 and 6.6.

Healthcare professionals who are immunocompromised or pregnant should not administer Imlygic and should not come into direct contact with the Imlygic injection site(s) or body fluids of treated patients (see sections 4.3 and 4.4).

Follow the instructions below to prepare and administer Imlygic to patients:

Pre-injection

- Thaw Imlygic vial(s) at room temperature. Thawed Imlygic may be stored prior to administration (see section 6.3).
- Draw the desired amount of Imlygic from the vial into a syringe using aseptic technique. A 22- to 26-gauge needle is recommended.
- The injection site may be treated with a topical anaesthetic agent. Injectable anaesthetic may be injected around the periphery of the lesion but should not be injected directly into the lesion.
- Clean the lesion and surrounding areas with an alcohol swab and let dry.

Injection

- Inject Imlygic intralesionally into cutaneous, subcutaneous, and/or nodal lesions that are visible, palpable or detectable by ultrasound guidance.
- Determine injection volume for each lesion using table 2 above.
- Using a single insertion point, inject Imlygic along multiple tracks as far as the radial reach of the needle allows within the lesion to achieve even and complete dispersion. Multiple insertion points may be used if a lesion is larger than the radial reach of the needle.

Cutaneous lesions




Figure 1.
Injection administration for cutaneous lesions

Subcutaneous lesions




Figure 2.
Injection administration for subcutaneous lesions

Nodal lesions




Figure 3.
Injection administration for nodal lesions

- Disperse Imlygic evenly and completely within the lesion by pulling the needle back without exiting the lesion. Redirect the needle as many times as necessary while injecting the remainder of the dose of Imlygic. Continue until the full dose is evenly and completely dispersed.
- When removing the needle, withdraw it from the lesion slowly to avoid leakage or splash back of Imlygic at the insertion point.
- Repeat these steps for other lesions that need to be injected. Use a new needle anytime the needle is completely removed from a lesion and each time a different lesion is injected.

Post-injection

- Apply pressure to the injection site with a sterile gauze for at least 30 seconds.
- Swab the injection site and surrounding area with alcohol, and cover the injected lesion with an absorbent pad and dry occlusive dressing.

Disposal

Dispose of all materials that have come in contact with Imlygic (e.g. vial, syringe, needle, any cotton or gauze) in accordance with local institutional procedures (see section 6.6).

4.3 Contraindications

- Patients with a history of hypersensitivity to talimogene laherparepvec or any of its excipients.
- Patients who are severely immunocompromised (e.g. patients with severe congenital or acquired cellular and/or humoral immune deficiency) (see section 4.4).

4.4 Special warnings and precautions for use

Previously treated patients

Efficacy data for Imlygic in the current second or later line treatment settings are limited.

Immunocompromised patients

Imlygic has not been studied in immunocompromised patients. Based on animal data, patients who are severely immunocompromised may be at an increased risk of disseminated herpetic infection and should not be treated with Imlygic (see sections 4.3 and 5.3). Disseminated herpetic infection may also occur in immunocompromised patients (such as those with HIV/AIDS, leukaemia, lymphoma, common variable immunodeficiency, or who require chronic high-dose steroids or other immunosuppressive agents). The risks and benefits of treatment should be considered before administering Imlygic to these patients.

Accidental exposure to Imlygic

Accidental exposure may lead to transmission of Imlygic and herpetic infection. Healthcare professionals and close contacts (e.g. household members, caregivers, sex partners or persons sharing the same bed) should avoid direct contact with injected lesions or body fluids of treated patients during the entirety of the treatment period and up to 30 days after the last treatment administration (see section 6.6). Accidental needle stick and splash-back have been reported in healthcare professionals during preparation and administration of Imlygic.

Close contacts who are pregnant or immunocompromised should not change the patient's dressing or clean their injection site. Pregnant women, neonates, and immunocompromised individuals should not be exposed to potentially contaminated materials.

Healthcare professionals should ensure that patients are able to comply with the requirement to cover injection sites with occlusive dressings (see section 6.6). Patients should also be advised to avoid touching or scratching injection sites as this could lead to inadvertent transfer of Imlygic to other areas of their body or to their close contacts.

Although it is not known if Imlygic could be transmitted through sexual contact, it is known that wild-type HSV-1 can be transmitted through sexual contact. Patients should be advised to use a latex condom during sexual contact to prevent possible transmission of Imlygic. Women of childbearing potential should be advised to use an effective method of contraception to prevent pregnancy during treatment with Imlygic (see section 4.6).

Caregivers should be advised to wear protective gloves when assisting patients in applying or changing occlusive dressings and to observe safety precautions for disposal of used dressings and cleaning materials (see sections 4.2 and 6.6).

In the event of an accidental exposure to Imlygic, exposed individuals should be advised to clean the affected area thoroughly with soap and water and/or a disinfectant. If signs or symptoms of herpetic infection develop, they should contact their healthcare professional. Talimogene laherparepvec is sensitive to acyclovir. In case suspected herpetic lesions occur, patients, close contacts or healthcare providers have the option of follow-up testing by the Marketing Authorisation Holder for further characterisation of the infection.

Herpetic infection in Imlygic-treated patients

In clinical studies, herpetic infections (including cold sores and herpes keratitis) have been reported in patients treated with Imlygic. Symptoms of a local or systemic infection possibly related to Imlygic are anticipated to be similar to symptoms caused by wild-type HSV-1 infections.

Individuals with wild-type HSV-1 infection are known to be at a lifelong risk for symptomatic herpetic infection due to reactivation of latent wild-type HSV-1. Symptomatic herpetic infection due to possible reactivation of Imlygic should be considered.

Patients who develop herpetic infections should be advised to follow standard hygienic practices to prevent viral transmission.

Talimogene laherparepvec is sensitive to acyclovir. The risks and benefits of Imlygic treatment should be considered before administering acyclovir or other anti-viral agents indicated for management of herpetic infection. These agents may interfere with the effectiveness of Imlygic if administered systemically or topically directly to the injection site.

Cellulitis at the injection site

Necrosis or ulceration of tumour tissue may occur following Imlygic treatment. Cellulitis and systemic bacterial infection have been reported. Careful wound care and infection precautions are recommended, particularly if tissue necrosis results in open wounds.

Impaired healing at the injection site

In clinical studies, impaired healing at the injection site has been reported. Imlygic may increase the risk of impaired healing in patients with underlying risk factors (e.g. previous radiation at the injection site, or lesions in poorly vascularised areas).

The risks and benefits of Imlygic should be considered before continuing treatment if persistent infection or delayed healing develops.

Immune-mediated events

In clinical studies, immune-mediated events including as glomerulonephritis, vasculitis, pneumonitis, worsening psoriasis, and vitiligo have been reported in patients treated with Imlygic.

The risks and benefits of Imlygic should be considered before initiating treatment in patients who have underlying autoimmune disease or before continuing treatment in patients who develop immune-mediated events.

Plasmacytoma at injection site

Plasmacytoma has been reported in proximity to the injection site after administration of Imlygic. The risks and benefits of Imlygic should be considered in patients with multiple myeloma or in whom plasmacytoma develops during treatment.

Obstructive airway disorder

Obstructive airway disorder has been reported following Imlygic treatment. Caution should be used when injecting lesions close to major airways.

HSV-1 seronegative patients

Patients who were HSV-1 seronegative at baseline were reported to have a greater incidence of pyrexia, chills, and influenza-like illness compared with those who were HSV-1 seropositive at baseline, especially within the period of the first 6 treatments (see section 4.8).

All patients

This medicinal product contains 80 mg sorbitol (E420) per 4 mL dose. The additive effect of concomitantly administered products containing sorbitol (or fructose) and dietary intake of sorbitol (or fructose) should be taken into account.

This medicinal product contains approximately 30 mg sodium per 4 mL dose, equivalent to 1.5% of the WHO recommended maximum daily intake of 2 g sodium for an adult.

Traceability of Imlygic

In order to improve the traceability of biological medicinal products, the tradename and the batch number of the administered product should be clearly recorded (or stated) in the patient file.

4.5 Interaction with other medicinal products and other forms of interaction

No interaction studies have been conducted with Imlygic. Acyclovir and other anti-viral agents may interfere with the effectiveness of Imlygic if administered systemically or topically directly to the injection site. Consider the risks and benefits of Imlygic treatment before administering acyclovir or other anti-viral agents indicated for management of herpetic infection.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential/contraception

Women of childbearing potential should be advised to use an effective method of contraception to prevent pregnancy during treatment with Imlygic.

All patients should be advised to use a latex condom during sexual contact to prevent possible transmission of Imlygic (see section 4.4).

Pregnancy

Adequate and well controlled studies with talimogene laherparepvec have not been conducted in pregnant women.

If a pregnant woman has an infection with wild type HSV-1 (primary or reactivation), there is potential for the virus to cross the placental barrier, and also a risk of transmission during birth due to viral shedding. Infections with wild-type HSV-1 have been associated with serious adverse effects, including multi-organ failure and death, if a foetus or neonate contracts the wild type herpes infection.

While there are no clinical data to date on talimogene laherparepvec infections in pregnant women, there could be a risk to the foetus or neonate if talimogene laherparepvec were to act in the same manner. No effects on embryo-foetal development have been observed in animal studies (see section 5.3). As a precautionary measure, it is preferable to avoid the use of talimogene laherparepvec during pregnancy.

Transplacental metastases of malignant melanoma can occur. Because talimogene laherparepvec is designed to enter and replicate in the tumour tissue, there could be a risk of foetal exposure to talimogene laherparepvec from tumour tissue that has crossed the placenta.

If Imlygic is used during pregnancy, or if the patient becomes pregnant while taking Imlygic, the patient should be apprised of the potential hazards to the foetus and/or neonate.

Breast-feeding

It is unknown whether talimogene laherparepvec is transferred into human milk. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from Imlygic therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

Fertility

No clinical studies have been performed to evaluate the effects of talimogene laherparepvec on fertility (see section 5.3).

4.7 Effects on ability to drive and use machines

Talimogene laherparepvec may have a minor influence on the ability to drive and use machines. Because of potential adverse reactions such as dizziness and confusional state (see section 4.8), patients should be advised to use caution when driving or operating machinery until they are certain that talimogene laherparepvec does not adversely affect them.

4.8 Undesirable effects

Summary of safety profile

The safety of Imlygic was evaluated in the pivotal study where 292 patients received at least 1 dose of Imlygic (see section 5.1). The median duration of exposure to Imlygic was 23 weeks (5.3 months). Twenty six (26) patients were exposed to Imlygic for at least one year.

The most commonly reported adverse reactions ($\geq 25\%$) in Imlygic-treated patients were fatigue (50.3%), chills (48.6%), pyrexia (42.8%), nausea (35.6%), influenza-like illness (30.5%), and injection site pain (27.7%). Overall, ninety eight per cent (98%) of these adverse reactions reported were mild or moderate in severity. The most common grade 3 or higher adverse reaction was cellulitis (2.1%) (see section 4.4).

Tabulated list of adverse reactions

Adverse reactions were determined based on clinical trials in patients with melanoma treated with Imlygic compared to GM-CSF and post-marketing experience. Incidence of adverse reactions are presented by system organ class and by frequency. Frequencies are defined as: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$) and uncommon ($\geq 1/1,000$ to $< 1/100$). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Table 3. Adverse reactions from clinical trials in patients with melanoma and post-marketing experience

Infections and infestations	
Common	Cellulitis*, Oral herpes
Uncommon	Incision site infection
Neoplasms benign, malignant and unspecified (including cysts and polyps)	
Common	Tumour pain, Infected neoplasm
Uncommon	Plasmacytoma at injection site*
Blood and lymphatic system disorders	
Very common	Oedema peripheral
Common	Anaemia
Immune system disorders	
Common	Immune-mediated events [†] *
Uncommon	Hypersensitivity
Metabolism and nutrition disorders	
Common	Dehydration
Nervous system disorders	
Very common	Headache
Common	Confusional state, Anxiety, Depression, Dizziness, Insomnia
Eye disorders	
Uncommon	Keratitis herpetic
Ear and labyrinth disorders	
Common	Ear pain
Cardiac disorders	
Common	Tachycardia
Vascular disorders	
Common	Deep vein thrombosis, Hypertension, Flushing
Respiratory, thoracic and mediastinal disorders	
Very common	Cough
Common	Dyspnoea, Oropharyngeal pain, Upper respiratory tract infection
Uncommon	Obstructive airways disorder
Gastrointestinal disorders	
Very common	Vomiting, Diarrhoea, Constipation, Nausea
Common	Abdominal pain, Abdominal discomfort
Skin and subcutaneous tissue disorders	
Common	Vitiligo, Rash, Dermatitis
Uncommon	Granulomatous dermatitis
Musculoskeletal and connective tissue disorders	
Very common	Myalgia, Arthralgia, Pain in extremity
Common	Back pain, Groin pain
General disorders and administration site conditions	
Very common	Influenza like illness*, Pyrexia, Chills, Fatigue, Pain, Injection site reactions [§]
Common	Malaise, Axillary pain
Investigations	
Common	Weight decreased
Injury, poisoning and procedural complications	
Common	Wound complication, Wound secretion, Contusion, Procedural pain

[§] Injection site reactions include: very common term of injection site pain, common terms of injection site erythema, injection site haemorrhage, injection site swelling, injection site reaction, injection site inflammation, secretion discharge, injection site discharge, uncommon term of injection site warmth.

[†] Immune mediated events include: uncommon terms of vasculitis, pneumonitis, worsening psoriasis and glomerulonephritis.

* See description of selected adverse reactions

Description of selected adverse reactions

Immune-mediated events

Immune-mediated events reported in the pivotal clinical study included a case of worsening psoriasis in a patient with a prior history of psoriasis, one case of pneumonitis in a patient with a prior history of autoimmune disease, one case of vasculitis, and two cases of glomerulonephritis of which one presented with acute renal failure.

Plasmacytoma

In clinical trials, one case of plasmacytoma at injection site was observed in a patient who was found to have multiple myeloma.

Cellulitis

In the pivotal clinical trial (study 005/05), events of cellulitis were recorded, some of them being considered as serious adverse events. However, none lead to permanent discontinuation of Imlygic treatment. Careful wound care and infection precautions are recommended, particularly if tissue necrosis results in open wounds.

Influenza-like symptoms

90% of patients treated with Imlygic experienced influenza-like symptoms. Pyrexia, chills, and influenza like illness, which can occur any time during Imlygic treatment, generally resolved within 72 hours. These events were reported more frequently within the period of the first 6 treatments, particularly in patients who were HSV-1 negative at baseline.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via [the national reporting system listed in Appendix V](#).

4.9 Overdose

There is no clinical experience with overdose with Imlygic. Doses up to 4 mL at a concentration of 10^8 PFU/mL every 2 weeks have been administered in clinical trials with no evidence of dose limiting toxicity. The maximum dose of Imlygic that can be safely administered has not been determined. In the event of a suspected overdose or inadvertent intravenous administration, the patient should be treated symptomatically, e.g. with acyclovir or other anti-viral agents (see section 4.4) and supportive measures instituted as required.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antineoplastic and immunomodulating agents, ATC code: L01XX51.

Mechanism of action

Talimogene laherparepvec is an oncolytic immunotherapy that is derived from HSV-1. Talimogene laherparepvec has been modified to replicate within tumours and to produce the immune stimulatory protein human GM-CSF. Talimogene laherparepvec causes the death of tumour cells and the release of tumour-derived antigens. It is thought that together with GM-CSF, it will promote a systemic anti-

tumour immune response and an effector T-cell response. Mice that had complete regression of their primary tumours following treatment were resistant to subsequent tumour rechallenge.

The modifications to talimogene laherparepvec from HSV-1 include deletion of ICP34.5 and ICP47. Whereas anti-viral immune responses defend normal cells following infection by talimogene laherparepvec, tumours have been shown to be susceptible to injury and cell death from ICP34.5-deficient HSV-1 viruses, including talimogene laherparepvec. Deletion of ICP47 prevents down-regulation of antigen presentation molecules and increases the expression of HSV US11 gene, thereby enhancing viral replication in tumour cells.

Clinical efficacy and safety

Study 005/05

The safety and efficacy of Imlygic monotherapy compared with subcutaneously administered GM-CSF were evaluated in a phase 3, multinational, open-label, and randomised clinical study of patients with stage IIIB, IIIC, and IV melanoma that was not considered to be surgically resectable. Previous systemic treatment for melanoma was allowed but not required. Patients with active brain metastases, bone metastases, extensive visceral disease, primary ocular or mucosal melanoma, evidence of immunosuppression, or receiving treatment with a systemic anti-herpetic agent were excluded from the study.

Patients were randomised in a 2:1 ratio to receive either Imlygic or GM-CSF (N = 436; 295 Imlygic, 141 GM-CSF). Imlygic was administered by intralesional injection at an initial concentration of 10^6 (1 million) PFU/mL on day 1, followed by a concentration of 10^8 (100 million) PFU/mL on day 21 and every 2 weeks thereafter at a dose of up to 4 mL. GM-CSF was administered subcutaneously at $125 \mu\text{g}/\text{m}^2$ delivered daily for 14 days followed by a 14-day rest period in repeating intervals.

To allow for delayed immune-mediated anti-tumour effects to occur, patients were treated for a minimum of 6 months or until there were no longer any injectable lesions. During this period, treatment was to continue despite an increase in size in existing lesion(s) and/or development of new lesion(s) unless the patient developed intolerable toxicity or the investigator believed that it was in the best interest of the patient to stop treatment or to be given other therapy for melanoma. After 6 months of treatment, patients were to continue treatment until clinically relevant disease progression (i.e. disease progression associated with a decline in performance status and/or alternative therapy was required in the opinion of the investigator). Patients experiencing a response at 12 months of treatment could continue treatment for up to an additional 6 months. The mean (SD) treatment duration for the intent-to-treat (ITT) population was 15.76 weeks (15.79) in the GM-CSF arm and 26.83 weeks (18.39) in the Imlygic arm. The primary endpoint was durable response rate (DRR) [defined as the percent of patients with complete response (CR) or partial response (PR) maintained continuously for a minimum of 6 months] per blinded central review. The secondary endpoints included overall survival (OS), overall response rate (ORR) [PR+CR], time to response, duration of response, and time to treatment failure (time from randomisation until the first episode of clinically relevant disease progression where there is no response achieved after the progression event, or until death).

The mean age was 63 (range: 22 to 94) years; 26.5% were over 65 years old and 23.3% were over 74 years old. The majority of patients, 98%, were caucasian. Male patients comprised 57% of study population and 70% of patients were baseline ECOG 0 performance status. Of the enrolled patients, 22% had stage IV M1c disease and 53% of patients had received prior therapy for melanoma such as chemotherapy and cytokine-based immunotherapy in addition to surgery, adjuvant therapy or radiation. Overall, 58% of all patients enrolled into the study were seropositive for wild-type HSV-1 at baseline and 32.6% were seronegative; the HSV-1 serostatus of the remaining 9.4% was unknown.

The difference in DRR between Imlygic and GM-CSF in the ITT population was statistically significant (see table 4) in favour of Imlygic.


Table 4. Summary of results for the ITT population from Imlygic study 005/05

	Study endpoint	Imlygic N = 295	GM-CSF N = 141
Durable response rate	Primary	16.3% (n = 48) (95% CI: 12.1, 20.5)	2.1% (n = 3) (95% CI: 0.0, 4.5)
		Odds ratio 8.9; (95% CI: 2.7, 29.2) P < 0.0001	
Overall response rate (% CR, % PR)	Secondary	26.4% (n = 78) (95% CI: 21.4%, 31.5%) (10.8% CR, 15.6% PR)	5.7% (n = 8) (95% CI: 1.9%, 9.5%) (0.7% CR, 5% PR)
Overall survival	Secondary	Median 23.3 (95% CI: 19.5, 29.6) months	Median 18.9 (95% CI: 16.0, 23.7) months
		HR: 0.79; (95% CI: 0.62, 1.00) p = 0.051	
Duration of response (ongoing response at last tumour evaluation)	Secondary	Not reached (Range: > 0.0 to > 16.8 months)	Median 2.8 months (Range: 1.2 to > 14.9 months)
		HR: 0.46; (95% CI: 0.35, 0.60)	
Time to response (median)	Secondary	4.1 months	3.7 months
Time to treatment failure (median)	Secondary	8.2 months (95% CI: 6.5, 9.9)	2.9 months (95% CI: 2.8, 4.0)
		HR: 0.42; (95% CI: 0.32, 0.54)	

Among the Imlygic-treated responders, 56 (72%) responses were still ongoing at the time of primary analysis. Of the responders, 42 (54%) experienced a $\geq 25\%$ increase in overall size of existing lesion(s) and/or developed a new lesion(s) prior to ultimately achieving a response.

In an analysis to evaluate systemic activity of Imlygic, 27 of 79 patients (34.2%) had a $\geq 50\%$ overall decrease in non-visceral lesions that were not injected with Imlygic, and 8 of 71 patients (11.3%) had a $\geq 50\%$ overall decrease in visceral lesions that were not injected with Imlygic.

Figure 4. Kaplan-Meier plot –overall survival (ITT population)



No overall differences in safety or efficacy were observed between elderly (≥ 65 years old) and younger adult patients.

Exploratory subgroups


Exploratory subgroup analyses for DRR and overall survival by stage of disease were also carried out (see figure 5 and table 5). While the pivotal study was not powered to evaluate efficacy in these individual subgroups, patients with no visceral disease derived greater benefit from Imlygic treatment than those with more advanced disease.

Table 5. Summary of results from exploratory subgroup analysis from Imlygic study 005/05

	DRR, (%)		ORR, (%)		OS (hazard ratio)
	Imlygic	GM-CSF	Imlygic	GM-CSF	Imlygic vs GM-CSF
Stage [§] IIIB/IIIC/ stage IVM1a (Imlygic, n = 163; GM-CSF, n = 86)	25.2	1.2	40.5	2.3	0.57, (95% CI: 0.40, 0.80);
Stage [§] IVM1B/ IVM1C (Imlygic, n = 131; GM-CSF, n = 55)	5.3	3.6	9.2	10.9	1.07, (95% CI: 0.75, 1.52);

[§] American Joint Committee on Cancer (AJCC) staging 6th edition.

Figure 5. Kaplan-Meier estimate of overall survival by randomised treatment arm for disease stage IIIB/IIIC/ stage IVM1a (exploratory subgroup analysis)



Censor indicated by vertical bar |
 NE = not estimable

Due to the exploratory nature of the analysis and based on the current evidence, it has not been established that Imlygic is associated with an effect on overall survival.

Paediatric population

The European Medicines Agency has deferred the obligation to submit the results of studies with Imlygic in one or more subsets of the paediatric population in melanoma (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

Talimogene laherparepvec is a genetically modified and replication-competent HSV-1 virus. Therefore, its pharmacokinetics and biodistribution are driven by the site of intralesional injection, tumour-selective replication, and release from tumour tissue.

Absorption

Cellular uptake of talimogene laherparepvec occurs through HSV-1 receptors on tumours and non-tumour cells following local injection into tumours. As talimogene laherparepvec is injected and replicates intratumorally, bioavailability and systemic concentration of talimogene laherparepvec are not predictive of drug substance activity and therefore have not been evaluated.

Metabolism/elimination

Talimogene laherparepvec is cleared through general host-defence mechanisms (e.g. autophagy, adaptive immune responses). Talimogene laherparepvec is degraded by typical endogenous protein

and DNA catabolic pathways. As with other wild-type HSV-1 infections, a latent pool of talimogene laherparepvec DNA may persist in neuronal cell bodies innervating the injection sites; therefore, the occurrence of latent infection with talimogene laherparepvec cannot be excluded.

Biodistribution (within the body) and viral shedding (excretion/secretion)

Talimogene laherparepvec DNA was quantified with a highly sensitive and specific quantitative Polymerase Chain Reaction (qPCR) assay which may not correlate with viral infectivity risk. Talimogene laherparepvec was also quantified in selected patient samples in clinical studies using viral infectivity assays at the injection sites and in some cases of potential herpetic lesions.

Clinical biodistribution, elimination, and shedding

The biodistribution and shedding of intralesionally administered talimogene laherparepvec are being investigated in a melanoma study. Interim results from 30 patients show that talimogene laherparepvec DNA was detected at transient and low concentrations in blood in 90% of patients and in urine in 20% of patients in the study. The proportion of patients with detectable talimogene laherparepvec DNA in blood and urine was highest during the second cycle. Talimogene laherparepvec DNA was detected in samples from injected lesions in approximately 90% of patients. However, only 14% of patients tested positive for infective virus by 50% Tissue Culture Infectious Dose (TCID₅₀) assay, all within 8 days of treatment administration. Seventeen percent of samples from the exterior of occlusive dressing tested positive for talimogene laherparepvec DNA but none tested positive for presence of infective virus. Among samples of oral mucosa, only 1 sample had detectable talimogene laherparepvec DNA during the study, but the sample did not test positive for presence of infective virus.

Pharmacokinetics in special populations

No pharmacokinetic studies using talimogene laherparepvec have been conducted in special populations.

5.3 Preclinical safety data

At doses up to 4×10^8 PFU/kg or 10^7 PFU/dose (60-fold over the highest proposed clinical dose), single or repeated doses of talimogene laherparepvec administered by SC, IV, or intratumoural injection were well tolerated in immunocompetent mice, rats, and dogs. No neuropathology or adverse neurological effects were observed. In an *in vivo* study of intracerebral injection, talimogene laherparepvec was 10,000-fold less neurovirulent as compared to the wild-type HSV-1 dose that results in death 50% of the time in mice.

Talimogene laherparepvec was injected into various xenograft tumours at doses up to 2×10^8 PFU/kg (30-fold over the highest proposed clinical dose) in immunodeficient mice (nude and SCID). Lethal systemic viral infection was observed in up to 20% of nude mice (primarily deficient in T lymphocyte function) and 100% of SCID mice (devoid of both T and B lymphocytes).

Across studies, fatal disseminated viral infection was observed in 14% of nude mice following treatment with talimogene laherparepvec at doses that are 10 to 100-fold higher than those that result in 100% lethality with wild-type HSV-1.

Mutagenicity

The genotoxic potential of talimogene laherparepvec has not been evaluated in long-term animal or human studies. Because wild-type HSV-1 does not integrate into the host genome, the risk of insertional mutagenesis with talimogene laherparepvec is negligible.

Carcinogenicity

The carcinogenic potential of talimogene laherparepvec has not been evaluated in long-term animal or human studies. However, available data for talimogene laherparepvec and wild-type HSV-1 do not indicate a carcinogenic risk in humans.

Reproductive and development toxicity

There were no impacts to male or female reproductive tissues following treatment of adult mice at doses up to 4×10^8 PFU/kg (60-fold higher, on a PFU/kg basis, compared to the maximum clinical dose). No effects on embryo-foetal development were observed when talimogene laherparepvec was administered during organogenesis to pregnant mice at doses up to 4×10^8 (400 million) PFU/kg (60-fold higher, on a PFU/kg basis, compared to the maximum clinical dose). Negligible amounts (< 0.001% of maternal blood levels) of talimogene laherparepvec DNA were found in foetal blood.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Di-sodium phosphate dihydrate
Sodium dihydrogen phosphate dihydrate
Sodium chloride
Myo-inositol
Sorbitol (E420)
Water for injections

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

Unopened vial

5 years.

Thawing Imlygic vials

- Before use, thaw frozen Imlygic vials at room temperature (20°C to 25°C) until Imlygic is liquid (approximately 30 minutes). Gently swirl. Do NOT shake.
- Vials should be thawed and stored in the original carton until administration in order to protect from light.

After thawing

- After thawing, administer Imlygic as soon as practically feasible.
- Thawed Imlygic is stable when stored at temperatures of 2°C up to 25°C protected from light in its original vial, in a syringe, or in the original vial followed by a syringe. Do not exceed the storage times specified in table 6 and table 7.
- If storing thawed Imlygic in the original vial followed by a syringe:
 - the same temperature range should be maintained throughout the duration of storage until administration.

- the storage time in the syringe at ambient temperature up to 25°C cannot exceed 2 hours for 10⁶ (1 million) PFU/mL and 4 hours for 10⁸ (100 million) PFU/mL (see table 6).
- the maximum cumulative storage time (storage time in vial plus storage time in syringe) cannot exceed the durations in table 7.
- Imlygic must not be refrozen once it has thawed. Discard any thawed Imlygic in the vial or syringe stored longer than the specified times below.

Table 6. Maximum storage time for thawed Imlygic in syringe

	10⁶ (1 million) PFU/mL	10⁸ (100 million) PFU/mL
2°C to 8°C	8 hours	8 hours
up to 25°C	2 hours	4 hours

Table 7. Maximum cumulative storage time (storage time in vial plus storage time in syringe) for thawed Imlygic

	10⁶ (1 million) PFU/mL	10⁸ (100 million) PFU/mL
2°C to 8°C	24 hours	1 week (7 days)
up to 25°C	12 hours	24 hours


6.4 Special precautions for storage

Store and transport frozen (-90°C to -70°C).
Store in the original carton in order to protect from light.

6.5 Nature and contents of container

Imlygic is provided as a one ml preservative-free solution in a single-use vial (cyclic olefin polymer plastic resin) with stopper (chlorobutyl elastomer) and seal (aluminium) with flip-off cap (polypropylene) in two different presentations:

Figure 6. Single-use vial permanently inserted into a clear copolyester plastic sleeve



OR

Figure 7. Single-use vial without a clear plastic sleeve



The vial cap is colour coded: 10⁶ (1 million) PFU/mL is light green and 10⁸ (100 million) PFU/mL is royal blue.

6.6 Special precautions for disposal and other handling

Follow local institutional guidelines for handling and administration, personal protective equipment, accidental spills, and waste disposal.

- Wear protective gown or laboratory coat, safety glasses, or face shield and gloves while preparing or administering Imlygic. Cover any exposed wounds before administering. Avoid contact with skin, eyes or mucous membranes.
- After administration, change gloves prior to applying occlusive dressings to injected lesions. Wipe the exterior of occlusive dressing with an alcohol wipe. It is recommended to keep injection sites covered with airtight and watertight dressings at all times, if possible. To minimise the risk of viral transmission, patients should keep their injection site covered for at least 8 days from the last treatment or longer if the injection site is weeping or oozing. Advise patients to apply dressing as instructed by the healthcare professional and to replace the dressing if it falls off.
- Dispose of all materials that have come in contact with Imlygic (e.g. vial, syringe, needle, any cotton or gauze) in accordance with local institutional procedures.

Accidental exposure

- In the event of an accidental occupational exposure to Imlygic (e.g. through a splash to the eyes or mucous membranes) during preparation or administration, flush with clean water for at least 15 minutes. In the event of exposure to broken skin or needle stick, clean the affected area thoroughly with soap and water and/or disinfectant.
- Treat all Imlygic spills with a virucidal agent and absorbent materials.
- Advise patients to place used dressings and cleaning materials in a sealed plastic bag as they may be potentially contaminated, and to dispose of the bag in household waste.

This medicine contains genetically modified organisms. Unused medicine must be disposed of in compliance with the institutional guidelines for genetically modified organisms or biohazardous waste, as appropriate.

7. MARKETING AUTHORISATION HOLDER

Amgen Europe B.V.
Minervum 7061
NL-4817 ZK Breda
The Netherlands

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/15/1064/001
EU/1/15/1064/002

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 16 December 2015

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>

ANNEX II

- A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer of the biological active substance

BioVex Inc. - Subsidiary of Amgen, Inc.
34 Commerce Way
Woburn
Massachusetts
01801
United States

Name and address of the manufacturers responsible for batch release

Amgen Europe B.V.
Minervum 7061
4817 ZK Breda
Netherlands

Amgen NV
Telecomlaan 5-7
1831 Diegem
Belgium

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

• **Periodic safety update reports**

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

The marketing authorisation holder shall submit the first periodic safety update report for this product within 6 months following authorisation.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

• **Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the Marketing Authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

- **Additional risk minimisation measures**

Prior to launch of IMLYGIC in each Member State the Marketing Authorisation Holder (MAH) must agree about the content and format of the educational and controlled distribution programme, including communication media, distribution modalities, and any other aspects of the programme, with the National Competent Authority.

The educational programme is aimed to inform about important risks associated with IMLYGIC:

- Herpetic infection occurring throughout the entire body (disseminated herpetic infection) in immunocompromised individuals (those with any congenital or acquired cellular and/or humoral immune deficiency, i.e. HIV/AIDS, leukaemia, lymphoma, common variable immunodeficiency, or those who require high-dose steroids or other immunosuppressive agents)
- Accidental exposure of Healthcare Professionals (HCPs) to IMLYGIC
- Spread of IMLYGIC to close contacts or healthcare providers after direct contact with injected lesions or body fluids
- Symptomatic herpetic infection due to latency and reactivation of IMLYGIC or herpes (wild-type HSV-1) in patients
- Patients with a weakened immune system (immunocompromised patients) treated with IMLYGIC and suffering from concomitant infection
- Combination with other therapies like chemotherapy or immunosuppressive agents
- Pregnant and lactating women

The MAH shall ensure that in each Member State where IMLYGIC is marketed, all healthcare professionals and patients/carers who are expected to prescribe, dispense and use IMLYGIC have access to/are provided with the following educational package:

- Physician educational material
- Patient information pack

The physician educational material should contain:

- The Summary of Product Characteristics
- Guide for healthcare professionals
- Patient alert card

- **Guide for healthcare professionals** shall contain the following key elements:

- Information on the risk of herpetic infection in patients treated by IMLYGIC
- Information on the risk of disseminated herpetic infection in immunocompromised individuals treated by IMLYGIC
- Recommendation regarding accidental exposure of IMLYGIC to HCPs:
- To always wear protective gown/laboratory coat, safety glasses and gloves while preparing or administering IMLYGIC;
- To avoid contact with skin, eyes, mucous membranes and ungloved direct contact with injected lesions or body fluids of treated patients;
- Instruction on first aid after accidental exposure;
- Immunocompromised and pregnant healthcare professionals should not prepare and administer IMLYGIC.
- Recommendation regarding the accidental transmission of IMLYGIC from patient to close contacts or HCPs:
- Instruction on how to behave after administration/accidental transmission and how and how often the dressing has to be changed and who should not change the dressing;
- Instructions to minimise the risk of exposure of blood and body fluids to close contacts for the duration of IMLYGIC treatment through 30 days after the last administration of IMLYGIC. The following activities should be avoided:
 - Sexual intercourse without a latex condom
 - Kissing if either party has an open mouth sore
 - Common usage of cutlery, crockery, and drinking vessels
 - Common usage of injection needles, razorblades and toothbrushes;

- Adequate waste disposal and decontamination, following the recommendations for disposal of biohazardous waste.
- Information on IMLYGIC use in pregnancy
- Instructions how to handle possible adverse events including providing of batch number when reporting adverse drug reactions
- **The patient alert card** shall contain the following key messages:
 - A warning message for HCPs treating the patient at any time, including in conditions of emergency, that the patient is using IMLYGIC
 - Contact details of the IMLYGIC prescriber
 - Details about IMLYGIC treatment start date, batch number, date administered, product manufacturer and license holder
 - Information of herpetic lesions
- The patient information pack should contain:
 - Patient information leaflet
 - A patient/carer and close contacts guide
- **The Patient/carer and close contacts guide** shall contain the following key messages:
 - A description of the important risks associated with the use of IMLYGIC;
 - Instruction on how to behave after administration and how and how often the dressing has to be changed and who should not change the dressing.
 - Information of the sign and symptoms of the risk of herpetic infection;
 - Information on IMLYGIC use in pregnancy;
 - Recommendation regarding the accidental transmission of IMLYGIC from patient to close contacts or HCPs:
 - Instructions to minimise the risk of exposure of blood and body fluids to close contacts for the duration of IMLYGIC treatment through 30 days after the last administration of IMLYGIC. The following activities should be avoided:
 - Sexual intercourse without a latex condom
 - Kissing if either party has an open mouth sore
 - Common usage of cutlery, crockery, and drinking vessels
 - Common usage of injection needles, razorblades and toothbrushes;
 - Adequate waste disposal and decontamination, following the recommendations for disposal of biohazardous waste.
 - Instruction on how to behave after accidental transmission.

The controlled distribution programme is aimed to manage the product supply chain to ensure that cold storage requirements are observed and to control the distribution of IMLYGIC to qualified centres and up to the patients.

The MAH shall ensure that in each Member State where IMLYGIC is marketed, a system aimed to control distribution to IMLYGIC beyond the level of control ensured by routine risk minimisation measures. The following requirements need to be fulfilled before the product is dispensed:

- Adequately trained and experienced HCPs in order to minimise the risk of occurrence of specified adverse drug reactions in patients, HCPs, and close contacts of the patients;
- Trained HCPs and support personnel regarding safe and appropriate storage, handling, and administration of IMLYGIC, and clinical follow-up for patients treated with IMLYGIC;
- Provide specified safety information to patients and communicate to patients the importance for sharing this information with family and caregivers;
- Trained HCPs to record batch number information in patient's charts and on the patient's alert card for all injections and to provide the batch number when reporting adverse drug reactions.

- **Obligation to conduct post-authorisation measures**

The MAH shall complete, within the stated timeframe, the below measures:

Description	Due date
The MAH should submit the preliminary results of Study 20120325 (a phase 2, multicentre, open-label, single-arm trial to evaluate the correlation between objective response rate and baseline intratumoral CD8+T-lymphocyte density in subjects with unresected stage IIIB to IVM1c melanoma treated with talimogene laherparepvec)	31st December, 2018
To submit the preliminary results from Study 20110266 (a phase 2, multicentre, randomised, open-label trial assessing the efficacy and safety of talimogene laherparepvec neoadjuvant treatment plus surgery vs surgery alone for resectable stage IIIB to IVM1a melanoma)	31th December, 2019
To provide preliminary efficacy results from the phase III part of the Study 20110265 (a multicentre trial evaluating the combination of talimogene laherparepvec with pembrolizumab)	30th June, 2019

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON

1. NAME OF THE MEDICINAL PRODUCT

Imlygic 10⁶ plaque forming units (PFU)/mL solution for injection
talimogene laherparepvec

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each vial contains 1 mL of 1 x 10⁶ (1 million) plaque forming units (PFU) of talimogene laherparepvec.

3. LIST OF EXCIPIENTS

Di-sodium phosphate dihydrate, sodium dihydrogen phosphate dihydrate, sodium chloride, myo-inositol, sorbitol (E420), water for injections.

4. PHARMACEUTICAL FORM AND CONTENTS

Solution for injection

1 vial.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Intralesional use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store and transport frozen at -90° to -70°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

This medicine contains genetically modified organisms.
Unused medicine must be disposed of in compliance with the institutional guidelines for genetically modified organisms or biohazardous waste.

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Amgen Europe B.V.
Minervum 7061
NL-4817 ZK Breda
The Netherlands

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/15/1064/001

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

VIAL LABEL

1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION

Imlygic 10⁶ PFU/mL injection
talimogene laherparepvec
Intralesional use

2. METHOD OF ADMINISTRATION

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT

1 mL

6. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON

1. NAME OF THE MEDICINAL PRODUCT

Imlygic 10⁸ plaque forming units (PFU)/mL solution for injection
talimogene laherparepvec

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each vial contains 1 mL of 1 x 10⁸ (100 million) plaque forming units (PFU) of talimogene laherparepvec.

3. LIST OF EXCIPIENTS

Di-sodium phosphate dihydrate, sodium dihydrogen phosphate dihydrate, sodium chloride, myo-inositol, sorbitol (E420), water for injections.

4. PHARMACEUTICAL FORM AND CONTENTS

Solution for injection
1 vial.

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Intralesional use.
Read the package leaflet before use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

Store and transport frozen at -90° to -70°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

This medicine contains genetically modified organisms.
Unused medicine must be disposed of in compliance with the institutional guidelines for genetically modified organisms or biohazardous waste.

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Amgen Europe B.V.
Minervum 7061
NL-4817 ZK Breda
The Netherlands

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/15/1064/002

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Justification for not including Braille accepted.

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS

VIAL LABEL

1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION

Imlygic 10⁸ PFU/mL injection
talimogene laherparepvec
Intralesional use

2. METHOD OF ADMINISTRATION

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT

1 mL

6. OTHER

B. PACKAGE LEAFLET

Package leaflet: Information for the patient

Imlygic 10⁶ plaque forming units (PFU)/mL solution for injection Imlygic 10⁸ plaque forming units (PFU)/mL solution for injection talimogene laherparepvec

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your healthcare professional (doctor or nurse).
- If you get any side effects, talk to your healthcare professional. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Imlygic is and what it is used for
2. What you need to know before and during Imlygic treatment
3. How Imlygic is given
4. Possible side effects
5. How Imlygic is stored
6. Contents of the pack and other information

1. What Imlygic is and what it is used for

Imlygic is used to treat adult patients with a type of skin cancer called melanoma that has spread in the skin or to the lymph nodes, when surgery is not an option.

The active ingredient of Imlygic is talimogene laherparepvec. This is a weakened form of herpes simplex virus type-1 (HSV-1), which is commonly called the cold sore virus. To get Imlygic from HSV-1, the virus has been changed so that it multiplies more effectively in tumours than in normal cells. This leads to destruction of infected tumour cells. Imlygic also works by helping your immune system to recognise and destroy tumours throughout your body.

2. What you need to know before and during Imlygic treatment

You will not be given Imlygic:

- if you are allergic to talimogene laherparepvec or any of the other ingredients of this medicine (listed in section 6).
- if your healthcare professional has told you that you have a severely weakened immune system.

Warnings and precautions

Talk to your healthcare professional before being given Imlygic.

Patients with weakened immune systems

Life-threatening herpes infection may occur in patients with weakened immune systems. Tell your healthcare professional if you have or have ever had a weakened immune system, if you have

HIV/AIDS, blood or bone marrow cancer, or if you are taking steroids or other medicines that suppress your immune system.

Accidental spread of Imlygic to yourself and others

Imlygic can be spread to other parts of your body or to other people through direct contact with your body fluids or injection sites.

You should do the following to avoid spreading Imlygic to other areas of your body or to your close contacts (close contacts include household members, caregivers, sex partners, or someone you share a bed with):

- Avoid direct contact between your injection sites or body fluids (e.g. blood and urine) and close contacts (e.g. use latex condoms when engaging in sexual activity, avoid kissing close contacts if either of you has an open mouth sore) while you are being treated with Imlygic and up to 30 days after your last dose.
- Avoid touching or scratching the injection sites.
- Keep injection sites covered with airtight and watertight dressings at all times. Apply the dressing as instructed by your healthcare professional. If the dressing comes loose or falls off, replace it immediately with a clean dressing.
- Place all used dressings and cleaning materials in a sealed plastic bag and throw them away in your household waste.

You should tell your close contacts to:

- Avoid direct contact with your body fluids or injection sites.
- Wear gloves while changing your dressing.

If your close contacts are accidentally exposed to Imlygic, they should clean the affected area on their body with soap and water and/or a disinfectant. If they develop signs or symptoms of herpes infection, you should ask them to contact their healthcare professional. If herpetic lesions (blisters or sores) are suspected, patients or close contacts have the option of follow-up testing by the Marketing Authorisation Holder for further characterisation of the infection. Please discuss with your Healthcare Professional.

Close contacts who are pregnant or who have a weakened immune system, and newborns

Ensure that your close contacts who are pregnant or who have a weakened immune system do not touch injection sites, used dressings and cleaning materials. Keep used dressings and cleaning materials away from newborns.

Herpes infection

Cold sores or a more serious herpes infection may occur during or after treatment with Imlygic. Signs and symptoms related to treatment with Imlygic may be the same as for herpes infections, and include pain, burning or tingling in a blister around the mouth or genitals, or on the fingers or ears, eye pain, light sensitivity, discharge from the eyes, or blurry vision, weakness in arms or legs, extreme drowsiness (feeling sleepy), and mental confusion. If you have these signs, you should follow standard hygiene practices to prevent viral transmission to others. If herpetic lesions (blisters or sores) are suspected, patients or close contacts have the option of follow-up testing by the Marketing Authorisation Holder for further characterisation of the infection. Please discuss with your Healthcare Professional.

Infection and delayed healing at Imlygic injection site

Imlygic may cause infection at the injection site. Signs and symptoms of infection include pain, redness, warmth, swelling, discharge or a sore (ulcer), fever, and chills. The injection site may take longer to heal than normal. You should tell your healthcare professional if you notice any of these symptoms.

Autoimmune reactions

Imlygic may cause autoimmune reactions (an over-reaction of the body's immune system). Some people taking Imlygic have developed inflammation in the kidneys (glomerulonephritis), narrowing or blockage of blood vessels (vasculitis), swelling of the lungs (pneumonitis), worsening skin scaling (psoriasis), and areas of skin without any colour (vitiligo). Inform your healthcare professional if you have a history of autoimmune disease.

Plasmacytoma

Imlygic may cause cancerous white blood cells to collect at or near the injection site (plasmacytoma). Inform your healthcare professional if you have a history of blood cancer including multiple myeloma.

Difficulty breathing

If you have a tumour in your neck, your healthcare professional may warn you that you might experience compression of your airways during treatment with Imlygic.

Patients with no prior herpes infection

If you have never had herpes infection in the past, you may be more likely to get fever, chills, and flu-like illnesses within the period of the first 6 treatments with Imlygic.

Children and adolescents

Imlygic is not recommended for children and adolescents since the effects of Imlygic in people younger than 18 years old are not known.

Other medicines and Imlygic

Tell your healthcare professional if you are taking, have recently taken or might take any other medicines, including medicines, such as acyclovir, to treat or prevent herpes infections. Acyclovir and other anti-viral treatments may decrease the effects of Imlygic.

Pregnancy and breast-feeding

Ask your healthcare professional for advice if you:

- think you may be pregnant
- or are planning to have a baby.

Your healthcare professional will determine if Imlygic is right for you.

If you are pregnant or breast-feeding, ask your healthcare professional for advice before being given this medicine. Imlygic may harm your unborn baby.

Women who are able to become pregnant should use effective contraception to avoid pregnancy during treatment with Imlygic. Talk to your healthcare professional about suitable methods of contraception.

It is not known whether Imlygic is excreted in breast milk. It is important to tell your healthcare professional if you are breast-feeding or plan to do so. They will then help you decide whether to stop breast-feeding, or whether to stop taking Imlygic, taking into account the benefit of breast-feeding to the baby and the benefit of Imlygic to you.

Driving and using machines

When you are being treated with Imlygic you may experience symptoms such as dizziness or confusion. This may impair your ability to drive or operate machinery. Use caution when driving or operating machinery until you are certain that Imlygic does not adversely affect you.

Imlygic contains sodium and sorbitol

This medicine contains approximately 30 mg sodium (main component of cooking/table salt) in each 4 mL dose. This is equivalent to 1.5% of the recommended maximum daily dietary intake of sodium for an adult.

This medicine contains 80 mg sorbitol in each 4 mL dose.

3. How Imlygic is given

Imlygic is given in a healthcare facility under the supervision of a healthcare professional. The initial recommended dose is up to 4 mL of Imlygic at a concentration of 10^6 (1 million) PFU/mL. Subsequent doses will be up to 4 mL of Imlygic at a concentration of 10^8 (100 million) PFU/mL.

Your healthcare professional will inject Imlygic directly into your tumour(s) with a needle and syringe. Your second injection will be given 3 weeks after the first injection. After that, you will receive injections every 2 weeks for as long as you have the tumour(s).

Your healthcare professional will decide which tumour(s) to inject and may not inject every tumour. Your existing tumour(s) may increase in size and new tumour(s) could appear while you are being treated with Imlygic.

You can expect to be treated with Imlygic for at least 6 months or longer.

If you miss a dose of Imlygic

It is important for you to keep all your appointments to receive Imlygic. If you miss an appointment, ask your healthcare professional when to schedule your next dose.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Infection caused by bacteria (cellulitis) at the injection site has been commonly seen in patients receiving Imlygic. Keeping wounds clean and dressed can help prevent these types of infections.

Very commonly, flu-like illnesses, fevers and chills have been seen in patients treated with Imlygic. These symptoms generally resolve within the first 72 hours after treatment.

The following side effects have been reported in patients receiving Imlygic:

Very common (may affect more than 1 in 10 people):

- Tissue swelling (peripheral oedema)
- Headache
- Cough
- Vomiting, diarrhoea, constipation, nausea
- Muscle pain (myalgia), painful/swollen joints (arthralgia), limb pain

- Flu-like illness, fever (pyrexia), chills, fatigue, pain
- Pain, redness, bleeding, swelling, inflammation, secretion, discharge, and warmth at the injection site

Common (may affect up to 1 in 10 people):

- Infection caused by bacteria (cellulitis), cold sores (oral herpes)
- Tumour pain, infected tumour
- Low red blood cell numbers (anaemia)
- Side effects related to the immune system such as narrowing or blockage of blood vessels (vasculitis), inflammation of the lungs (pneumonitis), worsening scaling of the skin (worsening psoriasis), and inflammation of kidneys (glomerulonephritis)
- Dehydration
- Confusion, anxiety, depression, dizziness, difficulty sleeping (insomnia)
- Pain in ear, throat, abdomen, groin, back and underarm
- Faster heart rate at rest (tachycardia)
- Blood clot within a deep vein (deep vein thrombosis), high blood pressure (hypertension), redness in the face (flushing)
- Shortness of breath (dyspnoea), upper respiratory infection
- Abdominal discomfort
- Areas of skin without any colour (vitiligo), rash, inflamed skin (dermatitis)
- Generally feeling unwell
- Weight loss
- Wound complication, secretion, bruising (contusion), pain after procedure

Uncommon (may affect up to 1 in 100 people):

- Incision site infection
- Cancerous white blood cells at or near the injection site (plasmacytoma).
- Eye infection caused by herpes (keratitis herpetic)
- Compressed airways (obstructive airways disorder)
- Allergic reaction (hypersensitivity)

Reporting of side effects

If you get any side effects, talk to your healthcare professional. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How Imlygic is stored

Imlygic will be stored by the healthcare professionals at your healthcare facility.

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label and carton after EXP. The expiry date refers to the last day of that month.

Store and transport frozen at -90°C to -70°C.

Store in the original carton in order to protect from light.

6. Contents of the pack and other information

What Imlygic contains

- The active substance is talimogene laherparepvec.
Each vial contains 1 extractable mL of solution at a nominal concentration of 1×10^6 (1 million) plaque forming units (PFU)/mL or 1×10^8 (100 million) PFU/mL.
- The other ingredients are di-sodium phosphate dihydrate, sodium dihydrogen phosphate dihydrate, sodium chloride, myo-inositol, sorbitol (E420), water for injections (see section 2).

What Imlygic looks like and contents of the pack

Imlygic is a clear to semi-translucent (10^6 PFU/mL) or semi-translucent to opaque (10^8 PFU/mL) liquid. It is supplied as a 1 mL preservative free solution in a single –use vial (cyclic olefin polymer plastic resin) with stopper (chlorobutyl elastomer) and seal (aluminium) with flip-off cap (polypropylene).

The vial cap is colour coded: 10^6 PFU/mL is light green and 10^8 PFU/mL is royal blue.

Marketing Authorisation Holder and Manufacturer

Amgen Europe B.V.
Minervum 7061
NL-4817 ZK Breda
The Netherlands

Marketing Authorisation Holder

Amgen Europe B.V.
Minervum 7061
NL-4817 ZK Breda
The Netherlands

Manufacturer

Amgen NV
Telecomlaan 5-7
1831 Diegem
Belgium

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

s.a. Amgen n.v.
Tel/Tél: +32 (0)2 7752711

Lietuva

Amgen Switzerland AG Vilniaus filialas
Tel: +370 5 219 7474

България

Амджен България ЕООД
Тел.: +359 (0)2 424 7440

Luxembourg/Luxemburg

s.a. Amgen
Belgique/Belgien
Tel/Tél: +32 (0)2 7752711

Česká republika

Amgen s.r.o.
Tel: +420 221 773 500

Magyarország

Amgen Kft.
Tel.: +36 1 35 44 700

Danmark

Amgen, filial af Amgen AB, Sverige
Tlf: +45 39617500

Deutschland

AMGEN GmbH
Tel.: +49 89 1490960

Eesti

Amgen Switzerland AG Vilniaus filialas
Tel: +372 586 09553

Ελλάδα

Amgen Ελλάς Φαρμακευτικά Ε.Π.Ε.
Τηλ.: +30 210 3447000

España

Amgen S.A.
Tel: +34 93 600 18 60

France

Amgen S.A.S.
Tél: +33 (0)9 69 363 363

Hrvatska

Amgen d.o.o.
Tel: +385 (0)1 562 57 20

Ireland

Amgen Limited
United Kingdom
Tel: +44 (0)1223 420305

Ísland

Vistor hf.
Sími: +354 535 7000

Italia

Amgen S.r.l.
Tel: +39 02 6241121

Κύπρος

C.A. Papaellinas Ltd
Τηλ.: +357 22741 741

Latvija

Amgen Switzerland AG Rīgas filiāle
Tel: +371 257 25888

Malta

Amgen B.V.
The Netherlands
Tel: +31 (0)76 5732500

Nederland

Amgen B.V.
Tel: +31 (0)76 5732500

Norge

Amgen AB
Tel: +47 23308000

Österreich

Amgen GmbH
Tel: +43 (0)1 50 217

Polska

Amgen Biotechnologia Sp. z o.o.
Tel.: +48 22 581 3000

Portugal

Amgen Biofarmacêutica, Lda.
Tel: +351 21 4220550

România

Amgen România SRL
Tel: +4021 527 3000

Slovenija

AMGEN zdravila d.o.o.
Tel: +386 (0)1 585 1767

Slovenská republika

Amgen Slovakia s.r.o.
Tel: +421 2 321 114 49

Suomi/Finland

Amgen AB, sivuliike Suomessa/Amgen AB, filial
i Finland
Puh/Tel: +358 (0)9 54900500

Sverige

Amgen AB
Tel: +46 (0)8 6951100

United Kingdom

Amgen Limited
Tel: +44 (0)1223 420305

This leaflet was last revised in

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site:
<http://www.ema.europa.eu>.

The following information is intended for healthcare professionals only:

Thawing Imlygic vials

- Before use, thaw frozen Imlygic vials at room temperature (20°C to 25°C) until Imlygic is liquid (approximately 30 minutes). Gently swirl. Do NOT shake.
- Vials should be thawed and stored in the original carton until administration in order to protect from light.

After thawing

- After thawing, administer Imlygic as soon as practically feasible.
- Thawed Imlygic is stable when stored at temperatures of 2°C up to 25°C protected from light in its original vial, in a syringe, or in the original vial followed by a syringe. Do not exceed the storage times specified in table 1 and table 2.
- If storing thawed Imlygic in the original vial followed by a syringe:
 - the same temperature range should be maintained throughout the duration of storage until administration.
 - the storage time in the syringe at ambient temperature up to 25°C cannot exceed 2 hours for 10⁶ (1 million) PFU/mL and 4 hours for 10⁸ (100 million) PFU/mL (see table 1).
 - the maximum cumulative storage time (storage time in vial plus storage time in syringe) cannot exceed the durations in table 2.
- Imlygic must not be refrozen once it has thawed. Discard any thawed Imlygic in the vial or syringe stored longer than the specified times below.

Table 1. Maximum storage time for thawed Imlygic in syringe

	10⁶ (1 million) PFU/mL	10⁸ (100 million) PFU/mL
2°C to 8°C	8 hours	8 hours
up to 25°C	2 hours	4 hours

Table 2. Maximum cumulative storage time (storage time in vial plus storage time in syringe) for thawed Imlygic

	10⁶ (1 million) PFU/mL	10⁸ (100 million) PFU/mL
2°C to 8°C	24 hours	1 week (7 days)
up to 25°C	12 hours	24 hours

This medicine contains genetically modified organisms. Unused medicine must be disposed of in compliance with the institutional guidelines for genetically modified organisms or biohazardous waste, as appropriate.