







While effective viral suppression with antiretroviral therapy has been proven to substantially reduce the risk of sexual transmission, a residual risk cannot be excluded. Precautions to prevent transmission should be taken in accordance with national guidelines.

The special warnings and precautions relevant to both lamivudine and zidovudine are included in this section. There are no additional precautions and warnings relevant to the combination Lamivudine/Zidovudine Teva.

It is recommended that separate preparations of lamivudine and zidovudine should be administered in cases where dosage adjustment is necessary (see section 4.2). In these cases the physician should refer to the individual prescribing information for these medicinal products.

The concomitant use of stavudine with zidovudine should be avoided (see section 4.5).

### Opportunistic infections

Patients receiving Lamivudine/Zidovudine Teva or any other antiretroviral therapy may continue to develop opportunistic infections and other complications of HIV infection. Therefore patients should remain under close clinical observation by physicians experienced in the treatment of HIV infection.

### Haematological adverse reactions

Anaemia, neutropenia and leukopenia (usually secondary to neutropenia) can be expected to occur in patients receiving zidovudine. These occurred more frequently at higher zidovudine dosages (1200-1500 mg/day) and in patients with poor bone marrow reserve prior to treatment, particularly with advanced HIV disease. Haematological parameters should therefore be carefully monitored (see section 4.3) in patients receiving Lamivudine/Zidovudine Teva. These haematological effects are not usually observed before four to six weeks therapy. For patients with advanced symptomatic HIV disease, it is generally recommended that blood tests are performed at least every two weeks for the first three months of therapy and at least monthly thereafter.

In patients with early HIV disease haematological adverse reactions are infrequent. Depending on the overall condition of the patient, blood tests may be performed less often, for example every one to three months. Additionally dosage adjustment of zidovudine may be required if severe anaemia or myelosuppression occurs during treatment with Lamivudine/Zidovudine Teva, or in patients with pre-existing bone marrow compromise e.g. haemoglobin <9 g/dL (5.59 mmol/L) or neutrophil count <1.0 x 10<sup>9</sup>/L (see section 4.2). As dosage adjustment of Lamivudine/Zidovudine Teva is not possible separate preparations of zidovudine and lamivudine should be used. Physicians should refer to the individual prescribing information for these medicinal products.

### Pancreatitis

Cases of pancreatitis have occurred rarely in patients treated with lamivudine and zidovudine. However, it is not clear whether these cases were due to the antiretroviral treatment or to the underlying HIV disease. Treatment with Lamivudine/Zidovudine Teva should be stopped immediately if clinical signs, symptoms or laboratory abnormalities suggestive of pancreatitis occur.

### Lactic acidosis

Lactic acidosis usually associated with hepatomegaly and hepatic steatosis has been reported with the use of zidovudine. Early symptoms (symptomatic hyperlactatemia) include benign digestive symptoms (nausea, vomiting and abdominal pain) non-specific malaise, loss of appetite, weight loss, respiratory symptoms (rapid and/or deep breathing) or neurological symptoms (including motor weakness).

Lactic acidosis has a high mortality and may be associated with pancreatitis, liver failure, or renal failure.

Lactic acidosis generally occurred after a few or several months of treatment.

Treatment with zidovudine should be discontinued if there is symptomatic hyperlactatemia and metabolic/lactic acidosis, progressive hepatomegaly, or rapidly elevating aminotransferase levels.

Caution should be exercised when administering zidovudine to any patient (particularly obese women) with hepatomegaly, hepatitis or other known risk factors for liver disease and hepatic steatosis (including certain medicinal products and alcohol). Patients co-infected with hepatitis C and treated with alpha interferon and ribavirin may constitute a special risk.

Patients at increased risk should be followed closely.

#### Mitochondrial dysfunction following exposure in utero

Nucleoside and nucleotide analogues may impact mitochondrial function to a variable degree, which is most pronounced with stavudine, didanosine and zidovudine. There have been reports of mitochondrial dysfunction in HIV-negative infants exposed *in utero* and/or post-natally to nucleoside analogues; these have predominantly concerned treatment with regimens containing zidovudine. The main adverse reactions reported are haematological disorders (anaemia, neutropenia) and metabolic disorders (hyperlactatemia, hyperlipasemia). These events have often been transitory. Late-onset neurological disorders have been reported rarely (hypertonia, convulsion, abnormal behaviour). Whether such neurological disorders are transient or permanent is currently unknown. These findings should be considered for any child exposed *in utero* to nucleoside and nucleotide analogues, who presents with severe clinical findings of unknown etiology particularly neurologic findings. These findings do not affect current national recommendations to use antiretroviral therapy in pregnant women to prevent vertical transmission of HIV.

#### Lipoatrophy

Treatment with zidovudine has been associated with loss of subcutaneous fat, which has been linked to mitochondrial toxicity. The incidence and severity of lipoatrophy are related to cumulative exposure. This fat loss, which is most evident in the face, limbs and buttocks, may not be reversible when switching to a zidovudine-free regimen. Patients should be regularly assessed for signs of lipoatrophy during therapy with zidovudine and zidovudine-containing products. Therapy should be switched to an alternative regimen if there is suspicion of lipoatrophy development.

#### Weight and metabolic parameters

An increase in weight and in levels of blood lipids and glucose may occur during antiretroviral therapy. Such changes may in part be linked to disease control and life style. For lipids, there is in some cases evidence for a treatment effect, while for weight gain there is no strong evidence relating this to any particular treatment. For monitoring of blood lipids and glucose reference is made to established HIV treatment guidelines. Lipid disorders should be managed as clinically appropriate.

#### Immune Reactivation Syndrome

In HIV-infected patients with severe immune deficiency at the time of institution of combination antiretroviral therapy (CART), an inflammatory reaction to asymptomatic or residual opportunistic pathogens may arise and cause serious clinical conditions, or aggravation of symptoms. Typically, such reactions have been observed within the first few weeks or months of initiation of CART. Relevant examples are cytomegalovirus retinitis, generalised and/or focal mycobacterium infections, and *Pneumocystis jirovecii pneumonia* (often referred to as PCP). Any inflammatory symptoms should be evaluated and treatment instituted when necessary. Autoimmune disorders (such as Graves'

disease and autoimmune hepatitis) have also been reported to occur in the setting of immune reactivation; however, the reported time to onset is more variable and these events can occur many months after initiation of treatment.

#### Liver disease

If lamivudine is being used concomitantly for the treatment of HIV and hepatitis B virus (HBV), additional information relating to the use of lamivudine in the treatment of hepatitis B infection is available in the lamivudine 100 mg SmPC.

The safety and efficacy of zidovudine has not been established in patients with significant underlying liver disorders.

Patients with chronic hepatitis B or C and treated with combination antiretroviral therapy are at an increased risk of severe and potentially fatal hepatic adverse events. In case of concomitant antiviral therapy for hepatitis B or C, please refer also to the relevant product information for these medicinal products.

If Lamivudine/Zidovudine Teva is discontinued in patients co-infected with HBV, periodic monitoring of both liver function tests and markers of HBV replication for 4 months is recommended, as withdrawal of lamivudine may result in an acute exacerbation of hepatitis.

Patients with pre-existing liver dysfunction, including chronic active hepatitis, have an increased frequency of liver function abnormalities during combination antiretroviral therapy, and should be monitored according to standard practice. If there is evidence of worsening liver disease in such patients, interruption or discontinuation of treatment must be considered.

#### Patients co-infected with hepatitis C virus

The concomitant use of ribavirin with zidovudine is not recommended due to an increased risk of anaemia (see section 4.5).

#### Osteonecrosis

Although the etiology is considered to be multifactorial (including corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index), cases of osteonecrosis have been reported particularly in patients with advanced HIV-disease and/or long-term exposure to combination antiretroviral therapy (CART). Patients should be advised to seek medical advice if they experience joint aches and pain, joint stiffness or difficulty in movement.

Lamivudine/Zidovudine Teva should not be taken with any other medicinal products containing lamivudine or medicinal products containing emtricitabine.

The combination of lamivudine with cladribine is not recommended (see section 4.5).

#### Administration in subjects with moderate renal impairment

Patients with a creatinine clearance between 30 and 49 mL/min receiving Lamivudine/Zidovudine Teva may experience a 1.6-to 3.3-fold higher lamivudine exposure (AUC) than patients with a creatinine clearance  $\geq 50$  mL/min. There are no safety data from randomized, controlled trials comparing Lamivudine/Zidovudine to the individual components in patients with a creatinine clearance between 30 and 49 mL/min who received dose-adjusted lamivudine. In the original lamivudine registrational trials in combination with zidovudine, higher lamivudine exposures were associated with higher rates of haematologic toxicities (neutropenia and anaemia), although discontinuations due to neutropenia or anaemia each occurred in <1% of subjects. Other lamivudine-related adverse events (such as gastro-intestinal and hepatic disorders) may occur.

Patients with a sustained creatinine clearance between 30 and 49 mL/min who receive Lamivudine/Zidovudine Teva should be monitored for lamivudine-related adverse events, notably

haematologic toxicities. If new or worsening neutropenia or anaemia develop, a dose adjustment of lamivudine, per lamivudine prescribing information, is indicated, which cannot be achieved with Lamivudine/Zidovudine Teva. Lamivudine/Zidovudine Teva should be discontinued and the individual components should be used to construct the treatment regimen.

#### Excipient(s)

#### *Sodium*

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Lamivudine/Zidovudine Teva contains lamivudine and zidovudine, therefore any interactions identified for these individually are relevant to Lamivudine/Zidovudine Teva. Clinical studies have shown that there are no clinically significant interactions between lamivudine and zidovudine.

Zidovudine is primarily metabolised by UGT enzymes; co-administration of inducers or inhibitors of UGT enzymes could alter zidovudine exposure. Lamivudine is cleared renally. Active renal secretion of lamivudine in the urine is mediated through organic cation transporters (OCTs); co-administration of lamivudine with OCT inhibitors or nephrotoxic drugs may increase lamivudine exposure.

Lamivudine and zidovudine are not significantly metabolised by cytochrome P<sub>450</sub> enzymes (such as CYP 3A4, CYP 2C9 or CYP 2D6) nor do they inhibit or induce this enzyme system. Therefore, there is little potential for interactions with antiretroviral protease inhibitors, non-nucleosides and other medicinal products metabolised by major P<sub>450</sub> enzymes.

Interaction studies have only been performed in adults. The list below should not be considered exhaustive but is representative of the classes studied.

<b>Drugs by Therapeutic Area</b>	<b>Interaction Geometric mean change (%) (Possible mechanism)</b>	<b>Recommendation concerning co- administration</b>
<b>ANTIRETROVIRAL MEDICINAL PRODUCTS</b>		
Didanosine/Lamivudine	Interaction not studied.	No dosage adjustment necessary.
Didanosine /Zidovudine	Interaction not studied.	
Stavudine/Lamivudine	Interaction not studied.	Combination not recommended.
Stavudine/Zidovudine	In vitro antagonism of anti-HIV activity between stavudine and zidovudine could result in decreased efficacy of both drugs.	
<b>ANTI-INFECTIVE PRODUCTS</b>		
Atovaquone/Lamivudine	Interaction not studied.	As only limited data available the clinical significance is unknown.
Atovaquone/Zidovudine (750 mg twice daily with food/200 mg thrice daily)	Zidovudine AUC ↑33% Atovaquone AUC ↔	
Clarithromycin/Lamivudine	Interaction not studied.	Separate administration of Lamivudine/Zidovudine Teva and clarithromycin by at least 2 hours
Clarithromycin/Zidovudine (500 mg twice daily/100 mg every 4 hours)	Zidovudine AUC ↓12%	

Trimethoprim/sulfamethoxazole (Co-trimoxazole)/Lamivudine (160 mg/800 mg once daily for 5 days/300 mg single dose)	Lamivudine: AUC ↑40% Trimethoprim: AUC ↔ Sulfamethoxazole: AUC ↔  (organic cation transporter inhibition)	No Lamivudine/Zidovudine Teva dosage adjustment necessary, unless patient has renal impairment (See Section 4.2).  When concomitant administration with co-trimoxazole is warranted, patients should be monitored clinically. High doses of trimethoprim/sulfamethoxazole for the treatment of <i>Pneumocystis jirovecii</i> pneumonia (PCP) and toxoplasmosis have not been studied and should be avoided.
Trimethoprim/sulfamethoxazole (Co-trimoxazole)/Zidovudine	Interaction not studied.	
<b>ANTIFUNGALS</b>		
Fluconazole/Lamivudine	Interaction not studied.	As only limited data are available the clinical significance is not known. Monitor for signs of zidovudine toxicity (see section 4.8).
Fluconazole/Zidovudine (400 mg once daily/200 mg thrice daily)	Zidovudine AUC ↑74%  (UGT inhibition)	
<b>ANTIMYCOBACTERIALS</b>		
Rifampicin/Lamivudine	Interaction not studied.	Insufficient data to recommend dosage adjustment.
Rifampicin/Zidovudine (600 mg once daily/200 mg thrice daily)	Zidovudine AUC ↓48%  (UGT induction)	
<b>ANTICONVULSANTS</b>		
Phenobarbital/Lamivudine	Interaction not studied.	Insufficient data to recommend dosage adjustment.
Phenobarbital/Zidovudine	Interaction not studied.  Potential to slightly decrease zidovudine plasma concentrations through UGT induction.	
Phenytoin/Lamivudine	Interaction not studied.	Monitor phenytoin concentrations.
Phenytoin/Zidovudine	Phenytoin AUC ↑↓	
Valproic acid/Lamivudine	Interaction not studied.	As only limited data are available the clinical significance is not known. Monitor for signs of zidovudine toxicity (see section 4.8).
Valproic acid/Zidovudine (250 mg or 500 mg thrice daily/100 mg thrice daily)	Zidovudine AUC ↑80%  (UGT inhibition)	
<b>ANTI-HISTAMINES (HISTAMINE H1 RECEPTOR ANTAGONISTS)</b>		
Ranitidine/Lamivudine	Interaction not studied.  Clinically significant interaction unlikely. Ranitidine eliminated only in part by renal organic cation transport system.	No dosage adjustment necessary.
Ranitidine/Zidovudine	Interaction not studied	



Cimetidine/Lamivudine	Interaction not studied.  Clinically significant interaction unlikely. Cimetidine eliminated only in part by renal organic cation transport system.	No dosage adjustment necessary.
Cimetidine/Zidovudine	Interaction not studied.	
<b>CYTOTOXICS</b>		
<b>Cladribine/Lamivudine</b>	Interaction not studied  <i>In vitro</i> lamivudine inhibits the intracellular phosphorylation of cladribine leading to a potential risk of cladribine loss of efficacy in case of combination in the clinical setting. Some clinical findings also support a possible interaction between lamivudine and cladribine	Therefore the concomitant use of lamivudine with cladribine is not recommended (see section 4.4)
<b>OPIOIDS</b>		
Methadone/Lamivudine	Interaction not studied.	As only limited data are available the clinical significance is not known. Monitor for signs of zidovudine toxicity (see section 4.8).  Methadone dosage adjustment unlikely in majority of patients; occasionally methadone re-titration may be required.
Methadone/Zidovudine (30 to 90 mg once daily/200 mg every 4 hours)	Zidovudine AUC ↑43% Methadone AUC ↔	
<b>URICOSURIC</b>		
Probenecid/Lamivudine	Interaction not studied.	As only limited data are available the clinical significance is not known. Monitor for signs of zidovudine toxicity (see section 4.8).
Probenecid/Zidovudine (500 mg four times daily/2 mg/kg thrice daily)	Zidovudine AUC ↑106%  (UGT inhibition)	
<b>MISCELLANEOUS</b>		
Sorbitol solution (3.2g , 10.2 g, 13.4 g)/ Lamivudine	Single dose lamivudine oral solution 300 mg  Lamivudine: AUC ↓ 14%; 32%; 36% Cmax ↓ 28%; 52%, 55%.	When possible, avoid chronic coadministration of Lamivudine/Zidovudine Teva with medicinal products containing sorbitol or other osmotic acting poly-alcohols or monosaccharide alcohols (e.g. xylitol, mannitol, lactitol, maltitol). Consider more frequent monitoring of HIV-1 viral load when chronic coadministration cannot be avoided.

Abbreviations: ↑ = Increase; ↓=decrease; ↔= no significant change; AUC=area under the concentration versus time curve; C<sub>max</sub>=maximum observed concentration; CL/F=apparent oral clearance

Exacerbation of anaemia due to ribavirin has been reported when zidovudine is part of the regimen used to treat HIV although the exact mechanism remains to be elucidated. The concomitant use of ribavirin with zidovudine is not recommended due to an increased risk of anaemia (see section 4.4).

Consideration should be given to replacing zidovudine in a combination ART regimen if this is already established. This would be particularly important in patients with a known history of zidovudine induced anaemia.

Concomitant treatment, especially acute therapy, with potentially nephrotoxic or myelosuppressive medicinal products (e.g. systemic pentamidine, dapsone, pyrimethamine, co-trimoxazole, amphotericin, flucytosine, ganciclovir, interferon, vincristine, vinblastine and doxorubicin) may also increase the risk of adverse reactions to zidovudine. If concomitant therapy with Lamivudine/Zidovudine Teva and any of these medicinal products is necessary then extra care should be taken in monitoring renal function and haematological parameters and, if required, the dosage of one or more agents should be reduced.

Limited data from clinical trials do not indicate a significantly increased risk of adverse reactions to zidovudine with cotrimoxazole (see interaction information above relating to lamivudine and co-trimoxazole), aerosolised pentamidine, pyrimethamine and acyclovir at doses used in prophylaxis.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

As a general rule, when deciding to use antiretroviral agents for the treatment of HIV infection in pregnant women and consequently for reducing the risk of HIV vertical transmission to the newborn, the animal data as well as the clinical experience in pregnant women should be taken into account. In the present case, the use in pregnant women of zidovudine, with subsequent treatment of the newborn infants, has been shown to reduce the rate of maternal-foetal transmission of HIV. A large amount of data on pregnant women taking lamivudine or zidovudine indicate no malformative toxicity (more than 3000 outcomes from first trimester exposure each of which over 2000 outcomes involved exposure to both lamivudine and zidovudine). The malformative risk is unlikely in humans based on the mentioned large amount of data.

The active ingredients of Lamivudine/Zidovudine Teva may inhibit cellular DNA replication and zidovudine has been shown to be transplacental carcinogen in one animal study (see section 5.3). The clinical relevance of these findings is unknown.

For patients co-infected with hepatitis who are being treated with lamivudine containing medicinal products such as Lamivudine/Zidovudine Teva and subsequently become pregnant, consideration should be given to the possibility of a recurrence of hepatitis on discontinuation of lamivudine.

Mitochondrial dysfunction: nucleoside and nucleotide analogues have been demonstrated in vitro and in vivo to cause a variable degree of mitochondrial damage. There have been reports of mitochondrial dysfunction in HIV-negative infants exposed in utero and/or post-natally to nucleoside analogues (see section 4.4).

##### Breast-feeding

Both lamivudine and zidovudine are excreted in breast milk at similar concentrations to those found in serum.

Based on more than 200 mother/child pairs treated for HIV, serum concentrations of lamivudine in breastfed infants of mothers treated for HIV are very low (< 4% of maternal serum concentrations) and progressively decrease to undetectable levels when breastfed infants reach 24 weeks of age. There are no data available on the safety of lamivudine when administered to babies less than three months old.

After administration of a single dose of 200 mg zidovudine to HIV-infected women, the mean concentration of zidovudine was similar in human milk and serum.

It is recommended that mothers infected by HIV do not breast-feed their infants under any circumstances in order to avoid transmission of HIV.

### Fertility

Neither zidovudine nor lamivudine have shown evidence of impairment of fertility in studies in male and female rats. There are no data on their effect on human female fertility. In men zidovudine has not been shown to affect sperm count, morphology or motility.

### **4.7 Effects on ability to drive and use machines**

No studies on the effects on the ability to drive and use machines have been performed.

### **4.8 Undesirable effects**

Adverse reactions have been reported during therapy for HIV disease with lamivudine and zidovudine separately or in combination. For many of these events, it is unclear whether they are related to lamivudine, zidovudine, the wide range of medicinal products used in the management of HIV disease, or as a result of the underlying disease process.

As Lamivudine/Zidovudine Teva contains lamivudine and zidovudine, the type and severity of adverse reactions associated with each of the compounds may be expected. There is no evidence of added toxicity following concurrent administration of the two compounds.

Cases of lactic acidosis, sometimes fatal, usually associated with severe hepatomegaly and hepatic steatosis, have been reported with the use of zidovudine (see section 4.4).

Treatment with zidovudine has been associated with loss of subcutaneous fat which is most evident in the face, limbs and buttocks. Patients receiving Lamivudine/Zidovudine Teva should be frequently examined and questioned for signs of lipoatrophy. When such development is found, treatment with Lamivudine/Zidovudine Teva should not be continued (see section 4.4).

Weight and levels of blood lipids and glucose may increase during antiretroviral therapy (see section 4.4).

In HIV-infected patients with severe immune deficiency at the time of initiation of combination antiretroviral therapy (CART), an inflammatory reaction to asymptomatic or residual opportunistic infections may arise. Autoimmune disorders (such as Graves' disease and autoimmune hepatitis) have also been reported to occur in the setting of immune reactivation; however, the reported time to onset is more variable and these events can occur many months after initiation of treatment (see section 4.4).

Cases of osteonecrosis have been reported, particularly in patients with generally acknowledged risk factors, advanced HIV disease or long-term exposure to CART. The frequency of this is unknown (see section 4.4).

### **Lamivudine**

The adverse reactions considered at least possibly related to the treatment are listed below by body system, organ class and absolute frequency. Frequencies are defined as very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1000$ ), very rare ( $< 1/10,000$ ). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

#### Blood and lymphatic systems disorders

*Uncommon:* Neutropenia and anaemia (both occasionally severe), thrombocytopenia

*Very rare:* Pure red cell aplasia

#### Metabolism and nutrition disorders

*Very Rare:* Lactic acidosis

#### Nervous system disorders

*Common:* Headache, insomnia

*Very rare:* Peripheral neuropathy (or paraesthesiae)

#### Respiratory, thoracic and mediastinal disorders

*Common:* Cough, nasal symptoms

#### Gastrointestinal disorders

*Common:* Nausea, vomiting, abdominal pain or cramps, diarrhoea

*Rare:* Pancreatitis, rises in serum amylase

#### Hepatobiliary disorders

*Uncommon:* Transient rises in liver enzymes (AST, ALT)

*Rare:* Hepatitis

#### Skin and subcutaneous tissue disorders

*Common:* Rash, alopecia

*Rare:* Angioedema

#### Musculoskeletal and connective tissue disorders

*Common:* Arthralgia, muscle disorders

*Rare:* Rhabdomyolysis

#### General disorders and administration site conditions

*Common:* Fatigue, malaise, fever

### **Zidovudine**

The adverse reactions profile appears similar for adults and adolescents. The most serious adverse reactions include anaemia (which may require transfusions), neutropenia and leukopenia. These occurred more frequently at higher dosages (1200-1500 mg/day) and in patients with advanced HIV disease (especially when there is poor bone marrow reserve prior to treatment), and particularly in patients with CD4 cell counts less than 100/mm<sup>3</sup> (see section 4.4).

The incidence of neutropenia was also increased in those patients whose neutrophil counts, haemoglobin levels and serum vitamin B<sub>12</sub> levels were low at the start of zidovudine therapy.

The adverse reactions considered at least possibly related to the treatment are listed below by body system, organ class and absolute frequency. Frequencies are defined as very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1000$ ), very rare ( $< 1/10,000$ ). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

#### Blood and lymphatic system disorders

*Common:* Anaemia, neutropenia and leukopenia

*Uncommon:* Thrombocytopenia and pancytopenia (with marrow hypoplasia)

*Rare:* Pure red cell aplasia

*Very rare:* Aplastic anaemia

#### Metabolism and nutrition disorders

*Rare:* Lactic acidosis in the absence of hypoxaemia, anorexia

#### Psychiatric disorders

*Rare:* Anxiety and depression

#### Nervous system disorders

*Very common:* Headache

*Common:* Dizziness

*Rare:* Insomnia, paraesthesiae, somnolence, loss of mental acuity, convulsions

#### Cardiac disorders

*Rare:* Cardiomyopathy

#### Respiratory, thoracic and mediastinal disorders

*Uncommon:* Dyspnoea

*Rare:* Cough

#### Gastrointestinal disorders

*Very common:* Nausea

*Common:* Vomiting, abdominal pain and diarrhoea

*Uncommon:* Flatulence

*Rare:* Oral mucosa pigmentation, taste perversion and dyspepsia. Pancreatitis

#### Hepatobiliary disorders

*Common:* Raised blood levels of liver enzymes and bilirubin

*Rare:* Liver disorders such as severe hepatomegaly with steatosis

#### Skin and subcutaneous tissue disorders

*Uncommon:* Rash and pruritus

*Rare:* Nail and skin pigmentation, urticaria and sweating

## Musculoskeletal and connective tissue disorders

*Common:* Myalgia

*Uncommon:* Myopathy

## Renal and urinary disorders

*Rare:* Urinary frequency

## Reproductive system and breast disorders

*Rare:* Gynaecomastia

## General disorders and administration site conditions

*Common:* Malaise

*Uncommon:* Fever, generalised pain and asthenia

*Rare:* Chills, chest pain and influenza-like syndrome

The available data from both placebo-controlled and open-label studies indicate that the incidence of nausea and other frequently reported clinical adverse events consistently decreases over time during the first few weeks of therapy with zidovudine.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

## **4.9 Overdose**

There is limited experience of overdosage with lamivudine/zidovudine. No specific symptoms or signs have been identified following acute overdose with zidovudine or lamivudine apart from those listed as undesirable effects.

If overdosage occurs the patient should be monitored for evidence of toxicity (see section 4.8), and standard supportive treatment applied as necessary. Since lamivudine is dialysable, continuous haemodialysis could be used in the treatment of overdosage, although this has not been studied. Haemodialysis and peritoneal dialysis appear to have a limited effect on elimination of zidovudine, but enhance the elimination of the glucuronide metabolite. For more details physicians should refer to the individual prescribing information for lamivudine and zidovudine.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Antivirals for treatment of HIV infections, combinations, ATC code: J05AR01

Lamivudine and zidovudine are nucleoside analogues which have activity against HIV. Additionally, lamivudine has activity against hepatitis B virus (HBV). Both medicinal products are metabolised intracellularly to their active moieties, lamivudine 5'-triphosphate (TP) and zidovudine 5'-TP respectively. Their main modes of action are as chain terminators of viral reverse transcription. Lamivudine-TP and zidovudine-TP have selective inhibitory activity against HIV-1 and HIV-2 replication *in vitro*; lamivudine is also active against zidovudine-resistant clinical isolates of HIV. No

antagonistic effects *in vitro* were seen with lamivudine and other antiretrovirals (tested agents: abacavir, didanosine and nevirapine). No antagonistic effects *in vitro* were seen with zidovudine and other antiretrovirals (tested agents: abacavir, didanosine and interferon-alpha).

HIV-1 resistance to lamivudine involves the development of a M184V amino acid change close to the active site of the viral reverse transcriptase (RT). This variant arises both *in vitro* and in HIV-1 infected patients treated with lamivudine-containing antiretroviral therapy. M184V mutants display greatly reduced susceptibility to lamivudine and show diminished viral replicative capacity *in vitro*. *In vitro* studies indicate that zidovudine-resistant virus isolates can become zidovudine sensitive when they simultaneously acquire resistance to lamivudine. The clinical relevance of such findings remains, however, not well defined.

*In vitro* data tend to suggest that the continuation of lamivudine in anti-retroviral regimen despite the development of M184V might provide residual anti-retroviral activity (likely through impaired viral fitness). The clinical relevance of these findings is not established. Indeed, the available clinical data are very limited and preclude any reliable conclusion in the field. In any case, initiation of susceptible nucleoside analogue reverse-transcriptase inhibitors (NRTIs) should always be preferred to maintenance of lamivudine therapy. Therefore, maintaining lamivudine therapy despite emergence of M184V mutation should only be considered in cases where no other active NRTIs are available.

Cross-resistance conferred by the M184V RT is limited within the nucleoside inhibitor class of antiretroviral agents. Zidovudine and stavudine maintain their antiretroviral activities against lamivudine-resistant HIV-1. Abacavir maintains its antiretroviral activities against lamivudine-resistant HIV-1 harbouring only the M184V mutation. The M184V RT mutant shows a <4-fold decrease in susceptibility to didanosine; the clinical significance of these findings is unknown. *In vitro* susceptibility testing has not been standardised and results may vary according to methodological factors.

Lamivudine demonstrates low cytotoxicity to peripheral blood lymphocytes, to established lymphocyte and monocyte-macrophage cell lines, and to a variety of bone marrow progenitor cells *in vitro*. Resistance to thymidine analogues (of which zidovudine is one) is well characterised and is conferred by the stepwise accumulation of up to six specific mutations in the HIV reverse transcriptase at codons 41, 67, 70, 210, 215 and 219. Viruses acquire phenotypic resistance to thymidine analogues through the combination of mutations at codons 41 and 215 or by the accumulation of at least four of the six mutations. These thymidine analogue mutations alone do not cause high-level cross-resistance to any of the other nucleosides, allowing for the subsequent use of any of the other approved reverse transcriptase inhibitors.

Two patterns of multi-drug resistance mutations, the first characterised by mutations in the HIV reverse transcriptase at codons 62, 75, 77, 116 and 151 and the second involving a T69S mutation plus a 6-base pair insert at the same position, result in phenotypic resistance to AZT as well as to the other approved NRTIs. Either of these two patterns of multinucleoside resistance mutations severely limits future therapeutic options.

### Clinical Experience

In clinical trials, lamivudine in combination with zidovudine has been shown to reduce HIV-1 viral load and increase CD4 cell count. Clinical end-point data indicate that lamivudine in combination with zidovudine, results in a significant reduction in the risk of disease progression and mortality.

Lamivudine and zidovudine have been widely used as components of antiretroviral combination therapy with other antiretroviral agents of the same class (NRTIs) or different classes (PIs, non-nucleoside reverse transcriptase inhibitors).

Multiple drug antiretroviral therapy containing lamivudine has been shown to be effective in antiretroviral-naïve patients as well as in patients presenting with viruses containing the M184V mutations.

Evidence from clinical studies shows that lamivudine plus zidovudine delays the emergence of zidovudine resistant isolates in individuals with no prior antiretroviral therapy. Subjects receiving lamivudine and zidovudine with or without additional concomitant antiretroviral therapies and who already present with the M184V mutant virus also experience a delay in the onset of mutations that confer resistance to zidovudine and stavudine (Thymidine Analogue Mutations; TAMs).

The relationship between *in vitro* susceptibility of HIV to lamivudine and zidovudine and clinical response to lamivudine/zidovudine containing therapy remains under investigation.

Lamivudine at a dose of 100 mg once daily has also been shown to be effective for the treatment of adult patients with chronic HBV infection (for details of clinical studies, see the prescribing information for lamivudine 100 mg). However, for the treatment of HIV infection only a 300 mg daily dose of lamivudine (in combination with other antiretroviral agents) has been shown to be efficacious.

Lamivudine has not been specifically investigated in HIV patients co-infected with HBV.

## 5.2 Pharmacokinetic properties

### Absorption

Lamivudine and zidovudine are well absorbed from the gastrointestinal tract. The bioavailability of oral lamivudine in adults is normally between 80-85 % and for zidovudine 60-70 %.

A bioequivalence study compared lamivudine/zidovudine with lamivudine 150 mg and zidovudine 300 mg tablets taken together. The effect of food on the rate and extent of absorption was also studied. Lamivudine/zidovudine was shown to be bioequivalent to lamivudine 150 mg and zidovudine 300 mg given as separate tablets, when administered to fasting subjects.

Following single dose lamivudine/zidovudine administration in healthy volunteers, mean (CV) lamivudine and zidovudine  $C_{max}$  values were 1.6 µg/mL (32 %) and 2.0 µg/mL (40 %), respectively and the corresponding values for AUC were 6.1 µg h/mL (20 %) and 2.4 µg h/mL (29 %) respectively. The median (range) lamivudine and zidovudine  $t_{max}$  values were 0.75 (0.50-2.00) hours and 0.50 (0.25-2.00) hours respectively. The extent of lamivudine and zidovudine absorption ( $AUC_{\infty}$ ) and estimates of half-life following administration of lamivudine/zidovudine with food were similar when compared to fasting subjects, although the rates of absorption ( $C_{max}$ ,  $t_{max}$ ) were slowed. Based on these data lamivudine/zidovudine may be administered with or without food.

Administration of crushed tablets with a small amount of semi-solid food or liquid would not be expected to have an impact on the pharmaceutical quality, and would therefore not be expected to alter the clinical effect. This conclusion is based on the physicochemical and pharmacokinetic data assuming that the patient crushes and transfers 100 % of the tablet and ingests immediately.

### Distribution

Intravenous studies with lamivudine and zidovudine showed that the mean apparent volume of distribution is 1.3 and 1.6 l/kg respectively. Lamivudine exhibits linear pharmacokinetics over the therapeutic dose range and displays limited binding to the major plasma protein albumin (<36 % serum albumin *in vitro*). Zidovudine plasma protein binding is 34 % to 38 %. Interactions involving binding site displacement are not anticipated with Lamivudine/Zidovudine Teva.

Data show that lamivudine and zidovudine penetrate the central nervous system (CNS) and reach the cerebrospinal fluid (CSF). The mean ratios of CSF/serum lamivudine and zidovudine concentrations



2-4 hours after oral administration were approximately 0.12 and 0.5 respectively. The true extent of CNS penetration of lamivudine and its relationship with any clinical efficacy is unknown.

### Biotransformation

Metabolism of lamivudine is a minor route of elimination. Lamivudine is predominately cleared unchanged by renal excretion. The likelihood of metabolic drug interactions with lamivudine is low due to the small extent of hepatic metabolism (5-10 %) and low plasma binding.

The 5'-glucuronide of zidovudine is the major metabolite in both plasma and urine, accounting for approximately 50-80 % of the administered dose eliminated by renal excretion. 3'-amino-3'-deoxythymidine (AMT) has been identified as a metabolite of zidovudine following intravenous dosing.

### Elimination

The observed lamivudine half-life of elimination is 18 to 19 hours. The mean systemic clearance of lamivudine is approximately 0.32 l/h/kg, with predominantly renal clearance (>70 %) via the organic cationic transport system. Studies in patients with renal impairment show lamivudine elimination is affected by renal dysfunction. Dose reduction is required for patients with creatinine clearance  $\leq 30$  mL/min (see section 4.2).

From studies with intravenous zidovudine, the mean terminal plasma half-life was 1.1 hours and the mean systemic clearance was 1.6 l/h/kg. Renal clearance of zidovudine is estimated to be 0.34 l/h/kg, indicating glomerular filtration and active tubular secretion by the kidneys. Zidovudine concentrations are increased in patients with advanced renal failure.

### Pharmacokinetics in children

In children over the age of 5-6 months, the pharmacokinetic profile of zidovudine is similar to that in adults. Zidovudine is well absorbed from the gut and at all dose levels studied in adults and children, the bioavailability was between 60-74 % with a mean of 65 %.  $C_{SSmax}$  levels were 4.45  $\mu$ M (1.19  $\mu$ g/mL) following a dose of 120 mg zidovudine (in solution)/m<sup>2</sup> body surface area and 7.7  $\mu$ M (2.06  $\mu$ g/mL) at 180 mg/m<sup>2</sup> body surface area. Dosages of 180 mg/m<sup>2</sup> four times daily in children produced similar systemic exposure (24 hour AUC 40.0 h  $\mu$ M or 10.7 h  $\mu$ g/mL) as doses of 200 mg six times daily in adults (40.7 h  $\mu$ M or 10.9 h  $\mu$ g/mL).

In six HIV-infected children from 2 to 13 years of age, zidovudine plasma pharmacokinetics were evaluated while subjects were receiving 120 mg/m<sup>2</sup> zidovudine three times daily and again after switching to 180 mg/m<sup>2</sup> twice daily. Systemic exposures (daily AUC and  $C_{max}$ ) in plasma from the twice daily regimen appeared equivalent to those from the same total daily dose given in three divided doses [Bergshoeff, 2004].

In general, lamivudine pharmacokinetics in paediatric patients are similar to adults. However, absolute bioavailability (approximately 55-65 %) was reduced in paediatric patients below 12 years of age. In addition, systemic clearance values were greater in younger paediatric patients and decreased with age, approaching adult values around 12 years of age. Due to these differences, the recommended dose for lamivudine in children (aged more than three months and weighing less than 30 kg) is 4 mg/kg twice a day. This dose will achieve an average AUC<sub>0-12</sub> ranging from approximately 3,800 to 5,300 ng h/mL. Recent findings indicate that exposure in children <6 years of age may be reduced by about 30 % compared with other age groups. Further data addressing this issue are currently awaited. At present, the available data do not suggest that lamivudine is less efficacious in this age group.

### Pharmacokinetics in pregnancy

The pharmacokinetics of lamivudine and zidovudine were similar to that of non-pregnant women.

### 5.3 Preclinical safety data

The clinically relevant effects of lamivudine and zidovudine in combination are anaemia, neutropenia and leukopenia.

#### Mutagenicity and carcinogenicity

Neither lamivudine nor zidovudine are mutagenic in bacterial tests, but consistent with other nucleoside analogues, inhibit cellular DNA replication in *in vitro* mammalian tests such as the mouse lymphoma assay.

Lamivudine has not shown any genotoxic activity in *in vivo* studies at doses that gave plasma concentrations up to 40-50 times higher than clinical plasma levels. Zidovudine showed clastogenic effects in an oral repeated dose micronucleus test in mice. Peripheral blood lymphocytes from acquired immune deficiency syndrome (AIDS) patients receiving zidovudine treatment have also been observed to contain higher numbers of chromosome breakages.

A pilot study has demonstrated that zidovudine is incorporated into leukocyte nuclear DNA of adults, including pregnant women, taking zidovudine as treatment for HIV-1 infection, or for the prevention of mother to child viral transmission. Zidovudine was also incorporated into DNA from cord blood leukocytes of infants from zidovudine-treated mothers. A transplacental genotoxicity study conducted in monkeys compared zidovudine alone with the combination of zidovudine and lamivudine at human-equivalent exposures. The study demonstrated that foetuses exposed *in utero* to the combination sustained a higher level of nucleoside analogue-DNA incorporation into multiple foetal organs, and showed evidence of more telomere shortening than in those exposed to zidovudine alone. The clinical significance of these findings is unknown.

The carcinogenic potential of a combination of lamivudine and zidovudine has not been tested.

In long-term oral carcinogenicity studies in rats and mice, lamivudine did not show any carcinogenic potential.

In oral carcinogenicity studies with zidovudine in mice and rats, late appearing vaginal epithelial tumours were observed. A subsequent intravaginal carcinogenicity study confirmed the hypothesis that the vaginal tumours were the result of long term local exposure of the rodent vaginal epithelium to high concentrations of unmetabolised zidovudine in urine. There were no other zidovudine-related tumours observed in either sex of either species.

In addition, two transplacental carcinogenicity studies have been conducted in mice. In one study, by the US National Cancer Institute, zidovudine was administered at maximum tolerated doses to pregnant mice from day 12 to 18 of gestation. One year post-natally, there was an increase in the incidence of tumours in the lung, liver and female reproductive tract of offspring exposed to the highest dose level (420 mg/kg term body weight).

In a second study, mice were administered zidovudine at doses up to 40 mg/kg for 24 months, with exposure beginning prenatally on gestation day 10. Treatment related findings were limited to late-occurring vaginal epithelial tumours, which were seen with a similar incidence and time of onset as in the standard oral carcinogenicity study. The second study thus provided no evidence that zidovudine acts as a transplacental carcinogen.

While the clinical relevance of these findings is unknown, these data suggest that a carcinogenic risk to humans is outweighed by the potential clinical benefit.

In reproductive toxicity studies lamivudine has demonstrated evidence of causing an increase in early embryonic deaths in the rabbit at relatively low systemic exposures, comparable to those achieved in

man, but not in the rat even at very high systemic exposure. Zidovudine had a similar effect in both species, but only at very high systemic exposures. Lamivudine was not teratogenic in animal studies. At maternally toxic doses, zidovudine given to rats during organogenesis resulted in an increased incidence of malformations, but no evidence of foetal abnormalities was observed at lower doses.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Cellulose, microcrystalline  
Sodium starch glycolate (Type A)  
Sodium stearyl fumarate

#### Tablet film-coat

Hypromellose 3cP  
Hypromellose 6cP  
Polysorbate 80  
Macrogol 400  
Titanium dioxide E171

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

30 months

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

Blisters:  
OPA/Alu/PVC Aluminium blisters

Containers:  
White opaque HDPE containers with white opaque polyethylene child resistant screw cap with Aluminium seal.

Each pack type contains 60 film-coated tablets.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

No special requirements for disposal.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

Teva B.V.  
Swensweg 5  
2031GA Haarlem  
The Netherlands

**8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/10/663/001  
EU/1/10/663/002

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 02 March 2011  
Date of latest renewal: 19 November 2015

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

Medicinal product no longer authorised

**ANNEX II**

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARDS TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

Medicinal product no longer authorised

## **A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturers responsible for batch release

Teva Pharmaceutical Works Private Limited Company  
Pallagi út 13  
Debrecen H-4042  
Hungary

Pharmachemie B.V.  
Swensweg 5  
2031 GA Haarlem  
The Netherlands

Teva Operations Poland Sp. z.o.o.  
80 Mogilska St.  
31-546 Krakow  
Poland

Merckle GmbH  
Ludwig-Merckle-Strasse 3  
89143 Blaubeuren  
Germany

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk management plan (RMP)**

Not applicable.

Medicinal product no longer authorised

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

**A. LABELLING**

Medicinal product no longer authorised



**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON – BLISTER PACK**

**1. NAME OF THE MEDICINAL PRODUCT**

Lamivudine/Zidovudine Teva 150 mg/300 mg film-coated tablets  
lamivudine/zidovudine

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains  
lamivudine 150 mg  
zidovudine 300 mg

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

60 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Teva B.V.  
Swensweg 5  
2031GA Haarlem  
The Netherlands

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/10/663/001  
EU/1/10/663/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Lamivudine/Zidovudine Teva

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTER FOIL**

**1. NAME OF THE MEDICINAL PRODUCT**

Lamivudine/Zidovudine Teva 150 mg/300 mg film-coated tablets  
lamivudine/zidovudine

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Teva B.V.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

Medicinal product no longer authorised

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON FOR THE CONTAINER**

**1. NAME OF THE MEDICINAL PRODUCT**

Lamivudine/Zidovudine Teva 150 mg/300 mg film-coated tablets  
lamivudine/zidovudine

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains  
lamivudine 150 mg  
zidovudine 300 mg

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

60 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Teva B.V.  
Swensweg 5  
2031GA Haarlem  
The Netherlands

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/10/663/001  
EU/1/10/663/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Lamivudine/Zidovudine Teva

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**CONTAINERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Lamivudine/Zidovudine Teva 150 mg/300 mg film-coated tablets  
lamivudine/zidovudine

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains  
lamivudine 150 mg  
zidovudine 300 mg

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

60 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Teva B.V.  
Swensweg 5  
2031GA Haarlem  
The Netherlands

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/10/663/001  
EU/1/10/663/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D-BARCODE**

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

Medicinal product no longer authorised

**B. PACKAGE LEAFLET**

Medicinal product no longer authorised



## Package leaflet: Information for the user

### Lamivudine/Zidovudine Teva 150 mg/300 mg film-coated tablets lamivudine/zidovudine

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet:

1. What Lamivudine/Zidovudine Teva is and what it is used for
2. What you need to know before you take Lamivudine/Zidovudine Teva
3. How to take Lamivudine/Zidovudine Teva
4. Possible side effects
5. How to store Lamivudine/Zidovudine Teva
6. Contents of the pack and other information

#### 1. What Lamivudine/Zidovudine Teva is and what it is used for

**Lamivudine/Zidovudine Teva is used to treat HIV (human immunodeficiency virus) infection in adults and children.**

Lamivudine/Zidovudine Teva contains two active ingredients that are used to treat HIV infection: lamivudine and zidovudine. Both of these belong to a group of anti-retroviral medicines called *nucleoside analogue reverse transcriptase inhibitors (NRTIs)*.

Lamivudine/Zidovudine Teva does not completely cure HIV infection; it reduces the amount of virus in your body, and keeps it at a low level. It also increases the CD4 cell count in your blood. CD4 cells are a type of white blood cells that are important in helping your body to fight infection.

Not everyone responds to treatment with Lamivudine/Zidovudine Teva in the same way. Your doctor will monitor the effectiveness of your treatment.

#### 2. What you need to know before you take Lamivudine/Zidovudine Teva

##### **Do not take Lamivudine/Zidovudine Teva**

- if you are allergic to lamivudine or zidovudine or any of the other ingredients of this medicine (listed in section 6).
- if you have a **very low red blood cell count (anaemia)** or a **very low white blood cell count (neutropenia)**.

**Check with your doctor** if you think any of these apply to you.

##### **Take special care with Lamivudine/Zidovudine Teva**

Some people taking Lamivudine/Zidovudine Teva or other combination treatments for HIV are more at risk of serious side effects. You need to be aware of the extra risks:

- if you have ever had **liver disease**, including hepatitis B or C (if you have hepatitis B infection, do not stop Lamivudine/Zidovudine Teva without your doctor's advice, as your hepatitis may come back)
- if you have **kidney disease**
- if you are seriously **overweight** (especially if you are a woman).

**Talk to your doctor if any of these apply to you.** Your doctor will decide if the active substances are suitable for you. You may need extra check-ups, including blood tests, while you are taking your medicine. See Section 4 for more information.

### **Look out for important symptoms**

Some people taking medicines for HIV infection develop other conditions, which can be serious. You need to know about important signs and symptoms to look out for while you are taking Lamivudine/Zidovudine Teva.

**Read the information 'Other possible side effects of combination therapy for HIV' in Section 4 of this leaflet.**

### **Protect other people**

HIV infection is spread by sexual contact with someone who has the infection, or by transfer of infected blood (for example, by sharing injection needles). You can still pass on HIV when taking this medicine, although the risk is lowered by effective antiretroviral therapy.

Discuss with your doctor the precautions needed to avoid infecting other people.

### **Other medicines and Lamivudine/Zidovudine Teva**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Remember to tell your doctor or pharmacist if you begin taking a new medicine while you are taking Lamivudine/Zidovudine Teva.

### **These medicines should not be used with Lamivudine/Zidovudine Teva:**

- other medicinal products containing lamivudine, to treat **HIV infection or hepatitis B infection**
- emtricitabine, to treat **HIV infection**
- stavudine or zalcitabine, to treat **HIV infection**
- ribavirin or injections of ganciclovir to treat **viral infections**
- high doses of **co-trimoxazole**, an antibiotic.
- cladribine, used to treat **hairy cell leukaemia**

**Tell your doctor** if you are being treated with any of these.

### **Some medicines can make it more likely that you'll have side effects, or make side effects worse.**

These include:

- sodium valproate, to treat **epilepsy**
- interferon, to treat **viral infections**
- pyrimethamine, to treat **malaria** and other parasitic infections
- dapsone, to prevent **pneumonia** and treat skin infections
- fluconazole or flucytosine, to treat **fungal infections** such as **candida**
- pentamidine or atovaquone to treat parasitic infections such as *Pneumocystis jirovecii* pneumonia (often referred to as **PCP**)
- amphotericin or co-trimoxazole, to treat **fungal and bacterial infections**
- probenecid, to treat **gout** and similar conditions, and given with some antibiotics to make them more effective
- **methadone**, used as a **heroin substitute**
- vincristine, vinblastine or doxorubicin, to treat **cancer**.

**Tell your doctor** if you are taking any of these.

### **Some medicines interact with Lamivudine/Zidovudine Teva**

These include:

- **clarithromycin**, an antibiotic  
if you are taking clarithromycin, take your dose at least 2 hours before or after you take your Lamivudine/Zidovudine Teva.
- **phenytoin**, for treating **epilepsy**.  
**Tell your doctor** if you are taking phenytoin. Your doctor may need to monitor you while you are taking Lamivudine/Zidovudine Teva.
- medicines (usually liquids) containing **sorbitol and other sugar alcohols** (such as xylitol, mannitol, lactitol or maltitol), if taken regularly.  
**Tell your doctor** or pharmacist if you are being treated with any of these.

### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

Lamivudine/Zidovudine Teva and similar medicines may cause side effects in unborn babies. If you have taken Lamivudine/Zidovudine Teva during your pregnancy, your doctor may request regular blood tests and other diagnostic tests to monitor the development of your child. In children whose mothers took NRTIs during pregnancy, the benefit from the protection against HIV outweighed the risk of side effects.

**Women who are HIV-positive must not breastfeed**, because HIV infection can be passed on to the baby in breast milk.

A small amount of the ingredients in Lamivudine/Zidovudine Teva can also pass into your breast-milk.

If you're breast-feeding, or thinking about breast-feeding:

**Talk to your doctor immediately.**

### **Driving and using machines**

**Lamivudine/Zidovudine Teva can make you dizzy** and have other side effects that make you less alert.

**Do not drive or operate machines** unless you are feeling well.

### **Lamivudine/Zidovudine Teva contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

## **3. How to take Lamivudine/Zidovudine Teva**

**Always take this medicine exactly as your doctor or pharmacist has told you.** Check with your doctor or pharmacist if you are not sure.

Swallow Lamivudine/Zidovudine Teva tablets with some water. Lamivudine/Zidovudine Teva can be taken with or without food.

If you cannot swallow the tablet whole, you may crush and combine them with a small amount of food or drink and take all the dose immediately.

**Stay in regular contact with your doctor**

Lamivudine/Zidovudine Teva helps to control your condition. You need to keep taking it every day to stop your illness getting worse. You may still develop other infections and illnesses linked to HIV infection.

**Keep in touch with your doctor, and do not stop taking Lamivudine/Zidovudine Teva** without your doctor's advice.

#### **The recommended dose is**

##### **Adults and adolescents 30 kg or more**

**The usual dose of Lamivudine/Zidovudine Teva is one tablet twice a day.**

Take the tablets at regular times, leaving approximately 12 hours between each tablet.

##### **Children who weigh between 21 and 30 kg**

The usual starting dose of Lamivudine/Zidovudine Teva is one half tablet ( $\frac{1}{2}$ ) taken in the morning and one whole tablet taken in the evening.

##### **Children who weigh between 14 and 21 kg**

The usual starting dose of Lamivudine/Zidovudine Teva is one half tablet ( $\frac{1}{2}$ ) taken in the morning and one half tablet ( $\frac{1}{2}$ ) taken in the evening.

For children who weigh less than 14 kg lamivudine and zidovudine (the ingredients of Lamivudine/Zidovudine Teva) should be taken separately.

##### **If you take more Lamivudine/Zidovudine Teva than you should**

If you accidentally take too much Lamivudine/Zidovudine Teva, tell your doctor or your pharmacist, or contact your nearest hospital emergency department for further advice.

##### **If you forget to take Lamivudine/Zidovudine Teva**

If you forget to take a dose, take it as soon as you remember. Then continue your treatment as before. Do not take a double dose to make up for a forgotten dose.

#### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

During HIV therapy there may be an increase in weight and in levels of blood lipids and glucose. This is partly linked to restored health and life style, and in the case of blood lipids sometimes to the HIV medicines themselves. Your doctor will test for these changes.

Treatment with Lamivudine/Zidovudine Teva often causes a loss of fat from legs, arms and face (lipoatrophy). This loss of body fat has been shown to be not fully reversible after discontinuation of zidovudine. Your doctor should monitor for signs of lipoatrophy. Tell your doctor if you notice any loss of fat from your legs, arms, and face. When these signs occur, Lamivudine/Zidovudine Teva should be stopped and your HIV treatment changed.

When you are being treated for HIV, it can be hard to tell whether a symptom is a side effect of Lamivudine/Zidovudine Teva or other medicines you are taking, or an effect of the HIV disease itself. **So it is very important to talk to your doctor about any changes in your health.**

**As well as the side effects listed below for Lamivudine/Zidovudine Teva,** other conditions can develop during combination therapy for HIV.

It is important to read the information later in this section under 'Other possible side effects of combination therapy for HIV'.

### Very common side effects

These may affect **more than 1 in 10** people:

- headache
- feeling sick (nausea).

### Common side effects

These may affect **up to 1 in 10** people:

- being sick (vomiting)
- diarrhoea
- stomach pains
- loss of appetite
- feeling dizzy
- tiredness, lack of energy
- fever (high temperature)
- general feeling of being unwell
- difficulty in sleeping (insomnia)
- muscle pain and discomfort
- joint pain
- cough
- irritated or runny nose
- skin rash
- hair loss (*alopecia*).

Common side effects that might show up in blood tests are:

- a low red blood cell count (*anaemia*) or low white blood cell count (*neutropenia* or *leukopenia*)
- an increase in the level of liver enzymes
- an increased amount in the blood of *bilirubin* (a substance produced in the liver) which may make your skin appear yellow.

### Uncommon side effects

These may affect **up to 1 in 100** people:

- feeling breathless
- wind (*flatulence*)
- itching
- muscle weakness.

An uncommon side effect that may show up in blood test is:

- a decrease in the number of cells involved in blood clotting (*thrombocytopenia*) or in all kinds of blood cells (*pancytopenia*).

### Rare side effects

These may affect **up to 1 in 1000** people:

- serious allergic reaction causing swelling of the face, tongue or throat which may cause difficulty in swallowing or breathing
- liver disorders, such as jaundice, enlarged liver or fatty liver, inflammation (*hepatitis*)
- lactic acidosis (excess lactic acid in the blood; *see the next section, 'Other possible side effects of combination therapy for HIV'*)
- inflammation of the pancreas (*pancreatitis*)
- chest pain; disease of the heart muscle (*cardiomyopathy*)
- fits (*convulsions*)

- feeling depressed or anxious, not being able to concentrate, feeling drowsy
- indigestion, taste disturbance
- changes in the colour of your nails, your skin or the skin inside your mouth
- a flu-like feeling – chills and sweating
- tingly feelings in the skin (pins and needles)
- sensation of weakness in the limbs
- breakdown of muscle tissue
- numbness
- passing urine more often
- enlarged breasts in men.

Rare side effects that may show up in blood tests are:

- an increase in an enzyme called amylase
- a failure of the bone marrow to produce new red blood cells (*pure red cell aplasia*).

### **Very rare side effects**

These may affect **up to 1 in 10,000** people

A very rare side effect that may show up in blood tests is:

- a failure of the bone marrow to produce new red or white blood cells (*aplastic anaemia*).

### **If you get side effects**

**Tell your doctor or pharmacist** if any of the side effects get severe or troublesome, or if you notice any side effects not listed in this leaflet.

### **Other possible side effects of combination therapy for HIV**

Combination therapy such as Lamivudine/Zidovudine Teva may cause other conditions to develop during HIV treatment.

### **Old infections may flare up**

People with advanced HIV infection (AIDS) have weak immune systems, and are more likely to develop serious infections (opportunistic infections). When these people start treatment, they may find that old, hidden infections flare up, causing signs and symptoms of inflammation. These symptoms are probably caused by the body's immune system becoming stronger, so that the body starts to fight these infections.

In addition to the opportunistic infections, autoimmune disorders (a condition that occurs when the immune system attacks healthy body tissue) may also occur after you start taking medicines for the treatment of your HIV infection. Autoimmune disorders may occur many months after the start of treatment. If you notice any symptoms of infection or other symptoms such as muscle weakness, weakness beginning in the hands and feet and moving up towards the trunk of the body, palpitations, tremor or hyperactivity, please inform your doctor immediately to seek necessary treatment.

If you get any symptoms of infection while you are taking Lamivudine/Zidovudine Teva:

**Tell your doctor immediately.** Do not take other medicines for the infection without your doctor's advice.

### **Lactic acidosis is a rare but serious side effect**

Some people taking Lamivudine/Zidovudine Teva develop a condition called lactic acidosis, together with an enlarged liver.

Lactic acidosis is caused by a build-up of lactic acid in the body. It is rare; if it happens, it usually develops after a few months of treatment. It can be life-threatening, causing failure of internal organs.

Lactic acidosis is more likely to develop in people who have liver disease, or in obese (very overweight) people, especially women.

**Signs of lactic acidosis include:**

- deep, rapid, difficult breathing
- drowsiness
- numbness or weakness in the limbs
- feeling sick (nausea), being sick (vomiting)
- stomach pain.

During your treatment, your doctor will monitor you for signs of lactic acidosis. If you have any of the symptoms listed above, or any other symptoms that worry you:

**See your doctor as soon as possible.**

**You may have problems with your bones**

Some people taking combination therapy for HIV develop a condition called osteonecrosis. With this condition, parts of the bone tissue die because of reduced blood supply to the bone. People may be more likely to get this condition:

- if they have been taking combination therapy for a long time
- if they are also taking anti-inflammatory medicines called corticosteroids
- if they drink alcohol
- if their immune systems are very weak
- if they are overweight.

**Signs of osteonecrosis include:**

- stiffness in the joints
- aches and pains (especially in the hip, knee or shoulder)
- difficulty moving.

If you notice any of these symptoms:

**Tell your doctor.**

**Other effects may show up in blood tests**

Combination therapy for HIV can also cause:

- increased levels of lactic acid in the blood, which on rare occasions can lead to lactic acidosis.

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Lamivudine/Zidovudine Teva**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the container or carton and blister after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Lamivudine/Zidovudine Teva contains**

- The active substances are lamivudine 150 mg and zidovudine 300 mg.
- The other ingredients are: Tablet core: Cellulose microcrystalline, sodium starch glycolate (Type A), sodium stearyl fumarate. Tablet film-coat: Hypromellose 3cP, hypromellose 6cP, polysorbate 80, macrogol 400, titanium dioxide E171.

### **What Lamivudine/Zidovudine Teva looks like and contents of the pack**

White, capsule shaped, biconvex, film-coated scored tablet – engraved with “L/Z” on one side and “150/300” on the other side.

The tablet is a breakable tablet.

Lamivudine/Zidovudine Teva is available in aluminium blisters or HDPE containers containing 60 tablets.

### **Marketing Authorisation Holder and Manufacturer**

#### **Marketing Authorisation Holder**

Teva B.V.  
Swensweg 5  
2031GA Haarlem  
The Netherlands

#### **Manufacturer**

Teva Pharmaceutical Works Private Limited Company  
Pallagi út 13  
Debrecen H-4042  
Hungary

Pharmachemie B.V.  
Swensweg 5  
2031 GA Haarlem  
The Netherlands

Teva Operations Poland Sp. z.o.o.  
80 Mogilska St.  
31-546 Krakow  
Poland

Merckle GmbH  
Ludwig-Merckle-Strasse 3  
89143 Blaubeuren  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:



**België/Belgique/Belgien**

Teva Pharma Belgium N.V./S.A./AG  
Tél/Tel: +32 38207373

**България**

Тева Фарма ЕАД  
Тел: +359 24899585

**Česká republika**

Teva Pharmaceuticals CR, s.r.o.  
Tel: +420 251007111

**Danmark**

Teva Denmark A/S  
Tlf: +45 44985511

**Deutschland**

ratiopharm GmbH  
Tel: +49 73140202

**Eesti**

UAB Teva Baltics Eesti filiaal  
Tel: +372 6610801

**Ελλάδα**

Specifar A.B.E.E.  
Τηλ: +30 2118805000

**España**

Teva Pharma, S.L.U.  
Tel: +34 913873280

**France**

Teva Santé  
Tél: +33 155917800

**Hrvatska**

Pliva Hrvatska d.o.o.  
Tel: +385 13720000

**Ireland**

Teva Pharmaceuticals Ireland  
Tel: +44 2075407117

**Ísland**

Teva Pharma Iceland ehf.  
Sími: +354 5503300

**Italia**

Teva Italia S.r.l.  
Tel: +39 028917981

**Lietuva**

UAB Teva Baltics  
Tel: +370 52660203

**Luxembourg/Luxemburg**

ratiopharm GmbH  
Allemagne/Deutschland  
Tél/Tel: +49 73140202

**Magyarország**

Teva Gyógyszergyár Zrt.  
Tel: +36 12886400

**Malta**

Teva Pharmaceuticals Ireland  
L-Irlanda  
Tel: +44 2075407117

**Nederland**

Teva Nederland B.V.  
Tel: +31 8000228400

**Norge**

Teva Norway AS  
Tlf: +47 66775590

**Österreich**

ratiopharm Arzneimittel Vertriebs-GmbH  
Tel: +43 1970070

**Polska**

Teva Pharmaceuticals Polska Sp. z o.o.  
Tel: +48 223459300

**Portugal**

Teva Pharma - Produtos Farmacêuticos,  
Lda.  
Tel: +351 214767550

**România**

Teva Pharmaceuticals S.R.L.  
Tel: +40 212306524

**Slovenija**

Pliva Ljubljana d.o.o.  
Tel: +386 15890390

**Slovenská republika**

TEVA Pharmaceuticals Slovakia s.r.o.  
Tel: +421 257267911

**Suomi/Finland**

Teva Finland Oy  
Puh/Tel: +358 201805900

**Κύπρος**

Specifar A.B.E.E.

Ελλάδα

Τηλ: +30 2118805000

**Sverige**

Teva Sweden AB

Tel: +46 42121100

**Latvija**

UAB Teva Baltics filiāle Latvijā

Tel: +371 67323666

**United Kingdom (Northern Ireland)**

Teva Pharmaceuticals Ireland

Ireland

Tel: +44 2075407117

**This leaflet was last revised in {MM/YYYY}.**

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu/>.

Medicinal product no longer authorised