

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## **1. NAME OF THE MEDICINAL PRODUCT**

Qarziba 4.5 mg/mL concentrate for solution for infusion

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

1 mL of concentrate contains 4.5 mg dinutuximab beta.  
Each vial contains 20 mg dinutuximab beta in 4.5 mL.

Dinutuximab beta is a mouse-human chimeric monoclonal IgG1 antibody produced in a mammalian cell line (CHO) by recombinant DNA technology.

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Concentrate for solution for infusion  
Colourless to slightly yellow liquid

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Qarziba is indicated for the treatment of high-risk neuroblastoma in patients aged 12 months and above, who have previously received induction chemotherapy and achieved at least a partial response, followed by myeloablative therapy and stem cell transplantation, as well as patients with history of relapsed or refractory neuroblastoma, with or without residual disease. Prior to the treatment of relapsed neuroblastoma, any actively progressing disease should be stabilised by other suitable measures.

In patients with a history of relapsed/refractory disease and in patients who have not achieved a complete response after first line therapy, Qarziba should be combined with interleukin-2 (IL-2).

### **4.2 Posology and method of administration**

Qarziba is restricted to hospital-use only and must be administered under the supervision of a physician experienced in the use of oncological therapies. It must be administered by a healthcare professional prepared to manage severe allergic reactions including anaphylaxis in an environment where full resuscitation services are immediately available.

#### Posology

Treatment with Qarziba consists of 5 consecutive courses, each course comprising 35 days. The individual dose is determined based on the body surface area and should be a total of 100 mg/m<sup>2</sup> per course.

Two modes of administration are possible:

- a continuous infusion over the first 10 days of each course (a total of 240 hours) at the daily dose of 10 mg/m<sup>2</sup>
- or five daily infusions of 20 mg/m<sup>2</sup> administered over 8 hours, on the first 5 days of each course

When IL-2 is combined with Qarziba, it should be administered as subcutaneous injections of 6×10<sup>6</sup> IU/m<sup>2</sup>/day, for 2 periods of 5 consecutive days, resulting in an overall dose of 60×10<sup>6</sup> IU/m<sup>2</sup> per course. The first 5-day course should start 7 days prior to the first infusion of dinutuximab beta and the second 5-day course should start concurrently with dinutuximab beta infusion (days 1 to 5 of each dinutuximab beta course).

Prior to starting each treatment course, the following clinical parameters should be evaluated and treatment should be delayed until these values are reached:

- pulse oximetry > 94% on room air
- adequate bone marrow function: absolute neutrophil count ≥ 500/μL, platelet count ≥ 20,000/μL, haemoglobin > 8.0 g/dL
- adequate liver function: alanine aminotransferase (ALT)/ aspartate aminotransferase (AST) < 5 times upper limit of normal (ULN)
- adequate renal function: creatinine clearance or glomerular filtration rate (GRF) > 60 mL/min/1.73 m<sup>2</sup>

#### *Dose modification of dinutuximab beta*

Based on the physician's evaluation of the severity of adverse drug reactions to dinutuximab beta, patients may undergo a dose reduction of 50% or a temporary interruption of the infusion. As a consequence, either the infusion period is prolonged or, if tolerated by the patient, the infusion rate may be increased up to 3 mL/h (continuous infusion), in order to administer the total dose.

#### **Recommended dose modifications for dinutuximab beta**

<b>Adverse reaction</b>	<b>Severity</b>	<b>Treatment modification</b>
<b>Any</b>	<b>Grade 1 – 2</b>	Decrease infusion rate to 50%, After resolution, resume infusion at original rate
Hypersensitivity reaction	e.g. hypotension	Interrupt infusion and administer supportive measures, After resolution, resume infusion at original rate
Dilated pupils with sluggish light reflex +/- photophobia		Interrupt infusion, After resolution, resume infusion at 50% rate
<b>Any</b>	<b>Grade ≥ 3</b>	Interrupt infusion and administer supportive measures, Resume infusion at 50% rate if ADR resolves or improves to Grade 1 – 2, After resolution, increase to original rate
	Recurrent	Discontinue infusion, Resume next day if ADR resolves
Hypersensitivity reaction	e.g. bronchospasm, angioedema	Interrupt infusion immediately and treat appropriately (see section 4.4), Resume treatment for subsequent courses
Capillary leak syndrome		Interrupt infusion and administer supportive measures, Resume at 50% rate if ADR resolves or improves to Grade 1 – 2

Adverse reaction	Severity	Treatment modification
Central neurotoxicity		Interrupt infusion immediately, rule out other influencing factors and treat appropriately. There is limited data available on resuming treatment and no recommendations can be made

Treatment with dinutuximab beta should be permanently discontinued if the following toxicities occur:

- grade 3 or 4 anaphylaxis
- prolonged grade 2 peripheral motor neuropathy
- grade 3 peripheral neuropathy
- grade 3 vision eye toxicity
- grade 4 hyponatremia (< 120 mEq/L) despite appropriate fluid management
- recurrent or grade 4 capillary leak syndrome (requires ventilator support)
- severe central neurotoxicity that includes grade 3 or 4 with substantial prolonged neurological deficit without any detectable reason, recurrent grade 1-3 neurotoxicity, and permanent neurological deficit
- all grades of posterior reversible encephalopathy syndrome and transverse myelitis

#### *Renal and hepatic impairment*

There are no data in patients with renal and hepatic impairment (see section 5.2).

#### *Paediatric population*

The safety and efficacy of Qarziba in children aged less than 12 months have not yet been established. No data are available.

#### Method of administration

Qarziba is for intravenous infusion. The solution should be administered via a peripheral or central intravenous line. Other intravenously co-administered agents should be delivered via a separate infusion line (see section 6.6).

For continuous infusions, the solution is administered at a rate of 2 mL per hour (48 mL per day) using an infusion pump.

For 8-hour daily infusions, the solution is administered at a rate of approximately 13 mL per hour.

Pre-medication should always be considered before starting each infusion (see section 4.4).

For instructions on dilution of the medicinal product before administration, see section 6.6.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Acute grade 3 or 4, or extensive chronic graft-versus-host disease (GvHD)

### **4.4 Special warnings and precautions for use**

#### Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

### Pain

Neuropathic pain usually occurs at the beginning of the treatment and premedication with analgesics, including intravenous opioids, prior to each infusion of dinutuximab beta is required. A triple therapy, including nonopioid analgesics (according to WHO guidelines), gabapentin and opioids, is recommended for pain treatment. The individual dose may vary widely.

#### *Nonopioid analgesics*

Nonopioid analgesics should be used permanently during the treatment, e.g. paracetamol or ibuprofen.

#### *Gabapentin*

The patient should be primed with 10 mg/kg/day, starting 3 days prior to dinutuximab beta infusion. The daily dose of gabapentin is increased to  $2 \times 10$  mg/kg/day orally, the next day and to  $3 \times 10$  mg/kg/day orally, the day before the onset of dinutuximab beta infusion and thereafter. The maximum single dose of gabapentin is 300 mg. This dosing schedule should be maintained for as long as required by the patient.

Oral gabapentin should be tapered off after weaning off intravenous morphine infusion, at the latest after dinutuximab beta infusion therapy has stopped.

#### *Opioids*

Treatment with opioids is standard with dinutuximab beta. The first infusion day and course usually requires a higher dose than subsequent days and courses.

- Before initiation of a continuous intravenous morphine infusion, a bolus infusion of 0.02 to 0.05 mg/kg/hour morphine should be started 2 hours before dinutuximab beta infusion.
- Subsequently, a dosing rate of 0.03 mg/kg/hour is recommended concomitantly with dinutuximab beta infusion.
- With daily infusions of dinutuximab beta, morphine infusion should be continued at a decreased rate (e.g. 0.01 mg/kg/h) for 4 hours after the end of dinutuximab beta infusion.
- With continuous infusion, in response to the patient's pain perception, it may be possible to wean off morphine over 5 days by progressively decreasing its dosing rate (e.g. to 0.02 mg/kg/hour, 0.01 mg/kg/hour, 0.005 mg/kg/hour).
- If continuous morphine infusion is required for more than 5 days, treatment should be gradually reduced by 20% per day after the last day of dinutuximab beta infusion.

After weaning off intravenous morphine, in case of severe neuropathic pain, oral morphine sulphate (0.2 to 0.4 mg/kg every 4 to 6 hours) can be administered on demand. For moderate neuropathic pain, oral tramadol may be administered.

### Hypersensitivity reactions

Severe infusion-related reactions, including cytokine release syndrome (CRS), anaphylactic and hypersensitivity reactions, may occur despite the use of premedication. Occurrence of a severe infusion related reaction (including CRS) requires immediate discontinuation of dinutuximab beta therapy and may necessitate emergency treatment.

Cytokine release syndrome frequently manifests itself within minutes to hours of initiating the first infusion and is characterised by systemic symptoms such as fever, hypotension and urticaria.

Anaphylactic reactions may occur as early as within a few minutes of the first infusion with dinutuximab beta and are commonly associated with bronchospasm and urticaria.

#### *Premedication*

Antihistamine premedication (e.g. diphenhydramine) should be administered by intravenous injection approximately 20 minutes before starting each dinutuximab beta infusion. It is recommended that antihistamine administration be repeated every 4 to 6 hours as required during dinutuximab infusion.

Patients should be closely monitored for anaphylaxis and allergic reactions, particularly during the first and second treatment course.

#### *Treatment of hypersensitivity reactions*

Intravenous antihistamine, epinephrine (adrenaline) and prednisolone for intravenous administration should be immediately available at the bedside during administration of dinutuximab beta to manage life-threatening allergic reactions. It is recommended that treatment for such reactions include prednisolone administered by intravenous bolus, and epinephrine administered by intravenous bolus every 3 to 5 minutes as necessary, according to clinical response. In case of bronchial and/or pulmonary hypersensitivity reaction, inhalation with epinephrine (adrenaline) is recommended and should be repeated every 2 hours, according to clinical response.

#### Capillary leak syndrome (CLS)

CLS is characterised by a loss of vascular tone and extravasation of plasma proteins and fluid into the extravascular space. CLS usually develops within hours after initiation of treatment, while clinical symptoms (i.e. hypotension, tachycardia) are reported to occur after 2 to 12 hours. Careful monitoring of circulatory and respiratory function is required.

#### Neurological disorders of the eye

Eye disorders may occur as dinutuximab beta binds to optic nerve cells. No dose modification is necessary in the case of an impaired visual accommodation that is correctable with eye glasses, as long as this is judged to be tolerable.

Treatment must be interrupted in patients who experience Grade 3 vision toxicity (i.e. subtotal vision loss per toxicity scale). In case of any eye problems, patients should be referred promptly to an ophthalmology specialist.

#### Peripheral neuropathy

Occasional occurrences of peripheral neuropathy have been reported with Qarziba. Cases of motor or sensory neuropathy lasting more than 4 days must be evaluated and non-inflammatory causes, such as disease progression, infections, metabolic syndromes and concomitant medication, should be excluded.

Treatment should be permanently discontinued in patients experiencing any objective prolonged weakness attributable to dinutuximab beta administration. For patients with moderate (Grade 2) neuropathy (motor with or without sensory), treatment should be interrupted and may be resumed after neurologic symptoms resolve.

#### Central neurotoxicity

Central neurotoxicity has been reported following treatment with Qarziba. If central neurotoxicity occurs the infusion should be interrupted immediately and the patient treated symptomatically, other influencing factors such as active infection, metastatic spread of neuroblastoma to CNS, neurotoxic concomitant medications should be ruled out.

Treatment with dinutuximab beta should be permanently discontinued following the occurrence of severe neurotoxicity that includes grade 3 or 4 central neurotoxicity with substantial prolonged neurological deficit without any detectable reason, recurrent grade 1-3 neurotoxicity and/or permanent neurological deficit and all grades of posterior reversible encephalopathy syndrome and transverse myelitis.

#### Systemic infections

Patients are likely to be immunocompromised as a result of prior therapies. As they typically have a central venous catheter in situ, they are at risk of developing systemic infection. Patients should have no evidence of systemic infection and any identified infection should be under control before starting therapy.

#### Haematologic toxicities

Occurrence of haematologic toxicities has been reported with Qarziba, such as erythropenia, thrombocytopenia or neutropenia. Grade 4 haematologic toxicities, improving to at least Grade 2 or baseline values by start of next treatment course, do not require dose modification.

#### Laboratory abnormalities

Regular monitoring of liver function and electrolytes is recommended.

#### Atypical haemolytic uraemic syndrome

Atypical haemolytic uraemic syndrome (aHUS) has been reported in patients who received dinutuximab beta, in some cases with fatal outcome. Signs and symptoms of aHUS should be monitored for. If aHUS is diagnosed, prompt treatment is required and dinutuximab beta should be permanently discontinued.

### **4.5 Interaction with other medicinal products and other forms of interaction**

No interaction studies have been performed. A risk for indirect reduction of CYP activity due to higher TNF- $\alpha$  and IL-6 levels and, therefore, interactions with concomitantly used medicinal products, cannot be excluded.

#### *Corticosteroids*

Due to their immunosuppressive activity, concomitant treatment with corticosteroids is not recommended within 2 weeks prior to the first treatment course until 1 week after the last treatment course with dinutuximab beta, except for life-threatening conditions.

#### *Vaccinations*

Vaccinations should be avoided during administration of dinutuximab beta until 10 weeks after the last treatment course, due to immune stimulation through dinutuximab beta and possible risk for rare neurological toxicities.

#### *Intravenous immunoglobulin*

Concomitant use of intravenous immunoglobulins is not recommended as they may interfere with dinutuximab beta-dependent cellular cytotoxicity.

### **4.6 Fertility, pregnancy and lactation**

#### Pregnancy

There are no data on pregnant women. No animal data are available on teratogenicity or embryotoxicity. Dinutuximab beta target (GD2) is expressed on neuronal tissues, especially during embryofetal development, and may cross the placenta; therefore, Qarziba may cause fetal harm when administered to pregnant women.

Qarziba should not be used during pregnancy.

#### Breast-feeding

There are no data on lactating women. It is unknown whether dinutuximab beta is excreted in human milk. Breast-feeding should be discontinued during treatment with Qarziba and for 6 months after the last dose.

#### Fertility

The effects of dinutuximab beta on fertility in humans are unknown. In animals, dedicated fertility studies have not been conducted, but no adverse effects on reproductive organs were observed in toxicity studies performed in Guinea pig and cynomolgous monkey.

Qarziba should not be used in women of childbearing potential not using contraception. It is recommended that women of childbearing potential use contraception for 6 months after discontinuation of treatment with dinutuximab beta.

#### 4.7 Effects on ability to drive and use machines

Dinutuximab beta has major influence on the ability to drive and use machines. Patients should not use or drive machines during treatment with dinutuximab beta.

#### 4.8 Undesirable effects

##### Summary of the safety profile

The safety of dinutuximab beta has been evaluated in 791 patients with high-risk and relapsed/refractory neuroblastoma, who received it as a continuous infusion (212) or as repeated daily infusions (416). It was combined with 13-cis retinoic in most patients and with IL-2 in 307 patients.

The most common adverse reactions were pyrexia (86%) and pain (57%) that occurred despite analgesic treatment. Other frequent adverse reactions were hypersensitivity (74.1%), vomiting (55%), diarrhoea (52%), capillary leak syndrome (36%), anaemia (49%), neutropenia (46%), thrombocytopenia (42%) and hypotension (41%).

##### Tabulated list of adverse reactions

Adverse reactions reported in clinical trials and post-marketing are listed by system organ class and by frequency and summarised in the table below. These adverse reactions are presented by MedDRA system organ class and frequency. Frequency categories are defined as: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ) and not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

System organ class	Very common	Common	Uncommon	Not known
Infections and infestations	infection (including pneumonia, skin infection, herpes virus infection, myelitis, encephalomyelitis), device related infection	sepsis		
Blood and lymphatic system disorders	anaemia leukopenia, neutropenia, thrombocytopenia	lymphopenia	disseminated intravascular coagulation, eosinophilia	Atypical haemolytic uraemic syndrome
Immune system disorders	hypersensitivity, cytokine release syndrome	anaphylactic reaction	serum sickness	
Metabolism and nutrition disorders	fluid retention	decreased appetite, hypoalbuminaemia, hyponatraemia, hypokalaemia, hypophosphataemia, hypomagnesaemia, hypocalcaemia, dehydration		
Psychiatric disorders		agitation, anxiety		
Nervous system disorders	headache	peripheral neuropathy, seizure, paraesthesia, dizziness, tremor	intracranial pressure increased,	



<b>System organ class</b>	<b>Very common</b>	<b>Common</b>	<b>Uncommon</b>	<b>Not known</b>
			posterior reversible encephalopathy syndrome	
Eye disorders	mydriasis, pupillonia, eye oedema (eyelid, periorbital)	ophthalmoplegia, papilloedema, accommodation disorder, blurred vision, photophobia		
Cardiac disorders	tachycardia	cardiac failure, left ventricular dysfunction, pericardial effusion		
Vascular disorders	hypotension, capillary leak syndrome	hypertension	hypovolaemic shock, veno-occlusive disease	
Respiratory, thoracic and mediastinal disorders	hypoxia, cough	bronchospasm, dyspnoea, respiratory failure, lung infiltration, pulmonary oedema, pleural effusion, tachypnoea, laryngospasm		
Gastrointestinal disorders	vomiting, diarrhoea, constipation, stomatitis	nausea, lip oedema, ascites, abdominal distension, ileus, dry lips	enterocolitis	
Hepatobiliary disorders			hepatocellular injury	
Skin and subcutaneous tissue disorders	pruritus, rash, urticaria	dermatitis (including exfoliative) , erythema, dry skin, hyperhidrosis, petechiae, photosensitivity reaction		
Musculoskeletal and connective tissue disorders		muscle spasms		
Renal and urinary disorders		oliguria, urinary retention, hyperphosphaturia, haematuria, proteinuria	renal failure	

System organ class	Very common	Common	Uncommon	Not known
General disorders and administration site conditions	pyrexia, chills, pain*, peripheral oedema, face oedema	injection site reaction		
Investigations	increased weight, increased transaminases, increased gamma glutamyltransferase, increased blood bilirubin increased blood creatinine	decreased weight, decreased glomerular filtration rate, hypertriglyceridaemia, prolonged activated partial thromboplastin time, prolonged prothrombin time, prolonged thrombin time		

\*includes abdominal pain, pain in extremity, oropharyngeal pain, and Back pain reported in >10% of patients. In addition, other common pain types reported were arthralgia, injection site pain, musculoskeletal pain, bone pain, chest pain, and neck pain.

### Description of selected adverse reactions

#### *Hypersensitivity*

The most frequent hypersensitivity reactions included hypotension (42.2%), urticaria (7%) and bronchospasm (1%). Cytokine release syndrome was also reported in 32% of the patients. Serious anaphylactic reactions occurred in 3.5% of the patients.

#### *Pain*

Pain typically occurs during the first infusion of dinutuximab beta and decreases over the treatment courses. Most commonly, patients reported abdominal pain, pain in the extremities, back pain, chest pain, or arthralgia.

#### *Capillary leak syndrome (CLS)*

Overall, 10% of CLS were severe (grade 3-4) and their frequency decreased over the treatment courses.

#### *Eye problems*

These included impaired visual accommodation that is correctable with eye glasses, as well as mydriasis (2%), periorbital oedema and eyelid oedema (3%), blurred vision (3%) or photophobia (3%), which were usually reversible after treatment discontinuation. Severe eye disorders were also reported including ophthalmoplegia (2%) and optic atrophy.

#### *Peripheral neuropathy*

Both motor and sensory peripheral neuropathies have been reported, overall in 9% of the patients. Most events were of grade 1-2 and resolved.

#### *Central Neurotoxicity*

Reports of central neurotoxicity and severe neurotoxicity have been received including posterior reversible encephalopathy syndrome (0.7%) and seizures (1.7%).

### *Safety profile with and without IL-2*

The combination of Qarziba with IL-2 increases the risk of adverse drug reactions compared to Qarziba without IL-2, especially for pyrexia (94% vs. 80%), CLS (45% vs. 20%), pain related to dinutuximab beta (70% vs. 62%), hypotension (44% vs. 27%), and peripheral neuropathy (9% vs. 5%), respectively.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

## **4.9 Overdose**

No cases of dinutuximab beta overdose have been reported.

In the case of overdose, patients should be carefully observed for signs or symptoms of adverse reactions and supportive care administered, as appropriate.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Antineoplastic agents, monoclonal antibodies, ATC code: L01FX06

#### Mechanism of action

Dinutuximab beta is a chimeric monoclonal IgG1 antibody that is specifically directed against the carbohydrate moiety of disialoganglioside 2 (GD2), which is overexpressed on neuroblastoma cells.

#### Pharmacodynamic effects

Dinutuximab beta has been shown *in vitro* to bind to neuroblastoma cell lines known to express GD2 and to induce both complement dependent cytotoxicity (CDC) and antibody dependent cell-mediated cytotoxicity (ADCC). In the presence of human effector cells, including peripheral blood nuclear cells and granulocytes from normal human donors, dinutuximab beta was found to mediate the lysis of human neuroblastoma and melanoma cell lines in a dose-dependent manner. Additionally, *in vivo* studies demonstrated that dinutuximab beta could suppress liver metastasis in a syngeneic liver metastasis mouse model.

Neurotoxicity associated to dinutuximab beta is likely due to the induction of mechanical allodynia that may be mediated by the reactivity of dinutuximab beta with the GD2 antigen located on the surface of peripheral nerve fibres and myelin.

#### Clinical efficacy

The efficacy of dinutuximab beta has been evaluated in a randomised controlled trial comparing the administration of dinutuximab beta with or without IL-2 in the first-line treatment of patients with high-risk neuroblastoma and in two single-arm studies in the relapsed/refractory setting.

#### *Relapsed and refractory patients*

In a compassionate use programme (study 1), 54 patients received 10 mg/m<sup>2</sup>/day dinutuximab beta given by continuous 10-day intravenous infusion in a 5-week treatment course, concurrently with subcutaneous IL-2 (6×10<sup>6</sup> IU/m<sup>2</sup>/day given on days 1-5 and 8-12 of each course) and followed by oral 13-cis-RA treatment (160 mg/m<sup>2</sup>/day for 14 days per course). The same treatment regimen was used in a Phase II study (study 2), which enrolled 44 patients.

Overall, these 98 patients had primary refractory neuroblastoma (40) or relapsed neuroblastoma (49) with an additional 9 patients enrolled after first-line therapy. These were 61 boys and 37 girls, aged 1 to 26 years (median 5 years). Most had an initial diagnosis of INSS stage 4 disease without MYCN

amplification (16% of the subjects had MYCN amplified tumours and in 14% this information was missing). Most patients with relapsed disease were enrolled after their first relapse and the median time from diagnosis to first relapse was about 14 months. Treatment of disease before immunotherapy included intensive chemotherapy regimen followed by autologous stem cell transplantation (ASCT), radiotherapy, and surgery. At baseline, 72 patients had measurable disease and 26 patients had no detectable disease.

Survival rates (event-free survival, overall survival) are presented by type of disease in Table 1. The overall response rate (complete response plus partial response) in patients with evidence of disease at baseline was 36% (95% confidence interval [25; 48]) and was more favourable in patients with refractory disease (41% [23; 57]) than in patients with relapsed disease (29% [15; 46]).

Table 1: Event-free survival (EFS) and overall survival (OS) rates in relapsed and refractory patients

		<b>Study 1 N=29</b>	<b>Study 2 N=19</b>	<b>Study 1 N=15</b>	<b>Study 2 N=25</b>
		<b>Relapsed patients</b>		<b>Refractory patients</b>	
EFS	1 year	45%	42%	58%	60%
	2 years	31%	37%	29%	56%
OS	1 year	90%	74%	93%	100%
	2 years	69%	42%	70%	78%

#### *First-line patients who received autologous stem cell transplantation*

In study 3, patients with high-risk neuroblastoma were enrolled after they had received induction chemotherapy and achieved at least a partial response, then myeloablative therapy and stem cell transplantation. Patients with progressive disease were excluded. Dinutuximab beta was administered at a dose of 20 mg/m<sup>2</sup>/day on 5 consecutive days, given by 8-hour intravenous infusion in a 5-week treatment course, and was combined with 13-cis-RA and with or without additional subcutaneous IL-2 at the same posologies as in the previous studies.

A total of 370 patients were randomised and received treatment. These included 64% male and 36% female patients with a median age of 3 years (0.6 to 20); 89% had a tumour INSS stage 4 and MYCN amplification was reported in 44% of the cases. The primary efficacy endpoint was 3-year EFS and secondary endpoint was OS. EFS and OS rates are presented in Tables 2 and 3 according to the evidence of disease at baseline.

For patients without evidence of disease at baseline, addition of IL-2 did not improve EFS and OS.

Table 2: Event-free survival (EFS) and overall survival (OS) rates [95% confidence interval] in patients without evidence of disease at baseline (complete response to initial treatment)

<b>Efficacy</b>	<b>without IL-2 N=104</b>			<b>with IL-2 N=107</b>		
	<b>1 year</b>	<b>2 year</b>	<b>3 year</b>	<b>1 year</b>	<b>2 year</b>	<b>3 year</b>
EFS	77% [67; 84]	67% [57; 75]	62% [51; 71]	73% [63; 80]	70% [60; 77]	66% [56; 75]
OS	89% [81; 94]	78% [68; 85]	71% [60; 80]	89% [81; 93]	78% [68; 85]	72% [61; 80]

Table 3: Event-free survival (EFS) and overall survival (OS) rates [95% confidence interval] in patients with evidence of disease at baseline (no complete response to initial treatment)

Efficacy	without IL-2 N=73			with IL-2 N=76		
	1 year	2 year	3 year	1 year	2 year	3 year
EFS	67% [55; 76]	58% [45; 69]	46% [33; 58]	72% [60; 81]	62% [49; 72]	54% [41; 65]
OS	83% [72; 90]	73% [61; 82]	54% [40; 66]	86% [75; 92]	71% [58; 80]	63% [50; 74]

#### Immunogenicity

In 3 clinical studies the appearance of ADA was 57.1% (112/196) in subjects being classed as ADA positive on the basis of having at least one measurable ADA response over the course of treatment. Neutralising antibody activity was observed in 63.5% (54/85) of the ADA-positive subjects in 2 studies. There was an overall trend of lower dinutuximab beta concentration with increasing ADA titre, (low, medium and high). In 16.8% of subjects (33/196) with a high ADA titre, the reduction in dinutuximab beta concentration impacted on pharmacodynamic responses. Based on the available data it is not possible to determine a quantitative association between ADA titre and impact on efficacy.

No clear associations between ADA response and relevant Selected Safety Events were observed.

From an efficacy and safety perspective, there is no rationale for adjusting or stopping treatment on the basis of measured ADA responses.

#### Paediatric population

The European Medicines Agency has deferred the obligation to submit the results of studies with Qarziba in one or more subsets of the paediatric population in neuroblastoma (see section 4.2 for information on paediatric use).

This medicinal product has been authorised under ‘exceptional circumstances’.

This means that for ethical reasons it has not been possible to obtain complete information on this medicinal product. The European Medicines Agency will review any new information which may become available every year and this SmPC will be updated as necessary.

## **5.2 Pharmacokinetic properties**

Dinutuximab beta has been investigated using short-term infusions (STI - five days of eight-hour infusions at 20 mg/m<sup>2</sup>/day) and long-term infusions (LTI - ten days of continuous infusion at 100 mg/m<sup>2</sup>).

#### Absorption

Dinutuximab beta is administered as an intravenous infusion. The maximum concentration (mean (± SD)) at the end of the long-term infusion was 11.2 (± 3.3) mg/L. Other routes of administration have not been investigated.

#### Distribution

The population mean (±SD) estimate for the central volume of distribution was 2.04 (± 1.05) L and for the peripheral volume of distribution 2.65 (±1.01) L.

#### Biotransformation

The metabolism of dinutuximab beta has not been investigated. As a protein, dinutuximab beta is expected to be metabolised to small peptides and individual amino acids by ubiquitous proteolytic enzymes.

### Elimination

The clearance after the LTI was 0.72 ( $\pm$  0.24) L/d/m<sup>2</sup>. The accumulation ratio for C<sub>max</sub> was 1.13 ( $\pm$  0.54) after 5 LTI courses (mean ( $\pm$ SD)). The apparent terminal elimination t<sub>1/2</sub> was 8.7 ( $\pm$  2.6) days (mean ( $\pm$  SD)). The clearance of dinutuximab beta increased in the presence of high anti-drug antibody titres regardless of neutralising activity. (See Immunogenicity in Section 5.1).

### Linearity/non-linearity

Variations in dose of the first infusion in Study 2 revealed a dose-proportional increase in exposure (AUC<sub>∞</sub>) up to the recommended dose of 100 mg/m<sup>2</sup> per course for 10 days.

### Specific populations

The age of patients ranged from 1 to 27 years (median 6 years). Body weight ranged from 9 to 75 kg (median 18.5 kg) and body-surface area ranged from 0.44 to 1.94 m<sup>2</sup> (median 0.75 m<sup>2</sup>). A two-compartment population-PK model with first-order elimination from the central compartment was developed using the data from 224 patients in four studies (STI 30 patients, LTI 194 patients). Volume and clearance parameters increased across the ranges with increasing body size. Body weight and ADA titre were covariates for clearance while body weight, age and IL-2 co-administration were covariates for volume of distribution.

### Age

The population pharmacokinetic analyses showed comparable exposure to dinutuximab beta in patients of all ages studied when dosed at 100 mg/m<sup>2</sup>.

### Gender

The population pharmacokinetic analysis with 89 female (40%) and 135 male (60%) patients showed no clinically meaningful effect of gender on dinutuximab beta pharmacokinetics.

### Race

Since the PK analysis population was predominantly Caucasian (92.9%) race was not formally examined as a potential PK covariate.

### Weight

Dosing on the basis of body surface area provides consistent exposure across populations.

### Renal impairment

No formal studies have been conducted in patients with renal impairment. Renal function was not a significant covariate in population pharmacokinetic analyses that included patients with normal renal function and mild renal impairment.

### Hepatic impairment

No formal studies have been conducted in patients with hepatic impairment. Subjects with ALT >3x ULN had comparable pharmacokinetics as subjects with ALT ≤3x ULN.

## **5.3 Preclinical safety data**

### General toxicology

Dinutuximab beta has been administered to male and female juvenile Guinea pigs, as well as male and female young cynomolgus monkeys, as repeat-dose regimens that exceeded the recommended clinical dose. Findings of note included changes (decrease) in thymus weight as well as bone marrow changes (atrophy affecting myeloid and erythroid precursor cell lines). The bone marrow changes were slight to severe and recovered after cessation of dosing. No effects on cardiovascular functions (ECG, blood pressure) were observed in monkeys.

#### Other

No non-clinical studies to evaluate the potential of dinutuximab beta to cause carcinogenicity, genotoxicity or developmental and reproductive toxicity have been conducted. In the repeat-dose toxicity studies in Guinea pigs and cynomolgus monkeys, no adverse effects of dinutuximab beta were observed on reproductive organs at exposure levels above clinical levels.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Histidine  
Sucrose  
Polysorbate 20  
Water for injections  
Hydrochloric acid (for pH adjustment)

### **6.2 Incompatibilities**

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

### **6.3 Shelf life**

#### Unopened vial

4 years

#### Diluted solution (solution for infusion)

Chemical and physical in-use stability has been demonstrated for up to 48 hours at 25 °C (50 mL syringe) and for up to 7 days at 37 °C (250 mL infusion bag), after cumulative storage in a refrigerator (2 °C – 8 °C) for 72 hours (see section 6.6).

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would not normally be longer than 24 hours at 2 to 8 °C, unless dilution has taken place in controlled and validated aseptic conditions.

### **6.4 Special precautions for storage**

Store in a refrigerator (2 °C – 8 °C).

Keep the vial in the outer carton in order to protect from light.

For storage conditions after dilution of the medicinal product, see section 6.3.

### **6.5 Nature and contents of container**

Clear Type I glass vial (6 mL) with a halobutyl rubber stopper and aluminium flip-off cap, containing a minimum extractable volume of 4.5 mL concentrate for solution for infusion.

Each carton contains 1 vial.

### **6.6 Special precautions for disposal and other handling**

The solution for infusion must be prepared under aseptic conditions. The solution must not be exposed to direct sunlight or heat.

The patient specific daily dose of Qarziba is calculated based on body surface area (see section 4.2). Qarziba should be diluted aseptically to the patient specific concentration/dose with sodium chloride 9 mg/mL (0.9%) solution for infusion containing 1% human albumin (e.g. 5 mL of human albumin 20% per 100 mL sodium chloride solution).

For continuous infusions, the solution for infusion can be prepared freshly on a daily basis, or sufficient for up to 5 days of continuous infusion. The daily dose is 10 mg/m<sup>2</sup>. The amount of solution to be infused per day (within a treatment course of 10 consecutive days) should be 48 mL; with 240 mL for a 5-day dose. It is recommended to prepare 50 mL solution in a 50 mL syringe, or 250 mL in an infusion bag suitable for the employed infusion pump, i.e. an overfill of 2 mL (syringe) or 10 mL (infusion bag) to allow for dead volumes of the infusion systems.

For repeated daily 8-hour infusions, the daily dose is 20 mg/m<sup>2</sup> and the calculated dose should be diluted in 100 mL sodium chloride 9 mg/mL (0.9%) containing 1% human albumin.

The solution for infusion should be administered via a peripheral or central intravenous line. Other intravenously co-administered agents should be delivered via a separate infusion line. The container should be inspected visually for particulates prior to administration. It is recommended that a 0.22 micrometre in-line filter is used during infusion.

For continuous infusions, any medical device suitable for infusion at a rate of 2 mL per hour can be used, e.g. syringe infusion pumps/infusors, electronic ambulatory infusion pumps. Note that elastomeric pumps are not considered suitable in combination with in-line filters.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Recordati Netherlands B.V.  
Beechavenue 54,  
1119PW Schiphol-Rijk  
Netherlands

## **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/17/1191/001

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 08 May 2017  
Date of latest renewal: 06 January 2022

## **10. DATE OF REVISION OF THE TEXT**

{MM/YYYY}

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.



## **ANNEX II**

- A. MANUFACTURER(S) OF THE BIOLOGICAL ACTIVE SUBSTANCE(S) AND MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**
- E. SPECIFIC OBLIGATION TO COMPLETE POST-AUTHORISATION MEASURES FOR THE MARKETING AUTHORISATION UNDER EXCEPTIONAL CIRCUMSTANCES**

**A. MANUFACTURER(S) OF THE BIOLOGICAL ACTIVE SUBSTANCE(S) AND MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer(s) of the biological active substance(s)

Rentschler Biopharma SE  
Erwin-Rentschler-Strasse 21  
Laupheim, Baden-Wuerttemberg  
88471  
Germany

Name and address of the manufacturer(s) responsible for batch release

Patheon Italia S.P.A.  
Via Morolense,  
5 – 03013 Ferentino  
Italy

**B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

**C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.  
The marketing authorisation holder (MAH) shall submit the first PSUR for this product within 6 months following authorisation.

**D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

**E. SPECIFIC OBLIGATION TO COMPLETE POST-AUTHORISATION MEASURES FOR THE MARKETING AUTHORISATION UNDER EXCEPTIONAL CIRCUMSTANCES**

This being a marketing authorisation under exceptional circumstances and pursuant to Article 14(8) of Regulation (EC) No 726/2004, the MAH shall complete, within the stated timeframe, the following measures:

<b>Description</b>	<b>Due date</b>
Non-interventional post-authorisation safety study (PASS): In order to collect data on pain and its management, effect on peripheral and central nervous system, including visual impairment, long-term safety and long-term effectiveness, the MAH should submit the results of a study based on data deriving from a registry of patients with high-risk neuroblastoma.	Annual reports to be submitted

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****CARTON****1. NAME OF THE MEDICINAL PRODUCT**

Qarziba 4.5 mg/mL concentrate for solution for infusion  
dinutuximab beta

**2. STATEMENT OF ACTIVE SUBSTANCE**

1 mL of concentrate contains 4.5 mg dinutuximab beta.  
Each 4.5 mL vial contains 20 mg dinutuximab beta.

**3. LIST OF EXCIPIENTS**

Histidine, sucrose, polysorbate 20, water for injections, hydrochloric acid.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Concentrate for solution for infusion  
1 vial  
20 mg/4.5 mL

**5. METHOD AND ROUTE OF ADMINISTRATION**

Read the package leaflet before use.  
Intravenous use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator.  
Keep the vial in the outer carton in order to protect from light.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Recordati Netherlands B.V.  
Beechavenue 54,  
1119PW Schiphol-Rijk  
Netherlands

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/17/1191/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Qarziba

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

<b>MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS</b> <b>VIAL LABEL</b>
--

<b>1. NAME OF THE MEDICINAL PRODUCT AND ROUTE OF ADMINISTRATION</b>
---

Qarziba 4.5 mg/mL concentrate for solution for infusion  
dinutuximab beta  
Intravenous use

<b>2. METHOD OF ADMINISTRATION</b>
------------------------------------

<b>3. EXPIRY DATE</b>
-----------------------

EXP

<b>4. BATCH NUMBER</b>
------------------------

Lot

<b>5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT</b>
--

20 mg/4.5 mL

<b>6. OTHER</b>
-----------------



## **B. PACKAGE LEAFLET**

## Package leaflet: Information for the user

### Qarziba 4.5 mg/mL concentrate for solution for infusion dinutuximab beta

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Qarziba is and what it is used for
2. What you need to know before you use Qarziba
3. How to use Qarziba
4. Possible side effects
5. How to store Qarziba
6. Contents of the pack and other information

#### 1. What Qarziba is and what it is used for

Qarziba contains dinutuximab beta, which belongs to a group of medicines called ‘monoclonal antibodies’. These are proteins, which specifically recognise and bind to other unique proteins in the body. Dinutuximab beta binds to the molecule known as disialoganglioside 2 (GD2), which is present on cancer cells, and this activates the body’s immune system, causing it to attack the cancer cells.

Qarziba is **used to treat neuroblastoma** that has a high risk of coming back after a series of treatments, which include a stem cell transplantation for rebuilding the immune system. It is also used to treat neuroblastoma that has come back (relapsed) or could not be completely treated with previous therapies.

Prior to the treatment of relapsed neuroblastoma, your treating physician will stabilise any actively progressing disease by other suitable measures.

Your doctor will further decide whether the co-administration of a second medicine, interleukin-2, is necessary for the treatment of your cancer.

Neuroblastoma is a type of cancer that grows from abnormal nerve cells in the body, in particular in the glands above the kidneys. It is one of the most common cancers in infancy.

It is used for patients aged 12 months and above.

#### 2. What you need to know before you use Qarziba

##### Do not use Qarziba if you

- are **allergic** to dinutuximab beta or any of the other ingredients of this medicine (listed in section 6)
- have acute grade 3 or 4, or extensive long-lasting graft-versus-host disease

This disease is a reaction in which **cells of transplanted tissue attack cells of the recipient**.

## **Warnings and precautions**

Before receiving Qarziba, you will have blood tests to check your liver, lung, renal and bone marrow functions.

You might notice the following when you first receive Qarziba and during the course of treatment:

- **pain**  
Pain is one of the most common side effects of Qarziba. It usually occurs at the beginning of infusion. Therefore, your doctor will give you an appropriate pain treatment starting 3 days before and continuing during use of Qarziba.
- **allergic reactions or other infusion-related reactions**  
Tell your doctor or nurse if you have any kind of reaction during or after the infusion, such as:
  - fever, shivering and/or low blood pressure
  - difficulties in breathing
  - skin rash, hivesYou will receive appropriate treatment to prevent these reactions and be closely monitored for these symptoms during infusion of Qarziba.
- **leakage from small blood vessels (capillary leak syndrome)**  
Leakage of blood components from small blood vessels may cause rapid swelling in arms, legs and other parts of the body. Rapid drop in blood pressure, light-headedness and breathing difficulties are further signs.
- **eye problems**  
You may notice changes to your vision.
- **problems with your nerves**  
You may notice numbness, tingling or burning in your hands, feet, legs or arms, reduced sensation or weakness with movement.
- **spinal cord and brain problems (central nervous system, CNS)**
- Tell your doctor or nurse if you have any kind of CNS symptoms, such as: substantial prolonged neurological deficit without apparent reason such as muscle weakness or loss of muscle strength in the legs (or arms), or mobility problems or unusual sensations and numbness. Persistent or sudden onset of a headache, or progressive loss of memory and cognitive ability, subtle personality changes, inability to concentrate, lethargy, and progressive loss of consciousness.
- **symptoms of kidney failure**  
Tell your doctor or nurse if you notice an altered frequency or absence of urination.

Tell your doctor immediately if you notice any of these problems.

Your doctor may decide to stop your treatment if you have any of the problems mentioned here. In some cases your treatment may be able to start again after a break or at a slower rate, but sometimes it may need to be stopped completely.

Your doctor will do blood tests and may do eye tests while you are taking this medicine.

## **Children**

This medicine should not be given to children under 12 months because there is insufficient experience in this age group.

## **Other medicines and Qarziba**

Tell your doctor if you are using, have recently used or might use any other medicines.

Do not use **medicines that suppress the immune system** from 2 weeks before the first dose of Qarziba until 1 week after the last treatment course, unless prescribed by your doctor. Examples of

medicines that suppress the immune system are corticosteroids used to reduce inflammation or prevent organ transplant rejection.

Avoid **vaccinations** during treatment with Qarziba and for 10 weeks afterwards.

### **Pregnancy, breast-feeding and fertility**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

Talk to your doctor before you receive Qarziba if you are of childbearing age. It is recommended to use contraception for 6 months after discontinuation of treatment with Qarziba. You may only use Qarziba if your doctor assesses that benefits outweigh risks for a foetus.

Tell your doctor if you are breast-feeding. Do not breast-feed during treatment with Qarziba and for 6 months after the last dose. It is not known if the medicine can pass into breast-milk.

### **Driving and using machines**

Qarziba has several side effects that may affect your ability to drive and use machines. Do not perform these activities if your ability to concentrate and react is affected.

## **3. How to use Qarziba**

A doctor experienced in the use of medicines to treat cancer will supervise your treatment. It will be given to you by a doctor or nurse while you are in hospital. It is given into one of your veins (intravenous infusion) usually by using special tubes (catheters) and a pump. During and after the infusion, you will be checked regularly for infusion-related side effects.

Qarziba will be given to you in five treatment courses of 35 days and the infusion will last 5 or 10 days in the beginning of each course. The recommended dose is **100 mg** dinutuximab beta **per square metre of body surface per treatment course**. The doctor will calculate your body surface area from your height and weight.

If your doctor considers co-administration of interleukin-2, it will be given twice, by injection under the skin, each time for 5 consecutive days (before and during treatment with Qarziba).

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Tell your doctor or nurse immediately** if you have any of the following:

**Very common** (may affect more than 1 in 10 people):

- rapid swelling of arms, legs and other body parts, rapid drop in blood pressure, light-headedness and breathing difficulties (capillary leak syndrome)
- pain in the stomach, throat, chest, face, hands, feet, legs, arms, back, neck, joint, or muscles
- allergic reactions and cytokine release syndrome with symptoms such as face or throat swelling, breathing difficulties, dizziness, hives, rapid or noticeable heartbeat, low blood pressure, hives, rash, fever, or nausea

Other side effects and their frequencies include:

**Very common** (may affect more than 1 in 10 people):

- fever, chills
- vomiting, diarrhoea, constipation
- inflammation of the mouth and lips (stomatitis)

- cough
- itching, rash
- low blood pressure, increased heartbeat
- oxygen deficiency
- tissue swelling (in the face, lip, around the eye, in the lower limbs)
- increased weight
- infection, in particular infection associated with the catheter that delivers the medicine
- headache
- dilated pupils or abnormal pupil reactions
- abnormal blood or urine tests (blood cells and other components, liver function, renal function)

**Common** (may affect up to 1 in 10 people):

- life-threatening infection (sepsis)
- fits
- agitation, anxiety
- nerve disorder in the arms and/or legs (with abnormal sensations or weakness), light-headedness, trembling, muscle spasms
- paralysis of eye muscles, blurred vision, light sensitivity, swelling in the retina
- high blood pressure
- cardiac failure, fluid around the heart
- respiratory failure, fluid in the lungs
- sudden constriction of the airways (bronchospasm, laryngospasm), rapid breathing
- decreased appetite, nausea, abdominal distension, accumulation of fluid in the abdominal cavity
- injection-site reactions, skin problems such as reddening, dry skin, eczema, excessive sweating, reaction to light
- unable to pass urine or passing reduced urine volume
- decreased weight, loss of fluids (dehydration)

**Uncommon** (may affect up to 1 in 100 people):

- shock due to decreased body fluid volume
- formation of blood clots in the small blood vessels (disseminated intravascular coagulation)
- a type of allergy (serum sickness) with fever, rash, joint inflammation
- a brain disorder characterised by headache, confusion, seizures and loss of vision (posterior reversible encephalopathy syndrome)
- inflammation of the intestine, injury to the liver
- kidney failure
- a condition in which some of the small veins in the liver are obstructed (veno-occlusive disease)

**Not known** (frequency cannot be estimated from the available data):

- extreme tiredness and shortness of breath (which may be due to a low number of red blood cells), bleeding and bruising (which may be due to a low number of blood platelets) and kidney disease where you pass little or no urine (atypical haemolytic uraemic syndrome)

**Reporting of side effects**

If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects, you can help provide more information on the safety of this medicine.

**5. How to store Qarziba**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label and the carton after EXP. The expiry date refers to the last day of that month.

Store in a refrigerator (2 °C – 8 °C). Keep the vial in the outer carton in order to protect from light.

Once opened, Qarziba is intended for immediate use.

## **6. Contents of the pack and other information**

### **What Qarziba contains**

- The active substance is dinutuximab beta.  
1 mL concentrate contains 4.5 mg dinutuximab beta. Each vial contains 20 mg dinutuximab beta in 4.5 mL.
- The other ingredients are histidine, sucrose, polysorbate 20, water for injections, hydrochloric acid (for pH adjustment).

### **What Qarziba looks like and contents of the pack**

Qarziba is colourless to slightly yellow liquid, provided in a clear glass vial with a rubber stopper and aluminium seal.

Each carton contains 1 vial.

- **Marketing Authorisation Holder**  
Recordati Netherlands B.V.  
Beechavenue 54,  
1119PW Schiphol-Rijk  
Netherlands
- **Manufacturer**  
Patheon Italia S.P.A.  
Via Morolense,  
5 – 03013 Ferentino  
Italy

**This leaflet was last revised in {MM/YYYY}.**

### **Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency website: <http://www.ema.europa.eu>. There are also links to other websites about rare diseases and treatments.

---

### **The following information is intended for healthcare professionals only:**

Qarziba is restricted to hospital-use only and must be administered under the supervision of a physician experienced in the use of oncological therapies. It must be administered by a healthcare professional prepared to manage severe allergic reactions including anaphylaxis in an environment where full resuscitation services are immediately available.

#### Posology

Treatment with dinutuximab beta consists of 5 consecutive courses, each course comprising 35 days. The individual dose is determined based on the body surface area and should be a total of 100 mg/m<sup>2</sup> per course.

Two modes of administration are possible:

- a continuous infusion over the first 10 days of each course (a total of 240 hours) at the daily dose of  $10 \text{ mg/m}^2$
- or five daily infusions of  $20 \text{ mg/m}^2$  administered over 8 hours, on the first 5 days of each course

If IL-2 is combined with dinutuximab beta, it should be given as subcutaneous injections for 5 consecutive days twice during each course. First 5-day treatment should start 7 days prior to first dinutuximab beta infusion. Second 5-day treatment with IL-2 should start concurrently with dinutuximab beta infusion (days 1 to 5 of each course). IL-2 is administered as  $6 \times 10^6 \text{ IU/m}^2/\text{day}$ , resulting in an overall dose of  $60 \times 10^6 \text{ IU/m}^2/\text{course}$ .

#### Preparation of the infusion

The solution for infusion must be prepared under aseptic conditions. The solution must not be exposed to direct sunlight or heat.

The patient-specific daily dose of Qarziba is calculated based on body surface area. Qarziba should be diluted aseptically to the patient-specific concentration/dose with sodium chloride  $9 \text{ mg/mL}$  (0.9%) solution for infusion, containing 1% human albumin (e.g. 5 mL of human albumin 20% per 100 mL sodium chloride solution).

- For continuous infusions, the solution for infusion can be prepared freshly on a daily basis, or sufficient for up to 5 days of continuous infusion. The daily dose is  $10 \text{ mg/m}^2$ . The amount of solution to be infused per day (within a treatment course of 10 consecutive days) should be 48 mL; with 240 mL for a 5-day dose. It is recommended to prepare 50 mL solution in a 50 mL syringe, or 250 mL in an infusion bag suitable for the employed infusion pump, i.e. an overfill of 2 mL (syringe) or 10 mL (infusion bag) to allow for dead volumes of the infusion systems.
- For repeated daily infusions, the daily dose is  $20 \text{ mg/m}^2$  and the calculated dose should be diluted in 100 mL sodium chloride  $9 \text{ mg/mL}$  (0.9%) containing 1% human albumin.

#### Administration of the infusion

The solution for infusion should be administered via a peripheral or central intravenous line. Other intravenously co-administered agents should be delivered via a separate infusion line. The container should be inspected visually for particulates prior to administration. It is recommended that a 0.22 micrometre in-line filter is used during infusion.

For continuous infusions, any medical device suitable for infusion at a rate of 2 mL per hour can be applied, e.g. syringe infusion pumps/infusors, electronic ambulatory infusion pumps. Note that elastomeric pumps are not considered suitable in combination with in-line filters.

#### Storage of the diluted solution

Chemical and physical in-use stability has been demonstrated for up to 48 hours at  $25^\circ\text{C}$  (50 mL syringe) and for up to 7 days at  $37^\circ\text{C}$  (250 mL infusion bag), after cumulative storage in a refrigerator ( $2^\circ\text{C} - 8^\circ\text{C}$ ) for 72 hours.

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would not normally be longer than 24 hours at  $2$  to  $8^\circ\text{C}$ , unless dilution has taken place in controlled and validated aseptic conditions.

#### Disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.