

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## 1. NAME OF THE MEDICINAL PRODUCT

Ranluspec 10 mg/mL solution for injection

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One mL contains 10 mg ranibizumab\*. Each vial contains 2.3 mg of ranibizumab in 0.23 mL solution. This provides a usable amount to deliver a single dose of 0.05 mL containing 0.5 mg ranibizumab to adult patients.

\*Ranibizumab is a humanised monoclonal antibody fragment produced in *Escherichia coli* cells by recombinant DNA technology.

### Excipient with known effect

Each mL contains 0.1 mg polysorbate 20.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Solution for injection.

Clear to slightly opalescent, colourless to pale brownish aqueous solution.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Ranluspec is indicated in adults for:

- The treatment of neovascular (wet) age-related macular degeneration (AMD)
- The treatment of visual impairment due to diabetic macular oedema (DME)
- The treatment of proliferative diabetic retinopathy (PDR)
- The treatment of visual impairment due to macular oedema secondary to retinal vein occlusion (branch RVO or central RVO)
- The treatment of visual impairment due to choroidal neovascularisation (CNV)

### 4.2 Posology and method of administration

Ranluspec must be administered by a qualified ophthalmologist experienced in intravitreal injections.

#### Posology

The recommended dose for ranibizumab in adults is 0.5 mg given as a single intravitreal injection. This corresponds to an injection volume of 0.05 mL. The interval between two doses injected into the same eye should be at least four weeks.

Treatment in adults is initiated with one injection per month until maximum visual acuity is achieved and/or there are no signs of disease activity i.e. no change in visual acuity and in other signs and

symptoms of the disease under continued treatment. In patients with wet AMD, DME, PDR and RVO, initially, three or more consecutive, monthly injections may be needed.

Thereafter, monitoring and treatment intervals should be determined by the physician and should be based on disease activity, as assessed by visual acuity and/or anatomical parameters.

If, in the physician's opinion, visual and anatomical parameters indicate that the patient is not benefiting from continued treatment, Ranluspec should be discontinued.

Monitoring for disease activity may include clinical examination, functional testing or imaging techniques (e.g. optical coherence tomography or fluorescein angiography).

If patients are being treated according to a treat-and-extend regimen, once maximum visual acuity is achieved and/or there are no signs of disease activity, the treatment intervals can be extended stepwise until signs of disease activity or visual impairment recur. The treatment interval should be extended by no more than two weeks at a time for wet AMD and may be extended by up to one month at a time for DME. For PDR and RVO, treatment intervals may also be gradually extended, however there are insufficient data to conclude on the length of these intervals. If disease activity recurs, the treatment interval should be shortened accordingly.

The treatment of visual impairment due to CNV should be determined individually per patient based on disease activity. Some patients may only need one injection during the first 12 months; others may need more frequent treatment, including a monthly injection. For CNV secondary to pathologic myopia (PM), many patients may only need one or two injections during the first year (see section 5.1).

#### *Ranibizumab and laser photocoagulation in DME and in macular oedema secondary to BRVO*

There is some experience of ranibizumab administered concomitantly with laser photocoagulation (see section 5.1). When given on the same day, ranibizumab should be administered at least 30 minutes after laser photocoagulation. Ranibizumab can be administered in patients who have received previous laser photocoagulation.

#### *Ranibizumab and verteporfin photodynamic therapy in CNV secondary to PM*

There is no experience of concomitant administration of ranibizumab and verteporfin.

### Special populations

#### *Hepatic impairment*

Ranibizumab has not been studied in patients with hepatic impairment. However, no special considerations are needed in this population.

#### *Renal impairment*

Dose adjustment is not needed in patients with renal impairment (see section 5.2).

#### *Elderly*

No dose adjustment is required in the elderly. There is limited experience in patients older than 75 years with DME.

#### *Paediatric population*

The safety and efficacy of Ranluspec in children and adolescents below 18 years of age have not been established. Available data in adolescent patients aged 12 to 17 years with visual impairment due to CNV are described in section 5.1 but no recommendation on a posology can be made.

### Method of administration

Single-use vial for intravitreal use only.

Since the volume contained in the vial (0.23 mL) is greater than the recommended dose (0.05 mL for adults), a portion of the volume contained in the vial must be discarded prior to administration.

Ranluspec should be inspected visually for particulate matter and discoloration prior to administration.

The injection procedure should be carried out under aseptic conditions, which includes the use of surgical hand disinfection, sterile gloves, a sterile drape and a sterile eyelid speculum (or equivalent) and the availability of sterile paracentesis (if required). The patient's medical history for hypersensitivity reactions should be carefully evaluated prior to performing the intravitreal procedure (see section 4.4). Adequate anaesthesia and a broad-spectrum topical microbicide to disinfect the periocular skin, eyelid and ocular surface should be administered prior to the injection, in accordance with local practice.

In adults the injection needle should be inserted 3.5–4.0 mm posterior to the limbus into the vitreous cavity, avoiding the horizontal meridian and aiming towards the centre of the globe. The injection volume of 0.05 mL is then delivered; a different scleral site should be used for subsequent injections.

For instructions on preparation of the medicinal product before administration, see section 6.6.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Patients with active or suspected ocular or periocular infections.

Patients with active severe intraocular inflammation.

### **4.4 Special warnings and precautions for use**

#### Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

#### Intravitreal injection-related reactions

Intravitreal injections, including those with ranibizumab, have been associated with endophthalmitis, intraocular inflammation, rhegmatogenous retinal detachment, retinal tear and iatrogenic traumatic cataract (see section 4.8). Proper aseptic injection techniques must always be used when administering ranibizumab. In addition, patients should be monitored during the week following the injection to permit early treatment if an infection occurs. Patients should be instructed to report any symptoms suggestive of endophthalmitis or any of the abovementioned events without delay.

#### Intraocular pressure increases

In adults, transient increases in intraocular pressure (IOP) have been seen within 60 minutes of injection of ranibizumab. Sustained IOP increases have also been identified (see section 4.8). Both intraocular pressure and the perfusion of the optic nerve head must be monitored and managed appropriately.

Patients should be informed of the symptoms of these potential adverse reactions and instructed to inform their physician if they develop signs such as eye pain or increased discomfort, worsening eye redness, blurred or decreased vision, an increased number of small particles in their vision, or increased sensitivity to light (see section 4.8).

#### Bilateral treatment

Limited data on bilateral use of ranibizumab (including same-day administration) do not suggest an increased risk of systemic adverse events compared with unilateral treatment.

### Immunogenicity

There is a potential for immunogenicity with ranibizumab. Since there is a potential for an increased systemic exposure in subjects with DME, an increased risk for developing hypersensitivity in this patient population cannot be excluded. Patients should also be instructed to report if an intraocular inflammation increases in severity, which may be a clinical sign attributable to intraocular antibody formation.

### Concomitant use of other anti-VEGF (vascular endothelial growth factor)

Ranibizumab should not be administered concurrently with other anti-VEGF medicinal products (systemic or ocular).

### Withholding ranibizumab in adults

The dose should be withheld and treatment should not be resumed earlier than the next scheduled treatment in the event of:

- a decrease in best-corrected visual acuity (BCVA) of  $\geq 30$  letters compared with the last assessment of visual acuity;
- an intraocular pressure of  $\geq 30$  mmHg;
- a retinal break;
- a subretinal haemorrhage involving the centre of the fovea, or, if the size of the haemorrhage is  $\geq 50\%$ , of the total lesion area;
- performed or planned intraocular surgery within the previous or next 28 days.

### Retinal pigment epithelial tear

Risk factors associated with the development of a retinal pigment epithelial tear after anti-VEGF therapy for wet AMD and potentially also other forms of CNV, include a large and/or high pigment epithelial retinal detachment. When initiating ranibizumab therapy, caution should be used in patients with these risk factors for retinal pigment epithelial tears.

### Rhegmatogenous retinal detachment or macular holes in adults

Treatment should be discontinued in subjects with rhegmatogenous retinal detachment or stage 3 or 4 macular holes.

### Populations with limited data

There is only limited experience in the treatment of subjects with DME due to type I diabetes. Ranibizumab has not been studied in patients who have previously received intravitreal injections, in patients with active systemic infections, or in patients with concurrent eye conditions such as retinal detachment or macular hole. There is limited experience of treatment with ranibizumab in diabetic patients with an HbA1c over 108 mmol/mol (12%) and no experience in patients with uncontrolled hypertension. This lack of information should be considered by the physician when treating such patients.

There are insufficient data to conclude on the effect of ranibizumab in patients with RVO presenting irreversible ischaemic visual function loss.

In patients with PM, there are limited data on the effect of ranibizumab in patients who have previously undergone unsuccessful verteporfin photodynamic therapy (vPDT) treatment. Also, while a consistent effect was observed in subjects with subfoveal and juxtafoveal lesions, there are insufficient data to conclude on the effect of ranibizumab in PM subjects with extrafoveal lesions.

### Systemic effects following intravitreal use

Systemic adverse events including non-ocular haemorrhages and arterial thromboembolic events have been reported following intravitreal injection of VEGF inhibitors.

There are limited data on safety in the treatment of DME, macular oedema due to RVO and CNV secondary to PM patients with prior history of stroke or transient ischaemic attacks. Caution should be exercised when treating such patients (see section 4.8).

#### Ranluspec contains polysorbate 20

This medicinal product contains 0.005 mg of polysorbate 20 in each administered dose of 0.05 mL which is equivalent to 0.1 mg/mL. Polysorbates may cause allergic reactions.

### **4.5 Interaction with other medicinal products and other forms of interaction**

No formal interaction studies have been performed.

For the adjunctive use of vPDT and ranibizumab in wet AMD and PM, see section 5.1.

For the adjunctive use of laser photocoagulation and ranibizumab in DME and BRVO, see sections 4.2 and 5.1.

In clinical studies for the treatment of visual impairment due to DME, the outcome with regard to visual acuity or central retinal subfield thickness (CSFT) in patients treated with ranibizumab was not affected by concomitant treatment with thiazolidinediones.

### **4.6 Fertility, pregnancy and lactation**

#### Women of childbearing potential/contraception in females

Women of childbearing potential should use effective contraception during treatment.

#### Pregnancy

For ranibizumab no clinical data on exposed pregnancies are available. Studies in cynomolgus monkeys do not indicate direct or indirect harmful effects with respect to pregnancy or embryonal/foetal development (see section 5.3). The systemic exposure to ranibizumab is low after ocular administration, but due to its mechanism of action, ranibizumab must be regarded as potentially teratogenic and embryo-/foetotoxic. Therefore, ranibizumab should not be used during pregnancy unless the expected benefit outweighs the potential risk to the foetus. For women who wish to become pregnant and have been treated with ranibizumab, it is recommended to wait at least 3 months after the last dose of ranibizumab before conceiving a child.

#### Breast-feeding

Based on very limited data, ranibizumab may be excreted in human milk at low levels. The effect of ranibizumab on a breast-fed newborn/infant is unknown. As a precautionary measure, breast-feeding is not recommended during the use of ranibizumab.

#### Fertility

There are no data available on fertility.

### **4.7 Effects on ability to drive and use machines**

The treatment procedure may induce temporary visual disturbances, which may affect the ability to drive or use machines (see section 4.8). Patients who experience these signs must not drive or use machines until these temporary visual disturbances subside.

## 4.8 Undesirable effects

### Summary of the safety profile

The majority of adverse reactions reported following administration of ranibizumab are related to the intravitreal injection procedure.

The most frequently reported ocular adverse reactions following injection of ranibizumab are: eye pain, ocular hyperaemia, increased intraocular pressure, vitritis, vitreous detachment, retinal haemorrhage, visual disturbance, vitreous floaters, conjunctival haemorrhage, eye irritation, foreign body sensation in eyes, increased lacrimation, blepharitis, dry eye and eye pruritus.

The most frequently reported non-ocular adverse reactions are headache, nasopharyngitis and arthralgia.

Less frequently reported, but more serious, adverse reactions include endophthalmitis, blindness, retinal detachment, retinal tear and iatrogenic traumatic cataract (see section 4.4).

The adverse reactions experienced following administration of ranibizumab in clinical studies are summarised in the table below.

### Tabulated list of adverse reactions<sup>#</sup>

The adverse reactions are listed by system organ class and frequency using the following convention: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1\ 000$  to  $< 1/100$ ), rare ( $\geq 1/10\ 000$  to  $< 1/1\ 000$ ), very rare ( $< 1/10\ 000$ ), not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

#### Infections and infestations

<i>Very common</i>	Nasopharyngitis
<i>Common</i>	Urinary tract infection*

#### Blood and lymphatic system disorders

<i>Common</i>	Anaemia
---------------	---------

#### Immune system disorders

<i>Common</i>	Hypersensitivity
---------------	------------------

#### Psychiatric disorders

<i>Common</i>	Anxiety
---------------	---------

#### Nervous system disorders

<i>Very common</i>	Headache
--------------------	----------

#### Eye disorders

<i>Very common</i>	Vitritis, vitreous detachment, retinal haemorrhage, visual disturbance, eye pain, vitreous floaters, conjunctival haemorrhage, eye irritation, foreign body sensation in eyes, lacrimation increased, blepharitis, dry eye, ocular hyperaemia, eye pruritus.
--------------------	--

<i>Common</i>	Retinal degeneration, retinal disorder, retinal detachment, retinal tear, detachment of the retinal pigment epithelium, retinal pigment epithelium tear, visual acuity reduced, vitreous haemorrhage, vitreous disorder, uveitis, iritis, iridocyclitis, cataract, cataract subcapsular, posterior capsule opacification, punctate keratitis, corneal abrasion, anterior chamber flare, vision blurred, injection site haemorrhage, eye
---------------	---

*Uncommon* haemorrhage, conjunctivitis, conjunctivitis allergic, eye discharge, photopsia, photophobia, ocular discomfort, eyelid oedema, eyelid pain, conjunctival hyperaemia.  
Blindness, endophthalmitis, hypopyon, hyphaema, keratopathy, iris adhesion, corneal deposits, corneal oedema, corneal striae, injection site pain, injection site irritation, abnormal sensation in eye, eyelid irritation.

Respiratory, thoracic and mediastinal disorders

*Common* Cough

Gastrointestinal disorders

*Common* Nausea

Skin and subcutaneous tissue disorders

*Common* Allergic reactions (rash, urticaria, pruritus, erythema)

Musculoskeletal and connective tissue disorders

*Very common* Arthralgia

Investigations

*Very common* Intraocular pressure increased

# Adverse reactions were defined as adverse events (in at least 0.5 percentage points of patients) which occurred at a higher rate (at least 2 percentage points) in patients receiving treatment with ranibizumab 0.5 mg than in those receiving control treatment (sham or verteporfin PDT).

\* observed only in DME population

#### Product-class-related adverse reactions

In the wet AMD phase III studies, the overall frequency of non-ocular haemorrhages, an adverse event potentially related to systemic VEGF (vascular endothelial growth factor) inhibition, was slightly increased in ranibizumab-treated patients. However, there was no consistent pattern among the different haemorrhages. There is a theoretical risk of arterial thromboembolic events, including stroke and myocardial infarction, following intravitreal use of VEGF inhibitors. A low incidence rate of arterial thromboembolic events was observed in the ranibizumab clinical studies in patients with AMD, DME, PDR, RVO and CNV and there were no major differences between the groups treated with ranibizumab compared to control.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

## **4.9 Overdose**

Cases of accidental overdose have been reported from the clinical studies in wet AMD and post-marketing data. Adverse reactions associated with these reported cases were intraocular pressure increased, transient blindness, reduced visual acuity, corneal oedema, corneal pain, and eye pain. If an overdose occurs, intraocular pressure should be monitored and treated, if deemed necessary by the attending physician.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Ophthalmologicals, antineovascularisation agents, ATC code: S01LA04

Ranluspec is a biosimilar medicinal product. Detailed information is available on the website of the European Medicines Agency <https://www.ema.europa.eu/en>.

### Mechanism of action

Ranibizumab is a humanised recombinant monoclonal antibody fragment targeted against human vascular endothelial growth factor A (VEGF-A). It binds with high affinity to the VEGF-A isoforms (e.g. VEGF110, VEGF121 and VEGF165), thereby preventing binding of VEGF-A to its receptors VEGFR-1 and VEGFR-2. Binding of VEGF-A to its receptors leads to endothelial cell proliferation and neovascularisation, as well as vascular leakage, all of which are thought to contribute to the progression of the neovascular form of age-related macular degeneration, pathologic myopia and CNV or to visual impairment caused by either diabetic macular oedema or macular oedema secondary to RVO in adults.

### Clinical efficacy and safety

#### Treatment of wet AMD

In wet AMD, the clinical safety and efficacy of ranibizumab have been assessed in three randomised, double-masked, sham- or active-controlled studies of 24 months duration in patients with neovascular AMD. A total of 1,323 patients (879 active and 444 control) were enrolled in these studies.

In study FVF2598g (MARINA), 716 patients with minimally classic or occult with no classic lesions were randomised in a 1:1:1 ratio to receive monthly injections of ranibizumab 0.3 mg, ranibizumab 0.5 mg or sham.

In study FVF2587g (ANCHOR), 423 patients with predominantly classic CNV lesions were randomised in a 1:1:1 ratio to receive ranibizumab 0.3 mg monthly, ranibizumab 0.5 mg monthly or verteporfin PDT (at baseline and every 3 months thereafter if fluorescein angiography showed persistence or recurrence of vascular leakage).



Key outcome measures are summarised in Table 1 and Figure 1.

**Table 1 Outcomes at Month 12 and Month 24 in study FVF2598g (MARINA) and FVF2587g (ANCHOR)**

Outcome measure	Month	FVF2598g (MARINA)		FVF2587g (ANCHOR)	
		Sham (n = 238)	Ranibizumab 0.5 mg (n = 240)	Verteporfin PDT (n = 143)	Ranibizumab 0.5 mg (n = 140)
Loss of < 15 letters in visual acuity (%) <sup>a</sup> (maintenance of vision, primary endpoint)	Month 12	62%	95%	64%	96%
	Month 24	53%	90%	66%	90%
Gain of ≥ 15 letters in visual acuity (%) <sup>a</sup>	Month 12	5%	34%	6%	40%
	Month 24	4%	33%	6%	41%
Mean change in visual acuity (letters) (SD) <sup>a</sup>	Month 12	-10.5 (16.6)	+7.2 (14.4)	-9.5 (16.4)	+11.3 (14.6)
	Month 24	-14.9 (18.7)	+6.6 (16.5)	-9.8 (17.6)	+10.7 (16.5)

<sup>a</sup> p < 0.01

**Figure 1 Mean change in visual acuity from baseline to Month 24 in study FVF2598g (MARINA) and study FVF2587g (ANCHOR)**



- |   |  |
|---|--|
| <p><b>MARINA</b></p> <ul style="list-style-type: none"> <li>■ Ranibizumab 0.5 mg (n=240)</li> <li>○ Sham (n=238)</li> </ul> | <p><b>ANCHOR</b></p> <ul style="list-style-type: none"> <li>■ Ranibizumab 0.5 mg (n=140)</li> <li>● Verteporfin PDT (n=143)</li> </ul> |
|---|--|

Results from both studies indicated that continued ranibizumab treatment may also be of benefit in patients who lost  $\geq 15$  letters of best-corrected visual acuity (BCVA) in the first year of treatment.

Statistically significant patient-reported visual functioning benefits were observed in both MARINA and ANCHOR with ranibizumab treatment over the control group as measured by the NEI VFQ-25.

In study FVF3192g (PIER), 184 patients with all forms of neovascular AMD were randomised in a 1:1:1 ratio to receive ranibizumab 0.3 mg, ranibizumab 0.5 mg or sham injections once a month for 3 consecutive doses, followed by a dose administered once every 3 months. From Month 14 of the study, sham-treated patients were allowed to receive ranibizumab and from Month 19, more frequent treatments were possible. Patients treated with ranibizumab in PIER received a mean of 10 total treatments.

After an initial increase in visual acuity (following monthly dosing), on average, patients' visual acuity declined with quarterly dosing, returning to baseline at Month 12 and this effect was maintained in most ranibizumab-treated patients (82%) at Month 24. Limited data from sham subjects who later received ranibizumab suggested that early initiation of treatment may be associated with better

preservation of visual acuity.

Data from two studies (MONT BLANC, BPD952A2308 and DENALI, BPD952A2309) conducted post approval confirmed the efficacy of ranibizumab but did not demonstrate additional effect of the combined administration of verteporfin (Visudyne PDT) and ranibizumab compared to ranibizumab monotherapy.

Treatment of visual impairment due to CNV secondary to PM

The clinical safety and efficacy of ranibizumab in patients with visual impairment due to CNV in PM have been assessed based on the 12-month data of the double-masked, controlled pivotal study F2301 (RADIANCE). In this study 277 patients were randomised in a 2:2:1 ratio to the following arms:

- Group I (ranibizumab 0.5 mg, dosing regimen driven by “stability” criteria defined as no change in BCVA compared to two preceding monthly evaluations).
- Group II (ranibizumab 0.5 mg, dosing regimen driven by “disease activity” criteria defined as vision impairment attributable to intra- or subretinal fluid or active leakage due to the CNV lesion as assessed by optical coherence tomography and/or fluorescence angiography).
- Group III (vPDT - patients were allowed to receive ranibizumab treatment as of Month 3).

In Group II, which is the recommended posology (see section 4.2), 50.9% of patients required 1 or 2 injections, 34.5% required 3 to 5 injections and 14.7% required 6 to 12 injections over the 12-month study period. 62.9% of Group II patients did not require injections in the second 6 months of the study.

The key outcomes from RADIANCE are summarised in Table 2 and Figure 2.


**Table 2 Outcomes at Month 3 and 12 (RADIANCE)**

	<b>Group I Ranibizumab 0.5 mg “Vision stability” (n = 105)</b>	<b>Group II Ranibizumab 0.5 mg “Disease activity” (n = 116)</b>	<b>Group III vPDT<sup>b</sup> (n = 55)</b>
<b>Month 3</b>			
Mean average BCVA change from Month 1 to Month 3 compared to baseline <sup>a</sup> (letters)	+10.5	+10.6	+2.2
Proportion of patients who gained: ≥ 15 letters, or reached ≥ 84 letters in BCVA	38.1%	43.1%	14.5%
<b>Month 12</b>			
Number of injections up to Month 12:			
Mean	4.6	3.5	N/A
Median	4.0	2.5	N/A
Mean average BCVA change from Month 1 to Month 12 compared to baseline (letters)	+12.8	+12.5	N/A
Proportion of patients who gained: ≥ 15 letters, or reached ≥ 84 letters in BCVA	53.3%	51.7%	N/A

<sup>a</sup> p< 0.00001 comparison with vPDT control

<sup>b</sup> Comparative control up to Month 3. Patients randomised to vPDT were allowed to receive ranibizumab treatment as of Month 3 (in Group III, 38 patients received ranibizumab as of Month 3)

**Figure 2 Mean change from baseline BCVA over time to Month 12 (RADIANCE)**



The improvement of vision was accompanied by a reduction in central retinal thickness.

Patient-reported benefits were observed with ranibizumab treatment arms over vPDT (p-value < 0.05) in terms of improvement in the composite score and several subscales (general vision, near activities, mental health and dependency) of the NEI VFQ-25.

Treatment of visual impairment due to CNV (other than secondary to PM and wet AMD)

The clinical safety and efficacy of ranibizumab in patients with visual impairment due to CNV have been assessed based on the 12-month data of the double-masked, sham-controlled pivotal study G2301 (MINERVA). In this study 178 adult patients were randomised in a 2:1 ratio to receive:

- ranibizumab 0.5 mg at baseline, followed by an individualised dosing regimen driven by disease activity as assessed by visual acuity and/or anatomical parameters (e.g. visual acuity impairment, intra/sub-retinal fluid, haemorrhage or leakage);
- sham injection at baseline, followed by an individualised treatment regimen driven by disease activity.

At Month 2, all patients received open-label treatment with ranibizumab as needed.

Key outcome measures from MINERVA are summarised in Table 3 and Figure 3. An improvement of vision was observed and was accompanied by a reduction in central subfield thickness over the 12-month period.

The mean number of injections given over 12 months was 5.8 in the ranibizumab arm versus 5.4 in

those patients in the sham arm who were eligible to receive ranibizumab from Month 2 onwards. In the sham arm 7 out of 59 patients did not receive any treatment with ranibizumab in the study eye during the 12-month period.


**Table 3 Outcomes at Month 2 (MINERVA)**

	Ranibizumab 0.5 mg (n = 119)	Sham (n = 59)
Mean BCVA change from baseline to Month 2 <sup>a</sup>	9.5 letters	-0.4 letters
Patients gaining ≥ 15 letters from baseline or reaching 84 letters at Month 2	31.4%	12.3%
Patients not losing > 15 letters from baseline at Month 2	99.2%	94.7%
Reduction in CSFT <sup>b</sup> from baseline to Month 2 <sup>a</sup>	77 μm	-9.8 μm

<sup>a</sup> One-sided p < 0.001 comparison with sham control

<sup>b</sup> CSFT - central retinal subfield thickness

**Figure 3 Mean change from baseline BCVA over time to Month 12 (MINERVA)**



When comparing ranibizumab versus sham control at Month 2, a consistent treatment effect both overall and across baseline aetiology subgroups was observed:

**Table 4 Treatment effect overall and across baseline aetiology subgroups**

Overall and per baseline aetiology	Treatment effect over sham [letters]	Patient numbers [n] (treatment +sham)
Overall	9.9	178
Angioid streaks	14.6	27
Post-inflammatory retinochoroidopathy	6.5	28
Central serous chorioretinopathy	5.0	23
Idiopathic chorioretinopathy	11.4	63
Miscellaneous aetiologies <sup>a</sup>	10.6	37

<sup>a</sup> encompasses different aetiologies of low frequency of occurrence not included in the other subgroups

In the pivotal study G2301 (MINERVA), five adolescent patients aged 12 to 17 years with visual impairment secondary to CNV received open-label treatment with ranibizumab 0.5 mg at baseline followed by an individualised treatment regimen as for the adult population. BCVA improved from baseline to Month 12 in all five patients, ranging from 5 to 38 letters (mean of 16.6 letters). The improvement of vision was accompanied by a stabilisation or reduction in central subfield thickness over the 12-month period. The mean number of ranibizumab injections given in the study eye over 12 months was 3 (ranged from 2 to 5). Overall, ranibizumab treatment was well tolerated.

Treatment of visual impairment due to DME


The efficacy and safety of ranibizumab have been assessed in three randomised, controlled studies of at least 12 months duration. A total of 868 patients (708 active and 160 control) were enrolled in these studies.

In the phase II study D2201 (RESOLVE), 151 patients were treated with ranibizumab (6 mg/mL, n = 51, 10 mg/mL, n = 51) or sham (n = 49) by monthly intravitreal injections. The mean average change in BCVA from Month 1 to Month 12 compared to baseline was +7.8 (±7.72) letters in the pooled ranibizumab-treated patients (n = 102), compared to -0.1 (±9.77) letters for sham-treated patients; and the mean change in BCVA at Month 12 from baseline was 10.3 (±9.1) letters compared to -1.4 (±14.2) letters, respectively (p < 0.0001 for the treatment difference).

In the phase III study D2301 (RESTORE), 345 patients were randomised in a 1:1:1 ratio to receive ranibizumab 0.5 mg monotherapy and sham laser photocoagulation, combined ranibizumab 0.5 mg and laser photocoagulation or sham injection and laser photocoagulation. 240 patients, who had previously completed the 12-month RESTORE study, were enrolled in the open-label, multicentre 24-month extension (RESTORE Extension) study. Patients were treated with ranibizumab 0.5 mg *pro re nata* (PRN) in the same eye as the core study (D2301 RESTORE).

Key outcome measures are summarised in Table 5 (RESTORE and Extension) and Figure 4 (RESTORE).

**Figure 4 Mean change in visual acuity from baseline over time in study D2301 (RESTORE)**



BL=baseline; SE=standard error of mean

\* Difference in least square means,  $p < 0.0001/0.0004$  based on two-sided stratified Cochran-Mantel- Haenszel test

The effect at 12 months was consistent in most subgroups. However, subjects with a baseline BCVA > 73 letters and macular oedema with central retinal thickness < 300  $\mu\text{m}$  did not appear to benefit from treatment with ranibizumab compared to laser photocoagulation.

**Table 5 Outcomes at Month 12 in study D2301 (RESTORE) and at Month 36 in study D2301-E1 (RESTORE Extension)**

Outcome measures at Month 12 compared to baseline in study D2301 (RESTORE)	Ranibizumab 0.5 mg n = 115	Ranibizumab 0.5 mg + Laser n = 118	Laser n = 110
Mean average change in BCVA from Month 1 to Month 12 <sup>a</sup> ( $\pm$ SD)	6.1 (6.4) <sup>a</sup>	5.9 (7.9) <sup>a</sup>	0.8 (8.6)
Mean change in BCVA at Month 12 ( $\pm$ SD)	6.8 (8.3) <sup>a</sup>	6.4 (11.8) <sup>a</sup>	0.9 (11.4)
Gain of $\geq 15$ letters or BCVA $\geq 84$ letters at Month 12 (%)	22.6	22.9	8.2
Mean number of injections (Months 0-11)	7.0	6.8	7.3 (sham)
Outcome measure at Month 36 compared to D2301 (RESTORE) baseline in study D2301-E1 (RESTORE Extension)	Prior ranibizumab 0.5 mg n = 83	Prior ranibizumab 0.5 mg + laser n = 83	Prior laser n = 74
Mean change in BCVA at Month 24 (SD)	7.9 (9.0)	6.7 (7.9)	5.4 (9.0)
Mean change in BCVA at Month 36 (SD)	8.0 (10.1)	6.7 (9.6)	6.0 (9.4)
Gain of $\geq 15$ letters or BCVA $\geq 84$ letters at Month 36 (%)	27.7	30.1	21.6
Mean number of injections (Months 12-35)*	6.8	6.0	6.5

<sup>a</sup> $p < 0.0001$  for comparisons of ranibizumab arms vs. laser arm.

n in D2301-E1 (RESTORE Extension) is the number of patients with a value at both D2301 (RESTORE) baseline (Month 0) and at the Month 36 visit.

\* The proportion of patients who did not require any ranibizumab treatment during the extension phase was 19%, 25% and 20% in the prior ranibizumab, prior ranibizumab + laser and prior laser groups, respectively.

Statistically significant patient-reported benefits for most vision-related functions were observed with ranibizumab (with or without laser) treatment over the control group as measured by the NEI VFQ-25. For other subscales of this questionnaire no treatment differences could be established.

The long-term safety profile of ranibizumab observed in the 24-month extension study is consistent with the known ranibizumab safety profile.

In the phase IIIb study D2304 (RETAIN), 372 patients were randomised in 1:1:1 ratio to receive:

- ranibizumab 0.5 mg with concomitant laser photocoagulation on a treat-and-extend (TE) regimen,
- ranibizumab 0.5 mg monotherapy on a TE regimen,
- ranibizumab 0.5 mg monotherapy on a PRN regimen.

In all groups, ranibizumab was administered monthly until BCVA was stable for at least three consecutive monthly assessments. On TE, ranibizumab was administered at treatment intervals of 2-3 months. In all groups, monthly treatment was re-initiated upon a decrease in BCVA due to DME progression and continued until stable BCVA was reached again.

The number of scheduled treatment visits after the initial 3 injections, was 13 and 20 for the TE and PRN regimens, respectively. With both TE regimens, more than 70% of patients maintained their

BCVA with an average visit frequency of  $\geq 2$  months.

The key outcome measures are summarised in Table 6.

**Table 6 Outcomes in study D2304 (RETAIN)**

Outcome measure compared to baseline	TE ranibizumab 0.5 mg + laser n = 117	TE ranibizumab 0.5 mg alone n = 125	PRN ranibizumab 0.5 mg n = 117
Mean average change in BCVA from Month 1 to Month 12 (SD)	5.9 (5.5) <sup>a</sup>	6.1 (5.7) <sup>a</sup>	6.2 (6.0)
Mean average change in BCVA from Month 1 to Month 24 (SD)	6.8 (6.0)	6.6 (7.1)	7.0 (6.4)
Mean change in BCVA at Month 24 (SD)	8.3 (8.1)	6.5 (10.9)	8.1 (8.5)
Gain of $\geq 15$ letters or BCVA $\geq 84$ letters at Month 24(%)	25.6	28.0	30.8
Mean number of injections (months 0-23)	12.4	12.8	10.7

<sup>a</sup>p < 0.0001 for assessment of non-inferiority to PRN

In DME studies, the improvement in BCVA was accompanied by a reduction over time in mean CSFT in all the treatment groups.

#### Treatment of PDR

The clinical safety and efficacy of ranibizumab in patients with PDR have been assessed in Protocol S which evaluated the treatment with ranibizumab 0.5 mg intravitreal injections compared with panretinal photocoagulation (PRP). The primary endpoint was the mean visual acuity change at year 2. Additionally, change in diabetic retinopathy (DR) severity was assessed based on fundus photographs using the DR severity score (DRSS).

Protocol S was a multicentre, randomised, active-controlled, parallel-assignment, non-inferiority phase III study in which 305 patients (394 study eyes) with PDR with or without DME at baseline were enrolled. The study compared ranibizumab 0.5 mg intravitreal injections to standard treatment with PRP. A total of 191 eyes (48.5%) were randomised to ranibizumab 0.5 mg and 203 eyes (51.5%) eyes were randomised to PRP. A total of 88 eyes (22.3%) had baseline DME: 42 (22.0%) and 46 (22.7%) eyes in the ranibizumab and PRP groups, respectively.

In this study, the mean visual acuity change at year 2 was +2.7 letters in the ranibizumab group compared to -0.7 letters in the PRP group. The difference in least square means was 3.5 letters (95% CI: [0.2 to 6.7]).

At year 1, 41.8% of eyes experienced a  $\geq 2$ -step improvement in the DRSS when treated with ranibizumab (n = 189) compared to 14.6% of eyes treated with PRP (n = 199). The estimated difference between ranibizumab and laser was 27.4% (95% CI: [18.9, 35.9]).

**Table 7 DRSS improvement or worsening of  $\geq 2$  or  $\geq 3$  steps at year 1 in Protocol S (LOCF Method)**

Categorised change from baseline	Protocol S		
	Ranibizumab 0.5 mg (N = 189)	PRP (N = 199)	Difference in proportion (%), CI
$\geq 2$ -step improvement			
n (%)	79 (41.8%)	29 (14.6%)	27.4 (18.9, 35.9)
$\geq 3$ -step improvement			
n (%)	54 (28.6%)	6 (3.0%)	25.7 (18.9, 32.6)
$\geq 2$ -step worsening			
n (%)	3 (1.6%)	23 (11.6%)	-9.9 (-14.7, -5.2)
$\geq 3$ -step worsening			
n (%)	1 (0.5%)	8 (4.0%)	-3.4 (-6.3, -0.5)
DRSS = diabetic retinopathy severity score, n = number of patients who satisfied the condition at the visit, N = total number of study eyes.			

At year 1 in the ranibizumab-treated group in Protocol S,  $\geq 2$ -step improvement in DRSS was consistent in eyes without DME (39.9%) and with baseline DME (48.8%).

An analysis of year 2 data from Protocol S demonstrated that 42.3% (n = 80) of eyes in the ranibizumab-treated group had  $\geq 2$ -step improvement in DRSS from baseline compared with 23.1% (n = 46) of eyes in the PRP group. In the ranibizumab-treated group,  $\geq 2$ -step improvement in DRSS from baseline was observed in 58.5% (n = 24) of eyes with baseline DME and 37.8% (n = 56) of eyes without DME.

DRSS was also assessed in three separate active-controlled phase III DME studies (ranibizumab 0.5 mg PRN vs laser) that included a total of 875 patients, of whom approximately 75% were of Asian origin. In a meta-analysis of these studies, 48.4% of the 315 patients with gradable DRSS scores in the subgroup of patients with moderately severe non-proliferative DR (NPDR) or worse at baseline experienced a  $\geq 2$ -step improvement in the DRSS at Month 12 when treated with ranibizumab (n = 192) vs 14.6% of patients treated with laser (n = 123). The estimated difference between ranibizumab and laser was 29.9% (95% CI: [20.0, 39.7]). In the 405 DRSS gradable patients with moderate NPDR or better a  $\geq 2$ -step DRSS improvement was observed in 1.4% and 0.9% of the ranibizumab and laser groups, respectively.

#### Treatment of visual impairment due to macular oedema secondary to RVO

The clinical safety and efficacy of ranibizumab in patients with visual impairment due to macular oedema secondary to RVO have been assessed in the randomised, double-masked, controlled studies BRAVO and CRUISE that recruited subjects with BRVO (n = 397) and CRVO (n = 392), respectively. In both studies, subjects received either 0.3 mg or 0.5 mg ranibizumab or sham injections. After 6 months, patients in the sham-control arms switched to 0.5 mg ranibizumab.

Key outcome measures from BRAVO and CRUISE are summarised in Table 8 and Figures 5 and 6.


**Table 8 Outcomes at Month 6 and 12 (BRAVO and CRUISE)**

	BRAVO		CRUISE	
	Sham/ Ranibizumab 0.5 mg (n = 132)	Ranibizumab 0.5 mg (n = 131)	Sham/ Ranibizumab 0.5 mg (n = 130)	Ranibizumab 0.5 mg (n = 130)
Mean change in visual acuity at	7.3 (13.0)	18.3 (13.2)	0.8 (16.2)	14.9 (13.2)

Month 6 <sup>a</sup> (letters) (SD) (primary endpoint)				
Mean change in BCVA at Month 12 (letters) (SD)	12.1 (14.4)	18.3 (14.6)	7.3 (15.9)	13.9 (14.2)
Gain of ≥ 15 letters in visual acuity at Month 6 <sup>a</sup> (%)	28.8	61.1	16.9	47.7
Gain of ≥ 15 letters in visual acuity at Month 12 (%)	43.9	60.3	33.1	50.8
Proportion (%) receiving laser rescue over 12 months	61.4	34.4	NA	NA


<sup>a</sup>p < 0.0001 for both studies

**Figure 5 Mean change from baseline BCVA over time to Month 6 and Month 12 (BRAVO)**



BL=baseline; SE=standard error of mean

**Figure 6 Mean change from baseline BCVA over time to Month 6 and Month 12 (CRUISE)**



BL=baseline; SE=standard error of mean

In both studies, the improvement of vision was accompanied by a continuous and significant reduction in the macular oedema as measured by central retinal thickness.

In patients with CRVO (CRUISE and extension study HORIZON): Subjects treated with sham in the first 6 months who subsequently received ranibizumab did not achieve comparable gains in visual acuity by Month 24 (~6 letters) compared to subjects treated with ranibizumab from study start (~12 letters).

Statistically significant patient-reported benefits in subscales related to near and distance activity were observed with ranibizumab treatment over the control group as measured by the NEI VFQ-25.

The long-term (24 months) clinical safety and efficacy of ranibizumab in patients with visual impairment due to macular oedema secondary to RVO were assessed in the BRIGHTER (BRVO) and CRYSTAL (CRVO) studies. In both studies, subjects received a 0.5 mg ranibizumab PRN dosing regimen driven by individualised stabilisation criteria. BRIGHTER was a 3-arm randomised active-controlled study that compared 0.5 mg ranibizumab given as monotherapy or in combination with adjunctive laser photocoagulation to laser photocoagulation alone. After 6 months, subjects in the laser arm could receive 0.5 mg ranibizumab. CRYSTAL was a single-arm study with 0.5 mg ranibizumab monotherapy.

Key outcome measures from BRIGHTER and CRYSTAL are shown in Table 9.

**Table 9 Outcomes at Months 6 and 24 (BRIGHTER and CRYSTAL)**

	BRIGHTER			CRYSTAL
	Ranibizumab 0.5 mg N = 180	Ranibizumab 0.5 mg + Laser N = 178	Laser* N = 90	Ranibizumab 0.5 mg N = 356
Mean change in	+14.8	+14.8	+6.0	+12.0

	<b>BRIGHTER</b>			<b>CRYSTAL</b>
	Ranibizumab 0.5 mg N = 180	Ranibizumab 0.5 mg + Laser N = 178	Laser* N = 90	Ranibizumab 0.5 mg N = 356
BCVA at Month 6 <sup>a</sup> (letters) (SD)	(10.7)	(11.13)	(14.27)	(13.95)
Mean change in BCVA at Month 24 <sup>b</sup> (letters) (SD)	+15.5 (13.91)	+17.3 (12.61)	+11.6 (16.09)	+12.1 (18.60)
Gain of ≥ 15 letters in BCVA at Month 24 (%)	52.8	59.6	43.3	49.2
Mean number of injections (SD) (Months 0-23)	11.4 (5.81)	11.3 (6.02)	NA	13.1 (6.39)
<sup>a</sup>	p < 0.0001 for both comparisons in BRIGHTER at Month 6: Ranibizumab 0.5 mg vs Laser and Ranibizumab 0.5 mg + Laser vs Laser.			
<sup>b</sup>	p < 0.0001 for null hypothesis in CRYSTAL that the mean change at Month 24 from baseline is zero.			
*	Starting at Month 6 ranibizumab 0.5 mg treatment was allowed (24 patients were treated with laser only).			

In BRIGHTER, ranibizumab 0.5 mg with adjunctive laser therapy demonstrated non-inferiority versus ranibizumab monotherapy from baseline to Month 24 (95% CI -2.8, 1.4).

In both studies, a rapid and statistically significant decrease from baseline in central retinal subfield thickness was observed at Month 1. This effect was maintained up to Month 24.

The effect of ranibizumab treatment was similar irrespective of the presence of retinal ischaemia. In BRIGHTER, patients with ischaemia present (N = 46) or absent (N = 133) and treated with ranibizumab monotherapy had a mean change from baseline of +15.3 and +15.6 letters, respectively, at Month 24. In CRYSTAL, patients with ischaemia present (N = 53) or absent (N = 300) and treated with ranibizumab monotherapy had a mean change from baseline of +15.0 and +11.5 letters, respectively.

The effect in terms of visual improvement was observed in all patients treated with 0.5 mg ranibizumab monotherapy regardless of their disease duration in both BRIGHTER and CRYSTAL. In patients with < 3 months disease duration an increase in visual acuity of 13.3 and 10.0 letters was seen at Month 1; and 17.7 and 13.2 letters at Month 24 in BRIGHTER and CRYSTAL, respectively. The corresponding visual acuity gain in patients with ≥ 12 months disease duration was 8.6 and 8.4 letters in the respective studies. Treatment initiation at the time of diagnosis should be considered.

The long-term safety profile of ranibizumab observed in the 24-month studies is consistent with the known ranibizumab safety profile.

#### Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with the reference medicinal product containing ranibizumab in all subsets of the paediatric population in neovascular AMD, visual impairment due to DME, visual impairment due to macular oedema secondary to RVO, visual impairment due to CNV and diabetic retinopathy (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

Following monthly intravitreal administration of ranibizumab to patients with neovascular AMD,

serum concentrations of ranibizumab were generally low, with maximum levels ( $C_{\max}$ ) generally below the ranibizumab concentration necessary to inhibit the biological activity of VEGF by 50% (11-27 ng/mL, as assessed in an *in vitro* cellular proliferation assay).  $C_{\max}$  was dose proportional over the dose range of 0.05 to 1.0 mg/eye. Serum concentrations in a limited number of DME patients indicate that a slightly higher systemic exposure cannot be excluded compared to those observed in neovascular AMD patients. Serum ranibizumab concentrations in RVO patients were similar or slightly higher compared to those observed in neovascular AMD patients.

Based on analysis of population pharmacokinetics and disappearance of ranibizumab from serum for patients with neovascular AMD treated with the 0.5 mg dose, the average vitreous elimination half-life of ranibizumab is approximately 9 days. Upon monthly intravitreal administration of ranibizumab 0.5 mg/eye, serum ranibizumab  $C_{\max}$ , attained approximately 1 day after dosing, is predicted to generally range between 0.79 and 2.90 ng/mL, and  $C_{\min}$  is predicted to generally range between 0.07 and 0.49 ng/mL. Serum ranibizumab concentrations are predicted to be approximately 90,000-fold lower than vitreal ranibizumab concentrations.

Patients with renal impairment: No formal studies have been conducted to examine the pharmacokinetics of ranibizumab in patients with renal impairment. In a population pharmacokinetic analysis of neovascular AMD patients, 68% (136 of 200) of patients had renal impairment (46.5% mild [50-80 mL/min], 20% moderate [30-50 mL/min], and 1.5% severe [ $<30$  mL/min]). In RVO patients, 48.2% (253 of 525) had renal impairment (36.4% mild, 9.5% moderate and 2.3% severe). Systemic clearance was slightly lower, but this was not clinically significant.

Hepatic impairment: No formal studies have been conducted to examine the pharmacokinetics of ranibizumab in patients with hepatic impairment.

### 5.3 Preclinical safety data

Bilateral intravitreal administration of ranibizumab to cynomolgus monkeys at doses between 0.25 mg/eye and 2.0 mg/eye once every 2 weeks for up to 26 weeks resulted in dose-dependent ocular effects.

Intraocularly, there were dose-dependent increases in anterior chamber flare and cells with a peak 2 days after injection. The severity of the inflammatory response generally diminished with subsequent injections or during recovery. In the posterior segment, there were vitreal cell infiltration and floaters, which also tended to be dose-dependent and generally persisted to the end of the treatment period. In the 26-week study, the severity of the vitreous inflammation increased with the number of injections. However, evidence of reversibility was observed after recovery. The nature and timing of the posterior segment inflammation is suggestive of an immune-mediated antibody response, which may be clinically irrelevant. Cataract formation was observed in some animals after a relatively long period of intense inflammation, suggesting that the lens changes were secondary to severe inflammation. A transient increase in post-dose intraocular pressure was observed following intravitreal injections, irrespective of dose.

Microscopic ocular changes were related to inflammation and did not indicate degenerative processes. Granulomatous inflammatory changes were noted in the optic disc of some eyes. These posterior segment changes diminished, and in some instances resolved, during the recovery period.

Following intravitreal administration, no signs of systemic toxicity were detected. Serum and vitreous antibodies to ranibizumab were found in a subset of treated animals.

No carcinogenicity or mutagenicity data are available.

In pregnant monkeys, intravitreal ranibizumab treatment resulting in maximal systemic exposures 0.9-7-fold a worst case clinical exposure did not elicit developmental toxicity or teratogenicity, and had no effect on weight or structure of the placenta, although, based on its pharmacological effect ranibizumab should be regarded as potentially teratogenic and embryo-/foetotoxic.

The absence of ranibizumab-mediated effects on embryo-foetal development is plausibly related mainly to the inability of the Fab fragment to cross the placenta. Nevertheless, a case was described with high maternal ranibizumab serum levels and presence of ranibizumab in foetal serum, suggesting that the anti-ranibizumab antibody acted as (Fc region containing) carrier protein for ranibizumab, thereby decreasing its maternal serum clearance and enabling its placental transfer. As the embryo-foetal development investigations were performed in healthy pregnant animals and disease (such as diabetes) may modify the permeability of the placenta towards a Fab fragment, the study should be interpreted with caution.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

$\alpha,\alpha$ -trehalose dihydrate  
Histidine hydrochloride, monohydrate  
Histidine  
Polysorbate 20  
Water for injections

### **6.2 Incompatibilities**

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

3 years

### **6.4 Special precautions for storage**

Store in a refrigerator (2 °C - 8 °C). Do not freeze.  
Keep the vial in the outer carton in order to protect from light.  
Prior to use, the unopened vial may be kept at room temperature (25 °C) for up to 24 hours.

### **6.5 Nature and contents of container**

#### Vial pack

One vial (type I glass) with a stopper (chlorobutyl rubber) containing 0.23 mL sterile solution.  
Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal and other handling**

#### Vial pack

The vial is for single use only. After injection any unused product must be discarded. Any vial showing signs of damage or tampering must not be used. The sterility cannot be guaranteed unless the packaging seal remains intact.

For preparation and intravitreal injection the following medical devices for single use are needed:  
a 5  $\mu$ m filter needle (18G)  
a 1 mL sterile syringe (including a 0.05 mL mark) and an injection needle (30G x 1/2"), for adult patients  
These medical devices are not included within this pack.

To prepare Ranluspec for intravitreal administration **to adults**, please adhere to the following instructions:

Before withdrawal, remove the vial cap and clean the vial septum (e.g., with 70% alcohol swab).

Assemble a 5 µm filter needle (18G x 1½", 1.2 mm x 40 mm) onto a 1 mL syringe using aseptic technique. Push the blunt filter needle into the centre of the vial stopper until the needle touches the bottom edge of the vial.

Withdraw all the liquid from the vial, keeping the vial in an upright position, slightly inclined to ease complete withdrawal.

Ensure that the plunger rod is drawn sufficiently back when emptying the vial in order to completely empty the filter needle.

Leave the blunt filter needle in the vial and disconnect the syringe from the blunt filter needle. The filter needle should be discarded after withdrawal of the vial contents and should not be used for the intravitreal injection.

Aseptically and firmly assemble an injection needle (30G x ½", 0.3 mm x 13 mm) onto the syringe.

Carefully remove the cap from the injection needle without disconnecting the injection needle from the syringe.

Note: Grip at the hub of the injection needle while removing the cap.

Carefully expel the air along with the excess solution and adjust the dose to the 0.05 mL mark on the syringe. The syringe is ready for injection.

Note: Do not wipe the injection needle. Do not pull back on the plunger.

After injection, do not recap the needle or detach it from the syringe. Dispose of the used syringe together with the needle in a sharps disposal container or in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Lupin Europe GmbH  
Hanauer Landstraße 139-143  
60314 Frankfurt  
Germany

## **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/25/2012/001

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation:

Date of latest renewal:

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <https://www.ema.europa.eu>.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## 1. NAME OF THE MEDICINAL PRODUCT

Ranluspec 10 mg/mL solution for injection in pre-filled syringe

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One mL contains 10 mg ranibizumab\*. One pre-filled syringe contains 0.165 mL, equivalent to 1.65 mg ranibizumab. The extractable volume of one pre-filled syringe is 0.1 mL. This provides a usable amount to deliver a single dose of 0.05 mL containing 0.5 mg ranibizumab.

\*Ranibizumab is a humanised monoclonal antibody fragment produced in *Escherichia coli* cells by recombinant DNA technology.

### Excipient with known effect

Each mL contains 0.1 mg polysorbate 20.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Solution for injection.

Clear to slightly opalescent, colourless to pale brownish aqueous solution.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Ranluspec is indicated in adults for:

- The treatment of neovascular (wet) age-related macular degeneration (AMD)
- The treatment of visual impairment due to diabetic macular oedema (DME)
- The treatment of proliferative diabetic retinopathy (PDR)
- The treatment of visual impairment due to macular oedema secondary to retinal vein occlusion (branch RVO or central RVO)
- The treatment of visual impairment due to choroidal neovascularisation (CNV)

### 4.2 Posology and method of administration

Ranluspec must be administered by a qualified ophthalmologist experienced in intravitreal injections.

#### Posology

The recommended dose for ranibizumab is 0.5 mg given as a single intravitreal injection. This corresponds to an injection volume of 0.05 mL. The interval between two doses injected into the same eye should be at least four weeks.

Treatment in adults is initiated with one injection per month until maximum visual acuity is achieved and/or there are no signs of disease activity i.e. no change in visual acuity and in other signs and symptoms of the disease under continued treatment. In patients with wet AMD, DME, PDR and RVO,

initially, three or more consecutive, monthly injections may be needed.

Thereafter, monitoring and treatment intervals should be determined by the physician and should be based on disease activity, as assessed by visual acuity and/or anatomical parameters.

If, in the physician's opinion, visual and anatomical parameters indicate that the patient is not benefiting from continued treatment, Ranluspec should be discontinued.

Monitoring for disease activity may include clinical examination, functional testing or imaging techniques (e.g. optical coherence tomography or fluorescein angiography).

If patients are being treated according to a treat-and-extend regimen, once maximum visual acuity is achieved and/or there are no signs of disease activity, the treatment intervals can be extended stepwise until signs of disease activity or visual impairment recur. The treatment interval should be extended by no more than two weeks at a time for wet AMD and may be extended by up to one month at a time for DME. For PDR and RVO, treatment intervals may also be gradually extended, however there are insufficient data to conclude on the length of these intervals. If disease activity recurs, the treatment interval should be shortened accordingly.

The treatment of visual impairment due to CNV should be determined individually per patient based on disease activity. Some patients may only need one injection during the first 12 months; others may need more frequent treatment, including a monthly injection. For CNV secondary to pathologic myopia (PM), many patients may only need one or two injections during the first year (see section 5.1).

#### *Ranibizumab and laser photocoagulation in DME and in macular oedema secondary to BRVO*

There is some experience of ranibizumab administered concomitantly with laser photocoagulation (see section 5.1). When given on the same day, ranibizumab should be administered at least 30 minutes after laser photocoagulation. Ranibizumab can be administered in patients who have received previous laser photocoagulation.

#### *Ranibizumab and verteporfin photodynamic therapy in CNV secondary to PM*

There is no experience of concomitant administration of ranibizumab and verteporfin.

### Special populations

#### *Hepatic impairment*

Ranibizumab has not been studied in patients with hepatic impairment. However, no special considerations are needed in this population.

#### *Renal impairment*

Dose adjustment is not needed in patients with renal impairment (see section 5.2).

#### *Elderly*

No dose adjustment is required in the elderly. There is limited experience in patients older than 75 years with DME.

#### *Paediatric population*

The safety and efficacy of Ranluspec in children and adolescents below 18 years of age have not been established. Available data in adolescent patients aged 12 to 17 years with visual impairment due to CNV are described in section 5.1.

### Method of administration

Single-use pre-filled syringe for intravitreal use only. The pre-filled syringe contains more than the recommended dose of 0.5 mg. The extractable volume of the pre-filled syringe (0.1 mL) is not to be used in total. The excess volume should be expelled prior to injection. Injecting the entire volume of

the pre-filled syringe could result in overdose. To expel the air bubble along with the excess medicinal product, slowly push the plunger until the edge below the dome of the rubber stopper is aligned with the black dosing line on the syringe (equivalent to 0.05 mL, i.e., 0.5 mg ranibizumab).

Ranluspec should be inspected visually for particulate matter and discoloration prior to administration.

The injection procedure should be carried out under aseptic conditions, which includes the use of surgical hand disinfection, sterile gloves, a sterile drape and a sterile eyelid speculum (or equivalent) and the availability of sterile paracentesis (if required). The patient's medical history for hypersensitivity reactions should be carefully evaluated prior to performing the intravitreal procedure (see section 4.4). Adequate anaesthesia and a broad-spectrum topical microbicide to disinfect the periocular skin, eyelid and ocular surface should be administered prior to the injection, in accordance with local practice.

The injection needle should be inserted 3.5-4.0 mm posterior to the limbus into the vitreous cavity, avoiding the horizontal meridian and aiming towards the centre of the globe. The injection volume of 0.05 mL is then delivered; a different scleral site should be used for subsequent injections. Each pre-filled syringe should only be used for the treatment of a single eye.

For instructions on preparation of the medicinal product before administration, see section 6.6.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Patients with active or suspected ocular or periocular infections.

Patients with active severe intraocular inflammation.

### **4.4 Special warnings and precautions for use**

#### Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

#### Intravitreal injection-related reactions

Intravitreal injections, including those with ranibizumab, have been associated with endophthalmitis, intraocular inflammation, rhegmatogenous retinal detachment, retinal tear and iatrogenic traumatic cataract (see section 4.8). Proper aseptic injection techniques must always be used when administering ranibizumab. In addition, patients should be monitored during the week following the injection to permit early treatment if an infection occurs. Patients should be instructed to report any symptoms suggestive of endophthalmitis or any of the abovementioned events without delay.

#### Intraocular pressure increases

Transient increases in intraocular pressure (IOP) have been seen within 60 minutes of injection of ranibizumab. Sustained IOP increases have also been identified (see section 4.8). Both intraocular pressure and the perfusion of the optic nerve head must be monitored and managed appropriately.

Patients should be informed of the symptoms of these potential adverse reactions and instructed to inform their physician if they develop signs such as eye pain or increased discomfort, worsening eye redness, blurred or decreased vision, an increased number of small particles in their vision, or increased sensitivity to light (see section 4.8).

#### Bilateral treatment

Limited data on bilateral use of ranibizumab (including same-day administration) do not suggest an

increased risk of systemic adverse events compared with unilateral treatment.

### Immunogenicity

There is a potential for immunogenicity with ranibizumab. Since there is a potential for an increased systemic exposure in subjects with DME, an increased risk for developing hypersensitivity in this patient population cannot be excluded. Patients should also be instructed to report if an intraocular inflammation increases in severity, which may be a clinical sign attributable to intraocular antibody formation.

### Concomitant use of other anti-VEGF (vascular endothelial growth factor)

Ranibizumab should not be administered concurrently with other anti-VEGF medicinal products (systemic or ocular).

### Withholding ranibizumab

The dose should be withheld and treatment should not be resumed earlier than the next scheduled treatment in the event of:

- a decrease in best-corrected visual acuity (BCVA) of  $\geq 30$  letters compared with the last assessment of visual acuity;
- an intraocular pressure of  $\geq 30$  mmHg;
- a retinal break;
- a subretinal haemorrhage involving the centre of the fovea, or, if the size of the haemorrhage is  $\geq 50\%$ , of the total lesion area;
- performed or planned intraocular surgery within the previous or next 28 days.

### Retinal pigment epithelial tear

Risk factors associated with the development of a retinal pigment epithelial tear after anti-VEGF therapy for wet AMD and potentially also other forms of CNV, include a large and/or high pigment epithelial retinal detachment. When initiating ranibizumab therapy, caution should be used in patients with these risk factors for retinal pigment epithelial tears.

### Rhegmatogenous retinal detachment or macular holes

Treatment should be discontinued in subjects with rhegmatogenous retinal detachment or stage 3 or 4 macular holes.

### Populations with limited data

There is only limited experience in the treatment of subjects with DME due to type I diabetes. Ranibizumab has not been studied in patients who have previously received intravitreal injections, in patients with active systemic infections, or in patients with concurrent eye conditions such as retinal detachment or macular hole. There is limited experience of treatment with ranibizumab in diabetic patients with an HbA1c over 108 mmol/mol (12%) and no experience in patients with uncontrolled hypertension. This lack of information should be considered by the physician when treating such patients.

There are insufficient data to conclude on the effect of ranibizumab in patients with RVO presenting irreversible ischaemic visual function loss.

In patients with PM, there are limited data on the effect of ranibizumab in patients who have previously undergone unsuccessful verteporfin photodynamic therapy (vPDT) treatment. Also, while a consistent effect was observed in subjects with subfoveal and juxtafoveal lesions, there are insufficient data to conclude on the effect of ranibizumab in PM subjects with extrafoveal lesions.

### Systemic effects following intravitreal use

Systemic adverse events including non-ocular haemorrhages and arterial thromboembolic events have been reported following intravitreal injection of VEGF inhibitors.

There are limited data on safety in the treatment of DME, macular oedema due to RVO and CNV secondary to PM patients with prior history of stroke or transient ischaemic attacks. Caution should be exercised when treating such patients (see section 4.8).

### Ranluspec contains polysorbate 20

This medicinal product contains 0.005 mg of polysorbate 20 in each administered dose of 0.05 mL which is equivalent to 0.1 mg/mL. Polysorbates may cause allergic reactions.

## **4.5 Interaction with other medicinal products and other forms of interaction**

No formal interaction studies have been performed.

For the adjunctive use of vPDT and ranibizumab in wet AMD and PM, see section 5.1.

For the adjunctive use of laser photocoagulation and ranibizumab in DME and BRVO, see sections 4.2 and 5.1.

In clinical studies for the treatment of visual impairment due to DME, the outcome with regard to visual acuity or central retinal subfield thickness (CSFT) in patients treated with ranibizumab was not affected by concomitant treatment with thiazolidinediones.

## **4.6 Fertility, pregnancy and lactation**

### Women of childbearing potential/contraception in females

Women of childbearing potential should use effective contraception during treatment.

### Pregnancy

For ranibizumab no clinical data on exposed pregnancies are available. Studies in cynomolgus monkeys do not indicate direct or indirect harmful effects with respect to pregnancy or embryonal/foetal development (see section 5.3). The systemic exposure to ranibizumab is low after ocular administration, but due to its mechanism of action, ranibizumab must be regarded as potentially teratogenic and embryo-/foetotoxic. Therefore, ranibizumab should not be used during pregnancy unless the expected benefit outweighs the potential risk to the foetus. For women who wish to become pregnant and have been treated with ranibizumab, it is recommended to wait at least 3 months after the last dose of ranibizumab before conceiving a child.

### Breast-feeding

Based on very limited data, ranibizumab may be excreted in human milk at low levels. The effect of ranibizumab on a breast-fed newborn/infant is unknown. As a precautionary measure, breast-feeding is not recommended during the use of ranibizumab.

### Fertility

There are no data available on fertility.

## **4.7 Effects on ability to drive and use machines**

The treatment procedure may induce temporary visual disturbances, which may affect the ability to drive or use machines (see section 4.8). Patients who experience these signs must not drive or use

machines until these temporary visual disturbances subside.

## 4.8 Undesirable effects

### Summary of the safety profile

The majority of adverse reactions reported following administration of ranibizumab are related to the intravitreal injection procedure.

The most frequently reported ocular adverse reactions following injection of ranibizumab are: eye pain, ocular hyperaemia, increased intraocular pressure, vitritis, vitreous detachment, retinal haemorrhage, visual disturbance, vitreous floaters, conjunctival haemorrhage, eye irritation, foreign body sensation in eyes, increased lacrimation, blepharitis, dry eye and eye pruritus.

The most frequently reported non-ocular adverse reactions are headache, nasopharyngitis and arthralgia.

Less frequently reported, but more serious, adverse reactions include endophthalmitis, blindness, retinal detachment, retinal tear and iatrogenic traumatic cataract (see section 4.4).

The adverse reactions experienced following administration of ranibizumab in clinical studies are summarised in the table below.

### Tabulated list of adverse reactions<sup>#</sup>

The adverse reactions are listed by system organ class and frequency using the following convention: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1\ 000$  to  $< 1/100$ ), rare ( $\geq 1/10\ 000$  to  $< 1/1\ 000$ ), very rare ( $< 1/10\ 000$ ), not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

#### Infections and infestations

*Very common*

Nasopharyngitis

*Common*

Urinary tract infection\*

#### Blood and lymphatic system disorders

*Common*

Anaemia

#### Immune system disorders

*Common*

Hypersensitivity

#### Psychiatric disorders

*Common*

Anxiety

#### Nervous system disorders

*Very common*

Headache

#### Eye disorders

*Very common*

Vitritis, vitreous detachment, retinal haemorrhage, visual disturbance, eye pain, vitreous floaters, conjunctival haemorrhage, eye irritation, foreign body sensation in eyes, lacrimation increased, blepharitis, dry eye, ocular hyperaemia, eye pruritus.

*Common*

Retinal degeneration, retinal disorder, retinal detachment, retinal tear, detachment of the retinal pigment epithelium, retinal pigment epithelium tear, visual acuity reduced, vitreous haemorrhage, vitreous disorder, uveitis, iritis, iridocyclitis, cataract, cataract subcapsular, posterior capsule opacification, punctate keratitis, corneal abrasion,

*Uncommon* anterior chamber flare, vision blurred, injection site haemorrhage, eye haemorrhage, conjunctivitis, conjunctivitis allergic, eye discharge, photopsia, photophobia, ocular discomfort, eyelid oedema, eyelid pain, conjunctival hyperaemia.  
Blindness, endophthalmitis, hypopyon, hyphaema, keratopathy, iris adhesion, corneal deposits, corneal oedema, corneal striae, injection site pain, injection site irritation, abnormal sensation in eye, eyelid irritation.

Respiratory, thoracic and mediastinal disorders

*Common* Cough

Gastrointestinal disorders

*Common* Nausea

Skin and subcutaneous tissue disorders

*Common* Allergic reactions (rash, urticaria, pruritus, erythema)

Musculoskeletal and connective tissue disorders

*Very common* Arthralgia

Investigations

*Very common* Intraocular pressure increased

# Adverse reactions were defined as adverse events (in at least 0.5 percentage points of patients) which occurred at a higher rate (at least 2 percentage points) in patients receiving treatment with ranibizumab 0.5 mg than in those receiving control treatment (sham or verteporfin PDT).

\* observed only in DME population

#### Product-class-related adverse reactions

In the wet AMD phase III studies, the overall frequency of non-ocular haemorrhages, an adverse event potentially related to systemic VEGF (vascular endothelial growth factor) inhibition, was slightly increased in ranibizumab-treated patients. However, there was no consistent pattern among the different haemorrhages. There is a theoretical risk of arterial thromboembolic events, including stroke and myocardial infarction, following intravitreal use of VEGF inhibitors. A low incidence rate of arterial thromboembolic events was observed in the Ranibizumab clinical studies in patients with AMD, DME, PDR, RVO and CNV and there were no major differences between the groups treated with ranibizumab compared to control.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

## **4.9 Overdose**

Cases of accidental overdose have been reported from the clinical studies in wet AMD and post-marketing data. Adverse reactions associated with these reported cases were intraocular pressure increased, transient blindness, reduced visual acuity, corneal oedema, corneal pain, and eye pain. If an overdose occurs, intraocular pressure should be monitored and treated, if deemed necessary by the attending physician.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Ophthalmologicals, antineovascularisation agents, ATC code: S01LA04

Ranluspec is a biosimilar medicinal product. Detailed information is available on the website of the European Medicines Agency <https://www.ema.europa.eu/en>.

### Mechanism of action

Ranibizumab is a humanised recombinant monoclonal antibody fragment targeted against human vascular endothelial growth factor A (VEGF-A). It binds with high affinity to the VEGF-A isoforms (e.g. VEGF110, VEGF121 and VEGF165), thereby preventing binding of VEGF-A to its receptors VEGFR-1 and VEGFR-2. Binding of VEGF-A to its receptors leads to endothelial cell proliferation and neovascularisation, as well as vascular leakage, all of which are thought to contribute to the progression of the neovascular form of age-related macular degeneration, pathologic myopia and CNV or to visual impairment caused by either diabetic macular oedema or macular oedema secondary to RVO in adults.

### Clinical efficacy and safety

#### Treatment of wet AMD

In wet AMD, the clinical safety and efficacy of ranibizumab have been assessed in three randomised, double-masked, sham- or active-controlled studies of 24 months duration in patients with neovascular AMD. A total of 1,323 patients (879 active and 444 control) were enrolled in these studies.

In study FVF2598g (MARINA), 716 patients with minimally classic or occult with no classic lesions were randomised in a 1:1:1 ratio to receive monthly injections of ranibizumab 0.3 mg, ranibizumab 0.5 mg or sham.

In study FVF2587g (ANCHOR), 423 patients with predominantly classic CNV lesions were randomised in a 1:1:1 ratio to receive ranibizumab 0.3 mg monthly, ranibizumab 0.5 mg monthly or verteporfin PDT (at baseline and every 3 months thereafter if fluorescein angiography showed persistence or recurrence of vascular leakage).



Key outcome measures are summarised in Table 1 and Figure 1.

**Table 1 Outcomes at Month 12 and Month 24 in study FVF2598g (MARINA) and FVF2587g (ANCHOR)**

Outcome measure	Month	FVF2598g (MARINA)		FVF2587g (ANCHOR)	
		Sham (n = 238)	Ranibizumab 0.5 mg (n = 240)	Verteporfin PDT (n = 143)	Ranibizumab 0.5 mg (n = 140)
Loss of < 15 letters in visual acuity (%) <sup>a</sup> (maintenance of vision, primary endpoint)	Month 12	62%	95%	64%	96%
	Month 24	53%	90%	66%	90%
Gain of ≥ 15 letters in visual acuity (%) <sup>a</sup>	Month 12	5%	34%	6%	40%
	Month 24	4%	33%	6%	41%
Mean change in visual acuity (letters) (SD) <sup>a</sup>	Month 12	-10.5 (16.6)	+7.2 (14.4)	-9.5 (16.4)	+11.3 (14.6)
	Month 24	-14.9 (18.7)	+6.6 (16.5)	-9.8 (17.6)	+10.7 (16.5)

<sup>a</sup> p < 0.01

**Figure 1 Mean change in visual acuity from baseline to Month 24 in study FVF2598g (MARINA) and study FVF2587g (ANCHOR)**



- |   |  |
|---|--|
| <p><b>MARINA</b></p> <ul style="list-style-type: none"> <li>■ Ranibizumab 0.5 mg (n=240)</li> <li>○ Sham (n=238)</li> </ul> | <p><b>ANCHOR</b></p> <ul style="list-style-type: none"> <li>■ Ranibizumab 0.5 mg (n=140)</li> <li>● Verteporfin PDT (n=143)</li> </ul> |
|---|--|

Results from both studies indicated that continued ranibizumab treatment may also be of benefit in patients who lost  $\geq 15$  letters of best-corrected visual acuity (BCVA) in the first year of treatment.

Statistically significant patient-reported visual functioning benefits were observed in both MARINA and ANCHOR with ranibizumab treatment over the control group as measured by the NEI VFQ-25.

In study FVF3192g (PIER), 184 patients with all forms of neovascular AMD were randomised in a 1:1:1 ratio to receive ranibizumab 0.3 mg, ranibizumab 0.5 mg or sham injections once a month for 3 consecutive doses, followed by a dose administered once every 3 months. From Month 14 of the study, sham-treated patients were allowed to receive ranibizumab and from Month 19, more frequent treatments were possible. Patients treated with ranibizumab in PIER received a mean of 10 total treatments.

After an initial increase in visual acuity (following monthly dosing), on average, patients' visual acuity declined with quarterly dosing, returning to baseline at Month 12 and this effect was maintained in most ranibizumab-treated patients (82%) at Month 24. Limited data from sham subjects who later received ranibizumab suggested that early initiation of treatment may be associated with better

preservation of visual acuity.

Data from two studies (MONT BLANC, BPD952A2308 and DENALI, BPD952A2309) conducted post approval confirmed the efficacy of ranibizumab but did not demonstrate additional effect of the combined administration of verteporfin (Visudyne PDT) and ranibizumab compared to ranibizumab monotherapy.

Treatment of visual impairment due to CNV secondary to PM

The clinical safety and efficacy of ranibizumab in patients with visual impairment due to CNV in PM have been assessed based on the 12-month data of the double-masked, controlled pivotal study F2301 (RADIANCE). In this study 277 patients were randomised in a 2:2:1 ratio to the following arms:

- Group I (ranibizumab 0.5 mg, dosing regimen driven by “stability” criteria defined as no change in BCVA compared to two preceding monthly evaluations).
- Group II (ranibizumab 0.5 mg, dosing regimen driven by “disease activity” criteria defined as vision impairment attributable to intra- or subretinal fluid or active leakage due to the CNV lesion as assessed by optical coherence tomography and/or fluorescence angiography).
- Group III (vPDT - patients were allowed to receive ranibizumab treatment as of Month 3).

In Group II, which is the recommended posology (see section 4.2), 50.9% of patients required 1 or 2 injections, 34.5% required 3 to 5 injections and 14.7% required 6 to 12 injections over the 12-month study period. 62.9% of Group II patients did not require injections in the second 6 months of the study.

The key outcomes from RADIANCE are summarised in Table 2 and Figure 2.


**Table 2 Outcomes at Month 3 and 12 (RADIANCE)**

	<b>Group I Ranibizumab 0.5 mg “Vision stability” (n = 105)</b>	<b>Group II Ranibizumab 0.5 mg “Disease activity” (n = 116)</b>	<b>Group III vPDT<sup>b</sup> (n = 55)</b>
<b>Month 3</b>			
Mean average BCVA change from Month 1 to Month 3 compared to baseline <sup>a</sup> (letters)	+10.5	+10.6	+2.2
Proportion of patients who gained: ≥ 15 letters, or reached ≥ 84 letters in BCVA	38.1%	43.1%	14.5%
<b>Month 12</b>			
Number of injections up to Month 12:			
Mean	4.6	3.5	N/A
Median	4.0	2.5	N/A
Mean average BCVA change from Month 1 to Month 12 compared to baseline (letters)	+12.8	+12.5	N/A
Proportion of patients who gained: ≥ 15 letters, or reached ≥ 84 letters in BCVA	53.3%	51.7%	N/A

<sup>a</sup> p< 0.00001 comparison with vPDT control

<sup>b</sup> Comparative control up to Month 3. Patients randomised to vPDT were allowed to receive ranibizumab treatment as of Month 3 (in Group III, 38 patients received ranibizumab as of Month 3)

**Figure 2 Mean change from baseline BCVA over time to Month 12 (RADIANCE)**



The improvement of vision was accompanied by a reduction in central retinal thickness.

Patient-reported benefits were observed with ranibizumab treatment arms over vPDT (p-value < 0.05) in terms of improvement in the composite score and several subscales (general vision, near activities, mental health and dependency) of the NEI VFQ-25.

Treatment of visual impairment due to CNV (other than secondary to PM and wet AMD)

The clinical safety and efficacy of ranibizumab in patients with visual impairment due to CNV have been assessed based on the 12-month data of the double-masked, sham-controlled pivotal study G2301 (MINERVA). In this study 178 adult patients were randomised in a 2:1 ratio to receive:

- ranibizumab 0.5 mg at baseline, followed by an individualised dosing regimen driven by disease activity as assessed by visual acuity and/or anatomical parameters (e.g. visual acuity impairment, intra/sub-retinal fluid, haemorrhage or leakage);
- sham injection at baseline, followed by an individualised treatment regimen driven by disease activity.

At Month 2, all patients received open-label treatment with ranibizumab as needed.

Key outcome measures from MINERVA are summarised in Table 3 and Figure 3. An improvement of vision was observed and was accompanied by a reduction in central subfield thickness over the 12-month period.

The mean number of injections given over 12 months was 5.8 in the ranibizumab arm versus 5.4 in

those patients in the sham arm who were eligible to receive ranibizumab from Month 2 onwards. In the sham arm 7 out of 59 patients did not receive any treatment with ranibizumab in the study eye during the 12-month period.


**Table 3 Outcomes at Month 2 (MINERVA)**

	Ranibizumab 0.5 mg (n = 119)	Sham (n = 59)
Mean BCVA change from baseline to Month 2 <sup>a</sup>	9.5 letters	-0.4 letters
Patients gaining ≥ 15 letters from baseline or reaching 84 letters at Month 2	31.4%	12.3%
Patients not losing > 15 letters from baseline at Month 2	99.2%	94.7%
Reduction in CSFT <sup>b</sup> from baseline to Month 2 <sup>a</sup>	77 μm	-9.8 μm

<sup>a</sup> One-sided p< 0.001 comparison with sham control

<sup>b</sup> CSFT - central retinal subfield thickness

**Figure 3 Mean change from baseline BCVA over time to Month 12 (MINERVA)**



When comparing ranibizumab versus sham control at Month 2, a consistent treatment effect both overall and across baseline aetiology subgroups was observed:

**Table 4 Treatment effect overall and across baseline aetiology subgroups**

Overall and per baseline aetiology	Treatment effect over sham [letters]	Patient numbers [n] (treatment +sham)
Overall	9.9	178
Angioid streaks	14.6	27
Post-inflammatory retinochoroidopathy	6.5	28
Central serous chorioretinopathy	5.0	23
Idiopathic chorioretinopathy	11.4	63
Miscellaneous aetiologies <sup>a</sup>	10.6	37

<sup>a</sup> encompasses different aetiologies of low frequency of occurrence not included in the other subgroups

In the pivotal study G2301 (MINERVA), five adolescent patients aged 12 to 17 years with visual impairment secondary to CNV received open-label treatment with ranibizumab 0.5 mg at baseline followed by an individualised treatment regimen as for the adult population. BCVA improved from baseline to Month 12 in all five patients, ranging from 5 to 38 letters (mean of 16.6 letters). The improvement of vision was accompanied by a stabilisation or reduction in central subfield thickness over the 12-month period. The mean number of ranibizumab injections given in the study eye over 12 months was 3 (ranged from 2 to 5). Overall, ranibizumab treatment was well tolerated.

#### Treatment of visual impairment due to DME


The efficacy and safety of ranibizumab have been assessed in three randomised, controlled studies of at least 12 months duration. A total of 868 patients (708 active and 160 control) were enrolled in these studies.

In the phase II study D2201 (RESOLVE), 151 patients were treated with ranibizumab (6 mg/mL, n = 51, 10 mg/mL, n = 51) or sham (n = 49) by monthly intravitreal injections. The mean average change in BCVA from Month 1 to Month 12 compared to baseline was +7.8 ( $\pm 7.72$ ) letters in the pooled ranibizumab-treated patients (n = 102), compared to -0.1 ( $\pm 9.77$ ) letters for sham-treated patients; and the mean change in BCVA at Month 12 from baseline was 10.3 ( $\pm 9.1$ ) letters compared to -1.4 ( $\pm 14.2$ ) letters, respectively ( $p < 0.0001$  for the treatment difference).

In the phase III study D2301 (RESTORE), 345 patients were randomised in a 1:1:1 ratio to receive ranibizumab 0.5 mg monotherapy and sham laser photocoagulation, combined ranibizumab 0.5 mg and laser photocoagulation or sham injection and laser photocoagulation. 240 patients, who had previously completed the 12-month RESTORE study, were enrolled in the open-label, multicentre 24-month extension (RESTORE Extension) study. Patients were treated with ranibizumab 0.5 mg *pro re nata* (PRN) in the same eye as the core study (D2301 RESTORE).

Key outcome measures are summarised in Table 5 (RESTORE and Extension) and Figure 4 (RESTORE).

#### **Figure 4 Mean change in visual acuity from baseline over time in study D2301 (RESTORE)**



BL=baseline; SE=standard error of mean

\* Difference in least square means,  $p < 0.0001/0.0004$  based on two-sided stratified Cochran-Mantel-Haenszel test

The effect at 12 months was consistent in most subgroups. However, subjects with a baseline BCVA  $> 73$  letters and macular oedema with central retinal thickness  $< 300 \mu\text{m}$  did not appear to benefit from treatment with ranibizumab compared to laser photocoagulation.

**Table 5 Outcomes at Month 12 in study D2301 (RESTORE) and at Month 36 in study D2301-E1 (RESTORE Extension)**

Outcome measures at Month 12 compared to baseline in study D2301 (RESTORE)	Ranibizumab 0.5 mg n = 115	Ranibizumab 0.5 mg + Laser n = 118	Laser n = 110
Mean average change in BCVA from Month 1 to Month 12 <sup>a</sup> ( $\pm$ SD)	6.1 (6.4) <sup>a</sup>	5.9 (7.9) <sup>a</sup>	0.8 (8.6)
Mean change in BCVA at Month 12 ( $\pm$ SD)	6.8 (8.3) <sup>a</sup>	6.4 (11.8) <sup>a</sup>	0.9 (11.4)
Gain of $\geq 15$ letters or BCVA $\geq 84$ letters at Month 12 (%)	22.6	22.9	8.2
Mean number of injections (Months 0-11)	7.0	6.8	7.3 (sham)
Outcome measure at Month 36 compared to D2301 (RESTORE) baseline in study D2301-E1 (RESTORE Extension)	Prior ranibizumab 0.5 mg n = 83	Prior ranibizumab 0.5 mg + laser n = 83	Prior laser n = 74
Mean change in BCVA at Month 24 (SD)	7.9 (9.0)	6.7 (7.9)	5.4 (9.0)
Mean change in BCVA at Month 36 (SD)	8.0 (10.1)	6.7 (9.6)	6.0 (9.4)
Gain of $\geq 15$ letters or BCVA $\geq 84$ letters at Month 36 (%)	27.7	30.1	21.6
Mean number of injections (Months 12-35)*	6.8	6.0	6.5

<sup>a</sup> $p < 0.0001$  for comparisons of ranibizumab arms vs. laser arm.

n in D2301-E1 (RESTORE Extension) is the number of patients with a value at both D2301

(RESTORE) baseline (Month 0) and at the Month 36 visit.

\* The proportion of patients who did not require any ranibizumab treatment during the extension phase was 19%, 25% and 20% in the prior ranibizumab, prior ranibizumab + laser and prior laser groups, respectively.

Statistically significant patient-reported benefits for most vision-related functions were observed with ranibizumab (with or without laser) treatment over the control group as measured by the NEI VFQ-25. For other subscales of this questionnaire no treatment differences could be established.

The long-term safety profile of ranibizumab observed in the 24-month extension study is consistent with the known ranibizumab safety profile.

In the phase IIIb study D2304 (RETAIN), 372 patients were randomised in 1:1:1 ratio to receive:

- ranibizumab 0.5 mg with concomitant laser photocoagulation on a treat-and-extend (TE) regimen,
- ranibizumab 0.5 mg monotherapy on a TE regimen,
- ranibizumab 0.5 mg monotherapy on a PRN regimen.

In all groups, ranibizumab was administered monthly until BCVA was stable for at least three consecutive monthly assessments. On TE, ranibizumab was administered at treatment intervals of 2-3 months. In all groups, monthly treatment was re-initiated upon a decrease in BCVA due to DME progression and continued until stable BCVA was reached again.

The number of scheduled treatment visits after the initial 3 injections, was 13 and 20 for the TE and PRN regimens, respectively. With both TE regimens, more than 70% of patients maintained their BCVA with an average visit frequency of  $\geq 2$  months.

The key outcome measures are summarised in Table 6.

**Table 6 Outcomes in study D2304 (RETAIN)**

Outcome measure compared to baseline	TE ranibizumab 0.5 mg + laser n = 117	TE ranibizumab 0.5 mg alone n = 125	PRN ranibizumab 0.5 mg n = 117
Mean average change in BCVA from Month 1 to Month 12 (SD)	5.9 (5.5) <sup>a</sup>	6.1 (5.7) <sup>a</sup>	6.2 (6.0)
Mean average change in BCVA from Month 1 to Month 24 (SD)	6.8 (6.0)	6.6 (7.1)	7.0 (6.4)
Mean change in BCVA at Month 24 (SD)	8.3 (8.1)	6.5 (10.9)	8.1 (8.5)
Gain of $\geq 15$ letters or BCVA $\geq 84$ letters at Month 24(%)	25.6	28.0	30.8
Mean number of injections (months 0-23)	12.4	12.8	10.7

<sup>a</sup>p < 0.0001 for assessment of non-inferiority to PRN

In DME studies, the improvement in BCVA was accompanied by a reduction over time in mean CSFT in all the treatment groups.

#### Treatment of PDR

The clinical safety and efficacy of ranibizumab in patients with PDR have been assessed in Protocol S which evaluated the treatment with ranibizumab 0.5 mg intravitreal injections compared with panretinal photocoagulation (PRP). The primary endpoint was the mean visual acuity change at year 2.

Additionally, change in diabetic retinopathy (DR) severity was assessed based on fundus photographs using the DR severity score (DRSS).

Protocol S was a multicentre, randomised, active-controlled, parallel-assignment, non-inferiority phase III study in which 305 patients (394 study eyes) with PDR with or without DME at baseline were enrolled. The study compared ranibizumab 0.5 mg intravitreal injections to standard treatment with PRP. A total of 191 eyes (48.5%) were randomised to ranibizumab 0.5 mg and 203 eyes (51.5%) eyes were randomised to PRP. A total of 88 eyes (22.3%) had baseline DME: 42 (22.0%) and 46 (22.7%) eyes in the ranibizumab and PRP groups, respectively.

In this study, the mean visual acuity change at year 2 was +2.7 letters in the ranibizumab group compared to -0.7 letters in the PRP group. The difference in least square means was 3.5 letters (95% CI: [0.2 to 6.7]).

At year 1, 41.8% of eyes experienced a  $\geq 2$ -step improvement in the DRSS when treated with ranibizumab (n = 189) compared to 14.6% of eyes treated with PRP (n = 199). The estimated difference between ranibizumab and laser was 27.4% (95% CI: [18.9, 35.9]).

**Table 7 DRSS improvement or worsening of  $\geq 2$  or  $\geq 3$  steps at year 1 in Protocol S (LOCF Method)**

Categorised change from baseline	Protocol S		
	Ranibizumab 0.5 mg (N = 189)	PRP (N = 199)	Difference in proportion (%), CI
$\geq 2$ -step improvement			
n (%)	79 (41.8%)	29 (14.6%)	27.4 (18.9, 35.9)
$\geq 3$ -step improvement			
n (%)	54 (28.6%)	6 (3.0%)	25.7 (18.9, 32.6)
$\geq 2$ -step worsening			
n (%)	3 (1.6%)	23 (11.6%)	-9.9 (-14.7, -5.2)
$\geq 3$ -step worsening			
n (%)	1 (0.5%)	8 (4.0%)	-3.4 (-6.3, -0.5)
DRSS = diabetic retinopathy severity score, n = number of patients who satisfied the condition at the visit, N = total number of study eyes.			

At year 1 in the ranibizumab-treated group in Protocol S,  $\geq 2$ -step improvement in DRSS was consistent in eyes without DME (39.9%) and with baseline DME (48.8%).

An analysis of year 2 data from Protocol S demonstrated that 42.3% (n = 80) of eyes in the ranibizumab-treated group had  $\geq 2$ -step improvement in DRSS from baseline compared with 23.1% (n = 46) of eyes in the PRP group. In the ranibizumab-treated group,  $\geq 2$ -step improvement in DRSS from baseline was observed in 58.5% (n = 24) of eyes with baseline DME and 37.8% (n = 56) of eyes without DME.

DRSS was also assessed in three separate active-controlled phase III DME studies (ranibizumab 0.5 mg PRN vs laser) that included a total of 875 patients, of whom approximately 75% were of Asian origin. In a meta-analysis of these studies, 48.4% of the 315 patients with gradable DRSS scores in the subgroup of patients with moderately severe non-proliferative DR (NPDR) or worse at baseline experienced a  $\geq 2$ -step improvement in the DRSS at Month 12 when treated with ranibizumab (n = 192) vs 14.6% of patients treated with laser (n = 123). The estimated difference between ranibizumab and laser was 29.9% (95% CI: [20.0, 39.7]). In the 405 DRSS gradable patients with

moderate NPDR or better, a  $\geq 2$ - step DRSS improvement was observed in 1.4% and 0.9% of the ranibizumab and laser groups, respectively.

Treatment of visual impairment due to macular oedema secondary to RVO

The clinical safety and efficacy of ranibizumab in patients with visual impairment due to macular oedema secondary to RVO have been assessed in the randomised, double-masked, controlled studies BRAVO and CRUISE that recruited subjects with BRVO (n = 397) and CRVO (n = 392), respectively. In both studies, subjects received either 0.3 mg or 0.5 mg ranibizumab or sham injections. After 6 months, patients in the sham-control arms switched to 0.5 mg ranibizumab.


Key outcome measures from BRAVO and CRUISE are summarised in Table 8 and Figures 5 and 6.

**Table 8 Outcomes at Month 6 and 12 (BRAVO and CRUISE)**

	BRAVO		CRUISE	
	Sham/ Ranibizumab 0.5 mg (n = 132)	Ranibizumab 0.5 mg (n = 131)	Sham/ Ranibizumab 0.5 mg (n = 130)	Ranibizumab 0.5 mg (n = 130)
Mean change in visual acuity at Month 6 <sup>a</sup> (letters) (SD) (primary endpoint)	7.3 (13.0)	18.3 (13.2)	0.8 (16.2)	14.9 (13.2)
Mean change in BCVA at Month 12 (letters) (SD)	12.1 (14.4)	18.3 (14.6)	7.3 (15.9)	13.9 (14.2)
Gain of $\geq 15$ letters in visual acuity at Month 6 <sup>a</sup> (%)	28.8	61.1	16.9	47.7
Gain of $\geq 15$ letters in visual acuity at Month 12 (%)	43.9	60.3	33.1	50.8
Proportion (%) receiving laser rescue over 12 months	61.4	34.4	NA	NA


<sup>a</sup>p < 0.0001 for both studies

**Figure 5 Mean change from baseline BCVA over time to Month 6 and Month 12 (BRAVO)**



BL=baseline; SE=standard error of mean

Figure 6 Mean change from baseline BCVA over time to Month 6 and Month 12 (CRUISE)



BL=baseline; SE=standard error of mean

In both studies, the improvement of vision was accompanied by a continuous and significant reduction

in the macular oedema as measured by central retinal thickness.

In patients with CRVO (CRUISE and extension study HORIZON): Subjects treated with sham in the first 6 months who subsequently received ranibizumab did not achieve comparable gains in visual acuity by Month 24 (~6 letters) compared to subjects treated with ranibizumab from study start (~12 letters).

Statistically significant patient-reported benefits in subscales related to near and distance activity were observed with ranibizumab treatment over the control group as measured by the NEI VFQ-25.

The long-term (24 months) clinical safety and efficacy of ranibizumab in patients with visual impairment due to macular oedema secondary to RVO were assessed in the BRIGHTER (BRVO) and CRYSTAL (CRVO) studies. In both studies, subjects received a 0.5 mg ranibizumab PRN dosing regimen driven by individualised stabilisation criteria. BRIGHTER was a 3-arm randomised active-controlled study that compared 0.5 mg ranibizumab given as monotherapy or in combination with adjunctive laser photocoagulation to laser photocoagulation alone. After 6 months, subjects in the laser arm could receive 0.5 mg ranibizumab. CRYSTAL was a single-arm study with 0.5 mg ranibizumab monotherapy.

Key outcome measures from BRIGHTER and CRYSTAL are shown in Table 9.

**Table 9 Outcomes at Months 6 and 24 (BRIGHTER and CRYSTAL)**

	<b>BRIGHTER</b>			<b>CRYSTAL</b>
	Ranibizumab 0.5 mg N = 180	Ranibizumab 0.5 mg + Laser N = 178	Laser* N = 90	Ranibizumab 0.5 mg N = 356
Mean change in BCVA at Month 6 <sup>a</sup> (letters) (SD)	+14.8 (10.7)	+14.8 (11.13)	+6.0 (14.27)	+12.0 (13.95)
Mean change in BCVA at Month 24 <sup>b</sup> (letters) (SD)	+15.5 (13.91)	+17.3 (12.61)	+11.6 (16.09)	+12.1 (18.60)
Gain of ≥ 15 letters in BCVA at Month 24 (%)	52.8	59.6	43.3	49.2
Mean number of injections (SD) (Months 0-23)	11.4 (5.81)	11.3 (6.02)	NA	13.1 (6.39)
<sup>a</sup>	p< 0.0001 for both comparisons in BRIGHTER at Month 6: Ranibizumab 0.5 mg vs Laser and Ranibizumab 0.5 mg + Laser vs Laser.			
<sup>b</sup>	p< 0.0001 for null hypothesis in CRYSTAL that the mean change at Month 24 from baseline is zero.			
*	Starting at Month 6 ranibizumab 0.5 mg treatment was allowed (24 patients were treated with laser only).			

In BRIGHTER, ranibizumab 0.5 mg with adjunctive laser therapy demonstrated non-inferiority versus ranibizumab monotherapy from baseline to Month 24 (95% CI -2.8, 1.4).

In both studies, a rapid and statistically significant decrease from baseline in central retinal subfield thickness was observed at Month 1. This effect was maintained up to Month 24.

The effect of ranibizumab treatment was similar irrespective of the presence of retinal ischaemia. In BRIGHTER, patients with ischaemia present (N = 46) or absent (N = 133) and treated with ranibizumab monotherapy had a mean change from baseline of +15.3 and +15.6 letters, respectively, at Month 24. In CRYSTAL, patients with ischaemia present (N = 53) or absent (N = 300) and treated

with ranibizumab monotherapy had a mean change from baseline of +15.0 and +11.5 letters, respectively.

The effect in terms of visual improvement was observed in all patients treated with 0.5 mg ranibizumab monotherapy regardless of their disease duration in both BRIGHTER and CRYSTAL. In patients with < 3 months disease duration an increase in visual acuity of 13.3 and 10.0 letters was seen at Month 1; and 17.7 and 13.2 letters at Month 24 in BRIGHTER and CRYSTAL, respectively. The corresponding visual acuity gain in patients with  $\geq$  12 months disease duration was 8.6 and 8.4 letters in the respective studies. Treatment initiation at the time of diagnosis should be considered.

The long-term safety profile of ranibizumab observed in the 24-month studies is consistent with the known ranibizumab safety profile.

### Paediatric population

The safety and efficacy of ranibizumab 0.5 mg in pre-filled syringe have not been studied in paediatric patients.

The European Medicines Agency has waived the obligation to submit the results of studies with the reference medicinal product containing ranibizumab in all subsets of the paediatric population in neovascular AMD, visual impairment due to DME, visual impairment due to macular oedema secondary to RVO, visual impairment due to CNV and diabetic retinopathy (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

Following monthly intravitreal administration of ranibizumab to patients with neovascular AMD, serum concentrations of ranibizumab were generally low, with maximum levels ( $C_{max}$ ) generally below the ranibizumab concentration necessary to inhibit the biological activity of VEGF by 50% (11-27 ng/mL, as assessed in an *in vitro* cellular proliferation assay).  $C_{max}$  was dose proportional over the dose range of 0.05 to 1.0 mg/eye. Serum concentrations in a limited number of DME patients indicate that a slightly higher systemic exposure cannot be excluded compared to those observed in neovascular AMD patients. Serum ranibizumab concentrations in RVO patients were similar or slightly higher compared to those observed in neovascular AMD patients.

Based on analysis of population pharmacokinetics and disappearance of ranibizumab from serum for patients with neovascular AMD treated with the 0.5 mg dose, the average vitreous elimination half-life of ranibizumab is approximately 9 days. Upon monthly intravitreal administration of ranibizumab 0.5 mg/eye, serum ranibizumab  $C_{max}$ , attained approximately 1 day after dosing, is predicted to generally range between 0.79 and 2.90 ng/mL, and  $C_{min}$  is predicted to generally range between 0.07 and 0.49 ng/mL. Serum ranibizumab concentrations are predicted to be approximately 90,000-fold lower than vitreal ranibizumab concentrations.

Patients with renal impairment: No formal studies have been conducted to examine the pharmacokinetics of ranibizumab in patients with renal impairment. In a population pharmacokinetic analysis of neovascular AMD patients, 68% (136 of 200) of patients had renal impairment (46.5% mild [50-80 mL/min], 20% moderate [30-50 mL/min], and 1.5% severe [ $<$ 30 mL/min]). In RVO patients, 48.2% (253 of 525) had renal impairment (36.4% mild, 9.5% moderate and 2.3% severe). Systemic clearance was slightly lower, but this was not clinically significant.

Hepatic impairment: No formal studies have been conducted to examine the pharmacokinetics of ranibizumab in patients with hepatic impairment.

## **5.3 Preclinical safety data**

Bilateral intravitreal administration of ranibizumab to cynomolgus monkeys at doses between 0.25 mg/eye and 2.0 mg/eye once every 2 weeks for up to 26 weeks resulted in dose-dependent ocular

effects.

Intraocularly, there were dose-dependent increases in anterior chamber flare and cells with a peak 2 days after injection. The severity of the inflammatory response generally diminished with subsequent injections or during recovery. In the posterior segment, there were vitreal cell infiltration and floaters, which also tended to be dose-dependent and generally persisted to the end of the treatment period. In the 26-week study, the severity of the vitreous inflammation increased with the number of injections. However, evidence of reversibility was observed after recovery. The nature and timing of the posterior segment inflammation is suggestive of an immune-mediated antibody response, which may be clinically irrelevant. Cataract formation was observed in some animals after a relatively long period of intense inflammation, suggesting that the lens changes were secondary to severe inflammation. A transient increase in post-dose intraocular pressure was observed following intravitreal injections, irrespective of dose.

Microscopic ocular changes were related to inflammation and did not indicate degenerative processes. Granulomatous inflammatory changes were noted in the optic disc of some eyes. These posterior segment changes diminished, and in some instances resolved, during the recovery period.

Following intravitreal administration, no signs of systemic toxicity were detected. Serum and vitreous antibodies to ranibizumab were found in a subset of treated animals.

No carcinogenicity or mutagenicity data are available.

In pregnant monkeys, intravitreal ranibizumab treatment resulting in maximal systemic exposures 0.9-7-fold a worst case clinical exposure did not elicit developmental toxicity or teratogenicity, and had no effect on weight or structure of the placenta, although, based on its pharmacological effect ranibizumab should be regarded as potentially teratogenic and embryo-/foetotoxic.

The absence of ranibizumab-mediated effects on embryo-foetal development is plausibly related mainly to the inability of the Fab fragment to cross the placenta. Nevertheless, a case was described with high maternal ranibizumab serum levels and presence of ranibizumab in foetal serum, suggesting that the anti-ranibizumab antibody acted as (Fc region containing) carrier protein for ranibizumab, thereby decreasing its maternal serum clearance and enabling its placental transfer. As the embryo-foetal development investigations were performed in healthy pregnant animals and disease (such as diabetes) may modify the permeability of the placenta towards a Fab fragment, the study should be interpreted with caution.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

$\alpha,\alpha$ -trehalose dihydrate  
Histidine hydrochloride monohydrate  
Histidine  
Polysorbate 20  
Water for injections

### **6.2 Incompatibilities**

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

12 months

## 6.4 Special precautions for storage

Store in a refrigerator (2 °C - 8 °C). Do not freeze.

Keep the pre-filled syringe in its sealed tray in the carton in order to protect from light.

Prior to use, the unopened tray may be kept at room temperature (25 °C) for up to 24 hours.

## 6.5 Nature and contents of container

0.165 mL sterile solution in a pre-filled syringe (Cyclo olefin polymer) with a chlorobutyl rubber plunger stopper with i-coating™ (silicone resin coating) and a syringe cap consisting of a translucent, tamper-evident rigid seal with a chlorobutyl rubber tip cap including a Luer lock adapter. The pre-filled syringe has a plunger rod and a finger grip, and is packed in a sealed tray.

Pack size of one pre-filled syringe.


## 6.6 Special precautions for disposal and other handling



The pre-filled syringe is for single use only. The pre-filled syringe is sterile. Do not use the product if the packaging is damaged. The sterility of the pre-filled syringe cannot be guaranteed unless the tray remains sealed. Do not use the pre-filled syringe if the solution is discoloured, cloudy or contains particles.


The pre-filled syringe contains more than the recommended dose of 0.5 mg. The extractable volume of the pre-filled syringe (0.1 mL) is not to be used in total. The excess volume should be expelled prior to injection. Injecting the entire volume of the pre-filled syringe could result in overdose. To expel the air bubble along with the excess medicinal product, slowly push the plunger until the edge below the dome of the rubber stopper is aligned with the black dosing line on the syringe (equivalent to 0.05 mL, i.e., 0.5 mg ranibizumab).

For the intravitreal injection, a 30G × ½” sterile injection needle should be used.

To prepare Ranluspec for intravitreal administration, please adhere to the instructions for use.

Introduction	<ol style="list-style-type: none"><li>1. The prefilled syringe is sterile and is for single dose only. Do not use the product if the packaging is damaged or has been tampered with.</li><li>2. To prepare Ranluspec for intravitreal administration, please adhere to these instructions for use. Read all the instructions carefully before using the prefilled syringe.</li><li>3. The opening of the sealed tray and all subsequent steps should be done under aseptic conditions.</li><li>4. For the intravitreal injection, a 30-gauge × ½ inch sterile injection needle should be used. (Not provided).</li></ol> <p><b>Note: The dose must be set to 0.05 mL.</b></p>
Pre-filled syringe description	 <p><b>Figure 1 Ranluspec pre-filled syringe illustration</b></p>

<p><b>Step 1</b></p>	<p><b>PREPARATION:</b>          Make sure that your pack contains a sterile prefilled syringe in a sealed tray.          Peel the lid off the syringe tray and using aseptic technique, remove the syringe.</p>	
<p><b>Step 2</b></p>	<p><b>INSPECT SYRINGE:</b>          1. RANIBIZUMAB should be clear to slightly opalescent, colorless to pale brownish.  <b>2. Do not use the prefilled syringe if:</b>          -The syringe cap is detached from the Luer lock.          - The syringe is damaged.          - Particulates, cloudiness, or discoloration are visible.          3. If any of the above is not true, discard the pre-filled syringe and use a new one.  <b>Note: To avoid compromising the sterility of the product, do not pull back the plunger</b></p>	 <p><b>Figure 2 Ranluspec 0.5 mg prefilled syringe with a White finger grip.</b></p>
<p><b>Step 3</b></p>	<p><b>REMOVE SYRINGE CAP:</b>          Remove the syringe cap by rotating anti-clockwise movement. (See Figure 3).   <b>Note: Syringe cap to be opened once user is ready for the injection. Once syringe cap opened, Do Not recap, Syringe should be disposed if user is not ready for injection.</b></p>	 <p><b>Figure 3 Removal of treaded closure cap</b></p>
<p><b>Step 4</b></p>	<p><b>ATTACH NEEDLE:</b>          1. Attach a 30G × ½ inch sterile injection needle firmly onto the syringe by screwing it tightly onto the Luer lock (See Figure 4)          2. Carefully remove the needle cap by pulling it straight off.   <b>Note: Do not wipe the needle at any time.</b></p>	 <p><b>Figure 4 Needle attachment</b></p>
<p><b>Step 5</b></p>	<p><b>DISLodge AIR BUBBLES</b>          1. Hold the syringe with the needle pointing up.           2. If there are any air bubbles, gently tap the syringe with your finger until the bubbles rise to the top (See Figure 5).</p>	 <p><b>Figure 5 Removal of air bubble</b></p>

<p><b>Step 6</b></p>	<p><b>EXPEL AIR AND ADJUST DRUG DOSE.</b></p> <p>1. Hold the syringe at eye level, and carefully push the plunger rod until the edge below the dome of the rubber stopper is aligned with the 0.05 mL dose mark (See Figure 6).</p> <p><b>Note: The plunger rod is <u>attached</u> to the rubber stopper, to avoid air entrapment do not pull back the plunger rod.</b></p>	 <p><b>Figure 6 Dose adjustment</b></p>
<p><b>Step 7</b></p>	<p><b>INJECT:</b></p> <ol style="list-style-type: none"> <li>1. The injection procedure should be carried out under aseptic conditions.</li> <li>2. The injection needle should be inserted 3.5-4.0 mm posterior to the limbus into vitreous cavity, avoiding the horizontal meridian and aiming towards the centre of the globe.</li> <li>3. Inject slowly until rubber stopper reaches the bottom of the syringe to deliver the volume of 0.05 mL.</li> <li>4. After injection, <b>do not</b> re-cap the needle or detach it from the syringe. dispose of the used syringe together with the needle in a sharp disposal container or in accordance with local requirements.</li> </ol>	

## 7. MARKETING AUTHORISATION HOLDER

Lupin Europe GmbH  
Hanauer Landstraße 139-143  
60314 Frankfurt  
Germany

## 8. MARKETING AUTHORISATION NUMBER(S)

EU/1/25/2012/002

## 9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation:  
Date of latest renewal:

## 10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <https://www.ema.europa.eu>

## **ANNEX II**

- A. MANUFACTURER OF THE BIOLOGICAL ACTIVE  
SUBSTANCE AND MANUFACTURER RESPONSIBLE FOR  
BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY  
AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE  
MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO  
THE SAFE AND EFFECTIVE USE OF THE MEDICINAL  
PRODUCT**

**A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer of the biological active substance

Lupin Limited (Biotech Division)  
Gat No.-1156, Village: Ghotawade,  
Taluka Mulshi, District: Pune - 412 115,  
Maharashtra  
India

Name and address of the manufacturer responsible for batch release

Hormosan Pharma GmbH  
Hanauer Landstraße 139-143  
60314 Frankfurt  
Germany

**B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

**C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

• **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

**D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

• **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

• **Additional risk minimisation measures**

Prior to launch in each Member State the MAH shall agree the final educational material with the National Competent Authority.

The MAH shall ensure that, following discussions and agreements with the National Competent Authorities in each Member State where Ranluspec is marketed, at launch and after launch all

ophthalmological clinics where Ranluspec is expected to be used are provided with an up-to-date patient information pack.

The patient information pack should be provided in both the form of patient information booklets and an audio-CD that contain following key elements:

- Patient information leaflet
- How to prepare for Ranluspec treatment
- What are the steps following treatment with Ranluspec
- Key signs and symptoms of serious adverse events including increased intraocular pressure, intraocular inflammation, retinal detachment & retinal tear and infectious endophthalmitis
- When to seek urgent attention from the health care provider.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON VIAL**

**1. NAME OF THE MEDICINAL PRODUCT**

Ranluspec 10 mg/mL solution for injection  
ranibizumab

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

One mL contains 10 mg of ranibizumab. Vial containing 2.3 mg of ranibizumab.

**3. LIST OF EXCIPIENTS**

Excipients:  $\alpha,\alpha$ -trehalose dihydrate; histidine hydrochloride, monohydrate; histidine; polysorbate 20; water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Solution for injection

1 × 0.23 mL vial  
Single dose for adults: 0.5 mg/0.05 mL. Excess volume to be expelled.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Intravitreal use.  
Vial for single use only.  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator (2 °C – 8 °C).  
Do not freeze.

Keep the vial in the outer carton in order to protect from light.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Lupin Europe GmbH  
Hanauer Landstraße 139-143  
60314 Frankfurt  
Germany

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/25/2012/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN



**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**LABEL VIAL**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Ranluspec 10 mg/mL solution for injection  
ranibizumab  
Intravitreal use

**2. METHOD OF ADMINISTRATION**

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

2.3 mg/0.23 mL

**6. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON PRE-FILLED SYRINGE**

**1. NAME OF THE MEDICINAL PRODUCT**

Ranluspec 10 mg/mL solution for injection in pre-filled syringe  
ranibizumab

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each pre-filled syringe of 0.165 mL solution contains 1.65 mg of ranibizumab (10 mg/mL).

**3. LIST OF EXCIPIENTS**

Excipients:  $\alpha,\alpha$ -trehalose dihydrate; histidine hydrochloride, monohydrate; histidine; polysorbate 20; water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Solution for injection

1 pre-filled syringe of 0.165 mL.  
Single dose of 0.5 mg/0.05 mL.  
Excess volume should be expelled prior to injection.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For single use only. Upon opening of the sealed tray, proceed under aseptic conditions.  
Set dose to 0.05 mL dose mark.  
Read the package leaflet before use.  
Intravitreal use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator (2 °C – 8 °C).

Do not freeze.

Keep the pre-filled syringe in its sealed tray in the carton to protect from light.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Lupin Europe GmbH  
Hanauer Landstraße 139-143  
60314 Frankfurt  
Germany

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/25/2012/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS  
BLISTER FOIL PRE-FILLED SYRINGE**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Ranluspec 10 mg/mL solution for injection in pre-filled syringe  
ranibizumab  
Intravitreal use

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Lupin Europe GmbH

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

0.165 mL

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS  
LABEL PRE-FILLED SYRINGE**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF ADMINISTRATION**

Ranluspec 10 mg/mL solution for injection in pre-filled syringe  
ranibizumab  
Intravitreal use

**2. METHOD OF ADMINISTRATION**

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

0.165 mL

**B. PACKAGE LEAFLET**

## Package leaflet: Information for the patient

Ranluspec 10 mg/mL solution for injection  
ranibizumab

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

**Read all of this leaflet carefully before you are given this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What Ranluspec is and what it is used for
2. What you need to know before you are given Ranluspec
3. How Ranluspec is given
4. Possible side effects
5. How to store Ranluspec
6. Contents of the pack and other information

## 1. What Ranluspec is and what it is used for

### What Ranluspec is

Ranluspec is a solution which is injected into the eye. Ranluspec belongs to a group of medicines called antineovascularisation agents. It contains the active substance called ranibizumab.

### What Ranluspec is used for

Ranluspec is used in adults to treat several eye diseases causing vision impairment.

These diseases result from damage to the retina (light-sensitive layer at the back of the eye) caused by:

- Growth of leaky, abnormal blood vessels. This is observed in diseases such as age-related macular degeneration (AMD) and proliferative diabetic retinopathy (PDR, a disease caused by diabetes). It may also be associated with choroidal neovascularisation (CNV) due to pathologic myopia (PM), angioid streaks, central serous chorioretinopathy or inflammatory CNV.
- Macular oedema (swelling of the centre of the retina). This swelling can be caused by diabetes (a disease called diabetic macular oedema (DME)) or by the blockage of retinal veins of the retina (a disease called retinal vein occlusion (RVO)).

### How Ranluspec works

Ranluspec specifically recognises and binds to a protein called human vascular endothelial growth factor A (VEGF-A) present in the eye. In excess, VEGF-A causes abnormal blood vessel growth and swelling in the eye which can lead to impairment of vision in diseases like AMD, DME, PDR, RVO, PM and CNV. By binding to VEGF-A, Ranluspec can block its actions and prevent this abnormal growth and swelling.

In these diseases, Ranluspec can help to stabilise and in many cases improve your vision

## 2. What you need to know before you are given Ranluspec

### You must not receive Ranluspec

- If you are allergic to ranibizumab or any of the other ingredients of this medicine (listed in section 6).
- If you have an infection in or around your eye.
- If you have pain or redness (severe intraocular inflammation) in your eye.

### **Warnings and precautions**

Talk to your doctor before you are given Ranluspec.

- Ranluspec is given as an injection into the eye. Occasionally, an infection in the internal portion of the eye, pain or redness (inflammation), detachment or tear of one of the layers in the back of the eye (retinal detachment or tear and retinal pigment epithelial detachment or tear), or clouding of the lens (cataract) may occur after Ranluspec treatment. It is important to identify and treat such an infection or retinal detachment as soon as possible. Please tell your doctor immediately if you develop signs such as eye pain or increased discomfort, worsening eye redness, blurred or decreased vision, an increased number of small particles in your vision or increased sensitivity to light.
- In some patients the eye pressure may increase for a short period directly after the injection. This is something you may not notice, therefore your doctor may monitor this after each injection.
- Inform your doctor if you have a prior history of eye conditions or eye treatments, or if you have had a stroke or experienced transient signs of stroke (weakness or paralysis of limbs or face, difficulty speaking or understanding). This information will be taken into account to evaluate if Ranluspec is the appropriate treatment for you.

Please see section 4 (“Possible side effects”) for more detailed information on side effects that could occur during Ranluspec therapy.

### **Children and adolescents (below 18 years of age)**

The use of Ranluspec in children and adolescents has not been established and is therefore not recommended.

### **Other medicines and Ranluspec**

Tell your doctor if you are using, have recently used or might use any other medicines.

### **Pregnancy and breast-feeding**

- Women who could become pregnant must use effective contraception during treatment and for at least three further months after the last injection of Ranluspec.
- There is no experience of using Ranluspec in pregnant women. Ranluspec should not be used during pregnancy unless the potential benefit outweighs the potential risk to the unborn child. If you are pregnant, think you may be pregnant or planning to become pregnant, discuss this with your doctor before treatment with Ranluspec.
- Small amounts of Ranluspec may pass into breast milk, therefore Ranluspec is not recommended during breast-feeding. Ask your doctor or pharmacist for advice before Ranluspec treatment.

### **Driving and using machines**

After Ranluspec treatment you may experience some temporary vision blurring. If this happens, do not drive or use machines until this resolves.

### **This medicine contains polysorbate 20**

This medicine contains 0.005 mg of polysorbate 20 in each dose of 0.05 mL which is equivalent to 0.1 mg/mL. Polysorbates may cause allergic reactions. Tell your doctor if you have any known allergies.

## **3. How Ranluspec is given**

Ranluspec is administered as a single injection into your eye by your eye doctor under a local

anaesthetic. The usual dose of an injection is 0.05 mL (which contains 0.5 mg of ranibizumab). The interval between two doses injected into the same eye should be at least four weeks. All injections will be administered by your eye doctor.

Before the injection, your doctor will wash your eye carefully to prevent infection. Your doctor will also give you a local anaesthetic to reduce or prevent any pain you might have with the injection.

The treatment is started with one injection of Ranluspec per month. Your doctor will monitor the condition of your eye and, depending on how you respond to the treatment, will decide if and when you need to receive further treatment.

Detailed instructions for use are given at the end of the leaflet under “How to prepare and administer Ranluspec to adults”.

#### **Elderly (age 65 years and over)**

Ranluspec can be used for people of 65 years of age and over without dose adjustment.

#### **Before stopping Ranluspec treatment**

If you are considering stopping Ranluspec treatment, please go to your next appointment and discuss this with your doctor. Your doctor will advise you and decide how long you should be treated with Ranluspec.

If you have any further questions on the use of this medicine, ask your doctor.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The side effects associated with the administration of Ranluspec are either due to the medicine itself or the injection procedure and mostly affect the eye.

The most serious side effects are described below:

**Common serious side effects** (may affect up to 1 in 10 people): Detachment or tear of the layer in the back of the eye (retinal detachment or tear), resulting in flashes of light with floaters progressing to a temporary loss of sight, or a clouding of the lens (cataract).

**Uncommon serious side effects** (may affect up to 1 in 100 people): Blindness, infection of the eyeball (endophthalmitis) with inflammation of the inside of the eye.

The symptoms you might experience are pain or increased discomfort in your eye, worsening eye redness, blurred or decreased vision, an increased number of small particles in your vision or increased sensitivity to light. **Please tell your doctor immediately if you develop any of these side effects.**

The most frequently reported side effects are described below:

**Very common side effects** (may affect more than 1 in 10 people)

Visual side effects include: Inflammation of the eye, bleeding in the back of the eye (retinal bleeding), visual disturbances, eye pain, small particles or spots in your vision (floaters), bloodshot eye, eye irritation, a feeling of having something in the eye, increased tear production, inflammation or infection of the eyelid margins, dry eye, redness or itching of the eye and increased eye pressure.

Non-visual side effects include: Sore throat, nasal congestion, runny nose, headache and joint pain.

Other side effects which may occur following Ranluspec treatment are described below:

#### **Common side effects**

Visual side effects include: Decreased sharpness of vision, swelling of a section of the eye (uvea, cornea), inflammation of the cornea (front part of eye), small marks on the surface of the eye, blurred vision, bleeding at the site of injection, bleeding in the eye, discharge from the eye with itching, redness and swelling (conjunctivitis), light sensitivity, eye discomfort, swelling of the eyelid, eyelid

pain.

Non-visual side effects include: Urinary tract infection, low red blood cells count (with symptoms such as tiredness, breathlessness, dizziness, pale skin), anxiety, cough, nausea, allergic reactions like rash, hives, itching and skin reddening.

### **Uncommon side effects**

Visual side effects include: Inflammation and bleeding in the front part of the eye, sac of pus on the eye, changes of the central part of the eye surface, pain or irritation at the site of injection, abnormal sensation in the eye, irritation of the eyelid.

### **Reporting of side effects**

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly to the national reporting system listed in [Appendix V](#). By reporting side effects, you can help provide more information on the safety of this medicine.

## **5. How to store Ranluspec**

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and vial label after EXP. The expiry date refers to the last day of that month.
- Store in a refrigerator (2 °C – 8 °C). Do not freeze.
- Prior to use, the unopened vial may be kept at room temperature (25 °C) for up to 24 hours.
- Keep the vial in the outer carton in order to protect from light.
- Do not use any pack that is damaged.

## **6. Contents of the pack and other information**

### **What Ranluspec contains**

- The active substance is ranibizumab. Each mL contains 10 mg ranibizumab. Each vial contains 2.3 mg ranibizumab in 0.23 mL solution. This provides a suitable amount to deliver a single dose of 0.05 mL containing 0.5 mg ranibizumab.
- The other ingredients are  $\alpha,\alpha$ -trehalose dihydrate; histidine hydrochloride, monohydrate; histidine; polysorbate 20; water for injections.

### **What Ranluspec looks like and contents of the pack**

Ranluspec is a solution for injection in a vial (0.23 mL). The solution is clear to slightly opalescent, colourless to pale brownish and aqueous.

Ranluspec is available as a pack containing one glass vial of ranibizumab with chlorobutyl rubber stopper. The vial is for single use only.

### **Marketing Authorisation Holder**

Lupin Europe GmbH  
Hanauer Landstraße 139-143  
60314 Frankfurt  
Germany

### **Manufacturer**

Hormosan Pharma GmbH  
Hanauer Landstraße 139-143  
60314 Frankfurt  
Germany

**This leaflet was last revised in**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<https://www.ema.europa.eu>.

## **THE FOLLOWING INFORMATION IS INTENDED FOR HEALTHCARE PROFESSIONALS ONLY:**

Please also refer to section 3 “How Ranluspec is given”.

### **How to prepare and administer Ranluspec**

Single-use vial for intravitreal use only

Ranluspec must be administered by a qualified ophthalmologist experienced in intravitreal injections.

In wet AMD, in CNV, in PDR and in visual impairment due to DME or to macular oedema secondary to RVO the recommended dose for Ranluspec is 0.5 mg given as a single intravitreal injection. This corresponds to an injection volume of 0.05 mL. The interval between two doses injected into the same eye should be at least four weeks.

Treatment is initiated with one injection per month until maximum visual acuity is achieved and/or there are no signs of disease activity i.e. no change in visual acuity and in other signs and symptoms of the disease under continued treatment. In patients with wet AMD, DME, PDR and RVO, initially, three or more consecutive, monthly injections may be needed.

Thereafter, monitoring and treatment intervals should be determined by the physician and should be based on disease activity, as assessed by visual acuity and/or anatomical parameters.

If, in the physician’s opinion, visual and anatomical parameters indicate that the patient is not benefiting from continued treatment, Ranluspec should be discontinued.

Monitoring for disease activity may include clinical examination, functional testing or imaging techniques (e.g. optical coherence tomography or fluorescein angiography).

If patients are being treated according to a treat-and-extend regimen, once maximum visual acuity is achieved and/or there are no signs of disease activity, the treatment intervals can be extended stepwise until signs of disease activity or visual impairment recur. The treatment interval should be extended by no more than two weeks at a time for wet AMD and may be extended by up to one month at a time for DME. For PDR and RVO, treatment intervals may also be gradually extended, however there are insufficient data to conclude on the length of these intervals. If disease activity recurs, the treatment interval should be shortened accordingly.

The treatment of visual impairment due to CNV should be determined individually per patient based on disease activity. Some patients may only need one injection during the first 12 months; others may need more frequent treatment, including a monthly injection. For CNV secondary to pathologic myopia (PM), many patients may only need one or two injections during the first year.

#### *Ranibizumab and laser photocoagulation in DME and macular oedema secondary to BRVO*

There is some experience of ranibizumab administered concomitantly with laser photocoagulation. When given on the same day, ranibizumab should be administered at least 30 minutes after laser photocoagulation. Ranibizumab can be administered in patients who have received previous laser photocoagulation.

#### *Ranibizumab and verteporfin photodynamic therapy in CNV secondary to PM*

There is no experience of concomitant administration of ranibizumab and verteporfin.

Ranluspec should be inspected visually for particulate matter and discoloration prior to administration.

The injection procedure should be carried out under aseptic conditions, which includes the use of surgical hand disinfection, sterile gloves, a sterile drape and a sterile eyelid speculum (or equivalent) and the availability of sterile paracentesis (if required). The patient’s medical history for




hypersensitivity reactions should be carefully evaluated prior to performing the intravitreal procedure. Adequate anaesthesia and a broad-spectrum topical microbicide to disinfect the periocular skin, eyelid and ocular surface should be administered prior to the injection, in accordance with local practice.


The vial is for single use only. After injection any unused product must be discarded. Any vial showing signs of damage or tampering must not be used. The sterility cannot be guaranteed unless the packaging seal remains intact. For preparation and intravitreal injection the following medical devices for single use are needed:

- a 5 µm filter needle (18G × 1½", 1.2 mm x 40 mm)
- a 1 mL sterile syringe (including a 0.05 mL mark)
- an injection needle (30G × ½").

These medical devices are not included within the Ranluspec pack.

To prepare Ranluspec for intravitreal administration to adult patients, please adhere to the following instructions:

	<p>Before withdrawal, disinfect the outer part of the rubber stopper of the vial.</p> <p>Place a 5-micron filter needle (18-gauge × 1-1/2 inch) onto a 1 mL Luer lock syringe using aseptic technique.</p> <p>Push the filter needle into the center of the vial stopper until the needle touches the bottom edge of the vial.</p> <p>Withdraw all the liquid from the vial, keeping the vial in an upright position, slightly inclined to ease complete withdrawal</p>
	<p>Ensure that the plunger rod is drawn sufficiently back when emptying the vial in order to completely empty the filter needle.</p> <p>The filter needle should be discarded after withdrawal of the vial contents and must not be used for the intravitreal injection.</p>
	<p>Attach a 30-gauge × 1/2-inch sterile injection needle firmly onto the syringe by screwing it tightly onto the Luer lock.</p> <p>Carefully remove the needle cap by pulling it straight off. Do not wipe the needle at any time.</p> <p>Note: Grip at the hub of the injection needle while removing the cap.</p>

	<p>Hold the syringe with the needle pointing up. If there are any air bubbles, gently tap the syringe with your finger until the bubbles rise to the top.</p> <p>Note: Do not wipe the injection needle. Do not pull back on the plunger.</p>
	<p>Hold the syringe at eye level, and carefully push the plunger rod until the plunger tip is aligned with the line that marks 0.05 mL on the syringe.</p>

The injection needle should be inserted 3.5-4.0 mm posterior to the limbus into the vitreous cavity, avoiding the horizontal meridian and aiming towards the centre of the globe. The injection volume of 0.05 mL is then delivered; a different scleral site should be used for subsequent injections.

After injection, do not recap the needle or detach it from the syringe. Dispose of the used syringe together with the needle in a sharps disposal container or in accordance with local requirements.

## Package leaflet: Information for the patient

Ranluspec 10 mg/mL solution for injection in pre-filled syringe  
ranibizumab

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

**Read all of this leaflet carefully before you are given this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What Ranluspec is and what it is used for
2. What you need to know before you are given Ranluspec
3. How Ranluspec is given
4. Possible side effects
5. How to store Ranluspec
6. Contents of the pack and other information

## 1. What Ranluspec is and what it is used for

### What Ranluspec is

Ranluspec is a solution which is injected into the eye. Ranluspec belongs to a group of medicines called antineovascularisation agents. It contains the active substance called ranibizumab.

### What Ranluspec is used for

Ranluspec is used in adults to treat several eye diseases causing vision impairment.

These diseases result from damage to the retina (light-sensitive layer at the back of the eye) caused by:

- Growth of leaky, abnormal blood vessels. This is observed in diseases such as age-related macular degeneration (AMD) and proliferative diabetic retinopathy (PDR, a disease caused by diabetes). It may also be associated with choroidal neovascularisation (CNV) due to pathologic myopia (PM), angioid streaks, central serous chorioretinopathy or inflammatory CNV.
- Macular oedema (swelling of the centre of the retina). This swelling can be caused by diabetes (a disease called diabetic macular oedema (DME)) or by the blockage of retinal veins of the retina (a disease called retinal vein occlusion (RVO)).

### How Ranluspec works

Ranluspec specifically recognises and binds to a protein called human vascular endothelial growth factor A (VEGF-A) present in the eye. In excess, VEGF-A causes abnormal blood vessel growth and swelling in the eye which can lead to impairment of vision in diseases like AMD, DME, PDR, RVO, PM and CNV. By binding to VEGF-A, Ranluspec can block its actions and prevent this abnormal growth and swelling.

In these diseases, Ranluspec can help to stabilise and in many cases improve your vision.

## 2. What you need to know before you are given Ranluspec

### You must not receive Ranluspec

- If you are allergic to ranibizumab or any of the other ingredients of this medicine (listed in

section 6).

- If you have an infection in or around your eye.
- If you have pain or redness (severe intraocular inflammation) in your eye.

### **Warnings and precautions**

Talk to your doctor before you are given Ranluspec.

- Ranluspec is given as an injection into the eye. Occasionally, an infection in the internal portion of the eye, pain or redness (inflammation), detachment or tear of one of the layers in the back of the eye (retinal detachment or tear and retinal pigment epithelial detachment or tear), or clouding of the lens (cataract) may occur after Ranluspec treatment. It is important to identify and treat such an infection or retinal detachment as soon as possible. Please tell your doctor immediately if you develop signs such as eye pain or increased discomfort, worsening eye redness, blurred or decreased vision, an increased number of small particles in your vision or increased sensitivity to light.
- In some patients the eye pressure may increase for a short period directly after the injection. This is something you may not notice, therefore your doctor may monitor this after each injection.
- Inform your doctor if you have a prior history of eye conditions or eye treatments, or if you have had a stroke or experienced transient signs of stroke (weakness or paralysis of limbs or face, difficulty speaking or understanding). This information will be taken into account to evaluate if Ranluspec is the appropriate treatment for you.

Please see section 4 (“Possible side effects”) for more detailed information on side effects that could occur during Ranluspec therapy.

### **Children and adolescents (below 18 years of age)**

The use of Ranluspec in children and adolescents has not been established and is therefore not recommended.

### **Other medicines and Ranluspec**

Tell your doctor if you are using, have recently used or might use any other medicines.

### **Pregnancy and breast-feeding**

- Women who could become pregnant must use effective contraception during treatment and for at least three further months after the last injection of Ranluspec.
- There is no experience of using Ranluspec in pregnant women. Ranluspec should not be used during pregnancy unless the potential benefit outweighs the potential risk to the unborn child. If you are pregnant, think you may be pregnant or planning to become pregnant, discuss this with your doctor before treatment with Ranluspec.
- Small amounts of Ranluspec may pass into breast milk, therefore Ranluspec is not recommended during breast-feeding. Ask your doctor or pharmacist for advice before Ranluspec treatment.

### **Driving and using machines**

After Ranluspec treatment you may experience some temporary vision blurring. If this happens, do not drive or use machines until this resolves.

### **This medicine contains polysorbate 20**

This medicine contains 0.005 mg of polysorbate 20 in each dose of 0.05 mL which is equivalent to 0.1 mg/mL. Polysorbates may cause allergic reactions. Tell your doctor if you have any known allergies.

## **3. How Ranluspec is given**

Ranluspec is administered as a single injection into your eye by your eye doctor under a local anaesthetic. The usual dose of an injection is 0.05 mL (which contains 0.5 mg of ranibizumab). The

pre-filled syringe contains more than the recommended dose of 0.5 mg. The extractable volume is not to be used in total. The excess volume should be expelled prior to injection. Injecting the entire volume of the pre-filled syringe could result in overdose.

The interval between two doses injected into the same eye should be at least four weeks. All injections will be administered by your eye doctor.

Before the injection, your doctor will wash your eye carefully to prevent infection. Your doctor will also give you a local anaesthetic to reduce or prevent any pain you might have with the injection.

The treatment is started with one injection of Ranluspec per month. Your doctor will monitor the condition of your eye and, depending on how you respond to the treatment, will decide if and when you need to receive further treatment.

Detailed instructions for use are given at the end of the leaflet under “How to prepare and administer Ranluspec”.

#### **Elderly (age 65 years and over)**

Ranluspec can be used for people of 65 years of age and over without dose adjustment.

#### **Before stopping Ranluspec treatment**

If you are considering stopping Ranluspec treatment, please go to your next appointment and discuss this with your doctor. Your doctor will advise you and decide how long you should be treated with Ranluspec.

If you have any further questions on the use of this medicine, ask your doctor.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The side effects associated with the administration of Ranluspec are either due to the medicine itself or the injection procedure and mostly affect the eye.

The most serious side effects are described below:

**Common serious side effects** (may affect up to 1 in 10 people): Detachment or tear of the layer in the back of the eye (retinal detachment or tear), resulting in flashes of light with floaters progressing to a temporary loss of sight, or a clouding of the lens (cataract).

**Uncommon serious side effects** (may affect up to 1 in 100 people): Blindness, infection of the eyeball (endophthalmitis) with inflammation of the inside of the eye.

The symptoms you might experience are pain or increased discomfort in your eye, worsening eye redness, blurred or decreased vision, an increased number of small particles in your vision or increased sensitivity to light. **Please tell your doctor immediately if you develop any of these side effects.**

The most frequently reported side effects are described below:

**Very common side effects** (may affect more than 1 in 10 people)

Visual side effects include: Inflammation of the eye, bleeding in the back of the eye (retinal bleeding), visual disturbances, eye pain, small particles or spots in your vision (floaters), bloodshot eye, eye irritation, a feeling of having something in the eye, increased tear production, inflammation or infection of the eyelid margins, dry eye, redness or itching of the eye and increased eye pressure. Non-visual side effects include: Sore throat, nasal congestion, runny nose, headache and joint pain.

Other side effects which may occur following Ranluspec treatment are described below:

#### **Common side effects**

Visual side effects include: Decreased sharpness of vision, swelling of a section of the eye (uvea, cornea), inflammation of the cornea (front part of eye), small marks on the surface of the eye, blurred vision, bleeding at the site of injection, bleeding in the eye, discharge from the eye with itching, redness and swelling (conjunctivitis), light sensitivity, eye discomfort, swelling of the eyelid, eyelid pain.

Non-visual side effects include: Urinary tract infection, low red blood cells count (with symptoms such as tiredness, breathlessness, dizziness, pale skin), anxiety, cough, nausea, allergic reactions like rash, hives, itching and skin reddening.

#### **Uncommon side effects**

Visual side effects include: Inflammation and bleeding in the front part of the eye, sac of pus on the eye, changes of the central part of the eye surface, pain or irritation at the site of injection, abnormal sensation in the eye, irritation of the eyelid.

#### **Reporting of side effects**

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly the national reporting system listed in [Appendix V](#). By reporting side effects, you can help provide more information on the safety of this medicine.

### **5. How to store Ranluspec**

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and vial label after EXP. The expiry date refers to the last day of that month.
- Store in a refrigerator (2 °C – 8 °C). Do not freeze.
- Prior to use, the sealed tray may be kept at room temperature (25 °C) for up to 24 hours.
- Keep the pre-filled syringe in its unopened tray in the carton in order to protect from light.
- Do not use any pack that is damaged.

### **6. Contents of the pack and other information**

#### **What Ranluspec contains**

- The active substance is ranibizumab. Each mL contains 10 mg ranibizumab. One pre-filled syringe contains 0.165 mL, equivalent to 1.65 mg ranibizumab. This provides a usable amount to deliver a single dose of 0.05 mL containing 0.5 mg ranibizumab.
- The other ingredients are  $\alpha,\alpha$ -trehalose dihydrate; histidine hydrochloride, monohydrate; histidine; polysorbate 20; water for injections.

#### **What Ranluspec looks like and contents of the pack**

Ranluspec is a solution for injection in a pre-filled syringe. The pre-filled syringe contains 0.165 mL of a sterile, clear to slightly opalescent, colourless to pale brownish and aqueous solution. The pre-filled syringe contains more than the recommended dose of 0.5 mg. The extractable volume is not to be used in total. The excess volume should be expelled prior to injection. Injecting the entire volume of the pre-filled syringe could result in overdose.

Pack size of one pre-filled syringe, packed in a sealed tray. The pre-filled syringe is for single use only.

#### **Marketing Authorisation Holder**

Lupin Europe GmbH  
Hanauer Landstraße 139-143  
60314 Frankfurt  
Germany

**Manufacturer**

Hormosan Pharma GmbH  
Hanauer Landstraße 139-143  
60314 Frankfurt  
Germany

**This leaflet was last revised in****Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<https://www.ema.europa.eu>.

The following information is intended for healthcare professionals only:

Please also refer to section 3 “How Ranluspec is given”.

### **How to prepare and administer Ranluspec**

Single-use pre-filled syringe for intravitreal use only

Ranluspec must be administered by a qualified ophthalmologist experienced in intravitreal injections.

In wet AMD, in CNV, in PDR and in visual impairment due to DME or to macular oedema secondary to RVO the recommended dose for Ranluspec is 0.5 mg given as a single intravitreal injection. This corresponds to an injection volume of 0.05 mL. The interval between two doses injected into the same eye should be at least four weeks.

Treatment is initiated with one injection per month until maximum visual acuity is achieved and/or there are no signs of disease activity i.e. no change in visual acuity and in other signs and symptoms of the disease under continued treatment. In patients with wet AMD, DME, PDR and RVO, initially, three or more consecutive, monthly injections may be needed.

Thereafter, monitoring and treatment intervals should be determined by the physician and should be based on disease activity, as assessed by visual acuity and/or anatomical parameters.

If, in the physician’s opinion, visual and anatomical parameters indicate that the patient is not benefiting from continued treatment, Ranluspec should be discontinued.

Monitoring for disease activity may include clinical examination, functional testing or imaging techniques (e.g. optical coherence tomography or fluorescein angiography).

If patients are being treated according to a treat-and-extend regimen, once maximum visual acuity is achieved and/or there are no signs of disease activity, the treatment intervals can be extended stepwise until signs of disease activity or visual impairment recur. The treatment interval should be extended by no more than two weeks at a time for wet AMD and may be extended by up to one month at a time for DME. For PDR and RVO, treatment intervals may also be gradually extended, however there are insufficient data to conclude on the length of these intervals. If disease activity recurs, the treatment interval should be shortened accordingly.

The treatment of visual impairment due to CNV should be determined individually per patient based on disease activity. Some patients may only need one injection during the first 12 months; others may need more frequent treatment, including a monthly injection. For CNV secondary to pathologic myopia (PM), many patients may only need one or two injections during the first year.

#### *Ranibizumab and laser photocoagulation in DME and macular oedema secondary to BRVO*

There is some experience of ranibizumab administered concomitantly with laser photocoagulation. When given on the same day, ranibizumab should be administered at least 30 minutes after laser photocoagulation. Ranibizumab can be administered in patients who have received previous laser photocoagulation.

#### *Ranibizumab and verteporfin photodynamic therapy in CNV secondary to PM*

There is no experience of concomitant administration of ranibizumab and verteporfin.

Ranluspec should be inspected visually for particulate matter and discoloration prior to administration.

The injection procedure should be carried out under aseptic conditions, which includes the use of surgical hand disinfection, sterile gloves, a sterile drape and a sterile eyelid speculum (or equivalent) and the availability of sterile paracentesis (if required). The patient’s medical history for hypersensitivity reactions should be carefully evaluated prior to performing the intravitreal procedure.


Adequate anaesthesia and a broad-spectrum topical microbicide to disinfect the periocular skin, eyelid and ocular surface should be administered prior to the injection, in accordance with local practice.



The pre-filled syringe is for single use only. The pre-filled syringe is sterile. Do not use the product if the packaging is damaged. The sterility of the pre-filled syringe cannot be guaranteed unless the tray remains sealed. Do not use the pre-filled syringe if the solution is discoloured, cloudy or contains particles.


The pre-filled syringe contains more than the recommended dose of 0.5 mg. The extractable volume of the pre-filled syringe (0.1 mL) is not to be used in total. The excess volume should be expelled prior to injection. Injecting the entire volume of the pre-filled syringe could result in overdose. To expel the air bubble along with the excess medicinal product, slowly push the plunger until the edge below the dome of the rubber stopper is aligned with the black dosing line on the syringe (equivalent to 0.05 mL, i.e., 0.5 mg ranibizumab).

For the intravitreal injection, a 30G x ½" sterile injection needle should be used.

To prepare Ranluspec for intravitreal administration, please adhere to the instructions for use:

Introduction	<p>1. The prefilled syringe is sterile and is for single dose only. Do not use the product if the packaging is damaged or has been tampered with.</p> <p>2. To prepare Ranluspec for intravitreal administration, please adhere to these instructions for use. Read all the instructions carefully before using the prefilled syringe.</p> <p>3. The opening of the sealed tray and all subsequent steps should be done under aseptic conditions.</p> <p>4. For the intravitreal injection, a 30-gauge × ½ inch sterile injection needle should be used. (Not provided).</p> <p><b>Note: The dose must be set to 0.05 mL.</b></p>
Pre-filled syringe description	 <p><b>Figure 1 Ranluspec pre-filled syringe illustration</b></p>
Step 1	<p><b>PREPARATION:</b>          Make sure that your pack contains a sterile prefilled syringe in a sealed tray. Peel the lid off the syringe tray and using aseptic technique, remove the syringe.</p>
Step 2	<p><b>INSPECT SYRINGE:</b></p> <p>1. RANIBIZUMAB should be clear to slightly opalescent.</p> <p><b>2. Do not use the prefilled syringe if:</b></p> <ul style="list-style-type: none"> <li>-The syringe cap is detached from the Luer lock.</li> <li>- The syringe is damaged.</li> <li>- Particulates, cloudiness, or</li> </ul>  <p><b>Figure 2 Ranluspec 0.5 mg prefilled syringe with a White finger grip.</b></p>

	<p>discoloration are visible.</p> <p>3. If any of the above is not true, discard the pre-filled syringe and use a new one.</p> <p><b>Note: To avoid compromising the sterility of the product, do not pull back the plunger</b></p>	
<p><b>Step 3</b></p>	<p><b>REMOVE SYRINGE CAP:</b> Remove the syringe cap by rotating anti-clockwise movement. (See Figure 3).</p> <p><b>Note: Syringe cap to be opened once user is ready for the injection. Once syringe cap opened, Do Not recap, Syringe should be disposed if user is not ready for injection.</b></p>	 <p><b>Figure 3 Removal of threaded closure cap</b></p>
<p><b>Step 4</b></p>	<p><b>ATTACH NEEDLE:</b></p> <p>1. Attach a 30G × ½ inch sterile injection needle firmly onto the syringe by screwing it tightly onto the Luer lock (See Figure 4)</p> <p>2. Carefully remove the needle cap by pulling it straight off.</p> <p><b>Note: Do not wipe the needle at any time.</b></p>	 <p><b>Figure 4 Needle attachment</b></p>
<p><b>Step 5</b></p>	<p><b>DISLodge AIR BUBBLES</b></p> <p>1. Hold the syringe with the needle pointing up.</p> <p>2. If there are any air bubbles, gently tap the syringe with your finger until the bubbles rise to the top (See Figure 5).</p>	 <p><b>Figure 5 Removal of air bubble</b></p>

<p><b>Step 6</b></p>	<p><b>EXPEL AIR AND ADJUST DRUG DOSE.</b></p> <p>1. Hold the syringe at eye level, and carefully push the plunger rod until the edge below the dome of the rubber stopper is aligned with the 0.05 mL dose mark (See Figure 6).</p> <p><b>Note: The plunger rod is <u>attached</u> to the rubber stopper, to avoid air entrapment do not pull back the plunger rod.</b></p>	 <p><b>Figure 6 Dose adjustment</b></p>
<p><b>Step 7</b></p>	<p><b>INJECT:</b></p> <ol style="list-style-type: none"> <li>1. The injection procedure should be carried out under aseptic conditions.</li> <li>2. Insert the needle into the injection site.</li> <li>3. Inject slowly until rubber stopper reaches the bottom of the syringe to deliver the volume of 0.05 mL.</li> <li>4. After injection, do not re-cap the needle or detach it from the syringe. dispose of the used syringe together with the needle in a sharp disposal container or in accordance with local requirements.</li> </ol>	