

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Sabervel 75 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 75 mg of irbesartan

Excipient with known effect:

20 mg of lactose monohydrate per film-coated tablet

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet.

White, concave, round, film-coated tablet with 7 mm diameter.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Sabervel is indicated in adults for the treatment of essential hypertension.

It is also indicated for the treatment of renal disease in adult patients with hypertension and type 2 diabetes mellitus as part of an antihypertensive medicinal product regimen (see sections 4.3, 4.4, 4.5 and 5.1).

### 4.2 Posology and method of administration

#### Posology

The usual recommended initial and maintenance dose is 150 mg once daily, with or without food. Sabervel at a dose of 150 mg once daily generally provides a better 24 hour blood pressure control than 75 mg. However, initiation of therapy with 75 mg could be considered, particularly in haemodialysed patients and in the elderly over 75 years.

In patients insufficiently controlled with 150 mg once daily, the dose of Sabervel can be increased to 300 mg, or other antihypertensive agents can be added (see sections 4.3, 4.4, 4.5 and 5.1). In particular, the addition of a diuretic such as hydrochlorothiazide has been shown to have an additive effect with Sabervel (see section 4.5).

In hypertensive type 2 diabetic patients, therapy should be initiated at 150 mg irbesartan once daily and titrated up to 300 mg once daily as the preferred maintenance dose for treatment of renal disease. The demonstration of renal benefit of Sabervel in hypertensive type 2 diabetic patients is based on studies where irbesartan was used in addition to other antihypertensive agents, as needed, to reach target blood pressure (see sections 4.3, 4.4, 4.5 and 5.1).

#### Special Populations

Renal impairment: no dosage adjustment is necessary in patients with impaired renal function. A lower starting dose (75 mg) should be considered for patients undergoing haemodialysis (see section 4.4).

Hepatic impairment: no dosage adjustment is necessary in patients with mild to moderate hepatic

impairment. There is no clinical experience in patients with severe hepatic impairment.

Elderly patients: although consideration should be given to initiating therapy with 75 mg in patients over 75 years of age, dosage adjustment is not usually necessary for the elderly.

Paediatric population: the safety and efficacy of Sabervel in children aged 0 to 18 has not been established. Currently available data are described in section 4.8, 5.1 and 5.2 but no recommendation on a posology can be made.

#### Method of Administration

For oral use.

### **4.3 Contraindications**

Hypersensitivity to the active substance, or to any of the excipients (see section 6.1).  
Second and third trimesters of pregnancy (see sections 4.4 and 4.6).

The concomitant use of Sabervel with aliskiren-containing products is contraindicated in patients with diabetes mellitus or renal impairment ( $\text{GFR} < 60 \text{ ml/min/1.73 m}^2$ ) (see sections 4.5 and 5.1).

### **4.4 Special warnings and precautions for use**

Intravascular volume depletion: symptomatic hypotension, especially after the first dose, may occur in patients who are volume and/or sodium depleted by vigorous diuretic therapy, dietary salt restriction, diarrhoea or vomiting. Such conditions should be corrected before the administration of Sabervel.

Renovascular hypertension: there is an increased risk of severe hypotension and renal insufficiency when patients with bilateral renal artery stenosis or stenosis of the artery to a single functioning kidney are treated with medicinal products that affect the renin-angiotensin-aldosterone system. While this is not documented with Sabervel, a similar effect should be anticipated with angiotensin-II receptor antagonists.

Renal impairment and kidney transplantation: when Sabervel is used in patients with impaired renal function, a periodic monitoring of potassium and creatinine serum levels is recommended. There is no experience regarding the administration of Sabervel in patients with a recent kidney transplantation.

Hypertensive patients with type 2 diabetes and renal disease: the effects of irbesartan both on renal and cardiovascular events were not uniform across all subgroups, in an analysis carried out in the study with patients with advanced renal disease. In particular, they appeared less favourable in women and non-white subjects (see section 5.1).

#### Dual blockade of the renin-angiotensin-aldosterone system (RAAS)

There is evidence that the concomitant use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren increases the risk of hypotension, hyperkalaemia and decreased renal function (including acute renal failure). Dual blockade of RAAS through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is therefore not recommended (see sections 4.5 and 5.1).

If dual blockade therapy is considered absolutely necessary, this should only occur under specialist supervision and subject to frequent close monitoring of renal function, electrolytes and blood pressure. ACE-inhibitors and angiotensin II receptor blockers should not be used concomitantly in patients with diabetic nephropathy.

Hyperkalaemia: as with other medicinal products that affect the renin-angiotensin-aldosterone system, hyperkalaemia may occur during the treatment with Sabervel, especially in the presence of renal impairment, overt proteinuria due to diabetic renal disease, and/or heart failure. Close monitoring of serum potassium in patients at risk is recommended (see section 4.5).

Lithium: the combination of lithium and Sabervel is not recommended (see section 4.5).

Aortic and mitral valve stenosis, obstructive hypertrophic cardiomyopathy: as with other vasodilators, special caution is indicated in patients suffering from aortic or mitral stenosis, or obstructive hypertrophic cardiomyopathy.

Primary aldosteronism: patients with primary aldosteronism generally will not respond to antihypertensive medicinal products acting through inhibition of the renin-angiotensin system. Therefore, the use of Sabervel is not recommended.

General: in patients whose vascular tone and renal function depend predominantly on the activity of the renin-angiotensin-aldosterone system (e.g. patients with severe congestive heart failure or underlying renal disease, including renal artery stenosis), treatment with angiotensin converting enzyme inhibitors or angiotensin-II receptor antagonists that affect this system has been associated with acute hypotension, azotaemia, oliguria, or rarely acute renal failure. As with any antihypertensive agent, excessive blood pressure decrease in patients with ischaemic cardiopathy or ischaemic cardiovascular disease could result in a myocardial infarction or stroke.

As observed for angiotensin converting enzyme inhibitors, irbesartan and the other angiotensin antagonists are apparently less effective in lowering blood pressure in black people than in non-blacks, possibly because of higher prevalence of low-renin states in the black hypertensive population (see section 5.1).

Pregnancy: Angiotensin II Receptor Antagonists (AIIRAs) should not be initiated during pregnancy. Unless continued AIIRA therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIRAs should be stopped immediately, and, if appropriate, alternative therapy should be started (see sections 4.3 and 4.6).

Lactose: this medicinal product contains lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.

Paediatric population: irbesartan has been studied in paediatric populations aged 6 to 16 years old but the current data are insufficient to support an extension of the use in children until further data become available (see sections 4.8, 5.1 and 5.2).

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Diuretics and other antihypertensive agents: other antihypertensive agents may increase the hypotensive effects of irbesartan; however irbesartan has been safely administered with other antihypertensive agents, such as beta-blockers, long-acting calcium channel blockers, and thiazide diuretics. Prior treatment with high dose diuretics may result in volume depletion and a risk of hypotension when initiating therapy with Sabervel (see section 4.4).

Clinical trial data has shown that dual blockade of the renin-angiotensin-aldosterone-system (RAAS) through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is associated with a higher frequency of adverse events such as hypotension, hyperkalaemia and decreased renal function (including acute renal failure) compared to the use of a single RAAS-acting agent (see sections 4.3, 4.4 and 5.1).

Potassium supplements and potassium-sparing diuretics: based on experience with the use of other medicinal products that affect the renin-angiotensin system, concomitant use of potassium-sparing diuretics, potassium supplements, salt substitutes containing potassium or other medicinal products that may increase serum potassium levels (e.g. heparin) may lead to increases in serum potassium and is, therefore, not recommended (see section 4.4).

Lithium: reversible increases in serum lithium concentrations and toxicity have been reported during

concomitant administration of lithium with angiotensin converting enzyme inhibitors. Similar effects have been very rarely reported with irbesartan so far. Therefore, this combination is not recommended (see section 4.4). If the combination proves necessary, careful monitoring of serum lithium levels is recommended.

Non-steroidal anti-inflammatory drugs: when angiotensin II antagonists are administered simultaneously with non-steroidal anti-inflammatory drugs (i.e. selective COX-2 inhibitors, acetylsalicylic acid (> 3 g/day) and non-selective NSAIDs), attenuation of the antihypertensive effect may occur.

As with ACE inhibitors, concomitant use of angiotensin II antagonists and NSAIDs may lead to an increased risk of worsening of renal function, including possible acute renal failure, and an increase in serum potassium, especially in patients with poor pre-existing renal function. The combination should be administered with caution, especially in the elderly. Patients should be adequately hydrated and consideration should be given to monitoring renal function after initiation of concomitant therapy, and periodically thereafter.

Additional information on irbesartan interactions: in clinical studies, the pharmacokinetic of irbesartan is not affected by hydrochlorothiazide. Irbesartan is mainly metabolised by CYP2C9 and to a lesser extent by glucuronidation. No significant pharmacokinetic or pharmacodynamic interactions were observed when irbesartan was coadministered with warfarin, a medicinal product metabolised by CYP2C9. The effects of CYP2C9 inducers such as rifampicin on the pharmacokinetic of irbesartan have not been evaluated. The pharmacokinetic of digoxin was not altered by coadministration of irbesartan.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy:

The use of AIIRAs is not recommended during the first trimester of pregnancy (see section 4.4). The use of AIIRAs is contraindicated during the second and third trimesters of pregnancy (see sections 4.3 and 4.4).

Epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded. Whilst there is no controlled epidemiological data on the risk with Angiotensin II Receptor Antagonists (AIIRAs), similar risks may exist for this class of drugs. Unless continued AIIRA therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIRAs should be stopped immediately, and, if appropriate, alternative therapy should be started.

Exposure to AIIRA therapy during the second and third trimesters is known to induce human fetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia). (See section 5.3).

Should exposure to AIIRAs have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended.

Infants whose mothers have taken AIIRAs should be closely observed for hypotension (see sections 4.3 and 4.4).

##### Breast-feeding:

Because no information is available regarding the use of Sabervel during breast-feeding, Sabervel is not recommended and alternative treatments with better established safety profiles during breast-feeding are preferable, especially while nursing a newborn or preterm infant.

It is unknown whether irbesartan or its metabolites are excreted in human milk.

Available pharmacodynamic/toxicological data in rats have shown excretion of irbesartan or its

metabolites in milk (for details see 5.3).

### Fertility

Irbesartan had no effect upon fertility of treated rats and their offspring up to the dose levels inducing the first signs of parental toxicity (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

No studies on the effects on the ability to drive and use machines have been performed. Based on its pharmacodynamic properties, irbesartan is unlikely to affect this ability. When driving vehicles or operating machines, it should be taken into account that dizziness or weariness may occur during treatment.

### **4.8 Undesirable effects**

In placebo-controlled trials in patients with hypertension, the overall incidence of adverse events did not differ between the irbesartan (56.2%) and the placebo groups (56.5%). Discontinuation due to any clinical or laboratory adverse event was less frequent for irbesartan-treated patients (3.3%) than for placebo-treated patients (4.5%). The incidence of adverse events was not related to dose (in the recommended dose range), gender, age, race, or duration of treatment.

In diabetic hypertensive patients with microalbuminuria and normal renal function, orthostatic dizziness and orthostatic hypotension were reported in 0.5% of the patients (i.e., uncommon) but in excess of placebo.

The following table presents the adverse drug reactions that were reported in placebo-controlled trials in which 1,965 hypertensive patients received irbesartan. Terms marked with a star (\*) refer to the adverse reactions that were additionally reported in > 2% of diabetic hypertensive patients with chronic renal insufficiency and overt proteinuria and in excess of placebo.

The frequency of adverse reactions listed below is defined using the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Adverse reactions additionally reported from post-marketing experience are also listed. These adverse reactions are derived from spontaneous reports.

#### *Immune system disorders:*

Not known: hypersensitivity reactions such as angioedema, rash, urticaria

#### *Metabolism and nutrition disorders:*

Not known: hyperkalaemia

#### *Nervous system disorders:*

Common: dizziness, orthostatic dizziness\*

Not known: vertigo, headache

#### *Ear and labyrinth disorder:*

Not known: tinnitus

#### *Cardiac disorders:*

Uncommon: tachycardia

#### *Vascular disorders:*

Common: orthostatic hypotension\*

Uncommon: flushing

*Respiratory, thoracic and mediastinal disorders:*

Uncommon: cough

*Gastrointestinal disorders:*

Common: nausea/vomiting

Uncommon: diarrhoea, dyspepsia/heartburn

Not known: dysgeusia

*Hepatobiliary disorders:*

Uncommon: jaundice

Not known: hepatitis, abnormal liver function

*Skin and subcutaneous tissue disorders:*

Not known: leukocytoclastic vasculitis

*Musculoskeletal and connective tissue disorders:*

Common: musculoskeletal pain\*

Not known: arthralgia, myalgia (in some cases associated with increased plasma creatine kinase levels), muscle cramps

*Renal and urinary disorders:*

Not known: impaired renal function including cases of renal failure in patients at risk (see section 4.4)

*Reproductive system and breast disorders:*

Uncommon: sexual dysfunction

*General disorders and administration site conditions:*

Common: fatigue

Uncommon: chest pain

Investigations:

Very common: Hyperkalaemia\* occurred more often in diabetic patients treated with irbesartan than with placebo. In diabetic hypertensive patients with microalbuminuria and normal renal function, hyperkalaemia ( $\geq 5.5$  mEq/L) occurred in 29.4% of the patients in the irbesartan 300 mg group and 22% of the patients in the placebo group. In diabetic hypertensive patients with chronic renal insufficiency and overt proteinuria, hyperkalaemia ( $\geq 5.5$  mEq/L) occurred in 46.3% of the patients in the irbesartan group and 26.3% of the patients in the placebo group.

Common: significant increases in plasma creatine kinase were commonly observed (1.7%) in irbesartan treated subjects. None of these increases were associated with identifiable clinical musculoskeletal events. In 1.7% of hypertensive patients with advanced diabetic renal disease treated with irbesartan, a decrease in haemoglobin\*, which was not clinically significant, has been observed.

Paediatric population:

In a randomised trial of 318 hypertensive children and adolescents aged 6 to 16 years, the following adverse reactions occurred in the 3-week double-blind phase: headache (7.9%), hypotension (2.2%), dizziness (1.9%), cough (0.9%). In the 26-week open-label period of this trial the most frequent laboratory abnormalities observed were creatinine increases (6.5%) and elevated CK values in 2% of child recipients.

## 4.9 Overdose

Experience in adults exposed to doses of up to 900 mg/day for 8 weeks revealed no toxicity. The most likely manifestations of overdose are expected to be hypotension and tachycardia; bradycardia might also occur from overdose. No specific information is available on the treatment of overdose with Sabervel. The patient should be closely monitored, and the treatment should be symptomatic and supportive. Suggested measures include induction of emesis and/or gastric lavage. Activated charcoal may be useful in the treatment of overdose. Irbesartan is not removed by haemodialysis.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Angiotensin-II antagonists, plain.

ATC code: C09C A04.

Mechanism of action: Irbesartan is a potent, orally active, selective angiotensin-II receptor (type AT<sub>1</sub>) antagonist. It is expected to block all actions of angiotensin-II mediated by the AT<sub>1</sub> receptor, regardless of the source or route of synthesis of angiotensin-II. The selective antagonism of the angiotensin-II (AT<sub>1</sub>) receptors results in increases in plasma renin levels and angiotensin-II levels, and a decrease in plasma aldosterone concentration. Serum potassium levels are not significantly affected by irbesartan alone at the recommended doses. Irbesartan does not inhibit ACE (kininase-II), an enzyme which generates angiotensin-II and also degrades bradykinin into inactive metabolites.

Irbesartan does not require metabolic activation for its activity.

Clinical efficacy:

#### Hypertension

Irbesartan lowers blood pressure with minimal change in heart rate. The decrease in blood pressure is dose-related for once a day doses with a tendency towards plateau at doses above 300 mg. Doses of 150-300 mg once daily lower supine or seated blood pressures at trough (i.e. 24 hours after dosing) by an average of 8-13/5-8 mm Hg (systolic/diastolic) greater than those associated with placebo. Peak reduction of blood pressure is achieved within 3-6 hours after administration and the blood pressure lowering effect is maintained for at least 24 hours. At 24 hours the reduction of blood pressure was 60-70% of the corresponding peak diastolic and systolic responses at the recommended doses. Once daily dosing with 150 mg produced trough and mean 24 hour responses similar to twice daily dosing on the same total dose.

The blood pressure lowering effect of irbesartan is evident within 1-2 weeks, with the maximal effect occurring by 4-6 weeks after start of therapy. The antihypertensive effects are maintained during long term therapy. After withdrawal of therapy, blood pressure gradually returns toward baseline. Rebound hypertension has not been observed.

The blood pressure lowering effects of irbesartan and thiazide-type diuretics are additive. In patients not adequately controlled by irbesartan alone, the addition of a low dose of hydrochlorothiazide (12.5 mg) to irbesartan once daily results in a further placebo-adjusted blood pressure reduction at trough of 7-10/3-6 mm Hg (systolic/diastolic).

The efficacy of irbesartan is not influenced by age or gender. As is the case with other medicinal products that affect the renin-angiotensin system, black hypertensive patients have notably less response to irbesartan monotherapy. When irbesartan is administered concomitantly with a low dose of hydrochlorothiazide (e.g. 12.5 mg daily), the antihypertensive response in black patients approaches that of white patients.

There is no clinically important effect on serum uric acid or urinary uric acid secretion.

#### Paediatric population

Reduction of blood pressure with 0.5 mg/kg (low), 1.5 mg/kg (medium) and 4.5 mg/kg (high) target titrated doses of irbesartan was evaluated in 318 hypertensive or at risk (diabetic, family history of



hypertension) children and adolescents aged 6 to 16 years over a three week period. At the end of the three weeks the mean reduction from baseline in the primary efficacy variable, trough seated systolic blood pressure (SeSBP) was 11.7 mmHg (low dose), 9.3 mmHg (medium dose), 13.2 mmHg (high dose). No significant difference was apparent between these doses. Adjusted mean change of trough seated diastolic blood pressure (SeDBP) was as follows: 3.8 mmHg (low dose), 3.2 mmHg (medium dose), 5.6 mmHg (high dose). Over a subsequent two week period where patients were re-randomized to either active medicinal product or placebo, patients on placebo had increases of 2.4 and 2.0 mmHg in SeSBP and SeDBP compared to +0.1 and -0.3 mmHg changes respectively in those on all doses of irbesartan (see section 4.2).

#### Hypertension and type 2 diabetes with renal disease

The “Irbesartan Diabetic Nephropathy Trial (IDNT)” shows that irbesartan decreases the progression of renal disease in patients with chronic renal insufficiency and overt proteinuria. IDNT was a double blind, controlled, morbidity and mortality trial comparing irbesartan, amlodipine and placebo. In 1,715 hypertensive patients with type 2 diabetes, proteinuria  $\geq 900$  mg/day and serum creatinine ranging from 1.0-3.0 mg/dl, the long-term effects (mean 2.6 years) of irbesartan on the progression of renal disease and all-cause mortality were examined. Patients were titrated from 75 mg to a maintenance dose of 300 mg irbesartan, from 2.5 mg to 10 mg amlodipine, or placebo as tolerated.

Patients in all treatment groups typically received between 2 and 4 antihypertensive agents (e.g., diuretics, beta blockers, alpha blockers) to reach a predefined blood pressure goal of  $\leq 135/85$  mmHg or a 10 mmHg reduction in systolic pressure if baseline was  $> 160$  mmHg. Sixty per cent (60%) of patients in the placebo group reached this target blood pressure whereas this figure was 76% and 78% in the irbesartan and amlodipine groups respectively.

Irbesartan significantly reduced the relative risk in the primary combined endpoint of doubling serum creatinine, end-stage renal disease (ESRD) or all-cause mortality. Approximately 33% of patients in the irbesartan group reached the primary renal composite endpoint compared to 39% and 41% in the placebo and amlodipine groups [20% relative risk reduction versus placebo ( $p = 0.024$ ) and 23% relative risk reduction compared to amlodipine ( $p = 0.006$ )]. When the individual components of the primary endpoint were analysed, no effect in all cause mortality was observed, while a positive trend in the reduction in ESRD and a significant reduction in doubling of serum creatinine were observed.

Subgroups consisting of gender, race, age, duration of diabetes, baseline blood pressure, serum creatinine, and albumin excretion rate were assessed for treatment effect. In the female and black subgroups which represented 32% and 26% of the overall study population respectively, a renal benefit was not evident, although the confidence intervals do not exclude it. As for the secondary endpoint of fatal and non-fatal cardiovascular events, there was no difference among the three groups in the overall population, although an increased incidence of non-fatal MI was seen for women and a decreased incidence of non-fatal MI was seen in males in the irbesartan group versus the placebo-based regimen. An increased incidence of non-fatal MI and stroke was seen in females in the irbesartan-based regimen versus the amlodipine-based regimen, while hospitalization due to heart failure was reduced in the overall population. However, no proper explanation for these findings in women has been identified.

The study of the “Effects of Irbesartan on Microalbuminuria in Hypertensive Patients with type 2 Diabetes Mellitus (IRMA 2)” shows that irbesartan 300 mg delays progression to overt proteinuria in patients with microalbuminuria. IRMA 2 was a placebo-controlled double blind morbidity study in 590 patients with type 2 diabetes, microalbuminuria (30-300 mg/day) and normal renal function (serum creatinine  $\leq 1.5$  mg/dl in males and  $< 1.1$  mg/dl in females). The study examined the long-term effects (2 years) of irbesartan on the progression to clinical (overt) proteinuria (urinary albumin excretion rate (UAER)  $> 300$  mg/day, and an increase in UAER of at least 30% from baseline). The predefined blood pressure goal was  $\leq 135/85$  mmHg. Additional antihypertensive agents (excluding ACE inhibitors, angiotensin II receptor antagonists and dihydropyridine calcium blockers) were added as needed to help achieve the blood pressure goal. While similar blood pressure was achieved in all treatment groups, fewer subjects in the irbesartan 300 mg group (5.2%) than in the placebo (14.9%) or in the irbesartan 150 mg group (9.7%) reached the endpoint of overt proteinuria, demonstrating a 70% relative risk reduction versus placebo ( $p = 0.0004$ ) for the higher dose. An accompanying improvement in the glomerular filtration rate (GFR) was not observed during the first three months of

treatment. The slowing in the progression to clinical proteinuria was evident as early as three months and continued over the 2 year period. Regression to normoalbuminuria (< 30 mg/day) was more frequent in the irbesartan 300 mg group (34%) than in the placebo group (21%).

Two large randomised, controlled trials (ONTARGET (ONgoing Telmisartan Alone and in combination with Ramipril Global Endpoint Trial) and VA NEPHRON-D (The Veterans Affairs Nephropathy in Diabetes)) have examined the use of the combination of an ACE-inhibitor with an angiotensin II receptor blocker.

ONTARGET was a study conducted in patients with a history of cardiovascular or cerebrovascular disease, or type 2 diabetes mellitus accompanied by evidence of end-organ damage. VA NEPHRON-D was a study in patients with type 2 diabetes mellitus and diabetic nephropathy.

These studies have shown no significant beneficial effect on renal and/or cardiovascular outcomes and mortality, while an increased risk of hyperkalaemia, acute kidney injury and/or hypotension as compared to monotherapy was observed. Given their similar pharmacodynamic properties, these results are also relevant for other ACE-inhibitors and angiotensin II receptor blockers.

ACE-inhibitors and angiotensin II receptor blockers should therefore not be used concomitantly in patients with diabetic nephropathy.

ALTITUDE (Aliskiren Trial in Type 2 Diabetes Using Cardiovascular and Renal Disease Endpoints) was a study designed to test the benefit of adding aliskiren to a standard therapy of an ACE-inhibitor or an angiotensin II receptor blocker in patients with type 2 diabetes mellitus and chronic kidney disease, cardiovascular disease, or both. The study was terminated early because of an increased risk of adverse outcomes. Cardiovascular death and stroke were both numerically more frequent in the aliskiren group than in the placebo group and adverse events and serious adverse events of interest (hyperkalaemia, hypotension and renal dysfunction) were more frequently reported in the aliskiren group than in the placebo group.

## 5.2 Pharmacokinetic properties

After oral administration, irbesartan is well absorbed: studies of absolute bioavailability gave values of approximately 60-80%. Concomitant food intake does not significantly influence the bioavailability of irbesartan. Plasma protein binding is approximately 96%, with negligible binding to cellular blood components. The volume of distribution is 53 - 93 litres. Following oral or intravenous administration of <sup>14</sup>C irbesartan, 80-85% of the circulating plasma radioactivity is attributable to unchanged irbesartan. Irbesartan is metabolised by the liver via glucuronide conjugation and oxidation. The major circulating metabolite is irbesartan glucuronide (approximately 6%). *In vitro* studies indicate that irbesartan is primarily oxidised by the cytochrome P450 enzyme CYP2C9; isoenzyme CYP3A4 has negligible effect.

Irbesartan exhibits linear and dose proportional pharmacokinetics over the dose range of 10 to 600 mg. A less than proportional increase in oral absorption at doses beyond 600 mg (twice the maximal recommended dose) was observed; the mechanism for this is unknown. Peak plasma concentrations are attained at 1.5 - 2 hours after oral administration. The total body and renal clearance are 157 - 176 and 3 - 3.5 ml/min, respectively. The terminal elimination half-life of irbesartan is 11 - 15 hours. Steady-state plasma concentrations are attained within 3 days after initiation of a once-daily dosing regimen. Limited accumulation of irbesartan (< 20%) is observed in plasma upon repeated once-daily dosing. In a study, somewhat higher plasma concentrations of irbesartan were observed in female hypertensive patients. However, there was no difference in the half-life and accumulation of irbesartan. No dosage adjustment is necessary in female patients. Irbesartan AUC and C<sub>max</sub> values were also somewhat greater in elderly subjects (≥ 65 years) than those of young subjects (18 - 40 years). However the terminal half-life was not significantly altered. No dosage adjustment is necessary in elderly patients.

Irbesartan and its metabolites are eliminated by both biliary and renal pathways. After either oral or IV administration of <sup>14</sup>C irbesartan, about 20% of the radioactivity is recovered in the urine, and the remainder in the faeces. Less than 2% of the dose is excreted in the urine as unchanged irbesartan.

### Paediatric population

The pharmacokinetics of irbesartan were evaluated in 23 hypertensive children after the administration of single and multiple daily doses of irbesartan (2 mg/kg) up to a maximum daily dose of 150 mg for four weeks. Of those 23 children, 21 were evaluable for comparison of pharmacokinetics with adults (twelve children over 12 years, nine children between 6 and 12 years). Results showed that  $C_{max}$ , AUC and clearance rates were comparable to those observed in adult patients receiving 150 mg irbesartan daily. A limited accumulation of irbesartan (18%) in plasma was observed upon repeated once daily dosing.

Renal impairment: in patients with renal impairment or those undergoing haemodialysis, the pharmacokinetic parameters of irbesartan are not significantly altered. Irbesartan is not removed by haemodialysis.

Hepatic impairment: in patients with mild to moderate cirrhosis, the pharmacokinetic parameters of irbesartan are not significantly altered.

Studies have not been performed in patients with severe hepatic impairment.

### **5.3 Preclinical safety data**

There was no evidence of abnormal systemic or target organ toxicity at clinically relevant doses. In non-clinical safety studies, high doses of irbesartan ( $\geq 250$  mg/kg/day in rats and  $\geq 100$  mg/kg/day in macaques) caused a reduction of red blood cell parameters (erythrocytes, haemoglobin, haematocrit). At very high doses ( $\geq 500$  mg/kg/day) degenerative changes in the kidney (such as interstitial nephritis, tubular distension, basophilic tubules, increased plasma concentrations of urea and creatinine) were induced by irbesartan in the rat and the macaque and are considered secondary to the hypotensive effects of the medicinal product which led to decreased renal perfusion. Furthermore, irbesartan induced hyperplasia/hypertrophy of the juxtaglomerular cells (in rats at  $\geq 90$  mg/kg/day, in macaques at  $\geq 10$  mg/kg/day). All of these changes were considered to be caused by the pharmacological action of irbesartan. For therapeutic doses of irbesartan in humans, the hyperplasia/hypertrophy of the renal juxtaglomerular cells does not appear to have any relevance.

There was no evidence of mutagenicity, clastogenicity or carcinogenicity.

Fertility and reproductive performance were not affected in studies of male and female rats even at oral doses of irbesartan causing some parental toxicity (from 50 to 650 mg/kg/day), including mortality at the highest dose. No significant effects on the number of corpora lutea, implants, or live fetuses were observed. Irbesartan did not affect survival, development, or reproduction of offspring. Studies in animals indicate that the radiolabeled irbesartan is detected in rat and rabbit fetuses. Irbesartan is excreted in the milk of lactating rats.

Animal studies with irbesartan showed transient toxic effects (increased renal pelvic cavitation, hydroureter or subcutaneous oedema) in rat foetuses, which were resolved after birth. In rabbits, abortion or early resorption were noted at doses causing significant maternal toxicity, including mortality. No teratogenic effects were observed in the rat or rabbit.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Tablet core:

Lactose monohydrate  
Microcrystalline cellulose  
Croscarmellose sodium  
Silica colloidal anhydrous  
Hypromellose  
Magnesium stearate.

Film-coating:  
Hypromellose  
Titanium dioxide (E171)  
Macrogol 400

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

4 years

## **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

## **6.5 Nature and contents of container**

Cartons of 28, 56, 90 or 98 film-coated tablets in PVC/PVDC-aluminium/PVDC blisters.  
Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

Any unused product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Pharmathen S.A.  
6, Dervenakion str.,  
15351 Pallini  
Attiki, Greece  
Tel.: +30 210 66 65 067

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/12/765/001  
EU/1/12/765/002  
EU/1/12/765/007  
EU/1/12/765/010

## **9. DATE OF FIRST AUTHORISATION / RENEWAL OF THE AUTHORISATION**

Date of first Authorization: 13 April 2012

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>

## 1. NAME OF THE MEDICINAL PRODUCT

Sabervel 150 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 150 mg of irbesartan

Excipient with known effect:

40 mg of lactose monohydrate per film-coated tablet

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet.

White, concave, round, film-coated tablet with 9 mm diameter.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Sabervel is indicated in adults for the treatment of essential hypertension.

It is also indicated for the treatment of renal disease in adult patients with hypertension and type 2 diabetes mellitus as part of an antihypertensive medicinal product regimen (see sections 4.3, 4.4, 4.5 and 5.1).

### 4.2 Posology and method of administration

#### Posology

The usual recommended initial and maintenance dose is 150 mg once daily, with or without food. Sabervel at a dose of 150 mg once daily generally provides a better 24 hour blood pressure control than 75 mg. However, initiation of therapy with 75 mg could be considered, particularly in haemodialysed patients and in the elderly over 75 years.

In patients insufficiently controlled with 150 mg once daily, the dose of Sabervel can be increased to 300 mg, or other antihypertensive agents can be added (see sections 4.3, 4.4, 4.5 and 5.1). In particular, the addition of a diuretic such as hydrochlorothiazide has been shown to have an additive effect with Sabervel (see section 4.5).

In hypertensive type 2 diabetic patients, therapy should be initiated at 150 mg irbesartan once daily and titrated up to 300 mg once daily as the preferred maintenance dose for treatment of renal disease. The demonstration of renal benefit of Sabervel in hypertensive type 2 diabetic patients is based on studies where irbesartan was used in addition to other antihypertensive agents, as needed, to reach target blood pressure (see sections 4.3, 4.4, 4.5 and 5.1).

#### Special Populations

Renal impairment: no dosage adjustment is necessary in patients with impaired renal function. A lower starting dose (75 mg) should be considered for patients undergoing haemodialysis (see section 4.4).

Hepatic impairment: no dosage adjustment is necessary in patients with mild to moderate hepatic

impairment. There is no clinical experience in patients with severe hepatic impairment.

Elderly patients: although consideration should be given to initiating therapy with 75 mg in patients over 75 years of age, dosage adjustment is not usually necessary for the elderly.

Paediatric population: the safety and efficacy of Sabervel in children aged 0 to 18 has not been established. Currently available data are described in section 4.8, 5.1 and 5.2 but no recommendation on a posology can be made.

#### Method of Administration

For oral use.

### **4.3 Contraindications**

Hypersensitivity to the active substance, or to any of the excipients (see section 6.1).  
Second and third trimesters of pregnancy (see sections 4.4 and 4.6).

The concomitant use of Sabervel with aliskiren-containing products is contraindicated in patients with diabetes mellitus or renal impairment ( $\text{GFR} < 60 \text{ ml/min/1.73 m}^2$ ) (see sections 4.5 and 5.1).

### **4.4 Special warnings and precautions for use**

Intravascular volume depletion: symptomatic hypotension, especially after the first dose, may occur in patients who are volume and/or sodium depleted by vigorous diuretic therapy, dietary salt restriction, diarrhoea or vomiting. Such conditions should be corrected before the administration of Sabervel.

Renovascular hypertension: there is an increased risk of severe hypotension and renal insufficiency when patients with bilateral renal artery stenosis or stenosis of the artery to a single functioning kidney are treated with medicinal products that affect the renin-angiotensin-aldosterone system. While this is not documented with Sabervel, a similar effect should be anticipated with angiotensin-II receptor antagonists.

Renal impairment and kidney transplantation: when Sabervel is used in patients with impaired renal function, a periodic monitoring of potassium and creatinine serum levels is recommended. There is no experience regarding the administration of Sabervel in patients with a recent kidney transplantation.

Hypertensive patients with type 2 diabetes and renal disease: the effects of irbesartan both on renal and cardiovascular events were not uniform across all subgroups, in an analysis carried out in the study with patients with advanced renal disease. In particular, they appeared less favourable in women and non-white subjects (see section 5.1).

#### Dual blockade of the renin-angiotensin-aldosterone system (RAAS)

There is evidence that the concomitant use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren increases the risk of hypotension, hyperkalaemia and decreased renal function (including acute renal failure). Dual blockade of RAAS through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is therefore not recommended (see sections 4.5 and 5.1).

If dual blockade therapy is considered absolutely necessary, this should only occur under specialist supervision and subject to frequent close monitoring of renal function, electrolytes and blood pressure. ACE-inhibitors and angiotensin II receptor blockers should not be used concomitantly in patients with diabetic nephropathy.

Hyperkalaemia: as with other medicinal products that affect the renin-angiotensin-aldosterone system, hyperkalaemia may occur during the treatment with Sabervel, especially in the presence of renal impairment, overt proteinuria due to diabetic renal disease, and/or heart failure. Close monitoring of serum potassium in patients at risk is recommended (see section 4.5).

Lithium: the combination of lithium and Sabervel is not recommended (see section 4.5).

Aortic and mitral valve stenosis, obstructive hypertrophic cardiomyopathy: as with other vasodilators, special caution is indicated in patients suffering from aortic or mitral stenosis, or obstructive hypertrophic cardiomyopathy.

Primary aldosteronism: patients with primary aldosteronism generally will not respond to antihypertensive medicinal products acting through inhibition of the renin-angiotensin system. Therefore, the use of Sabervel is not recommended.

General: in patients whose vascular tone and renal function depend predominantly on the activity of the renin-angiotensin-aldosterone system (e.g. patients with severe congestive heart failure or underlying renal disease, including renal artery stenosis), treatment with angiotensin converting enzyme inhibitors or angiotensin-II receptor antagonists that affect this system has been associated with acute hypotension, azotaemia, oliguria, or rarely acute renal failure. As with any antihypertensive agent, excessive blood pressure decrease in patients with ischaemic cardiopathy or ischaemic cardiovascular disease could result in a myocardial infarction or stroke.

As observed for angiotensin converting enzyme inhibitors, irbesartan and the other angiotensin antagonists are apparently less effective in lowering blood pressure in black people than in non-blacks, possibly because of higher prevalence of low-renin states in the black hypertensive population (see section 5.1).

Pregnancy: Angiotensin II Receptor Antagonists (AIIRAs) should not be initiated during pregnancy. Unless continued AIIRA therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIRAs should be stopped immediately, and, if appropriate, alternative therapy should be started (see sections 4.3 and 4.6).

Lactose: this medicinal product contains lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.

Paediatric population: irbesartan has been studied in paediatric populations aged 6 to 16 years old but the current data are insufficient to support an extension of the use in children until further data become available (see sections 4.8, 5.1 and 5.2).

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Diuretics and other antihypertensive agents: other antihypertensive agents may increase the hypotensive effects of irbesartan; however irbesartan has been safely administered with other antihypertensive agents, such as beta-blockers, long-acting calcium channel blockers, and thiazide diuretics. Prior treatment with high dose diuretics may result in volume depletion and a risk of hypotension when initiating therapy with Sabervel (see section 4.4).

Clinical trial data has shown that dual blockade of the renin-angiotensin-aldosterone-system (RAAS) through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is associated with a higher frequency of adverse events such as hypotension, hyperkalaemia and decreased renal function (including acute renal failure) compared to the use of a single RAAS-acting agent (see sections 4.3, 4.4 and 5.1).

Potassium supplements and potassium-sparing diuretics: based on experience with the use of other medicinal products that affect the renin-angiotensin system, concomitant use of potassium-sparing diuretics, potassium supplements, salt substitutes containing potassium or other medicinal products that may increase serum potassium levels (e.g. heparin) may lead to increases in serum potassium and is, therefore, not recommended (see section 4.4).

Lithium: reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with angiotensin converting enzyme inhibitors. Similar effects have been very rarely reported with irbesartan so far. Therefore, this combination is not recommended (see section 4.4). If the combination proves necessary, careful monitoring of serum lithium levels is recommended.

Non-steroidal anti-inflammatory drugs: when angiotensin II antagonists are administered simultaneously with non-steroidal anti-inflammatory drugs (i.e. selective COX-2 inhibitors, acetylsalicylic acid (> 3 g/day) and non-selective NSAIDs), attenuation of the antihypertensive effect may occur.

As with ACE inhibitors, concomitant use of angiotensin II antagonists and NSAIDs may lead to an increased risk of worsening of renal function, including possible acute renal failure, and an increase in serum potassium, especially in patients with poor pre-existing renal function. The combination should be administered with caution, especially in the elderly. Patients should be adequately hydrated and consideration should be given to monitoring renal function after initiation of concomitant therapy, and periodically thereafter.

Additional information on irbesartan interactions: in clinical studies, the pharmacokinetic of irbesartan is not affected by hydrochlorothiazide. Irbesartan is mainly metabolised by CYP2C9 and to a lesser extent by glucuronidation. No significant pharmacokinetic or pharmacodynamic interactions were observed when irbesartan was coadministered with warfarin, a medicinal product metabolised by CYP2C9. The effects of CYP2C9 inducers such as rifampicin on the pharmacokinetic of irbesartan have not been evaluated. The pharmacokinetic of digoxin was not altered by coadministration of irbesartan.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy:

The use of AIIRAs is not recommended during the first trimester of pregnancy (see section 4.4). The use of AIIRAs is contraindicated during the second and third trimesters of pregnancy (see sections 4.3 and 4.4).

Epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded. Whilst there is no controlled epidemiological data on the risk with Angiotensin II Receptor Antagonists (AIIRAs), similar risks may exist for this class of drugs. Unless continued AIIRA therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIRAs should be stopped immediately, and, if appropriate, alternative therapy should be started.

Exposure to AIIRA therapy during the second and third trimesters is known to induce human fetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia). (See section 5.3).

Should exposure to AIIRAs have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended.

Infants whose mothers have taken AIIRAs should be closely observed for hypotension (see sections 4.3 and 4.4).

##### Breast-feeding:

Because no information is available regarding the use of Sabervel during breast-feeding, Sabervel is not recommended and alternative treatments with better established safety profiles during breast-feeding are preferable, especially while nursing a newborn or preterm infant.

It is unknown whether irbesartan or its metabolites are excreted in human milk.



Available pharmacodynamic/toxicological data in rats have shown excretion of irbesartan or its metabolites in milk (for details see 5.3).

### Fertility

Irbesartan had no effect upon fertility of treated rats and their offspring up to the dose levels inducing the first signs of parental toxicity (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

No studies on the effects on the ability to drive and use machines have been performed. Based on its pharmacodynamic properties, irbesartan is unlikely to affect this ability. When driving vehicles or operating machines, it should be taken into account that dizziness or weariness may occur during treatment.

### **4.8 Undesirable effects**

In placebo-controlled trials in patients with hypertension, the overall incidence of adverse events did not differ between the irbesartan (56.2%) and the placebo groups (56.5%). Discontinuation due to any clinical or laboratory adverse event was less frequent for irbesartan-treated patients (3.3%) than for placebo-treated patients (4.5%). The incidence of adverse events was not related to dose (in the recommended dose range), gender, age, race, or duration of treatment.

In diabetic hypertensive patients with microalbuminuria and normal renal function, orthostatic dizziness and orthostatic hypotension were reported in 0.5% of the patients (i.e., uncommon) but in excess of placebo.

The following table presents the adverse drug reactions that were reported in placebo-controlled trials in which 1,965 hypertensive patients received irbesartan. Terms marked with a star (\*) refer to the adverse reactions that were additionally reported in > 2% of diabetic hypertensive patients with chronic renal insufficiency and overt proteinuria and in excess of placebo.

The frequency of adverse reactions listed below is defined using the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Adverse reactions additionally reported from post-marketing experience are also listed. These adverse reactions are derived from spontaneous reports.

#### *Immune system disorders:*

Not known: hypersensitivity reactions such as angioedema, rash, urticaria

#### *Metabolism and nutrition disorders:*

Not known: hyperkalaemia

#### *Nervous system disorders:*

Common: dizziness, orthostatic dizziness\*

Not known: vertigo, headache

#### *Ear and labyrinth disorder:*

Not known: tinnitus

#### *Cardiac disorders:*

Uncommon: tachycardia

#### *Vascular disorders:*

Common: orthostatic hypotension\*  
Uncommon: flushing

*Respiratory, thoracic and mediastinal disorders:*

Uncommon: cough

*Gastrointestinal disorders:*

Common: nausea/vomiting  
Uncommon: diarrhoea, dyspepsia/heartburn  
Not known: dysgeusia

*Hepatobiliary disorders:*

Uncommon: jaundice  
Not known: hepatitis, abnormal liver function

*Skin and subcutaneous tissue disorders:*

Not known: leukocytoclastic vasculitis

*Musculoskeletal and connective tissue disorders:*

Common: musculoskeletal pain\*  
Not known: arthralgia, myalgia (in some cases associated with increased plasma creatine kinase levels), muscle cramps

*Renal and urinary disorders:*

Not known: impaired renal function including cases of renal failure in patients at risk (see section 4.4)

*Reproductive system and breast disorders:*

Uncommon: sexual dysfunction

*General disorders and administration site conditions:*

Common: fatigue  
Uncommon: chest pain

Investigations:

Very common: Hyperkalaemia\* occurred more often in diabetic patients treated with irbesartan than with placebo. In diabetic hypertensive patients with microalbuminuria and normal renal function, hyperkalaemia ( $\geq 5.5$  mEq/L) occurred in 29.4% of the patients in the irbesartan 300 mg group and 22% of the patients in the placebo group. In diabetic hypertensive patients with chronic renal insufficiency and overt proteinuria, hyperkalaemia ( $\geq 5.5$  mEq/L) occurred in 46.3% of the patients in the irbesartan group and 26.3% of the patients in the placebo group.

Common: significant increases in plasma creatine kinase were commonly observed (1.7%) in irbesartan treated subjects. None of these increases were associated with identifiable clinical musculoskeletal events. In 1.7% of hypertensive patients with advanced diabetic renal disease treated with irbesartan, a decrease in haemoglobin\*, which was not clinically significant, has been observed.

Paediatric population:

In a randomised trial of 318 hypertensive children and adolescents aged 6 to 16 years, the following adverse reactions occurred in the 3-week double-blind phase: headache (7.9%), hypotension (2.2%), dizziness (1.9%), cough (0.9%). In the 26-week open-label period of this trial the most frequent laboratory abnormalities observed were creatinine increases (6.5%) and elevated CK values in 2% of child recipients.

## 4.9 Overdose

Experience in adults exposed to doses of up to 900 mg/day for 8 weeks revealed no toxicity. The most likely manifestations of overdose are expected to be hypotension and tachycardia; bradycardia might also occur from overdose. No specific information is available on the treatment of overdose with Sabervel. The patient should be closely monitored, and the treatment should be symptomatic and supportive. Suggested measures include induction of emesis and/or gastric lavage. Activated charcoal may be useful in the treatment of overdose. Irbesartan is not removed by haemodialysis.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Angiotensin-II antagonists, plain.

ATC code: C09C A04.

Mechanism of action: Irbesartan is a potent, orally active, selective angiotensin-II receptor (type AT<sub>1</sub>) antagonist. It is expected to block all actions of angiotensin-II mediated by the AT<sub>1</sub> receptor, regardless of the source or route of synthesis of angiotensin-II. The selective antagonism of the angiotensin-II (AT<sub>1</sub>) receptors results in increases in plasma renin levels and angiotensin-II levels, and a decrease in plasma aldosterone concentration. Serum potassium levels are not significantly affected by irbesartan alone at the recommended doses. Irbesartan does not inhibit ACE (kininase-II), an enzyme which generates angiotensin-II and also degrades bradykinin into inactive metabolites.

Irbesartan does not require metabolic activation for its activity.

Clinical efficacy:

#### Hypertension

Irbesartan lowers blood pressure with minimal change in heart rate. The decrease in blood pressure is dose-related for once a day doses with a tendency towards plateau at doses above 300 mg. Doses of 150-300 mg once daily lower supine or seated blood pressures at trough (i.e. 24 hours after dosing) by an average of 8-13/5-8 mm Hg (systolic/diastolic) greater than those associated with placebo. Peak reduction of blood pressure is achieved within 3-6 hours after administration and the blood pressure lowering effect is maintained for at least 24 hours. At 24 hours the reduction of blood pressure was 60-70% of the corresponding peak diastolic and systolic responses at the recommended doses. Once daily dosing with 150 mg produced trough and mean 24 hour responses similar to twice daily dosing on the same total dose.

The blood pressure lowering effect of irbesartan is evident within 1-2 weeks, with the maximal effect occurring by 4-6 weeks after start of therapy. The antihypertensive effects are maintained during long term therapy. After withdrawal of therapy, blood pressure gradually returns toward baseline. Rebound hypertension has not been observed.

The blood pressure lowering effects of irbesartan and thiazide-type diuretics are additive. In patients not adequately controlled by irbesartan alone, the addition of a low dose of hydrochlorothiazide (12.5 mg) to irbesartan once daily results in a further placebo-adjusted blood pressure reduction at trough of 7-10/3-6 mm Hg (systolic/diastolic).

The efficacy of irbesartan is not influenced by age or gender. As is the case with other medicinal products that affect the renin-angiotensin system, black hypertensive patients have notably less response to irbesartan monotherapy. When irbesartan is administered concomitantly with a low dose of hydrochlorothiazide (e.g. 12.5 mg daily), the antihypertensive response in black patients approaches that of white patients.

There is no clinically important effect on serum uric acid or urinary uric acid secretion.

#### Paediatric population

Reduction of blood pressure with 0.5 mg/kg (low), 1.5 mg/kg (medium) and 4.5 mg/kg (high) target titrated doses of irbesartan was evaluated in 318 hypertensive or at risk (diabetic, family history of

hypertension) children and adolescents aged 6 to 16 years over a three week period. At the end of the three weeks the mean reduction from baseline in the primary efficacy variable, trough seated systolic blood pressure (SeSBP) was 11.7 mmHg (low dose), 9.3 mmHg (medium dose), 13.2 mmHg (high dose). No significant difference was apparent between these doses. Adjusted mean change of trough seated diastolic blood pressure (SeDBP) was as follows: 3.8 mmHg (low dose), 3.2 mmHg (medium dose), 5.6 mmHg (high dose). Over a subsequent two week period where patients were re-randomized to either active medicinal product or placebo, patients on placebo had increases of 2.4 and 2.0 mmHg in SeSBP and SeDBP compared to +0.1 and -0.3 mmHg changes respectively in those on all doses of irbesartan (see section 4.2).

#### Hypertension and type 2 diabetes with renal disease

The “Irbesartan Diabetic Nephropathy Trial (IDNT)” shows that irbesartan decreases the progression of renal disease in patients with chronic renal insufficiency and overt proteinuria. IDNT was a double blind, controlled, morbidity and mortality trial comparing irbesartan, amlodipine and placebo. In 1,715 hypertensive patients with type 2 diabetes, proteinuria  $\geq 900$  mg/day and serum creatinine ranging from 1.0-3.0 mg/dl, the long-term effects (mean 2.6 years) of irbesartan on the progression of renal disease and all-cause mortality were examined. Patients were titrated from 75 mg to a maintenance dose of 300 mg irbesartan, from 2.5 mg to 10 mg amlodipine, or placebo as tolerated.

Patients in all treatment groups typically received between 2 and 4 antihypertensive agents (e.g., diuretics, beta blockers, alpha blockers) to reach a predefined blood pressure goal of  $\leq 135/85$  mmHg or a 10 mmHg reduction in systolic pressure if baseline was  $> 160$  mmHg. Sixty per cent (60%) of patients in the placebo group reached this target blood pressure whereas this figure was 76% and 78% in the irbesartan and amlodipine groups respectively.

Irbesartan significantly reduced the relative risk in the primary combined endpoint of doubling serum creatinine, end-stage renal disease (ESRD) or all-cause mortality. Approximately 33% of patients in the irbesartan group reached the primary renal composite endpoint compared to 39% and 41% in the placebo and amlodipine groups [20% relative risk reduction versus placebo ( $p = 0.024$ ) and 23% relative risk reduction compared to amlodipine ( $p = 0.006$ )]. When the individual components of the primary endpoint were analysed, no effect in all cause mortality was observed, while a positive trend in the reduction in ESRD and a significant reduction in doubling of serum creatinine were observed.

Subgroups consisting of gender, race, age, duration of diabetes, baseline blood pressure, serum creatinine, and albumin excretion rate were assessed for treatment effect. In the female and black subgroups which represented 32% and 26% of the overall study population respectively, a renal benefit was not evident, although the confidence intervals do not exclude it. As for the secondary endpoint of fatal and non-fatal cardiovascular events, there was no difference among the three groups in the overall population, although an increased incidence of non-fatal MI was seen for women and a decreased incidence of non-fatal MI was seen in males in the irbesartan group versus the placebo-based regimen. An increased incidence of non-fatal MI and stroke was seen in females in the irbesartan-based regimen versus the amlodipine-based regimen, while hospitalization due to heart failure was reduced in the overall population. However, no proper explanation for these findings in women has been identified.

The study of the “Effects of Irbesartan on Microalbuminuria in Hypertensive Patients with type 2 Diabetes Mellitus (IRMA 2)” shows that irbesartan 300 mg delays progression to overt proteinuria in patients with microalbuminuria. IRMA 2 was a placebo-controlled double blind morbidity study in 590 patients with type 2 diabetes, microalbuminuria (30-300 mg/day) and normal renal function (serum creatinine  $\leq 1.5$  mg/dl in males and  $< 1.1$  mg/dl in females). The study examined the long-term effects (2 years) of irbesartan on the progression to clinical (overt) proteinuria (urinary albumin excretion rate (UAER)  $> 300$  mg/day, and an increase in UAER of at least 30% from baseline). The predefined blood pressure goal was  $\leq 135/85$  mmHg. Additional antihypertensive agents (excluding ACE inhibitors, angiotensin II receptor antagonists and dihydropyridine calcium blockers) were added as needed to help achieve the blood pressure goal. While similar blood pressure was achieved in all treatment groups, fewer subjects in the irbesartan 300 mg group (5.2%) than in the placebo (14.9%) or in the irbesartan 150 mg group (9.7%) reached the endpoint of overt proteinuria, demonstrating a 70% relative risk reduction versus placebo ( $p = 0.0004$ ) for the higher dose. An accompanying improvement in the glomerular filtration rate (GFR) was not observed during the first three months of

treatment. The slowing in the progression to clinical proteinuria was evident as early as three months and continued over the 2 year period. Regression to normoalbuminuria (< 30 mg/day) was more frequent in the irbesartan 300 mg group (34%) than in the placebo group (21%).

Two large randomised, controlled trials (ONTARGET (ONgoing Telmisartan Alone and in combination with Ramipril Global Endpoint Trial) and VA NEPHRON-D (The Veterans Affairs Nephropathy in Diabetes)) have examined the use of the combination of an ACE-inhibitor with an angiotensin II receptor blocker.

ONTARGET was a study conducted in patients with a history of cardiovascular or cerebrovascular disease, or type 2 diabetes mellitus accompanied by evidence of end-organ damage. VA NEPHRON-D was a study in patients with type 2 diabetes mellitus and diabetic nephropathy.

These studies have shown no significant beneficial effect on renal and/or cardiovascular outcomes and mortality, while an increased risk of hyperkalaemia, acute kidney injury and/or hypotension as compared to monotherapy was observed. Given their similar pharmacodynamic properties, these results are also relevant for other ACE-inhibitors and angiotensin II receptor blockers.

ACE-inhibitors and angiotensin II receptor blockers should therefore not be used concomitantly in patients with diabetic nephropathy.

ALTITUDE (Aliskiren Trial in Type 2 Diabetes Using Cardiovascular and Renal Disease Endpoints) was a study designed to test the benefit of adding aliskiren to a standard therapy of an ACE-inhibitor or an angiotensin II receptor blocker in patients with type 2 diabetes mellitus and chronic kidney disease, cardiovascular disease, or both. The study was terminated early because of an increased risk of adverse outcomes. Cardiovascular death and stroke were both numerically more frequent in the aliskiren group than in the placebo group and adverse events and serious adverse events of interest (hyperkalaemia, hypotension and renal dysfunction) were more frequently reported in the aliskiren group than in the placebo group.

## 5.2 Pharmacokinetic properties

After oral administration, irbesartan is well absorbed: studies of absolute bioavailability gave values of approximately 60-80%. Concomitant food intake does not significantly influence the bioavailability of irbesartan. Plasma protein binding is approximately 96%, with negligible binding to cellular blood components. The volume of distribution is 53 - 93 litres. Following oral or intravenous administration of <sup>14</sup>C irbesartan, 80-85% of the circulating plasma radioactivity is attributable to unchanged irbesartan. Irbesartan is metabolised by the liver via glucuronide conjugation and oxidation. The major circulating metabolite is irbesartan glucuronide (approximately 6%). *In vitro* studies indicate that irbesartan is primarily oxidised by the cytochrome P450 enzyme CYP2C9; isoenzyme CYP3A4 has negligible effect.

Irbesartan exhibits linear and dose proportional pharmacokinetics over the dose range of 10 to 600 mg. A less than proportional increase in oral absorption at doses beyond 600 mg (twice the maximal recommended dose) was observed; the mechanism for this is unknown. Peak plasma concentrations are attained at 1.5 - 2 hours after oral administration. The total body and renal clearance are 157 - 176 and 3 - 3.5 ml/min, respectively. The terminal elimination half-life of irbesartan is 11 - 15 hours. Steady-state plasma concentrations are attained within 3 days after initiation of a once-daily dosing regimen. Limited accumulation of irbesartan (< 20%) is observed in plasma upon repeated once-daily dosing. In a study, somewhat higher plasma concentrations of irbesartan were observed in female hypertensive patients. However, there was no difference in the half-life and accumulation of irbesartan. No dosage adjustment is necessary in female patients. Irbesartan AUC and C<sub>max</sub> values were also somewhat greater in elderly subjects (≥ 65 years) than those of young subjects (18 - 40 years). However the terminal half-life was not significantly altered. No dosage adjustment is necessary in elderly patients.

Irbesartan and its metabolites are eliminated by both biliary and renal pathways. After either oral or IV administration of <sup>14</sup>C irbesartan, about 20% of the radioactivity is recovered in the urine, and the remainder in the faeces. Less than 2% of the dose is excreted in the urine as unchanged irbesartan.

### Paediatric population

The pharmacokinetics of irbesartan were evaluated in 23 hypertensive children after the administration of single and multiple daily doses of irbesartan (2 mg/kg) up to a maximum daily dose of 150 mg for four weeks. Of those 23 children, 21 were evaluable for comparison of pharmacokinetics with adults (twelve children over 12 years, nine children between 6 and 12 years). Results showed that  $C_{max}$ , AUC and clearance rates were comparable to those observed in adult patients receiving 150 mg irbesartan daily. A limited accumulation of irbesartan (18%) in plasma was observed upon repeated once daily dosing.

Renal impairment: in patients with renal impairment or those undergoing haemodialysis, the pharmacokinetic parameters of irbesartan are not significantly altered. Irbesartan is not removed by haemodialysis.

Hepatic impairment: in patients with mild to moderate cirrhosis, the pharmacokinetic parameters of irbesartan are not significantly altered.

Studies have not been performed in patients with severe hepatic impairment.

### **5.3 Preclinical safety data**

There was no evidence of abnormal systemic or target organ toxicity at clinically relevant doses. In non-clinical safety studies, high doses of irbesartan ( $\geq 250$  mg/kg/day in rats and  $\geq 100$  mg/kg/day in macaques) caused a reduction of red blood cell parameters (erythrocytes, haemoglobin, haematocrit). At very high doses ( $\geq 500$  mg/kg/day) degenerative changes in the kidney (such as interstitial nephritis, tubular distension, basophilic tubules, increased plasma concentrations of urea and creatinine) were induced by irbesartan in the rat and the macaque and are considered secondary to the hypotensive effects of the medicinal product which led to decreased renal perfusion. Furthermore, irbesartan induced hyperplasia/hypertrophy of the juxtaglomerular cells (in rats at  $\geq 90$  mg/kg/day, in macaques at  $\geq 10$  mg/kg/day). All of these changes were considered to be caused by the pharmacological action of irbesartan. For therapeutic doses of irbesartan in humans, the hyperplasia/hypertrophy of the renal juxtaglomerular cells does not appear to have any relevance.

There was no evidence of mutagenicity, clastogenicity or carcinogenicity.

Fertility and reproductive performance were not affected in studies of male and female rats even at oral doses of irbesartan causing some parental toxicity (from 50 to 650 mg/kg/day), including mortality at the highest dose. No significant effects on the number of corpora lutea, implants, or live fetuses were observed. Irbesartan did not affect survival, development, or reproduction of offspring. Studies in animals indicate that the radiolabeled irbesartan is detected in rat and rabbit fetuses. Irbesartan is excreted in the milk of lactating rats.

Animal studies with irbesartan showed transient toxic effects (increased renal pelvic cavitation, hydroureter or subcutaneous oedema) in rat foetuses, which were resolved after birth. In rabbits, abortion or early resorption were noted at doses causing significant maternal toxicity, including mortality. No teratogenic effects were observed in the rat or rabbit.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Tablet core:

Lactose monohydrate  
Microcrystalline cellulose  
Croscarmellose sodium  
Silica colloidal anhydrous  
Hypromellose  
Magnesium stearate.

Film-coating:  
Hypromellose  
Titanium dioxide (E171)  
Macrogol 400

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

4 years

## **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

## **6.5 Nature and contents of container**

Cartons of 28, 56, 90 or 98 film-coated tablets in PVC/PVDC-aluminium/PVDC blisters.  
Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

Any unused product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Pharmathen S.A.  
6, Dervenakion str.,  
15351 Pallini  
Attiki, Greece  
Tel.: +30 210 66 65 067

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/12/765/003  
EU/1/12/765/004  
EU/1/12/765/008  
EU/1/12/765/011

## **9. DATE OF FIRST AUTHORISATION / RENEWAL OF THE AUTHORISATION**

Date of first Authorization: 13 April 2012

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>

## 1. NAME OF THE MEDICINAL PRODUCT

Sabervel 300 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 300 mg of irbesartan

Excipient with known effect:

80 mg of lactose monohydrate per film-coated tablet

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet.

White, concave, round, film-coated tablet with 11 mm diameter.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Sabervel is indicated in adults for the treatment of essential hypertension.

It is also indicated for the treatment of renal disease in adult patients with hypertension and type 2 diabetes mellitus as part of an antihypertensive medicinal product regimen (see sections 4.3, 4.4, 4.5 and 5.1).

### 4.2 Posology and method of administration

#### Posology

The usual recommended initial and maintenance dose is 150 mg once daily, with or without food. Sabervel at a dose of 150 mg once daily generally provides a better 24 hour blood pressure control than 75 mg. However, initiation of therapy with 75 mg could be considered, particularly in haemodialysed patients and in the elderly over 75 years.

In patients insufficiently controlled with 150 mg once daily, the dose of Sabervel can be increased to 300 mg, or other antihypertensive agents can be added (see sections 4.3, 4.4, 4.5 and 5.1). In particular, the addition of a diuretic such as hydrochlorothiazide has been shown to have an additive effect with Sabervel (see section 4.5).

In hypertensive type 2 diabetic patients, therapy should be initiated at 150 mg irbesartan once daily and titrated up to 300 mg once daily as the preferred maintenance dose for treatment of renal disease. The demonstration of renal benefit of Sabervel in hypertensive type 2 diabetic patients is based on studies where irbesartan was used in addition to other antihypertensive agents, as needed, to reach target blood pressure (see sections 4.3, 4.4, 4.5 and 5.1).

#### Special Populations

Renal impairment: no dosage adjustment is necessary in patients with impaired renal function. A lower starting dose (75 mg) should be considered for patients undergoing haemodialysis (see section 4.4).

Hepatic impairment: no dosage adjustment is necessary in patients with mild to moderate hepatic



impairment. There is no clinical experience in patients with severe hepatic impairment.

Elderly patients: although consideration should be given to initiating therapy with 75 mg in patients over 75 years of age, dosage adjustment is not usually necessary for the elderly.

Paediatric population: the safety and efficacy of Sabervel in children aged 0 to 18 has not been established. Currently available data are described in section 4.8, 5.1 and 5.2 but no recommendation on a posology can be made.

#### Method of Administration

For oral use.

### **4.3 Contraindications**

Hypersensitivity to the active substance, or to any of the excipients (see section 6.1).  
Second and third trimesters of pregnancy (see sections 4.4 and 4.6).

The concomitant use of Sabervel with aliskiren-containing products is contraindicated in patients with diabetes mellitus or renal impairment ( $\text{GFR} < 60 \text{ ml/min/1.73 m}^2$ ) (see sections 4.5 and 5.1).

### **4.4 Special warnings and precautions for use**

Intravascular volume depletion: symptomatic hypotension, especially after the first dose, may occur in patients who are volume and/or sodium depleted by vigorous diuretic therapy, dietary salt restriction, diarrhoea or vomiting. Such conditions should be corrected before the administration of Sabervel.

Renovascular hypertension: there is an increased risk of severe hypotension and renal insufficiency when patients with bilateral renal artery stenosis or stenosis of the artery to a single functioning kidney are treated with medicinal products that affect the renin-angiotensin-aldosterone system. While this is not documented with Sabervel, a similar effect should be anticipated with angiotensin-II receptor antagonists.

Renal impairment and kidney transplantation: when Sabervel is used in patients with impaired renal function, a periodic monitoring of potassium and creatinine serum levels is recommended. There is no experience regarding the administration of Sabervel in patients with a recent kidney transplantation.

Hypertensive patients with type 2 diabetes and renal disease: the effects of irbesartan both on renal and cardiovascular events were not uniform across all subgroups, in an analysis carried out in the study with patients with advanced renal disease. In particular, they appeared less favourable in women and non-white subjects (see section 5.1).

#### Dual blockade of the renin-angiotensin-aldosterone system (RAAS)

There is evidence that the concomitant use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren increases the risk of hypotension, hyperkalaemia and decreased renal function (including acute renal failure). Dual blockade of RAAS through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is therefore not recommended (see sections 4.5 and 5.1).

If dual blockade therapy is considered absolutely necessary, this should only occur under specialist supervision and subject to frequent close monitoring of renal function, electrolytes and blood pressure. ACE-inhibitors and angiotensin II receptor blockers should not be used concomitantly in patients with diabetic nephropathy.

Hyperkalaemia: as with other medicinal products that affect the renin-angiotensin-aldosterone system, hyperkalaemia may occur during the treatment with Sabervel, especially in the presence of renal impairment, overt proteinuria due to diabetic renal disease, and/or heart failure. Close monitoring of serum potassium in patients at risk is recommended (see section 4.5).

Lithium: the combination of lithium and Sabervel is not recommended (see section 4.5).

Aortic and mitral valve stenosis, obstructive hypertrophic cardiomyopathy: as with other vasodilators, special caution is indicated in patients suffering from aortic or mitral stenosis, or obstructive hypertrophic cardiomyopathy.

Primary aldosteronism: patients with primary aldosteronism generally will not respond to antihypertensive medicinal products acting through inhibition of the renin-angiotensin system. Therefore, the use of Sabervel is not recommended.

General: in patients whose vascular tone and renal function depend predominantly on the activity of the renin-angiotensin-aldosterone system (e.g. patients with severe congestive heart failure or underlying renal disease, including renal artery stenosis), treatment with angiotensin converting enzyme inhibitors or angiotensin-II receptor antagonists that affect this system has been associated with acute hypotension, azotaemia, oliguria, or rarely acute renal failure. As with any antihypertensive agent, excessive blood pressure decrease in patients with ischaemic cardiopathy or ischaemic cardiovascular disease could result in a myocardial infarction or stroke.

As observed for angiotensin converting enzyme inhibitors, irbesartan and the other angiotensin antagonists are apparently less effective in lowering blood pressure in black people than in non-blacks, possibly because of higher prevalence of low-renin states in the black hypertensive population (see section 5.1).

Pregnancy: Angiotensin II Receptor Antagonists (AIIRAs) should not be initiated during pregnancy. Unless continued AIIRA therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIRAs should be stopped immediately, and, if appropriate, alternative therapy should be started (see sections 4.3 and 4.6).

Lactose: this medicinal product contains lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product.

Paediatric population: irbesartan has been studied in paediatric populations aged 6 to 16 years old but the current data are insufficient to support an extension of the use in children until further data become available (see sections 4.8, 5.1 and 5.2).

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Diuretics and other antihypertensive agents: other antihypertensive agents may increase the hypotensive effects of irbesartan; however irbesartan has been safely administered with other antihypertensive agents, such as beta-blockers, long-acting calcium channel blockers, and thiazide diuretics. Prior treatment with high dose diuretics may result in volume depletion and a risk of hypotension when initiating therapy with Sabervel (see section 4.4).

Clinical trial data has shown that dual blockade of the renin-angiotensin-aldosterone-system (RAAS) through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is associated with a higher frequency of adverse events such as hypotension, hyperkalaemia and decreased renal function (including acute renal failure) compared to the use of a single RAAS-acting agent (see sections 4.3, 4.4 and 5.1).

Potassium supplements and potassium-sparing diuretics: based on experience with the use of other medicinal products that affect the renin-angiotensin system, concomitant use of potassium-sparing diuretics, potassium supplements, salt substitutes containing potassium or other medicinal products that may increase serum potassium levels (e.g. heparin) may lead to increases in serum potassium and is, therefore, not recommended (see section 4.4).

Lithium: reversible increases in serum lithium concentrations and toxicity have been reported during

concomitant administration of lithium with angiotensin converting enzyme inhibitors. Similar effects have been very rarely reported with irbesartan so far. Therefore, this combination is not recommended (see section 4.4). If the combination proves necessary, careful monitoring of serum lithium levels is recommended.

Non-steroidal anti-inflammatory drugs: when angiotensin II antagonists are administered simultaneously with non-steroidal anti-inflammatory drugs (i.e. selective COX-2 inhibitors, acetylsalicylic acid (> 3 g/day) and non-selective NSAIDs), attenuation of the antihypertensive effect may occur.

As with ACE inhibitors, concomitant use of angiotensin II antagonists and NSAIDs may lead to an increased risk of worsening of renal function, including possible acute renal failure, and an increase in serum potassium, especially in patients with poor pre-existing renal function. The combination should be administered with caution, especially in the elderly. Patients should be adequately hydrated and consideration should be given to monitoring renal function after initiation of concomitant therapy, and periodically thereafter.

Additional information on irbesartan interactions: in clinical studies, the pharmacokinetic of irbesartan is not affected by hydrochlorothiazide. Irbesartan is mainly metabolised by CYP2C9 and to a lesser extent by glucuronidation. No significant pharmacokinetic or pharmacodynamic interactions were observed when irbesartan was coadministered with warfarin, a medicinal product metabolised by CYP2C9. The effects of CYP2C9 inducers such as rifampicin on the pharmacokinetic of irbesartan have not been evaluated. The pharmacokinetic of digoxin was not altered by coadministration of irbesartan.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy:

The use of AIIRAs is not recommended during the first trimester of pregnancy (see section 4.4). The use of AIIRAs is contraindicated during the second and third trimesters of pregnancy (see sections 4.3 and 4.4).

Epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded. Whilst there is no controlled epidemiological data on the risk with Angiotensin II Receptor Antagonists (AIIRAs), similar risks may exist for this class of drugs. Unless continued AIIRA therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIRAs should be stopped immediately, and, if appropriate, alternative therapy should be started.

Exposure to AIIRA therapy during the second and third trimesters is known to induce human fetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia). (See section 5.3).

Should exposure to AIIRAs have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended.

Infants whose mothers have taken AIIRAs should be closely observed for hypotension (see sections 4.3 and 4.4).

##### Breast-feeding:

Because no information is available regarding the use of Sabervel during breast-feeding, Sabervel is not recommended and alternative treatments with better established safety profiles during breast-feeding are preferable, especially while nursing a newborn or preterm infant.

It is unknown whether irbesartan or its metabolites are excreted in human milk.

Available pharmacodynamic/toxicological data in rats have shown excretion of irbesartan or its

metabolites in milk (for details see 5.3).

### Fertility

Irbesartan had no effect upon fertility of treated rats and their offspring up to the dose levels inducing the first signs of parental toxicity (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

No studies on the effects on the ability to drive and use machines have been performed. Based on its pharmacodynamic properties, irbesartan is unlikely to affect this ability. When driving vehicles or operating machines, it should be taken into account that dizziness or weariness may occur during treatment.

### **4.8 Undesirable effects**

In placebo-controlled trials in patients with hypertension, the overall incidence of adverse events did not differ between the irbesartan (56.2%) and the placebo groups (56.5%). Discontinuation due to any clinical or laboratory adverse event was less frequent for irbesartan-treated patients (3.3%) than for placebo-treated patients (4.5%). The incidence of adverse events was not related to dose (in the recommended dose range), gender, age, race, or duration of treatment.

In diabetic hypertensive patients with microalbuminuria and normal renal function, orthostatic dizziness and orthostatic hypotension were reported in 0.5% of the patients (i.e., uncommon) but in excess of placebo.

The following table presents the adverse drug reactions that were reported in placebo-controlled trials in which 1,965 hypertensive patients received irbesartan. Terms marked with a star (\*) refer to the adverse reactions that were additionally reported in > 2% of diabetic hypertensive patients with chronic renal insufficiency and overt proteinuria and in excess of placebo.

The frequency of adverse reactions listed below is defined using the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Adverse reactions additionally reported from post-marketing experience are also listed. These adverse reactions are derived from spontaneous reports.

#### *Immune system disorders:*

Not known: hypersensitivity reactions such as angioedema, rash, urticaria

#### *Metabolism and nutrition disorders:*

Not known: hyperkalaemia

#### *Nervous system disorders:*

Common: dizziness, orthostatic dizziness\*

Not known: vertigo, headache

#### *Ear and labyrinth disorder:*

Not known: tinnitus

#### *Cardiac disorders:*

Uncommon: tachycardia

#### *Vascular disorders:*

Common: orthostatic hypotension\*

Uncommon: flushing

*Respiratory, thoracic and mediastinal disorders:*

Uncommon: cough

*Gastrointestinal disorders:*

Common: nausea/vomiting

Uncommon: diarrhoea, dyspepsia/heartburn

Not known: dysgeusia

*Hepatobiliary disorders:*

Uncommon: jaundice

Not known: hepatitis, abnormal liver function

*Skin and subcutaneous tissue disorders:*

Not known: leukocytoclastic vasculitis

*Musculoskeletal and connective tissue disorders:*

Common: musculoskeletal pain\*

Not known: arthralgia, myalgia (in some cases associated with increased plasma creatine kinase levels), muscle cramps

*Renal and urinary disorders:*

Not known: impaired renal function including cases of renal failure in patients at risk (see section 4.4)

*Reproductive system and breast disorders:*

Uncommon: sexual dysfunction

*General disorders and administration site conditions:*

Common: fatigue

Uncommon: chest pain

Investigations:

Very common: Hyperkalaemia\* occurred more often in diabetic patients treated with irbesartan than with placebo. In diabetic hypertensive patients with microalbuminuria and normal renal function, hyperkalaemia ( $\geq 5.5$  mEq/L) occurred in 29.4% of the patients in the irbesartan 300 mg group and 22% of the patients in the placebo group. In diabetic hypertensive patients with chronic renal insufficiency and overt proteinuria, hyperkalaemia ( $\geq 5.5$  mEq/L) occurred in 46.3% of the patients in the irbesartan group and 26.3% of the patients in the placebo group.

Common: significant increases in plasma creatine kinase were commonly observed (1.7%) in irbesartan treated subjects. None of these increases were associated with identifiable clinical musculoskeletal events. In 1.7% of hypertensive patients with advanced diabetic renal disease treated with irbesartan, a decrease in haemoglobin\*, which was not clinically significant, has been observed.

Paediatric population:

In a randomised trial of 318 hypertensive children and adolescents aged 6 to 16 years, the following adverse reactions occurred in the 3-week double-blind phase: headache (7.9%), hypotension (2.2%), dizziness (1.9%), cough (0.9%). In the 26-week open-label period of this trial the most frequent laboratory abnormalities observed were creatinine increases (6.5%) and elevated CK values in 2% of child recipients.

## 4.9 Overdose

Experience in adults exposed to doses of up to 900 mg/day for 8 weeks revealed no toxicity. The most likely manifestations of overdose are expected to be hypotension and tachycardia; bradycardia might also occur from overdose. No specific information is available on the treatment of overdose with Sabervel. The patient should be closely monitored, and the treatment should be symptomatic and supportive. Suggested measures include induction of emesis and/or gastric lavage. Activated charcoal may be useful in the treatment of overdose. Irbesartan is not removed by haemodialysis.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Angiotensin-II antagonists, plain.

ATC code: C09C A04.

Mechanism of action: Irbesartan is a potent, orally active, selective angiotensin-II receptor (type AT<sub>1</sub>) antagonist. It is expected to block all actions of angiotensin-II mediated by the AT<sub>1</sub> receptor, regardless of the source or route of synthesis of angiotensin-II. The selective antagonism of the angiotensin-II (AT<sub>1</sub>) receptors results in increases in plasma renin levels and angiotensin-II levels, and a decrease in plasma aldosterone concentration. Serum potassium levels are not significantly affected by irbesartan alone at the recommended doses. Irbesartan does not inhibit ACE (kininase-II), an enzyme which generates angiotensin-II and also degrades bradykinin into inactive metabolites.

Irbesartan does not require metabolic activation for its activity.

Clinical efficacy:

#### Hypertension

Irbesartan lowers blood pressure with minimal change in heart rate. The decrease in blood pressure is dose-related for once a day doses with a tendency towards plateau at doses above 300 mg. Doses of 150-300 mg once daily lower supine or seated blood pressures at trough (i.e. 24 hours after dosing) by an average of 8-13/5-8 mm Hg (systolic/diastolic) greater than those associated with placebo. Peak reduction of blood pressure is achieved within 3-6 hours after administration and the blood pressure lowering effect is maintained for at least 24 hours. At 24 hours the reduction of blood pressure was 60-70% of the corresponding peak diastolic and systolic responses at the recommended doses. Once daily dosing with 150 mg produced trough and mean 24 hour responses similar to twice daily dosing on the same total dose.

The blood pressure lowering effect of irbesartan is evident within 1-2 weeks, with the maximal effect occurring by 4-6 weeks after start of therapy. The antihypertensive effects are maintained during long term therapy. After withdrawal of therapy, blood pressure gradually returns toward baseline. Rebound hypertension has not been observed.

The blood pressure lowering effects of irbesartan and thiazide-type diuretics are additive. In patients not adequately controlled by irbesartan alone, the addition of a low dose of hydrochlorothiazide (12.5 mg) to irbesartan once daily results in a further placebo-adjusted blood pressure reduction at trough of 7-10/3-6 mm Hg (systolic/diastolic).

The efficacy of irbesartan is not influenced by age or gender. As is the case with other medicinal products that affect the renin-angiotensin system, black hypertensive patients have notably less response to irbesartan monotherapy. When irbesartan is administered concomitantly with a low dose of hydrochlorothiazide (e.g. 12.5 mg daily), the antihypertensive response in black patients approaches that of white patients.

There is no clinically important effect on serum uric acid or urinary uric acid secretion.

#### Paediatric population

Reduction of blood pressure with 0.5 mg/kg (low), 1.5 mg/kg (medium) and 4.5 mg/kg (high) target titrated doses of irbesartan was evaluated in 318 hypertensive or at risk (diabetic, family history of

hypertension) children and adolescents aged 6 to 16 years over a three week period. At the end of the three weeks the mean reduction from baseline in the primary efficacy variable, trough seated systolic blood pressure (SeSBP) was 11.7 mmHg (low dose), 9.3 mmHg (medium dose), 13.2 mmHg (high dose). No significant difference was apparent between these doses. Adjusted mean change of trough seated diastolic blood pressure (SeDBP) was as follows: 3.8 mmHg (low dose), 3.2 mmHg (medium dose), 5.6 mmHg (high dose). Over a subsequent two week period where patients were re-randomized to either active medicinal product or placebo, patients on placebo had increases of 2.4 and 2.0 mmHg in SeSBP and SeDBP compared to +0.1 and -0.3 mmHg changes respectively in those on all doses of irbesartan (see section 4.2).

#### Hypertension and type 2 diabetes with renal disease

The “Irbesartan Diabetic Nephropathy Trial (IDNT)” shows that irbesartan decreases the progression of renal disease in patients with chronic renal insufficiency and overt proteinuria. IDNT was a double blind, controlled, morbidity and mortality trial comparing irbesartan, amlodipine and placebo. In 1,715 hypertensive patients with type 2 diabetes, proteinuria  $\geq$  900 mg/day and serum creatinine ranging from 1.0-3.0 mg/dl, the long-term effects (mean 2.6 years) of irbesartan on the progression of renal disease and all-cause mortality were examined. Patients were titrated from 75 mg to a maintenance dose of 300 mg irbesartan, from 2.5 mg to 10 mg amlodipine, or placebo as tolerated.

Patients in all treatment groups typically received between 2 and 4 antihypertensive agents (e.g., diuretics, beta blockers, alpha blockers) to reach a predefined blood pressure goal of  $\leq$  135/85 mmHg or a 10 mmHg reduction in systolic pressure if baseline was  $>$  160 mmHg. Sixty per cent (60%) of patients in the placebo group reached this target blood pressure whereas this figure was 76% and 78% in the irbesartan and amlodipine groups respectively.

Irbesartan significantly reduced the relative risk in the primary combined endpoint of doubling serum creatinine, end-stage renal disease (ESRD) or all-cause mortality. Approximately 33% of patients in the irbesartan group reached the primary renal composite endpoint compared to 39% and 41% in the placebo and amlodipine groups [20% relative risk reduction versus placebo ( $p = 0.024$ ) and 23% relative risk reduction compared to amlodipine ( $p = 0.006$ )]. When the individual components of the primary endpoint were analysed, no effect in all cause mortality was observed, while a positive trend in the reduction in ESRD and a significant reduction in doubling of serum creatinine were observed.

Subgroups consisting of gender, race, age, duration of diabetes, baseline blood pressure, serum creatinine, and albumin excretion rate were assessed for treatment effect. In the female and black subgroups which represented 32% and 26% of the overall study population respectively, a renal benefit was not evident, although the confidence intervals do not exclude it. As for the secondary endpoint of fatal and non-fatal cardiovascular events, there was no difference among the three groups in the overall population, although an increased incidence of non-fatal MI was seen for women and a decreased incidence of non-fatal MI was seen in males in the irbesartan group versus the placebo-based regimen. An increased incidence of non-fatal MI and stroke was seen in females in the irbesartan-based regimen versus the amlodipine-based regimen, while hospitalization due to heart failure was reduced in the overall population. However, no proper explanation for these findings in women has been identified.

The study of the “Effects of Irbesartan on Microalbuminuria in Hypertensive Patients with type 2 Diabetes Mellitus (IRMA 2)” shows that irbesartan 300 mg delays progression to overt proteinuria in patients with microalbuminuria. IRMA 2 was a placebo-controlled double blind morbidity study in 590 patients with type 2 diabetes, microalbuminuria (30-300 mg/day) and normal renal function (serum creatinine  $\leq$  1.5 mg/dl in males and  $<$  1.1 mg/dl in females). The study examined the long-term effects (2 years) of irbesartan on the progression to clinical (overt) proteinuria (urinary albumin excretion rate (UAER)  $>$  300 mg/day, and an increase in UAER of at least 30% from baseline). The predefined blood pressure goal was  $\leq$  135/85 mmHg. Additional antihypertensive agents (excluding ACE inhibitors, angiotensin II receptor antagonists and dihydropyridine calcium blockers) were added as needed to help achieve the blood pressure goal. While similar blood pressure was achieved in all treatment groups, fewer subjects in the irbesartan 300 mg group (5.2%) than in the placebo (14.9%) or in the irbesartan 150 mg group (9.7%) reached the endpoint of overt proteinuria, demonstrating a 70% relative risk reduction versus placebo ( $p = 0.0004$ ) for the higher dose. An accompanying improvement in the glomerular filtration rate (GFR) was not observed during the first three months of

treatment. The slowing in the progression to clinical proteinuria was evident as early as three months and continued over the 2 year period. Regression to normoalbuminuria (< 30 mg/day) was more frequent in the irbesartan 300 mg group (34%) than in the placebo group (21%).

Two large randomised, controlled trials (ONTARGET (ONgoing Telmisartan Alone and in combination with Ramipril Global Endpoint Trial) and VA NEPHRON-D (The Veterans Affairs Nephropathy in Diabetes)) have examined the use of the combination of an ACE-inhibitor with an angiotensin II receptor blocker.

ONTARGET was a study conducted in patients with a history of cardiovascular or cerebrovascular disease, or type 2 diabetes mellitus accompanied by evidence of end-organ damage. VA NEPHRON-D was a study in patients with type 2 diabetes mellitus and diabetic nephropathy.

These studies have shown no significant beneficial effect on renal and/or cardiovascular outcomes and mortality, while an increased risk of hyperkalaemia, acute kidney injury and/or hypotension as compared to monotherapy was observed. Given their similar pharmacodynamic properties, these results are also relevant for other ACE-inhibitors and angiotensin II receptor blockers.

ACE-inhibitors and angiotensin II receptor blockers should therefore not be used concomitantly in patients with diabetic nephropathy.

ALTITUDE (Aliskiren Trial in Type 2 Diabetes Using Cardiovascular and Renal Disease Endpoints) was a study designed to test the benefit of adding aliskiren to a standard therapy of an ACE-inhibitor or an angiotensin II receptor blocker in patients with type 2 diabetes mellitus and chronic kidney disease, cardiovascular disease, or both. The study was terminated early because of an increased risk of adverse outcomes. Cardiovascular death and stroke were both numerically more frequent in the aliskiren group than in the placebo group and adverse events and serious adverse events of interest (hyperkalaemia, hypotension and renal dysfunction) were more frequently reported in the aliskiren group than in the placebo group.

## 5.2 Pharmacokinetic properties

After oral administration, irbesartan is well absorbed: studies of absolute bioavailability gave values of approximately 60-80%. Concomitant food intake does not significantly influence the bioavailability of irbesartan. Plasma protein binding is approximately 96%, with negligible binding to cellular blood components. The volume of distribution is 53 - 93 litres. Following oral or intravenous administration of <sup>14</sup>C irbesartan, 80-85% of the circulating plasma radioactivity is attributable to unchanged irbesartan. Irbesartan is metabolised by the liver via glucuronide conjugation and oxidation. The major circulating metabolite is irbesartan glucuronide (approximately 6%). *In vitro* studies indicate that irbesartan is primarily oxidised by the cytochrome P450 enzyme CYP2C9; isoenzyme CYP3A4 has negligible effect.

Irbesartan exhibits linear and dose proportional pharmacokinetics over the dose range of 10 to 600 mg. A less than proportional increase in oral absorption at doses beyond 600 mg (twice the maximal recommended dose) was observed; the mechanism for this is unknown. Peak plasma concentrations are attained at 1.5 - 2 hours after oral administration. The total body and renal clearance are 157 - 176 and 3 - 3.5 ml/min, respectively. The terminal elimination half-life of irbesartan is 11 - 15 hours. Steady-state plasma concentrations are attained within 3 days after initiation of a once-daily dosing regimen. Limited accumulation of irbesartan (< 20%) is observed in plasma upon repeated once-daily dosing. In a study, somewhat higher plasma concentrations of irbesartan were observed in female hypertensive patients. However, there was no difference in the half-life and accumulation of irbesartan. No dosage adjustment is necessary in female patients. Irbesartan AUC and C<sub>max</sub> values were also somewhat greater in elderly subjects (≥ 65 years) than those of young subjects (18 - 40 years). However the terminal half-life was not significantly altered. No dosage adjustment is necessary in elderly patients.

Irbesartan and its metabolites are eliminated by both biliary and renal pathways. After either oral or IV administration of <sup>14</sup>C irbesartan, about 20% of the radioactivity is recovered in the urine, and the remainder in the faeces. Less than 2% of the dose is excreted in the urine as unchanged irbesartan.

### Paediatric population



The pharmacokinetics of irbesartan were evaluated in 23 hypertensive children after the administration of single and multiple daily doses of irbesartan (2 mg/kg) up to a maximum daily dose of 150 mg for four weeks. Of those 23 children, 21 were evaluable for comparison of pharmacokinetics with adults (twelve children over 12 years, nine children between 6 and 12 years). Results showed that C<sub>max</sub>, AUC and clearance rates were comparable to those observed in adult patients receiving 150 mg irbesartan daily. A limited accumulation of irbesartan (18%) in plasma was observed upon repeated once daily dosing.

Renal impairment: in patients with renal impairment or those undergoing haemodialysis, the pharmacokinetic parameters of irbesartan are not significantly altered. Irbesartan is not removed by haemodialysis.

Hepatic impairment: in patients with mild to moderate cirrhosis, the pharmacokinetic parameters of irbesartan are not significantly altered.

Studies have not been performed in patients with severe hepatic impairment.

### **5.3 Preclinical safety data**

There was no evidence of abnormal systemic or target organ toxicity at clinically relevant doses. In non-clinical safety studies, high doses of irbesartan ( $\geq 250$  mg/kg/day in rats and  $\geq 100$  mg/kg/day in macaques) caused a reduction of red blood cell parameters (erythrocytes, haemoglobin, haematocrit). At very high doses ( $\geq 500$  mg/kg/day) degenerative changes in the kidney (such as interstitial nephritis, tubular distension, basophilic tubules, increased plasma concentrations of urea and creatinine) were induced by irbesartan in the rat and the macaque and are considered secondary to the hypotensive effects of the medicinal product which led to decreased renal perfusion. Furthermore, irbesartan induced hyperplasia/hypertrophy of the juxtaglomerular cells (in rats at  $\geq 90$  mg/kg/day, in macaques at  $\geq 10$  mg/kg/day). All of these changes were considered to be caused by the pharmacological action of irbesartan. For therapeutic doses of irbesartan in humans, the hyperplasia/hypertrophy of the renal juxtaglomerular cells does not appear to have any relevance.

There was no evidence of mutagenicity, clastogenicity or carcinogenicity.

Fertility and reproductive performance were not affected in studies of male and female rats even at oral doses of irbesartan causing some parental toxicity (from 50 to 650 mg/kg/day), including mortality at the highest dose. No significant effects on the number of corpora lutea, implants, or live fetuses were observed. Irbesartan did not affect survival, development, or reproduction of offspring. Studies in animals indicate that the radiolabeled irbesartan is detected in rat and rabbit fetuses. Irbesartan is excreted in the milk of lactating rats.

Animal studies with irbesartan showed transient toxic effects (increased renal pelvic cavitation, hydroureter or subcutaneous oedema) in rat foetuses, which were resolved after birth. In rabbits, abortion or early resorption were noted at doses causing significant maternal toxicity, including mortality. No teratogenic effects were observed in the rat or rabbit.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Tablet core:

Lactose monohydrate  
Microcrystalline cellulose  
Croscarmellose sodium  
Silica colloidal anhydrous  
Hypromellose  
Magnesium stearate.

Film-coating:  
Hypromellose  
Titanium dioxide (E171)  
Macrogol 400

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

4 years

## **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

## **6.5 Nature and contents of container**

Cartons of 28, 56, 90 or 98 film-coated tablets in PVC/PVDC-aluminium/PVDC blisters.  
Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

Any unused product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Pharmathen S.A.  
6, Dervenakion str.,  
15351 Pallini  
Attiki, Greece  
Tel.: +30 210 66 65 067

## **8. MARKETING AUTHORISATION NUMBERS**

EU/1/12/765/005  
EU/1/12/765/006  
EU/1/12/765/009  
EU/1/12/765/012

## **9. DATE OF FIRST AUTHORISATION / RENEWAL OF THE AUTHORISATION**

Date of first Authorization: 13 April 2012

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu/>

## **ANNEX II**

- A. MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

## **A. MANUFACTURERS RESPONSIBLE FOR BATCH RELEASE**

### Name and address of the manufacturers responsible for batch release

J. Uriach y Compañía, S.A.  
Av. Camí Reial, 51-57  
ES-08184 – Palau-solità i Plegamans  
Barcelona  
Spain

Pharmathen S.A.  
6 Dervenakion str.  
15351 Pallini Attiki  
Greece

Pharmathen International S.A.  
Industrial Park Sapes,  
Rodopi Prefecture, Block No 5,  
Rodopi 69300,  
Greece

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to medical prescription.

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

### Pharmacovigilance system

The MAH must ensure that the system of pharmacovigilance presented in Module 1.8.1. of the Marketing Authorisation is in place and functioning before and whilst the medicinal product is on the market.

### Risk Management Plan (RMP)

Not applicable

### PSURs

The PSUR submission schedule should follow the PSUR submission schedule for the reference medicinal product.

## **• CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

Not applicable.

•

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Sabervel 75 mg film-coated tablets  
irbesartan

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each tablet contains 75 mg irbesartan

**3. LIST OF EXCIPIENTS**

Excipients: also contains lactose monohydrate.  
See leaflet for further information

**4. PHARMACEUTICAL FORM AND CONTENTS**

28 film-coated tablets  
56 film-coated tablets  
90 film-coated tablets  
98 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use. Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Pharmathen S.A.  
6, Dervenakion str.,  
153 51 Pallini  
Attiki, Greece

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/12/765/001  
EU/1/12/765/002  
EU/1/12/765/007  
EU/1/12/765/010

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Sabervel 75 mg



**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**1. NAME OF THE MEDICINAL PRODUCT**

Sabervel 75 mg film-coated tablets  
irbesartan

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Pharmathen S.A.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Sabervel 150 mg film-coated tablets  
irbesartan

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each tablet contains 150 mg irbesartan

**3. LIST OF EXCIPIENTS**

Excipients: also contains lactose monohydrate.  
See leaflet for further information

**4. PHARMACEUTICAL FORM AND CONTENTS**

28 film-coated tablets  
56 film-coated tablets  
90 film-coated tablets  
98 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use. Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Pharmathen S.A.  
6, Dervenakion str.,  
153 51 Pallini  
Attiki, Greece

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/12/765/003  
EU/1/12/765/004  
EU/1/12/765/008  
EU/1/12/765/011

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Sabervel 150 mg

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**1. NAME OF THE MEDICINAL PRODUCT**

Sabervel 150 mg film-coated tablets  
irbesartan

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Pharmathen S.A.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Sabervel 300 mg film-coated tablets  
irbesartan

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each tablet contains 300 mg irbesartan

**3. LIST OF EXCIPIENTS**

Excipients: also contains lactose monohydrate.  
See leaflet for further information

**4. PHARMACEUTICAL FORM AND CONTENTS**

28 film-coated tablets  
56 film-coated tablets  
90 film-coated tablets  
98 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use. Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Pharmathen S.A.  
6, Dervenakion str.,  
153 51 Pallini  
Attiki, Greece

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/12/765/005  
EU/1/12/765/006  
EU/1/12/765/009  
EU/1/12/765/012

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription.

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Sabervel 300 mg

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**1. NAME OF THE MEDICINAL PRODUCT**

Sabervel 300 mg film-coated tablets  
irbesartan

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Pharmathen S.A.

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

## **B. PACKAGE LEAFLET**



## PACKAGE LEAFLET: INFORMATION FOR THE USER

### Sabervel 75 mg film-coated tablets irbesartan

**Read all of this leaflet carefully before you start taking this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

#### **In this leaflet:**

1. What Sabervel is and what it is used for
2. Before you take Sabervel
3. How to take Sabervel
4. Possible side effects
5. How to store Sabervel
6. Further information

## **1. WHAT SABERVEL IS AND WHAT IT IS USED FOR**

Sabervel belongs to a group of medicines known as angiotensin-II receptor antagonists. Angiotensin-II is a substance produced in the body which binds to receptors in blood vessels causing them to tighten. This results in an increase in blood pressure. Sabervel prevents the binding of angiotensin-II to these receptors, causing the blood vessels to relax and the blood pressure to lower. Sabervel slows the decrease of kidney function in patients with high blood pressure and type 2 diabetes.

Sabervel is used in adult patients

- to treat high blood pressure (*essential hypertension*)
- to protect the kidney in patients with high blood pressure, type 2 diabetes and laboratory evidence of impaired kidney function.

## **2. BEFORE YOU TAKE SABERVEL**

### **Do not take Sabervel**

- if you are **allergic** (hypersensitive) to irbesartan or any other ingredients of Sabervel
- if you are **more than 3 months pregnant**. (It is also better to avoid Sabervel in early pregnancy – see pregnancy section)
- if you have diabetes or impaired kidney function and you are treated with a blood pressure lowering medicine containing aliskiren.

**Warnings and precautions** Tell your doctor if any of the following apply to you:

- if you get **excessive vomiting or diarrhoea**
- if you suffer from **kidney problems**
- if you suffer from **heart problems**
- if you receive Sabervel for **diabetic kidney disease**. In this case your doctor may perform regular blood tests, especially for measuring blood potassium levels in case of poor kidney function
- if you are **going to have an operation** (surgery) or **be given anaesthetics**.
- if you are taking any of the following medicines used to treat high blood pressure:  
- an ACE-inhibitor (for example enalapril, lisinopril, ramipril), in particular if you have diabetes-related kidney problems.

- aliskiren

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (e.g. potassium) in your blood at regular intervals.

See also information under the heading “Do not take Sabervel”.

You must tell your doctor if you think you are (or might become) pregnant. Sabervel is not recommended in early pregnancy, and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see pregnancy section).

### **Use in Children**

This medicine should not be used in children and adolescents (under 18 years) because the safety and efficacy have not yet been fully established.

### **Other medicines and Sabervel**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Your doctor may need to change your dose and/or to take other precautions:

If you are taking an ACE-inhibitor or aliskiren (see also information under the headings “Do not take Sabervel” and “Warnings and precautions”).

### **You may need to have blood checks if you take:**

- potassium supplements
- salt substitutes containing potassium
- potassium-sparing medicines (such as certain diuretics)
- medicines containing lithium

If you take certain painkillers, called non-steroidal anti-inflammatory drugs, the effect of irbesartan may be reduced.

### **Taking Sabervel with food and drink**

Sabervel can be taken with or without food.

### **Pregnancy and breast-feeding**

#### **Pregnancy**

You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking Sabervel before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Sabervel. Sabervel is not recommended in early pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

#### **Breast-feeding**

Tell your doctor if you are breast-feeding or about to start breast-feeding. Sabervel is not recommended for mothers who are breast-feeding, and your doctor may choose another treatment for you if you wish to breast-feed, especially if your baby is newborn, or was born prematurely.

### **Driving and using machines**

No studies on the effects on the ability to drive and use machines have been performed. Sabervel is unlikely to affect your ability to drive or use machines. However, occasionally dizziness or weariness may occur during treatment of high blood pressure. If you experience these, talk to your doctor before attempting to drive or use machines.

### **Important information about some of the ingredients of Sabervel**

**Sabervel contains lactose.** If you have been told by your doctor that you have an intolerance to some sugars (e.g. lactose), contact your doctor before taking this medicine.

### 3. HOW TO TAKE SABERVEL

Always take Sabervel exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

#### Method of administration

Sabervel is for **oral use**. Swallow the tablets with a sufficient amount of fluid (e.g. one glass of water). You can take Sabervel with or without food. Try to take your daily dose at about the same time each day. It is important that you continue to take Sabervel until your doctor tells you otherwise.

- **Patients with high blood pressure**  
The usual dose is 150 mg once a day. The dose may later be increased to 300 mg once daily depending on blood pressure response.
- **Patients with high blood pressure and type 2 diabetes with kidney disease**  
In patients with high blood pressure and type 2 diabetes, 300 mg once daily is the preferred maintenance dose for the treatment of associated kidney disease.

The doctor may advise a lower dose, especially when starting treatment in certain patients such as those on **haemodialysis**, or those **over the age of 75 years**.

The maximal blood pressure lowering effect should be reached 4-6 weeks after beginning treatment.

#### If you take more Sabervel than you should:

If you accidentally take too many tablets, contact your doctor immediately.

#### Children should not take Sabervel

Sabervel should not be given to children under 18 years of age. If a child swallows some tablets, contact your doctor immediately.

#### If you forget to take Sabervel:

If you accidentally miss a daily dose, just take the next dose as normal. Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

### 4. POSSIBLE SIDE EFFECTS

Like all medicines, Sabervel can cause side effects, although not everybody gets them. Some of these effects may be serious and may require medical attention.

As with similar medicines, rare cases of allergic skin reactions (rash, urticaria), as well as localised swelling of the face, lips and/or tongue have been reported in patients taking irbesartan. If you get any of these symptoms or get short of breath, **stop taking Sabervel and contact your doctor immediately**.

The frequency of the side effects listed below is defined using the following convention:

Very common: at least 1 in 10 patients or more

Common: at least 1 in 100 and less than 1 in 10 patients

Uncommon: at least 1 in 1000 and less than 1 in 100 patients  
Side effects reported in clinical studies for patients treated with Irbesartan were:

- Very common : if you suffer from high blood pressure and type 2 diabetes with kidney disease, blood tests may show an increased level of potassium.

- Common : dizziness, feeling sick/vomiting, fatigue and blood tests may show raised levels of an enzyme that measures the muscle and heart function (creatine kinase enzyme). In patients with high blood pressure and type 2 diabetes with kidney disease, dizziness when getting up from a lying or sitting position, low blood pressure when getting up from a lying or sitting position, pain in joints or muscles and decreased levels of a protein in the red blood cells (haemoglobin) were also reported.
- Uncommon : heart rate increased, flushing, cough, diarrhoea, indigestion/heartburn, sexual dysfunction (problems with sexual performance), chest pain.

Some side effects have been reported since marketing of Irbesartan Side effects where the frequency is not known are: feeling of spinning, headache, taste disturbance, ringing in the ears, muscle cramps, pain in joints and muscles, abnormal liver function, increased blood potassium levels, impaired kidney function, and inflammation of small blood vessels mainly affecting the skin (a condition known as leukocytoclastic vasculitis).

Uncommon cases of jaundice (yellowing of the skin and/or whites of the eyes) have also been reported.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## 5. HOW TO STORE SABERVEL

Keep out of the sight and reach of children.

Do not use Sabervel after the expiry date which is stated on the carton and on the blister after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## 6. FURTHER INFORMATION

### What Sabervel contains

- The active substance is irbesartan. Each tablet of Sabervel 75 mg contains 75 mg irbesartan.
- The other ingredients are:  
Tablet core: lactose monohydrate, microcrystalline cellulose, croscarmellose sodium, silica colloidal anhydrous, hypromellose, magnesium stearate,  
Film-coating: hypromellose, titanium dioxide (E171), macrogol 400

### What Sabervel looks like and contents of the pack

Sabervel 75 mg film-coated tablets are white, concave, round, film-coated tablets with 7 mm diameter.

Sabervel 75mg film-coated tablets are supplied in blister packs of 28, 56, 90 or 98 film-coated tablets.

Not all pack sizes may be marketed.

### Marketing Authorisation Holder:

Pharmathen S.A,  
6 Dervenakion Str.,  
15351 Pallini, Attiki, Greece

**Manufacturer:**

J. Uriach y Compañía, S.A.  
Av. Camí Reial, 51-57  
08184 – Palau-solità i Plegamans, Barcelona, Spain

Pharmathen S.A.,  
6 Dervenakion Str.,  
15351 Pallini, Attiki, Greece

Pharmathen International S.A.  
Industrial Park Sapes,  
Rodopi Prefecture, Block No 5, Rodopi 69300, Greece

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder.

**België/Belgique/Belgien**

Pharmathen S.A.  
Tél/Tel: +30 210 66 65 067

**Lietuva**

Portfarma ehf.  
Tel: +354 534 4030

**България**

Pharmathen S.A.  
Тел.: +30 210 66 65 067

**Luxembourg/Luxemburg**

Portfarma ehf.  
Tel: +354 534 4030

**Česká republika**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Magyarország**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Danmark**

Portfarma ehf.  
Tlf: +354 534 4030

**Malta**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Deutschland**

Glenmark Arzneimittel GmbH  
Tel: +49 8142 44392 0

**Nederland**

Glenmark Generics B.V.  
Tel: 0031 20 5226030

**Eesti (Estonia)**

Portfarma ehf.  
Tel: +354 534 4030

**Norge**

Pharmathen S.A.  
Tlf: +30 210 66 65 067

**Ελλάδα**

Pharmathen Hellas S.A.  
Τηλ: +30 210 66 04 300

**Österreich**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**España**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Polska**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**France**

Pharmathen S.A.  
Tél: +30 210 66 65 067

**Portugal**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Hrvatska**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**România**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Ireland**

Aspire Pharma Limited  
Tel: +44(0)1730 234527

**Ísland**

Portfarma ehf.  
Tel: +354 534 4030

**Italia**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Κύπρος**

C.V. MEDILINE LTD  
Τηλ: +357 25761699

**Latvija**

Portfarma ehf.  
Tel: +354 534 4030

**Slovenija**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Slovenská republika**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Suomi/Finland**

Pharmathen S.A.  
Puh/Tel: +30 210 66 65 067

**Sverige**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**United Kingdom**

Aspire Pharma Limited  
Tel: +44(0)1730 234527

**This leaflet was last approved in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu/>

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### Sabervel 150 mg film-coated tablets irbesartan

#### **Read all of this leaflet carefully before you start taking this medicine .**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

#### **In this leaflet:**

1. What Sabervel is and what it is used for
2. Before you take Sabervel
3. How to take Sabervel
4. Possible side effects
5. How to store Sabervel
6. Further information

## **1. WHAT SABERVEL IS AND WHAT IT IS USED FOR**

Sabervel belongs to a group of medicines known as angiotensin-II receptor antagonists. Angiotensin-II is a substance produced in the body which binds to receptors in blood vessels causing them to tighten. This results in an increase in blood pressure. Sabervel prevents the binding of angiotensin-II to these receptors, causing the blood vessels to relax and the blood pressure to lower. Sabervel slows the decrease of kidney function in patients with high blood pressure and type 2 diabetes.

Sabervel is used in adult patients

- to treat high blood pressure (*essential hypertension*)
- to protect the kidney in patients with high blood pressure, type 2 diabetes and laboratory evidence of impaired kidney function.

## **2. BEFORE YOU TAKE SABERVEL**

### **Do not take Sabervel**

- if you are **allergic** (hypersensitive) to irbesartan or any other ingredients of Sabervel
- if you are **more than 3 months pregnant**. (It is also better to avoid Sabervel in early pregnancy – see pregnancy section)
- if you have diabetes or impaired kidney function and you are treated with a blood pressure lowering medicine containing aliskiren.

### **Warning and precautions**

**Tell your doctor** if any of the following apply to you:

- if you get **excessive vomiting or diarrhoea**
- if you suffer from **kidney problems**
- if you suffer from **heart problems**
- if you receive Sabervel for **diabetic kidney disease**. In this case your doctor may perform regular blood tests, especially for measuring blood potassium levels in case of poor kidney function
- if you are **going to have an operation** (surgery) or **be given anaesthetics**.
- if you are taking any of the following medicines used to treat high blood pressure:  
- an ACE-inhibitor (for example enalapril, lisinopril, ramipril), in particular if you have

diabetes-related kidney problems.  
- aliskiren

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (e.g. potassium) in your blood at regular intervals.  
See also information under the heading “Do not take Sabervel”.

You must tell your doctor if you think you are (or might become) pregnant. Sabervel is not recommended in early pregnancy, and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see pregnancy section).

### **Use in Children**

This medicine should not be used in children and adolescents (under 18 years) because the safety and efficacy have not yet been fully established.

### **Other medicines and Sabervel**

Tell your doctor or pharmacist if you are taking , have recently taken or might take any other medicines.

Your doctor may need to change your dose and/or to take other precautions:

If you are taking an ACE-inhibitor or aliskiren (see also information under the headings “Do not take Sabervel” and “Warnings and precautions”).

### **You may need to have blood checks if you take :**

- potassium supplements
- salt substitutes containing potassium
- potassium-sparing medicines (such as certain diuretics)
- medicines containing lithium

If you take certain painkillers, called non-steroidal anti-inflammatory drugs, the effect of irbesartan may be reduced.

### **Taking Sabervel with food and drink**

Sabervel can be taken with or without food.

### **Pregnancy and breast-feeding**

#### **Pregnancy**

You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking Sabervel before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Sabervel. Sabervel is not recommended in early pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

#### **Breast-feeding**

Tell your doctor if you are breast-feeding or about to start breast-feeding. Sabervel is not recommended for mothers who are breast-feeding, and your doctor may choose another treatment for you if you wish to breast-feed, especially if your baby is newborn, or was born prematurely.

### **Driving and using machines**

No studies on the effects on the ability to drive and use machines have been performed. Sabervel is unlikely to affect your ability to drive or use machines. However, occasionally dizziness or weariness may occur during treatment of high blood pressure. If you experience these, talk to your doctor before attempting to drive or use machines.

### **Important information about some of the ingredients of Sabervel**

**Sabervel contains lactose.** If you have been told by your doctor that you have an intolerance to some sugars (e.g. lactose), contact your doctor before taking this medicine.



### 3. HOW TO TAKE SABERVEL

Always take Sabervel exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

#### Method of administration

Sabervel is for **oral use**. Swallow the tablets with a sufficient amount of fluid (e.g. one glass of water). You can take Sabervel with or without food. Try to take your daily dose at about the same time each day. It is important that you continue to take Sabervel until your doctor tells you otherwise.

- **Patients with high blood pressure**  
The usual dose is 150 mg once a day. The dose may later be increased to 300 mg once daily depending on blood pressure response.
- **Patients with high blood pressure and type 2 diabetes with kidney disease**  
In patients with high blood pressure and type 2 diabetes, 300 mg once daily is the preferred maintenance dose for the treatment of associated kidney disease.

The doctor may advise a lower dose, especially when starting treatment in certain patients such as those on **haemodialysis**, or those **over the age of 75 years**.

The maximal blood pressure lowering effect should be reached 4-6 weeks after beginning treatment.

#### If you take more Sabervel than you should:

If you accidentally take too many tablets, contact your doctor immediately.

#### Children should not take Sabervel

Sabervel should not be given to children under 18 years of age. If a child swallows some tablets, contact your doctor immediately.

#### If you forget to take Sabervel:

If you accidentally miss a daily dose, just take the next dose as normal. Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

### 4. POSSIBLE SIDE EFFECTS

Like all medicines, Sabervel can cause side effects, although not everybody gets them. Some of these effects may be serious and may require medical attention.

As with similar medicines, rare cases of allergic skin reactions (rash, urticaria), as well as localised swelling of the face, lips and/or tongue have been reported in patients taking irbesartan. If you get any of these symptoms or get short of breath, **stop taking Sabervel and contact your doctor immediately**.

The frequency of the side effects listed below is defined using the following convention:

Very common: at least 1 in 10 patients or more

Common: at least 1 in 100 and less than 1 in 10 patients

Uncommon: at least 1 in 1000 and less than 1 in 100 patients

Side effects reported in clinical studies for patients treated with Irbesartan were:

- Very common : if you suffer from high blood pressure and type 2 diabetes with kidney disease, blood tests may show an increased level of potassium.

- Common : dizziness, feeling sick/vomiting, fatigue and blood tests may show raised levels of an enzyme that measures the muscle and heart function (creatine kinase enzyme). In patients with high blood pressure and type 2 diabetes with kidney disease, dizziness when getting up from a lying or sitting position, low blood pressure when getting up from a lying or sitting position, pain in joints or muscles and decreased levels of a protein in the red blood cells (haemoglobin) were also reported.
- Uncommon : heart rate increased, flushing, cough, diarrhoea, indigestion/heartburn, sexual dysfunction (problems with sexual performance), chest pain.

Some side effects have been reported since marketing of Irbesartan Side effects where the frequency is not known are: feeling of spinning, headache, taste disturbance, ringing in the ears, muscle cramps, pain in joints and muscles, abnormal liver function, increased blood potassium levels, impaired kidney function, and inflammation of small blood vessels mainly affecting the skin (a condition known as leukocytoclastic vasculitis).

Uncommon cases of jaundice (yellowing of the skin and/or whites of the eyes) have also been reported.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## 5. HOW TO STORE SABERVEL

Keep out of the sight and reach of children.

Do not use Sabervel after the expiry date which is stated on the carton and on the blister after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## 6. FURTHER INFORMATION

### What Sabervel contains

- The active substance is irbesartan. Each tablet of Sabervel 150 mg contains 150 mg irbesartan.
- The other ingredients are:  
Tablet core: lactose monohydrate, microcrystalline cellulose, croscarmellose sodium, silica colloidal anhydrous, hypromellose, magnesium stearate,  
Film-coating: hypromellose, titanium dioxide (E171), macrogol 400

### What Sabervel looks like and contents of the pack

Sabervel 150 mg film-coated tablets are white, concave, round, film-coated tablets with 9 mm diameter.

Sabervel 150mg film-coated tablets are supplied in blister packs of 28, 56, 90 or 98 film-coated tablets.

Not all pack sizes may be marketed.

### Marketing Authorisation Holder:

Pharmathen S.A,  
6 Dervenakion Str.,

15351 Pallini, Attiki, Greece

**Manufacturer:**

J. Uriach y Compañía, S.A.  
Av. Camí Reial, 51-57  
08184 – Palau-solità i Plegamans, Barcelona, Spain

Pharmathen S.A.,  
6 Dervenakion Str.,  
15351 Pallini, Attiki, Greece

Pharmathen International S.A.  
Industrial Park Sapes,  
Rodopi Prefecture, Block No 5, Rodopi 69300, Greece

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder.

**België/Belgique/Belgien**

Pharmathen S.A.  
Tél/Tel: +30 210 66 65 067

**Lietuva**

Portfarma ehf.  
Tel: +354 534 4030

**България**

Pharmathen S.A.  
Тел.: +30 210 66 65 067

**Luxembourg/Luxemburg**

Portfarma ehf.  
Tel: +354 534 4030

**Česká republika**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Magyarország**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Danmark**

Portfarma ehf.  
Tlf: +354 534 4030

**Malta**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Deutschland**

Glenmark Arzneimittel GmbH  
Tel: +49 8142 44392 0

**Nederland**

Glenmark Generics B.V.  
Tel: 0031 20 5226030

**Eesti (Estonia)**

Portfarma ehf.  
Tel: +354 534 4030

**Norge**

Pharmathen S.A.  
Tlf: +30 210 66 65 067

**Ελλάδα**

Pharmathen Hellas S.A.  
Τηλ: +30 210 66 04 300

**Österreich**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**España**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Polska**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**France**

Pharmathen S.A.  
Tél: +30 210 66 65 067

**Portugal**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Hrvatska**

Pharmathen S.A.

**România**

Pharmathen S.A.

Tel: +30 210 66 65 067

**Ireland**

Aspire Pharma Limited  
Tel: +44(0)1730 234527

**Ísland**

Portfarma ehf.  
Tel: +354 534 4030

**Italia**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Κύπρος**

C.V. MEDILINE LTD  
Τηλ: +357 25761699

**Latvija**

Portfarma ehf.  
Tel: +354 534 4030

Tel: +30 210 66 65 067

**Slovenija**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Slovenská republika**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Suomi/Finland**

Pharmathen S.A.  
Puh/Tel: +30 210 66 65 067

**Sverige**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**United Kingdom**

Aspire Pharma Limited  
Tel: +44(0)1730 234527

**This leaflet was last approved in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu/>

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### Sabervel 300 mg film-coated tablets irbesartan

#### **Read all of this leaflet carefully before you start taking this medicine .**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

#### **In this leaflet:**

1. What Sabervel is and what it is used for
2. Before you take Sabervel
3. How to take Sabervel
4. Possible side effects
5. How to store Sabervel
6. Further information

### **1. WHAT SABERVEL IS AND WHAT IT IS USED FOR**

Sabervel belongs to a group of medicines known as angiotensin-II receptor antagonists. Angiotensin-II is a substance produced in the body which binds to receptors in blood vessels causing them to tighten. This results in an increase in blood pressure. Sabervel prevents the binding of angiotensin-II to these receptors, causing the blood vessels to relax and the blood pressure to lower. Sabervel slows the decrease of kidney function in patients with high blood pressure and type 2 diabetes.

Sabervel is used in adult patients

- to treat high blood pressure (*essential hypertension*)
- to protect the kidney in patients with high blood pressure, type 2 diabetes and laboratory evidence of impaired kidney function.

### **2. BEFORE YOU TAKE SABERVEL**

#### **Do not take Sabervel**

- if you are **allergic** (hypersensitive) to irbesartan or any other ingredients of Sabervel
- if you are **more than 3 months pregnant**. (It is also better to avoid Sabervel in early pregnancy – see pregnancy section)
- if you have diabetes or impaired kidney function and you are treated with a blood pressure lowering medicine containing aliskiren.

#### **Warning and precautions**

**Tell your doctor** if any of the following apply to you:

- if you get **excessive vomiting or diarrhoea**
- if you suffer from **kidney problems**
- if you suffer from **heart problems**
- if you receive Sabervel for **diabetic kidney disease**. In this case your doctor may perform regular blood tests, especially for measuring blood potassium levels in case of poor kidney function
- if you are **going to have an operation** (surgery) or **be given anaesthetics**.
- if you are taking any of the following medicines used to treat high blood pressure:  
- an ACE-inhibitor (for example enalapril, lisinopril, ramipril), in particular if you have

diabetes-related kidney problems.  
- aliskiren

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (e.g. potassium) in your blood at regular intervals.  
See also information under the heading “Do not take Sabervel”.

You must tell your doctor if you think you are (or might become) pregnant. Sabervel is not recommended in early pregnancy, and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see pregnancy section).

### **Use in Children**

This medicine should not be used in children and adolescents (under 18 years) because the safety and efficacy have not yet been fully established.

### **Other medicines and Sabervel**

Tell your doctor or pharmacist if you are taking , have recently taken or might take any other medicines.

Your doctor may need to change your dose and/or to take other precautions:

If you are taking an ACE-inhibitor or aliskiren (see also information under the headings “Do not take Sabervel” and “Warnings and precautions”).

### **You may need to have blood checks if you take:**

- potassium supplements
- salt substitutes containing potassium
- potassium-sparing medicines (such as certain diuretics)
- medicines containing lithium

If you take certain painkillers, called non-steroidal anti-inflammatory drugs, the effect of irbesartan may be reduced.

### **Taking Sabervel with food and drink**

Sabervel can be taken with or without food.

### **Pregnancy and breast-feeding**

#### **Pregnancy**

You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking Sabervel before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Sabervel. Sabervel is not recommended in early pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

#### **Breast-feeding**

Tell your doctor if you are breast-feeding or about to start breast-feeding. Sabervel is not recommended for mothers who are breast-feeding, and your doctor may choose another treatment for you if you wish to breast-feed, especially if your baby is newborn, or was born prematurely.

### **Driving and using machines**

No studies on the effects on the ability to drive and use machines have been performed. Sabervel is unlikely to affect your ability to drive or use machines. However, occasionally dizziness or weariness may occur during treatment of high blood pressure. If you experience these, talk to your doctor before attempting to drive or use machines.

### **Important information about some of the ingredients of Sabervel**

**Sabervel contains lactose.** If you have been told by your doctor that you have an intolerance to some

sugars (e.g. lactose), contact your doctor before taking this medicine.

### 3. HOW TO TAKE SABERVEL

Always take Sabervel exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

#### Method of administration

Sabervel is for **oral use**. Swallow the tablets with a sufficient amount of fluid (e.g. one glass of water). You can take Sabervel with or without food. Try to take your daily dose at about the same time each day. It is important that you continue to take Sabervel until your doctor tells you otherwise.

- **Patients with high blood pressure**  
The usual dose is 150 mg once a day. The dose may later be increased to 300 mg once daily depending on blood pressure response.
- **Patients with high blood pressure and type 2 diabetes with kidney disease**  
In patients with high blood pressure and type 2 diabetes, 300 mg once daily is the preferred maintenance dose for the treatment of associated kidney disease.

The doctor may advise a lower dose, especially when starting treatment in certain patients such as those on **haemodialysis**, or those **over the age of 75 years**.

The maximal blood pressure lowering effect should be reached 4-6 weeks after beginning treatment.

#### If you take more Sabervel than you should:

If you accidentally take too many tablets, contact your doctor immediately.

#### Children should not take Sabervel

Sabervel should not be given to children under 18 years of age. If a child swallows some tablets, contact your doctor immediately.

#### If you forget to take Sabervel:

If you accidentally miss a daily dose, just take the next dose as normal. Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

### 4. POSSIBLE SIDE EFFECTS

Like all medicines, Sabervel can cause side effects, although not everybody gets them. Some of these effects may be serious and may require medical attention.

As with similar medicines, rare cases of allergic skin reactions (rash, urticaria), as well as localised swelling of the face, lips and/or tongue have been reported in patients taking irbesartan. If you get any of these symptoms or get short of breath, **stop taking Sabervel and contact your doctor immediately**.

The frequency of the side effects listed below is defined using the following convention:

Very common: at least 1 in 10 patients or more

Common: at least 1 in 100 and less than 1 in 10 patients

Uncommon: at least 1 in 1000 and less than 1 in 100 patients

Side effects reported in clinical studies for patients treated with Irbesartan were:

- Very common : if you suffer from high blood pressure and type 2 diabetes with kidney disease, blood tests may show an increased level of potassium.
- Common : dizziness, feeling sick/vomiting, fatigue and blood tests may show raised levels of an enzyme that measures the muscle and heart function (creatine kinase enzyme). In patients with high blood pressure and type 2 diabetes with kidney disease, dizziness when getting up from a lying or sitting position, low blood pressure when getting up from a lying or sitting position, pain in joints or muscles and decreased levels of a protein in the red blood cells (haemoglobin) were also reported.
- Uncommon : heart rate increased, flushing, cough, diarrhoea, indigestion/heartburn, sexual dysfunction (problems with sexual performance), chest pain.

Some side effects have been reported since marketing of Irbesartan Side effects where the frequency is not known are: feeling of spinning, headache, taste disturbance, ringing in the ears, muscle cramps, pain in joints and muscles, abnormal liver function, increased blood potassium levels, impaired kidney function, and inflammation of small blood vessels mainly affecting the skin (a condition known as leukocytoclastic vasculitis).

Uncommon cases of jaundice (yellowing of the skin and/or whites of the eyes) have also been reported.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## 5. HOW TO STORE SABERVEL

Keep out of the sight and reach of children.

Do not use Sabervel after the expiry date which is stated on the carton and on the blister after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## 6. FURTHER INFORMATION

### What Sabervel contains

- The active substance is irbesartan. Each tablet of Sabervel 300 mg contains 300 mg irbesartan.
- The other ingredients are:  
Tablet core: lactose monohydrate, microcrystalline cellulose, croscarmellose sodium, silica colloidal anhydrous, hypromellose, magnesium stearate,  
Film-coating: hypromellose, titanium dioxide (E171), macrogol 400

### What Sabervel looks like and contents of the pack

Sabervel 300 mg film-coated tablets are white, concave, round, film-coated tablets with 11 mm diameter.

Sabervel 300mg film-coated tablets are supplied in blister packs of 28, 56, 90 or 98 film-coated tablets.

Not all pack sizes may be marketed.



**Marketing Authorisation Holder:**

Pharmathen S.A,  
6 Dervenakion Str.,  
15351 Pallini, Attiki, Greece

**Manufacturer:**

J. Uriach y Compañía, S.A.  
Av. Camí Reial, 51-57  
08184 – Palau-solità i Plegamans, Barcelona, Spain

Pharmathen S.A,  
6 Dervenakion Str.,  
15351 Pallini, Attiki, Greece

Pharmathen International S.A.  
Industrial Park Sapes,  
Rodopi Prefecture, Block No 5, Rodopi 69300, Greece

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder.

**België/Belgique/Belgien**

Pharmathen S.A.  
Tél/Tel: +30 210 66 65 067

**Lietuva**

Portfarma ehf.  
Tel: +354 534 4030

**България**

Pharmathen S.A.  
Тел.: +30 210 66 65 067

**Luxembourg/Luxemburg**

Portfarma ehf.  
Tel: +354 534 4030

**Česká republika**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Magyarország**

Pharmathen S.A.  
Tel.: +30 210 66 65 067

**Danmark**

Portfarma ehf.  
Tlf: +354 534 4030

**Malta**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Deutschland**

Glenmark Arzneimittel GmbH  
Tel: +49 8142 44392 0

**Nederland**

Glenmark Generics B.V.  
Tel: 0031 20 5226030

**Eesti (Estonia)**

Portfarma ehf.  
Tel: +354 534 4030

**Norge**

Pharmathen S.A.  
Tlf: +30 210 66 65 067

**Ελλάδα**

Pharmathen Hellas S.A.  
Τηλ: +30 210 66 04 300

**Österreich**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**España**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Polska**

Pharmathen S.A.  
Tel.: +30 210 66 65 067

**France**

Pharmathen S.A.  
Tél: +30 210 66 65 067

**Portugal**

Pharmathen S.A.  
Tel: +30 210 66 65 067

**Hrvatska**  
Pharmathen S.A.  
Tel: +30 210 66 65 067

**Ireland**  
Aspire Pharma Limited  
Tel: +44(0)1730 234527

**Ísland**  
Portfarma ehf.  
Tel: +354 534 4030

**Italia**  
Pharmathen S.A.  
Tel: +30 210 66 65 067

**Κύπρος**  
C.V. MEDILINE LTD  
Τηλ: +357 25761699

**Latvija**  
Portfarma ehf.  
Tel: +354 534 4030

**România**  
Pharmathen S.A.  
Tel: +30 210 66 65 067

**Slovenija**  
Pharmathen S.A.  
Tel: +30 210 66 65 067

**Slovenská republika**  
Pharmathen S.A.  
Tel: +30 210 66 65 067

**Suomi/Finland**  
Pharmathen S.A.  
Puh/Tel: +30 210 66 65 067

**Sverige**  
Pharmathen S.A.  
Tel: +30 210 66 65 067

**United Kingdom**  
Aspire Pharma Limited  
Tel: +44(0)1730 234527

**This leaflet was last approved in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu/>